


$$f(z) = \frac{(z - p_1)(z - p_2)}{(z - q_1)(z - q_2)},$$

$$\text{ord}_{p_1} f = \text{ord}_{p_2} f = +1, \quad \text{ord}_{q_1} f = \text{ord}_{q_2} f = -1$$

$$\frac{df}{f} = \frac{dz}{z - p_1} + \frac{dz}{z - p_2} - \frac{dz}{z - q_1} - \frac{dz}{z - q_2}$$

$$\Rightarrow \#Z - \#P = 2 - 2 = 0$$

$w = f(z)$ -plane



$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{(\zeta - p_1)^{n+1}} d\zeta,$$

$$f(\zeta) = \frac{(\zeta - p_1)(\zeta - p_2)}{(\zeta - q_1)(\zeta - q_2)}.$$

At $z_0 = p_1$: $a_0 = f(p_1) = 0$,

$$a_1 = f'(p_1) = \frac{p_1 - p_2}{(p_1 - q_1)(p_1 - q_2)},$$

$$\left. \frac{1}{2} \left(\frac{d}{dz} \left(\frac{z - p_2}{(z - q_1)(z - q_2)} \right) \right) \right|_{z=p_1} = \frac{(p_1 - q_1)(p_1 - q_2) - (p_1 - p_2)[2p_1 - (q_1 + q_2)]}{2(p_1 - q_1)^2(p_1 - q_2)^2}.$$

$$\text{wind}(f(C), 0) = \#Z_C - \#P_C = 2 - 2 = 0 \Rightarrow \oint_C \frac{f'(z)}{f(z)} dz = 0.$$