

Advanced Calculus II

Ji, Yong-hyeon

March 31, 2025

We cover the following topics in this note.

- Convergence of Sequences
- Inequality Rule for Absolute Values
- Limit Theorem (Algebraic Property of Limit of Sequence)

Sequence

Definition. Let $X \subseteq \mathbb{R}$. A **sequence** is a function

$$f : \mathbb{N} \rightarrow X (\subseteq \mathbb{R}), \quad n \mapsto f(n) := a_n.$$

Instead of using function notation $f(n)$, the values of the sequence are denoted by $\{a_n\}_{n=1}^{\infty}$, where $a_n = f(n)$ is called n -th term of the sequence.

Remark. A sequence in $X \subseteq \mathbb{R}$ is a function

$$a : \mathbb{N} \rightarrow X, \quad n \mapsto a_n,$$

where $a_n \in X$ for all $n \in \mathbb{N}$. We sometimes write

$$\{a_n\}, \quad \{a_n\}_{n=1}^{\infty}, \quad \{a_n\}_{n \in \mathbb{N}}, \quad (a_n)_{n \in \mathbb{N}}, \quad \text{or} \quad \langle a_n \rangle_{n \in \mathbb{N}}.$$

Convergence of Sequence

Definition. A real sequence $\{a_n\}_{n=1}^{\infty}$ ($\subseteq \mathbb{R}$) is said to **converge** to $L \in \mathbb{R}$ if and only if

$$\forall \varepsilon > 0, \exists N_{\varepsilon} \in \mathbb{N} \text{ such that } [n \geq N_{\varepsilon} \implies |a_n - L| < \varepsilon].$$

Remark. The real number $L \in \mathbb{R}$ is called **the limit**¹. When a sequence $\{a_n\}_{n=1}^{\infty}$ has the limit L , we will use the notation

$$\lim_{n \rightarrow \infty} a_n = L \quad \text{or} \quad a_n \rightarrow L \text{ as } n \rightarrow \infty.$$


That is,

$$\lim_{n \rightarrow \infty} a_n = L \iff \forall \varepsilon > 0 : \exists N \in \mathbb{N} : [n \geq N \implies |a_n - L| < \varepsilon].$$

Note. If a sequence has a limit, we say that the sequence is **convergent**; if it has no limit, we say that the sequence is **divergent**.

Example. Consider the sequence defined by $a_n = 1/n$ for each $n \in \mathbb{N}$. Prove that

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$



Proof. Let $\varepsilon > 0$. By the Archimedean property, we obtain

$$\exists N_{\varepsilon} \in \mathbb{N} \quad \text{s.t.} \quad 1 < \varepsilon \cdot N_{\varepsilon}, \text{ i.e., } \frac{1}{N_{\varepsilon}} < \varepsilon.$$


Assume that $n \geq N_{\varepsilon}$ then

$$|a_n - 0| = \left| \frac{1}{n} \right| = \frac{1}{n} \leq \frac{1}{N_{\varepsilon}} < \varepsilon.$$


Hence $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$. □

¹The limit of a sequence is unique. See **Theorem 4**

Example. Consider the sequence defined by $b_n = 1 - (-1)^n$ for all $n \in \mathbb{N}$. Prove that b_n does not converge.



Proof.



□

Absolute Value in Reals

Definition. Let $x \in \mathbb{R}$. A **absolute value** $|x|$ of x is defined by

$$|x| := \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Remark. For $x \in \mathbb{R}$,

$$|x| = \begin{cases} x & : x > 0 \\ 0 & : x = 0 \\ -x & : x < 0 \end{cases}$$

Proposition 1. Let $x, y \in \mathbb{R}$.

$$(1) |x| = +\sqrt{x^2}$$

$$(2) |x| \geq 0$$

$$(3) |x| = 0 \Leftrightarrow x = 0$$

$$(4) |x| = |-x|$$

$$(5) |xy| = |x||y|$$

(6) (Fundamental Theorem of Absolute Values) For $c \geq 0$, we have

$$|x| \leq c \iff -c \leq x \leq c$$

$$(7) -|x| \leq x \leq |x|$$

Proof. (1) If $(x \geq 0)$ then $|x| = x = \sqrt{x^2}$. Similarly if $x < 0$ then $|x| = -x = \sqrt{x^2}$.

$$(2) |x| = \begin{cases} x \geq 0 & : x \geq 0 \\ -x > 0 & : x < 0 \end{cases} \geq 0.$$

(3) (\Leftarrow) If $x = 0$ then $|x| = x = 0$.


(\Rightarrow) Let $|x| = 0$. Suppose that $x \neq 0$.

$$(i) \ x > 0 \implies |x| = x > 0 \nlessdot$$

$$(ii) \ x < 0 \implies |x| = -x > 0 \nlessdot$$

Thus x must be zero.

$$(4) \quad |-x| = \begin{cases} -x & : -x \geq 0 \text{ (i.e., } x \leq 0) \\ -(-x) = x & : -x < 0 \text{ (i.e., } x > 0) \end{cases} = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases} = |x|.$$



$$(5) \quad |xy| = \begin{cases} xy = |x||y| & : x \geq 0, y \geq 0 \\ -xy = x(-y) = |x||y| & : x \geq 0, y < 0 \\ -xy = (-x)y = |x||y| & : x < 0, y \geq 0 \\ xy = (-x)(-y) = |x||y| & : x < 0, y < 0 \end{cases}$$

(6) (\Rightarrow) Let $|x| \leq c$.

(i) $x \geq 0 \Rightarrow x = |x| \leq c$, i.e., $-c \leq 0 \leq x \leq c$.

(ii) $x < 0 \Rightarrow -x = |x| \leq c$, i.e., $-c \leq x < 0 \leq c$.

Thus, $-c \leq x \leq c$.

(\Leftarrow) Let $-c \leq x \leq c$.

(i) $x \geq 0 \Rightarrow |x| = x \leq c$.


(ii) $x < 0 \Rightarrow |x| = -x \leq c$.

Thus, $|x| \leq c$.

Key equivalence:

$$|x| \leq c \iff -c \leq x \leq c$$

Interval: $-c \leq x \leq c$



Absolute: $|x| \leq c$

(7) Let $c = |x|$, where $c \geq 0$. By (6), thus, the result follows.

□

Triangle Inequality

Proposition 2. Let $x, y \in \mathbb{R}$.

$$(1) |x + y| \leq |x| + |y|$$

$$(2) \left| |x| - |y| \right| \leq |x - y|.$$

Proof. (1) By (7) of **Proposition 1**, we have

$$-|x| \leq x \leq |x|, \quad -|y| \leq y \leq |y|.$$

Then

$$\begin{array}{ccccccc} -|x| & \leq & x & \leq & |x| \\ + & & & & \\ -|y| & \leq & y & \leq & |y| \\ \hline -(|x| + |y|) & \leq & x + y & \leq & |x| + |y| \end{array}$$

Thus, we have $|x + y| \leq |x| + |y|$.

(2)

(i) Note that

$$\begin{aligned} |x| &= |x - y + y| \\ &\leq |x - y| + |y| \quad \text{by (1) of Proposition 2} \end{aligned}$$

$$\text{Thus } |x| - |y| \leq |x - y|.$$

(ii) Note that

$$\begin{aligned} |y| &= |x - (x - y)| \\ &\leq |x| + |-(x - y)| \quad \text{by (1) of Proposition 2} \\ &= |x| + |x - y| \quad \text{by (4) of Proposition 1} \end{aligned}$$

$$\text{Therefore } -|x - y| \leq |x| - |y|.$$

By (i) and (ii), we know

$$-|x - y| \leq |x| - |y| \leq |x - y|, \quad \text{i.e., } \left| |x| - |y| \right| \leq |x - y|.$$

□


Boundedness of Sequence

Definition. Let $\{a_n\}_{n=1}^{\infty} (\subseteq \mathbb{R})$ is a sequence. $\{a_n\}$ is said to be **bounded** if

$$\exists M \in \mathbb{R} \text{ such that } \forall n \in \mathbb{N}, |a_n| \leq M.$$

Proposition 3. A convergent sequence is bounded.

Proof.



□

Note. We have established that if the limit of a sequence a_n exists as n approaches infinity, then there exists a real number M such that $|a_n| \leq M$ for all n :

$$\exists A \in \mathbb{R} \text{ s.t. } A = \lim_{n \rightarrow \infty} a_n \implies \exists M \in \mathbb{R} \text{ s.t. } \forall n \in \mathbb{N}, |a_n| \leq M.$$

However, the converse is not necessarily true:

$$\exists A \in \mathbb{R} \text{ s.t. } A = \lim_{n \rightarrow \infty} a_n \not\Leftarrow \exists M \in \mathbb{R} \text{ s.t. } \forall n \in \mathbb{N}, |a_n| \leq M.$$

To illustrate, consider the sequence $\{a_n\} = 1 - (-1)^n$. This sequence is bounded, yet it does not converge, serving as a counterexample. Furthermore, we note the following important theorems:

1. Monotone Convergence Theorem:

- (i) If a sequence $\{a_n\}$ is bounded above and monotone increasing, then it converges.
- (ii) If a sequence $\{a_n\}$ is bounded below and monotone decreasing, then it converges.

2. Bolzano-Weierstrass Theorem: Every bounded sequence of real numbers has a convergent subsequence. That is, if there exists a real number M such that $|a_n| < M$ for all n , then there exists a convergent subsequence $\{a_{n_k}\}$ of $\{a_n\}$.

Limit Theorem (Algebraic Property of Limit of Sequence)

Theorem. Let $\lim_{n \rightarrow \infty} a_n = \alpha$, $\lim_{n \rightarrow \infty} b_n = \beta$, and $k \in \mathbb{R}$. Then

- (1) $\lim_{n \rightarrow \infty} ka_n = k\alpha = k \lim_{n \rightarrow \infty} a_n$.
- (2) $\lim_{n \rightarrow \infty} a_n \pm b_n = \alpha \pm \beta = \lim_{n \rightarrow \infty} a_n \pm \lim_{n \rightarrow \infty} b_n$.
- (3) $\lim_{n \rightarrow \infty} a_n b_n = \alpha\beta = \lim_{n \rightarrow \infty} a_n \lim_{n \rightarrow \infty} b_n$.
- (4) $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\alpha}{\beta} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n}$. (Here, $\beta \neq 0$ and $b_n \neq 0$)


Proof. Let $\varepsilon > 0$.

(1) If $k = 0$, it is trivial. Let $k \neq 0$. Since $\lim_{n \rightarrow \infty} a_n = \alpha$, we know

$$\exists N \in \mathbb{N} \text{ such that } \left[n \geq N \implies |a_n - \alpha| < \frac{\varepsilon}{|k|} \right] \quad (*)$$

Thus, if $n \geq N$ then

$$\begin{aligned} |ka_n - k\alpha| &= |k(a_n - \alpha)| \\ &= |k||a_n - \alpha| \quad \because |xy| = |x||y| \\ &< |k| \cdot \frac{\varepsilon}{|k|} \quad \text{by } (*) \\ &= \varepsilon. \end{aligned}$$



(2)

(3)

(4)

□

Uniqueness of Limits

Theorem 4. *The limit of a sequence is unique.*

Proof.

□