

Quotient Rings, Spectra, and Cyclic Codes

A comparative lecture note based on $\mathbb{Z}$, $\mathbb{C}[x]$, and $\mathbb{F}_q[x]/\langle x^n - 1 \rangle$

Contents

1	Introduction	2
2	The ideal correspondence for a quotient ring	2
3	First model: the quotient $\mathbb{Z} \rightarrow \mathbb{Z}/(n)$	3
3.1	Ideals of $\mathbb{Z}$	3
4	Second model: the quotient $\mathbb{C}[x] \rightarrow \mathbb{C}[x]/\langle g(x) \rangle$	4
4.1	Ideals of $\mathbb{C}[x]$	4
5	The geometric meaning: spectra and closed subspaces	5
5.1	From quotients to closed immersions	5
5.2	Points and residue fields	5
6	The spectrum of $\mathbb{Z}$	6
6.1	Residue fields	6
6.2	Example: $\text{Spec}(\mathbb{Z}/(12))$	6
7	The spectrum of $\mathbb{C}[x]$	7
7.1	Residue fields	7
7.2	Example: $\text{Spec}(\mathbb{C}[x]/(g(x)))$	7
8	The cyclic-code quotient ring	8
9	From the quotient ring to the vector space $\mathbb{F}_q^n$	8
9.1	Multiplication by x becomes cyclic shift	9
10	Cyclic codes as ideals	9
11	Dimension and basis	10
12	Parity-check polynomial	11
13	Geometric picture of the cyclic-code ring	11
14	A small worked example	12

15 Full comparison tables	12
16 Final summary	14
17 Elements of a ring as functions on $\text{Spec}(R)$	15
17.1 The general construction	15
17.2 The case $R = \mathbb{C}[x]$	15
17.3 The case $R = \mathbb{Z}$	16
17.4 Closed-point versions of the coproduct picture	16
17.5 Quotients as restricting the domain of definition	17
17.6 The closed-point evaluation section for $\mathbb{C}[x]$	19
17.7 The closed-point evaluation section for $\mathbb{Z}$	19
17.8 The closed-point evaluation section for the cyclic-code ring	20

1 Introduction

The purpose of these notes is to explain, in one unified framework, the following three themes:

- (1) the ideal correspondence between a ring R and a quotient R/I ;
- (2) the geometric meaning of this correspondence through spectra, especially for

$$R = \mathbb{Z} \quad \text{and} \quad R = \mathbb{C}[x];$$

- (3) the application of the same idea to cyclic codes through the quotient ring

$$\mathbb{F}_q[x]/\langle x^n - 1 \rangle.$$

The basic algebraic fact is simple:

Passing from R to R/I means forcing every element of I to become zero. Therefore only those ideals of R that already contain I survive in the quotient.

In coding theory, the quotient

$$\mathbb{F}_q[x]/\langle x^n - 1 \rangle$$

has an additional structure: as an $\mathbb{F}_q$ -vector space it is naturally identified with $\mathbb{F}_q^n$, and multiplication by x becomes cyclic shift of coordinates. Because of this, ideals of the quotient ring become cyclic codes.

2 The ideal correspondence for a quotient ring

We begin with the universal algebraic statement.

Theorem 2.1 (Ideal correspondence). *Let R be a ring, let $I \triangleleft R$ be an ideal, and let*

$$\pi : R \rightarrow R/I$$

be the quotient map. Then the assignment

$$J' \mapsto \pi^{-1}(J')$$

defines an inclusion-preserving bijection

$$\{\text{ideals of } R/I\} \longleftrightarrow \{\text{ideals } J \triangleleft R \text{ such that } I \subseteq J\}.$$

Its inverse is given by

$$J \mapsto J/I.$$

Proof. Let $J' \triangleleft R/I$. Then $\pi^{-1}(J')$ is an ideal of R . Since $\pi(I) = 0$, we have

$$I \subseteq \pi^{-1}(J').$$

Conversely, if $J \triangleleft R$ and $I \subseteq J$, then

$$J/I = \{\bar{x} \in R/I : x \in J\}$$

is an ideal of R/I .

Now we verify that the two constructions are inverse.

First, let $J \triangleleft R$ with $I \subseteq J$. Then

$$\pi^{-1}(J/I) = J.$$

Indeed, $\pi(x) \in J/I$ if and only if $x \in J$.

Second, let $J' \triangleleft R/I$. Then

$$\pi^{-1}(J')/I = J'.$$

Indeed, every element of J' is represented by an element of $\pi^{-1}(J')$, and conversely every class from $\pi^{-1}(J')$ lies in J' .

Thus the correspondence is bijective. Inclusion-preservation is immediate. □

Remark 2.2. A useful slogan is

ideals of R/I are exactly ideals of R containing I .

3 First model: the quotient $\mathbb{Z} \rightarrow \mathbb{Z}/(n)$

We now specialize to $R = \mathbb{Z}$.

3.1 Ideals of $\mathbb{Z}$

Every ideal of $\mathbb{Z}$ has the form

$$(m) = m\mathbb{Z}$$

for a unique integer $m \geq 0$.

Let us consider the quotient

$$\pi : \mathbb{Z} \rightarrow \mathbb{Z}/(n).$$

Proposition 3.1. *The ideals of $\mathbb{Z}/(n)$ are in bijection with the ideals $(d) \subseteq \mathbb{Z}$ satisfying*

$$(n) \subseteq (d).$$

Equivalently, they are in bijection with the positive divisors d of n .

Proof. By Theorem 2.1, the ideals of $\mathbb{Z}/(n)$ correspond to the ideals of $\mathbb{Z}$ containing (n) . Since every ideal of $\mathbb{Z}$ is of the form (d) , we must determine when

$$(n) \subseteq (d).$$

This happens if and only if $d \mid n$. □

Example 3.2. Take $n = 12$. Then the positive divisors of 12 are

$$1, 2, 3, 4, 6, 12.$$

Hence the ideals of $\mathbb{Z}/(12)$ are

$$(1)/(12), (2)/(12), (3)/(12), (4)/(12), (6)/(12), (12)/(12).$$

For example,

$$(4)/(12) = \{\bar{0}, \bar{4}, \bar{8}\} \subseteq \mathbb{Z}/(12).$$

Thus quotienting by (12) preserves exactly the ideal structure compatible with the relation $12 = 0$.

4 Second model: the quotient $\mathbb{C}[x] \rightarrow \mathbb{C}[x]/\langle g(x) \rangle$

We now turn to the polynomial ring $\mathbb{C}[x]$.

4.1 Ideals of $\mathbb{C}[x]$

Since $\mathbb{C}[x]$ is a principal ideal domain, every ideal is of the form

$$(f(x))$$

for some polynomial $f(x) \in \mathbb{C}[x]$.

Let

$$\pi : \mathbb{C}[x] \rightarrow \mathbb{C}[x]/\langle g(x) \rangle$$

be the quotient map.

Proposition 4.1. *The ideals of $\mathbb{C}[x]/\langle g(x) \rangle$ are in bijection with the ideals $(f(x)) \subseteq \mathbb{C}[x]$ satisfying*

$$(g(x)) \subseteq (f(x)).$$

Equivalently, they are in bijection with the divisors of $g(x)$, up to multiplication by a nonzero constant.

Proof. By Theorem 2.1, the ideals of $\mathbb{C}[x]/\langle g \rangle$ correspond to the ideals of $\mathbb{C}[x]$ containing (g) . Since every ideal is principal, we consider ideals (f) such that

$$(g) \subseteq (f).$$

In a PID, this is equivalent to $f \mid g$. □

Example 4.2. Let

$$g(x) = x^2(x - 1).$$

Then the monic divisors of $g(x)$ are

$$1, \quad x, \quad x^2, \quad x - 1, \quad x(x - 1), \quad x^2(x - 1).$$

Therefore the ideals of

$$\mathbb{C}[x]/\langle x^2(x - 1) \rangle$$

are precisely

$$(1)/(g), (x)/(g), (x^2)/(g), (x - 1)/(g), (x(x - 1))/(g), (g)/(g).$$

This is the exact polynomial analogue of the integer case:

divisors of $n \longleftrightarrow$ ideals of $\mathbb{Z}/(n)$,	divisors of $g(x) \longleftrightarrow$ ideals of $\mathbb{C}[x]/(g(x))$.
--	---

5 The geometric meaning: spectra and closed subspaces

We now reinterpret the quotient construction geometrically.

5.1 From quotients to closed immersions

The quotient map

$$R \rightarrow R/I$$

induces a map

$$\mathrm{Spec}(R/I) \rightarrow \mathrm{Spec}(R).$$

Its image is the closed subset

$$V(I) = \{\mathfrak{p} \in \mathrm{Spec}(R) : I \subseteq \mathfrak{p}\}.$$

Thus there is a natural homeomorphism

$$\mathrm{Spec}(R/I) \cong V(I) \subseteq \mathrm{Spec}(R).$$

Remark 5.1. The quotient ring R/I is therefore the coordinate ring of the closed subspace cut out by the equation $I = 0$.

5.2 Points and residue fields

Let $x \in \mathrm{Spec}(R)$, corresponding to a prime ideal $\mathfrak{p}_x$. Its residue field is

$$\kappa(x) = \mathrm{Frac}(R_{\mathfrak{p}_x}/\mathfrak{p}_x R_{\mathfrak{p}_x}).$$

If $\mathfrak{p}_x$ is maximal, then

$$\kappa(x) \cong R/\mathfrak{p}_x.$$

There is a canonical morphism

$$\mathrm{Spec}(\kappa(x)) \rightarrow \mathrm{Spec}(R)$$

whose image is the point x .

Remark 5.2. It is essential to distinguish two kinds of maps:

(a) the quotient-induced map

$$\mathrm{Spec}(R/I) \rightarrow \mathrm{Spec}(R),$$

which cuts out a closed subspace;

(b) the residue-field map

$$\mathrm{Spec}(\kappa(x)) \rightarrow \mathrm{Spec}(R),$$

which picks out a single point.

6 The spectrum of $\mathbb{Z}$

The prime ideals of $\mathbb{Z}$ are:

$$(0) \quad \text{and} \quad (p) \text{ for each prime } p.$$

Thus $\mathrm{Spec}(\mathbb{Z})$ has one generic point (0) and one closed point (p) for each prime p .

6.1 Residue fields

At the generic point,

$$\kappa((0)) = \mathbb{Q}.$$

At a closed point (p) ,

$$\kappa((p)) \cong \mathbb{Z}/(p) \cong \mathbb{F}_p.$$

Therefore each prime p gives a morphism

$$\mathrm{Spec}(\mathbb{F}_p) \rightarrow \mathrm{Spec}(\mathbb{Z}).$$

Remark 6.1. The coproduct

$$\coprod_{p \text{ prime}} \mathrm{Spec}(\mathbb{F}_p) \rightarrow \mathrm{Spec}(\mathbb{Z})$$

captures all closed points, but it omits the generic point. The full family of point maps is

$$\coprod_{x \in \mathrm{Spec}(\mathbb{Z})} \mathrm{Spec}(\kappa(x)) \rightarrow \mathrm{Spec}(\mathbb{Z}).$$

6.2 Example: $\mathrm{Spec}(\mathbb{Z}/(12))$

Now let

$$\mathbb{Z} \rightarrow \mathbb{Z}/(12).$$

Then

$$\mathrm{Spec}(\mathbb{Z}/(12)) \cong V(12) \subseteq \mathrm{Spec}(\mathbb{Z}).$$

Set-theoretically,

$$V(12) = \{(2), (3)\},$$

because the prime ideals containing (12) are exactly (2) and (3) .

Thus quotienting by 12 cuts out the closed points corresponding to the prime divisors of 12.

7 The spectrum of $\mathbb{C}[x]$

The prime ideals of $\mathbb{C}[x]$ are:

$$(0) \quad \text{and} \quad (x - \alpha) \text{ for each } \alpha \in \mathbb{C}.$$

Thus $\text{Spec}(\mathbb{C}[x])$ has one generic point (0) and one closed point $(x - \alpha)$ for each complex number α .

7.1 Residue fields

At the generic point,

$$\kappa((0)) = \mathbb{C}(x).$$

At a closed point $(x - \alpha)$,

$$\kappa((x - \alpha)) \cong \mathbb{C}[x]/(x - \alpha) \cong \mathbb{C}.$$

Thus each $\alpha \in \mathbb{C}$ gives a morphism

$$\text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(\mathbb{C}[x]).$$

Equivalently,

$$\coprod_{\alpha \in \mathbb{C}} \text{Spec}(\mathbb{C}[x]/(x - \alpha)) \rightarrow \text{Spec}(\mathbb{C}[x])$$

describes all closed points.

Remark 7.1. Again this closed-point coproduct omits the generic point. The complete family is

$$\coprod_{x \in \text{Spec}(\mathbb{C}[x])} \text{Spec}(\kappa(x)) \rightarrow \text{Spec}(\mathbb{C}[x]).$$

7.2 Example: $\text{Spec}(\mathbb{C}[x]/(g(x)))$

Let

$$\mathbb{C}[x] \rightarrow \mathbb{C}[x]/(g(x)).$$

Then

$$\text{Spec}(\mathbb{C}[x]/(g(x))) \cong V(g) \subseteq \text{Spec}(\mathbb{C}[x]).$$

If

$$g(x) = \prod_{i=1}^r (x - \alpha_i)^{e_i},$$

then set-theoretically

$$V(g) = \{(x - \alpha_1), \dots, (x - \alpha_r)\}.$$

Thus quotienting by (g) cuts out the roots of $g(x)$.

Proposition 7.2 (Squarefree case). *If*

$$g(x) = \prod_{i=1}^r (x - \alpha_i)$$

has distinct roots, then

$$\mathbb{C}[x]/(g(x)) \cong \prod_{i=1}^r \mathbb{C}[x]/(x - \alpha_i) \cong \prod_{i=1}^r \mathbb{C}.$$

Consequently,

$$\text{Spec}(\mathbb{C}[x]/(g(x))) \cong \prod_{i=1}^r \text{Spec}(\mathbb{C}).$$

Proof. The ideals $(x - \alpha_i)$ are pairwise comaximal. The Chinese remainder theorem gives

$$\mathbb{C}[x]/\left(\prod_{i=1}^r (x - \alpha_i)\right) \cong \prod_{i=1}^r \mathbb{C}[x]/(x - \alpha_i).$$

Each factor is isomorphic to $\mathbb{C}$. □

8 The cyclic-code quotient ring

We now pass to the finite-field setting relevant to coding theory.

Fix a finite field $\mathbb{F}_q$ and a positive integer n . Define

$$A := \mathbb{F}_q[x]/\langle x^n - 1 \rangle.$$

Proposition 8.1. *The ideals of A are exactly the ideals*

$$\langle \overline{g(x)} \rangle \subseteq A$$

where $g(x)$ runs through the monic divisors of $x^n - 1$.

Proof. By Theorem 2.1, the ideals of A correspond to ideals of $\mathbb{F}_q[x]$ containing $(x^n - 1)$. Since $\mathbb{F}_q[x]$ is a PID, every ideal is of the form $(g(x))$, and

$$(x^n - 1) \subseteq (g(x)) \iff g(x) \mid x^n - 1.$$

□

This is formally identical to the earlier two examples:

$$\begin{aligned} \text{ideals of } \mathbb{Z}/(n) &\leftrightarrow \text{divisors of } n, \\ \text{ideals of } \mathbb{C}[x]/(g) &\leftrightarrow \text{divisors of } g, \\ \text{ideals of } \mathbb{F}_q[x]/(x^n - 1) &\leftrightarrow \text{divisors of } x^n - 1. \end{aligned}$$

9 From the quotient ring to the vector space $\mathbb{F}_q^n$

Every residue class in A has a unique representative of degree $< n$:

$$a_0 + a_1x + \cdots + a_{n-1}x^{n-1}.$$

Hence there is a natural $\mathbb{F}_q$ -linear isomorphism

$$\Phi : A \rightarrow \mathbb{F}_q^n, \quad \Phi\left(\overline{a_0 + a_1x + \cdots + a_{n-1}x^{n-1}}\right) = (a_0, a_1, \dots, a_{n-1}).$$

Therefore

$$A \cong \mathbb{F}_q^n$$

as $\mathbb{F}_q$ -vector spaces.

9.1 Multiplication by x becomes cyclic shift

Let

$$f(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}.$$

Then in A ,

$$xf(x) = a_0x + a_1x^2 + \cdots + a_{n-2}x^{n-1} + a_{n-1}x^n.$$

Since

$$x^n \equiv 1 \pmod{x^n - 1},$$

we obtain

$$xf(x) \equiv a_{n-1} + a_0x + \cdots + a_{n-2}x^{n-1}.$$

Thus under Φ , multiplication by x becomes the cyclic shift

$$(a_0, a_1, \dots, a_{n-1}) \mapsto (a_{n-1}, a_0, \dots, a_{n-2}).$$

10 Cyclic codes as ideals

Definition 10.1. A *cyclic code* of length n over $\mathbb{F}_q$ is an $\mathbb{F}_q$ -linear subspace

$$C \subseteq \mathbb{F}_q^n$$

such that for every

$$(c_0, c_1, \dots, c_{n-1}) \in C,$$

its cyclic shift

$$(c_{n-1}, c_0, \dots, c_{n-2})$$

also belongs to C .

Theorem 10.2 (Cyclic codes are ideals). *Under the identification*

$$\Phi : \mathbb{F}_q[x]/\langle x^n - 1 \rangle \xrightarrow{\sim} \mathbb{F}_q^n,$$

the cyclic codes of length n over $\mathbb{F}_q$ are exactly the ideals of

$$\mathbb{F}_q[x]/\langle x^n - 1 \rangle.$$

Proof. Let $C \subseteq \mathbb{F}_q^n$ be a cyclic code. Since C is an $\mathbb{F}_q$ -subspace, $\Phi^{-1}(C) \subseteq A$ is closed under addition and scalar multiplication. Since C is closed under cyclic shift and multiplication by x corresponds to cyclic shift, $\Phi^{-1}(C)$ is closed under multiplication by x .

Now let

$$a(x) = a_0 + a_1x + \cdots + a_mx^m \in \mathbb{F}_q[x]$$

and let $c \in \Phi^{-1}(C)$. Then

$$a(x)c = a_0c + a_1(xc) + \cdots + a_m(x^m c).$$

Each term lies in $\Phi^{-1}(C)$, so $a(x)c \in \Phi^{-1}(C)$. Hence $\Phi^{-1}(C)$ is an ideal of A .

Conversely, if $J \triangleleft A$ is an ideal, then J is an $\mathbb{F}_q$ -subspace and is stable under multiplication by x . Under Φ , this means $\Phi(J) \subseteq \mathbb{F}_q^n$ is an $\mathbb{F}_q$ -subspace closed under cyclic shift, i.e. a cyclic code. $\square$

Combining Theorem 10.2 with Proposition 8.1, we obtain:

Corollary 10.3 (Classification of cyclic codes). *Every cyclic code of length n over $\mathbb{F}_q$ has the form*

$$C = \langle \overline{g(x)} \rangle \subseteq \mathbb{F}_q[x]/\langle x^n - 1 \rangle$$

for a unique monic divisor $g(x) \mid x^n - 1$.

The polynomial $g(x)$ is called the *generator polynomial* of the code.

11 Dimension and basis

Let

$$C = \langle \overline{g(x)} \rangle \subseteq \mathbb{F}_q[x]/\langle x^n - 1 \rangle, \quad x^n - 1 = g(x)h(x).$$

Theorem 11.1. *If $\deg g = r$, then*

$$\dim_{\mathbb{F}_q} C = n - r.$$

Moreover, the vectors corresponding to

$$\overline{g(x)}, \overline{xg(x)}, \overline{x^2g(x)}, \dots, \overline{x^{n-r-1}g(x)}$$

form a basis of C .

Proof. Consider the $\mathbb{F}_q$ -linear map

$$\psi : \{m(x) \in \mathbb{F}_q[x] : \deg m < n - r\} \rightarrow C, \quad m(x) \mapsto \overline{m(x)g(x)}.$$

First, ψ is injective. Indeed, if $\overline{m(x)g(x)} = 0$, then

$$x^n - 1 = g(x)h(x) \mid m(x)g(x).$$

Since $\mathbb{F}_q[x]$ is a domain, this implies $h(x) \mid m(x)$. But

$$\deg m < n - r = \deg h,$$

so $m(x) = 0$.

Second, ψ is surjective. Any element of C has the form $\overline{a(x)g(x)}$. Divide $a(x)$ by $h(x)$:

$$a(x) = q(x)h(x) + r(x), \quad \deg r < \deg h = n - r.$$

Then

$$a(x)g(x) = q(x)h(x)g(x) + r(x)g(x) = q(x)(x^n - 1) + r(x)g(x),$$

hence

$$\overline{a(x)g(x)} = \overline{r(x)g(x)}.$$

So ψ is surjective.

Therefore ψ is an isomorphism. Its domain has basis

$$1, x, \dots, x^{n-r-1},$$

and transporting that basis through ψ gives the desired basis of C . □

12 Parity-check polynomial

The complementary factor

$$h(x) = \frac{x^n - 1}{g(x)}$$

is called the *parity-check polynomial*.

Theorem 12.1 (Kernel description). *Let*

$$\mu_h : A \rightarrow A, \quad f \mapsto fh.$$

Then

$$C = \text{Ker}(\mu_h).$$

Equivalently,

$$C = \{ c(x) \in A : c(x)h(x) = 0 \}.$$

Proof. If $c(x) \in C$, then

$$c(x) = m(x)g(x)$$

for some $m(x) \in \mathbb{F}_q[x]$. Hence in A ,

$$c(x)h(x) = m(x)g(x)h(x) = m(x)(x^n - 1) = 0.$$

So $C \subseteq \text{Ker}(\mu_h)$.

Conversely, suppose $c(x)h(x) = 0$ in A . Then

$$x^n - 1 = g(x)h(x) \mid c(x)h(x)$$

in $\mathbb{F}_q[x]$. Since $\mathbb{F}_q[x]$ is a domain and $h(x) \neq 0$, it follows that

$$g(x) \mid c(x).$$

Hence $c(x) \in C$. □

Remark 12.2. There is a useful symmetry:

$$C = \mathfrak{I}(\mu_g) = \text{Ker}(\mu_h).$$

Thus $g(x)$ describes the code as an image, while $h(x)$ describes it as a kernel.

13 Geometric picture of the cyclic-code ring

The quotient map

$$\mathbb{F}_q[x] \rightarrow \mathbb{F}_q[x] / \langle x^n - 1 \rangle$$

induces a closed immersion

$$\text{Spec}(\mathbb{F}_q[x] / \langle x^n - 1 \rangle) \hookrightarrow \text{Spec}(\mathbb{F}_q[x]),$$

whose image is

$$V(x^n - 1).$$

If

$$x^n - 1 = \prod_{i=1}^r f_i(x)^{e_i}$$

is the factorization into monic irreducibles, then the points of this closed subspace are the primes

$$(f_1), \dots, (f_r).$$

Remark 13.1. Topologically, only the distinct irreducible factors matter. Scheme-theoretically, the exponents e_i remain visible through nilpotents.

If $x^n - 1$ is squarefree, then the Chinese remainder theorem gives

$$\mathbb{F}_q[x]/\langle x^n - 1 \rangle \cong \prod_{i=1}^r \mathbb{F}_q[x]/\langle f_i(x) \rangle.$$

Therefore

$$\text{Spec}(\mathbb{F}_q[x]/\langle x^n - 1 \rangle) \cong \prod_{i=1}^r \text{Spec}(\mathbb{F}_q[x]/\langle f_i(x) \rangle).$$

This is the exact finite-field analogue of Proposition 7.2.

14 A small worked example

Take $q = 2$ and $n = 3$. Then

$$x^3 - 1 = x^3 + 1 = (x + 1)(x^2 + x + 1)$$

in $\mathbb{F}_2[x]$.

The monic divisors of $x^3 - 1$ are

$$1, \quad x + 1, \quad x^2 + x + 1, \quad x^3 + 1.$$

Hence there are four cyclic codes of length 3 over $\mathbb{F}_2$.

(1) $g(x) = 1$. Then

$$C = \langle \bar{1} \rangle = \mathbb{F}_2[x]/\langle x^3 - 1 \rangle \cong \mathbb{F}_2^3.$$

(2) $g(x) = x + 1$. Then $\deg g = 1$, so $\dim C = 2$. This is the even-weight code:

$$\{000, 011, 101, 110\}.$$

(3) $g(x) = x^2 + x + 1$. Then $\deg g = 2$, so $\dim C = 1$. This is the repetition code:

$$\{000, 111\}.$$

(4) $g(x) = x^3 + 1 = x^3 - 1$. Then

$$C = 0.$$

15 Full comparison tables

We now collect the three worlds side by side.

Table A: ring-theoretic comparison

Theme	$\mathbb{Z}$	$\mathbb{C}[x]$	$\mathbb{F}_q[x]$ / cyclic-code case
Ambient ring R	$\mathbb{Z}$	$\mathbb{C}[x]$	$\mathbb{F}_q[x]$
Typical quotient R/I	$\mathbb{Z}/(n)$	$\mathbb{C}[x]/(g(x))$	$\mathbb{F}_q[x]/(x^n - 1)$
Ideals of R	(m)	$(f(x))$	$(f(x))$
Ideals of R/I correspond to	ideals of $\mathbb{Z}$ containing (n)	ideals of $\mathbb{C}[x]$ containing (g)	ideals of $\mathbb{F}_q[x]$ containing $(x^n - 1)$
Equivalent divisibility condition	$d \mid n$	$f \mid g$	$g \mid x^n - 1$
Quotient ideals classified by	divisors of n	divisors of $g(x)$	divisors of $x^n - 1$
Meaning of quotient	force $n = 0$	force $g(x) = 0$	force $x^n = 1$

Table B: spectral comparison

Theme	$\text{Spec}(\mathbb{Z})$	$\text{Spec}(\mathbb{C}[x])$	$\text{Spec}(\mathbb{F}_q[x])$
Generic point	(0)	(0)	(0)
Closed points	$(p), p \text{ prime}$	$(x - \alpha), \alpha \in \mathbb{C}$	$(f), f \text{ irreducible}$
Residue field at generic point	$\mathbb{Q}$	$\mathbb{C}(x)$	$\mathbb{F}_q(x)$
Residue field at a closed point	$\mathbb{F}_p$	$\mathbb{C}$	$\mathbb{F}_q[x]/(f)$
Closed-point coproduct map	$\coprod_p \text{Spec}(\mathbb{F}_p) \rightarrow \text{Spec}(\mathbb{Z})$	$\coprod_{\alpha \in \mathbb{C}} \text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(\mathbb{C}[x])$	$\coprod_{f \text{ irr}} \text{Spec}(\mathbb{F}_q[x]/(f)) \rightarrow \text{Spec}(\mathbb{F}_q[x])$
What this coproduct sees	all closed points	all closed points	all closed points
What it misses	generic point	generic point	generic point

Table C: quotients as closed subspaces

Theme	$\mathbb{Z}/(n)$	$\mathbb{C}[x]/(g)$	$\mathbb{F}_q[x]/(x^n - 1)$
Closed subset $V(I)$	$V(n)$	$V(g)$	$V(x^n - 1)$
Set-theoretic description	prime divisors of n	roots of g	irreducible factors of $x^n - 1$
Squarefree decomposition	CRT over pairwise coprime prime powers	$\mathbb{C}[x]/(g) \cong \prod \mathbb{C}$ if g squarefree	$\mathbb{F}_q[x]/(x^n - 1) \cong \prod \mathbb{F}_q[x]/(f_i)$ if square-free
Topology sees	distinct prime divisors	distinct roots	distinct irreducible factors
Scheme structure remembers	prime powers	repeated roots	repeated irreducible powers

Table D: coding-theoretic interpretation

Theme	Cyclic-code meaning
Ring	$A = \mathbb{F}_q[x]/(x^n - 1)$
Vector-space identification	$A \cong \mathbb{F}_q^n$
Typical element of A	$a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$
Corresponding vector	$(a_0, a_1, \dots, a_{n-1})$
Multiplication by x	cyclic shift of coordinates
Ideal of A	$\mathbb{F}_q$ -subspace stable under cyclic shift
Coding name	cyclic code
Classification	monic divisors $g(x) \mid x^n - 1$
Code generated by g	$C = \langle \overline{g(x)} \rangle \subseteq A$
Generator polynomial	$g(x)$
Parity-check polynomial	$h(x) = \frac{x^n - 1}{g(x)}$
Image description	$C = \Im(\mu_g)$
Kernel description	$C = \text{Ker}(\mu_h)$

16 Final summary

The three main examples are governed by the same principle.

- In $\mathbb{Z}/(n)$, ideals are controlled by divisors of n .
- In $\mathbb{C}[x]/(g(x))$, ideals are controlled by divisors of $g(x)$.
- In $\mathbb{F}_q[x]/(x^n - 1)$, ideals are controlled by divisors of $x^n - 1$.

The only extra feature in the third case is that

$$\mathbb{F}_q[x]/(x^n - 1) \cong \mathbb{F}_q^n$$

as a vector space, and multiplication by x becomes cyclic shift. Therefore the ideals of the quotient ring are exactly the cyclic codes.

cyclic code = ideal of $\mathbb{F}_q[x]/\langle x^n - 1 \rangle$ = subspace of $\mathbb{F}_q^n$ invariant under cyclic shift.

17 Elements of a ring as functions on $\mathrm{Spec}(R)$

A point that is often left implicit is the following:

An element of a ring R is not, in general, a function with values in one fixed field. Rather, it is a function whose value at a point $x \in \mathrm{Spec}(R)$ lies in the residue field $\kappa(x)$.

17.1 The general construction

Let R be a ring and let $a \in R$. For each point

$$x \in \mathrm{Spec}(R),$$

corresponding to a prime ideal $\mathfrak{p}_x$, the image of a in the residue field

$$\kappa(x)$$

is denoted by

$$a(x) \in \kappa(x).$$

Thus a defines a pointwise rule

$$x \longmapsto a(x) \in \kappa(x).$$

Equivalently, a determines a section of the projection

$$\coprod_{x \in \mathrm{Spec}(R)} \kappa(x) \longrightarrow \mathrm{Spec}(R).$$

Remark 17.1. So the correct geometric statement is:

$$\boxed{a \in R \quad \rightsquigarrow \quad (x \mapsto a(x) \in \kappa(x)).}$$

In particular, an element of R is not naturally a function

$$\mathrm{Spec}(R) \rightarrow R$$

or even a function to a single field. Its codomain varies from point to point.

17.2 The case $R = \mathbb{C}[x]$

Let

$$f(x) \in \mathbb{C}[x].$$

Then f defines a function on $\mathrm{Spec}(\mathbb{C}[x])$ in the above sense.

- At a closed point $(x - \alpha)$, one has

$$f \longmapsto f(\alpha) \in \mathbb{C} \cong \kappa((x - \alpha)).$$

- At the generic point (0) , one has

$$f \longmapsto f(x) \in \mathbb{C}(x) = \kappa((0)).$$

Thus a polynomial is a regular function on $\text{Spec}(\mathbb{C}[x])$, whose value at each closed point is the usual complex number $f(\alpha)$, but whose value at the generic point lies in the rational function field $\mathbb{C}(x)$.

Remark 17.2. If one restricts only to closed points, then one recovers the usual classical picture

$$\alpha \mapsto f(\alpha), \quad \alpha \in \mathbb{C}.$$

So the usual polynomial function on $\mathbb{C}$ is only the closed-point part of the full Spec-picture.

17.3 The case $R = \mathbb{Z}$

Now let $n \in \mathbb{Z}$. Then n defines a function on $\text{Spec}(\mathbb{Z})$ in the same sense.

- At a closed point (p) , where p is prime,

$$n \mapsto n \bmod p \in \mathbb{F}_p \cong \kappa((p)).$$

- At the generic point (0) ,

$$n \mapsto n \in \mathbb{Q} = \kappa((0)).$$

So an integer may be viewed as a function assigning to each prime p its residue modulo p , together with its value in the generic fiber $\mathbb{Q}$.

17.4 Closed-point versions of the coproduct picture

The previous discussion explains the precise meaning of expressions such as

$$\coprod_{\alpha \in \mathbb{C}} \mathbb{C}[x]/(x - \alpha) \quad \text{and} \quad \coprod_{p \text{ prime}} \mathbb{Z}/(p).$$

These should be interpreted geometrically as

$$\coprod_{\alpha \in \mathbb{C}} \text{Spec}(\mathbb{C}[x]/(x - \alpha)) \rightarrow \text{Spec}(\mathbb{C}[x]),$$

and

$$\coprod_{p \text{ prime}} \text{Spec}(\mathbb{Z}/(p)) \rightarrow \text{Spec}(\mathbb{Z}),$$

which recover all closed points.

At the level of values, an element of the ring restricts to a section over the closed points:

$$f \mapsto (\alpha \mapsto f(\alpha) \in \mathbb{C}[x]/(x - \alpha) \cong \mathbb{C}),$$

and

$$n \mapsto (p \mapsto n \bmod p \in \mathbb{Z}/(p) \cong \mathbb{F}_p).$$

However, these closed-point pictures do *not* include the generic point. The full pointwise picture requires the residue fields at *all* points:

$$\coprod_{x \in \text{Spec}(R)} \text{Spec}(\kappa(x)) \rightarrow \text{Spec}(R).$$

17.5 Quotients as restricting the domain of definition

Let

$$\pi : R \rightarrow R/I.$$

Then an element of R/I is a regular function on

$$\mathrm{Spec}(R/I) \cong V(I) \subseteq \mathrm{Spec}(R).$$

In other words, quotienting by I means restricting from all of $\mathrm{Spec}(R)$ to the closed subspace on which the elements of I vanish.

Example 17.3. Let

$$g(x) = \prod_{i=1}^r (x - \alpha_i)$$

with distinct roots. Then

$$\mathbb{C}[x]/(g(x)) \cong \prod_{i=1}^r \mathbb{C},$$

so an element of $\mathbb{C}[x]/(g(x))$ may be viewed literally as a tuple

$$(c_1, \dots, c_r), \quad c_i \in \mathbb{C},$$

that is, as a function on the finite set of roots $\{\alpha_1, \dots, \alpha_r\}$.

Example 17.4. For

$$A = \mathbb{F}_q[x]/(x^n - 1),$$

an element of A is a regular function on

$$V(x^n - 1) \subseteq \mathrm{Spec}(\mathbb{F}_q[x]).$$

At the same time, every element of A has a unique representative

$$a_0 + a_1x + \dots + a_{n-1}x^{n-1},$$

so it also corresponds to a vector

$$(a_0, a_1, \dots, a_{n-1}) \in \mathbb{F}_q^n.$$

This is the bridge from algebraic geometry to coding theory.

Table B: spectral comparison

Theme	$\mathrm{Spec}(\mathbb{Z})$	$\mathrm{Spec}(\mathbb{C}[x])$	$\mathrm{Spec}(\mathbb{F}_q[x])$
Generic point	(0)	(0)	(0)
Closed points	(p) , p prime	$(x - \alpha)$, $\alpha \in \mathbb{C}$	(f) , f irreducible
Residue field at generic point	$\mathbb{Q}$	$\mathbb{C}(x)$	$\mathbb{F}_q(x)$
Residue field at a closed point	$\mathbb{F}_p$	$\mathbb{C}$	$\mathbb{F}_q[x]/(f)$
Closed-point coproduct map	$\coprod_p \mathrm{Spec}(\mathbb{F}_p) \rightarrow \mathrm{Spec}(\mathbb{Z})$	$\rightarrow \coprod_{\alpha \in \mathbb{C}} \mathrm{Spec}(\mathbb{C}) \rightarrow \mathrm{Spec}(\mathbb{C}[x])$	$\rightarrow \coprod_{f \text{ irr}} \mathrm{Spec}(\mathbb{F}_q[x]/(f)) \rightarrow \mathrm{Spec}(\mathbb{F}_q[x])$

What this coproduct sees	all closed points	all closed points	all closed points
What it misses	generic point	generic point	generic point
Element of the ring as a function	$n \mapsto ((p) \mapsto n \bmod p, (0) \mapsto n \in \mathbb{Q})$	$f \mapsto ((x - \alpha) \mapsto f(\alpha), (0) \mapsto f(x) \in \mathbb{C}(x))$	$a(x) \mapsto ((f) \mapsto a(x) \bmod f, (0) \mapsto a(x) \in \mathbb{F}_q(x))$
Closed-point value of an element	$n \mapsto ((p) \mapsto n \bmod p)$	$f(x) \mapsto ((x - \alpha) \mapsto f(\alpha))$	$\overline{a(x)} \mapsto (\mathfrak{m}_i \mapsto a(x) \bmod f_i(x))$

Table D: coding-theoretic interpretation

Theme	Cyclic-code meaning
Ring	$A = \mathbb{F}_q[x]/(x^n - 1)$
Geometric meaning of A	coordinate ring of the closed subscheme $V(x^n - 1) \subseteq \text{Spec}(\mathbb{F}_q[x])$
Element of A as a function	a regular function on $V(x^n - 1)$, i.e. a section $x \mapsto a(x) \in \kappa(x)$ over points of $V(x^n - 1)$
Vector-space identification	$A \cong \mathbb{F}_q^n$
Typical element of A	$a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$
Corresponding vector	$(a_0, a_1, \dots, a_{n-1})$
Multiplication by x	cyclic shift of coordinates
Ideal of A	$\mathbb{F}_q$ -subspace stable under cyclic shift
Coding name	cyclic code
Classification	monic divisors $g(x) \mid x^n - 1$
Code generated by g	$C = \langle \overline{g(x)} \rangle \subseteq A$
Generator polynomial	$g(x)$
Parity-check polynomial	$h(x) = \frac{x^n - 1}{g(x)}$
Image description	$C = \mathfrak{S}(\mu_g)$
Kernel description	$C = \text{Ker}(\mu_h)$
Element of A as a closed-point function	$\overline{a(x)} \mapsto (\mathfrak{m}_i \mapsto a(x) \bmod f_i(x) \in \mathbb{F}_q[x]/(f_i))$

17.6 The closed-point evaluation section for $\mathbb{C}[x]$

Let $f \in \mathbb{C}[x]$. For each closed point $(x - \alpha) \in \text{Spec}(\mathbb{C}[x])$, the image of f in the residue field at that point is

$$f \bmod (x - \alpha) \in \mathbb{C}[x]/(x - \alpha) \cong \mathbb{C}.$$

Thus f determines a section over the closed points

$$s_f : \{\text{closed points of } \text{Spec}(\mathbb{C}[x])\} \longrightarrow \prod_{\alpha \in \mathbb{C}} \mathbb{C}[x]/(x - \alpha),$$

defined by

$$s_f((x - \alpha)) = f \bmod (x - \alpha).$$

Under the natural identification

$$\mathbb{C}[x]/(x - \alpha) \cong \mathbb{C},$$

this is exactly the usual evaluation map

$$(x - \alpha) \longmapsto f(\alpha).$$

However, this description only involves the closed points of $\text{Spec}(\mathbb{C}[x])$. To include the generic point (0) , one must pass to the full residue-field picture:

$$\tilde{f} : \text{Spec}(\mathbb{C}[x]) \longrightarrow \prod_{x \in \text{Spec}(\mathbb{C}[x])} \kappa(x), \quad x \longmapsto f(x) \in \kappa(x).$$

Here

$$\tilde{f}((x - \alpha)) = f(\alpha) \in \mathbb{C}, \quad \tilde{f}((0)) = f(x) \in \mathbb{C}(x).$$

Thus the usual polynomial function $\alpha \mapsto f(\alpha)$ is the closed-point part of the full regular-function interpretation of f on $\text{Spec}(\mathbb{C}[x])$.

17.7 The closed-point evaluation section for $\mathbb{Z}$

Let $n \in \mathbb{Z}$. For each closed point $(p) \in \text{Spec}(\mathbb{Z})$, where p is prime, the image of n in the residue field is

$$n \bmod p \in \mathbb{Z}/(p) \cong \mathbb{F}_p.$$

Thus n determines a section over the closed points

$$s_n : \{\text{closed points of } \text{Spec}(\mathbb{Z})\} \longrightarrow \prod_{p \text{ prime}} \mathbb{Z}/(p),$$

defined by

$$s_n((p)) = n \bmod p.$$

To include the generic point (0) , one uses the full residue-field picture

$$\tilde{n} : \text{Spec}(\mathbb{Z}) \longrightarrow \prod_{x \in \text{Spec}(\mathbb{Z})} \kappa(x), \quad x \longmapsto n(x) \in \kappa(x),$$

where

$$\tilde{n}((p)) = n \bmod p \in \mathbb{F}_p, \quad \tilde{n}((0)) = n \in \mathbb{Q}.$$

So an integer may be viewed as a function assigning to each prime p its residue modulo p , together with its value in the generic fiber $\mathbb{Q}$.

17.8 The closed-point evaluation section for the cyclic-code ring

Let

$$A = \mathbb{F}_q[x]/\langle x^n - 1 \rangle.$$

Factor

$$x^n - 1 = \prod_{i=1}^r f_i(x)^{e_i}$$

into distinct monic irreducible polynomials $f_i(x) \in \mathbb{F}_q[x]$. Then the closed points of $\text{Spec}(A)$ are the prime ideals

$$\mathfrak{m}_i = \langle f_i(x) \rangle / \langle x^n - 1 \rangle \subseteq A.$$

The residue field at $\mathfrak{m}_i$ is

$$\kappa(\mathfrak{m}_i) \cong A/\mathfrak{m}_i \cong \mathbb{F}_q[x]/\langle f_i(x) \rangle.$$

Now let $\overline{a(x)} \in A$. Then $\overline{a(x)}$ determines a section over the closed points

$$s_{\overline{a}} : \{\mathfrak{m}_1, \dots, \mathfrak{m}_r\} \longrightarrow \prod_{i=1}^r \mathbb{F}_q[x]/\langle f_i(x) \rangle,$$

defined by

$$s_{\overline{a}}(\mathfrak{m}_i) = a(x) \bmod f_i(x).$$

Thus an element of the cyclic-code ring may be viewed as a function assigning to each closed point of $V(x^n - 1)$ its residue modulo the corresponding irreducible factor $f_i(x)$.

More generally, the full regular-function interpretation is

$$\tilde{a} : \text{Spec}(A) \longrightarrow \prod_{y \in \text{Spec}(A)} \kappa(y), \quad y \longmapsto \overline{a(x)}(y) \in \kappa(y).$$

If $x^n - 1$ is squarefree, then by the Chinese remainder theorem,

$$A \cong \prod_{i=1}^r \mathbb{F}_q[x]/\langle f_i(x) \rangle.$$

Hence in that case an element of A is literally a tuple

$$(a(x) \bmod f_1(x), \dots, a(x) \bmod f_r(x)).$$

At the same time, every element of A has a unique representative

$$a_0 + a_1x + \dots + a_{n-1}x^{n-1},$$

so as an $\mathbb{F}_q$ -vector space,

$$A \cong \mathbb{F}_q^n.$$

Under this identification, multiplication by x becomes cyclic shift of coordinates. Therefore the ideals of A are exactly the cyclic codes of length n over $\mathbb{F}_q$.

18 Regular functions versus tangent-bundle sections

In the previous sections, we interpreted an element $a \in R$ as a pointwise assignment

$$x \longmapsto a(x) \in \kappa(x), \quad x \in \operatorname{Spec}(R),$$

where $\kappa(x)$ is the residue field at the point x .

It is important to understand that these are *not* sections of the tangent bundle. Rather, they are sections of the *structure sheaf*.

The tangent-bundle analogue exists, but it is a different object.

18.1 The manifold analogy

Let M be a smooth manifold. A smooth function

$$f \in C^\infty(M)$$

assigns to each point $p \in M$ a scalar value

$$f(p) \in \mathbb{R}.$$

This is a section of the trivial line bundle of scalar-valued functions.

By contrast, a vector field

$$v \in \Gamma(M, TM)$$

assigns to each point $p \in M$ a tangent vector

$$v(p) \in T_p M.$$

Thus a vector field is a section of the tangent bundle.

So already in differential geometry we distinguish two kinds of sections:

$$\text{function: } p \mapsto f(p) \in \mathbb{R}, \quad \text{vector field: } p \mapsto v(p) \in T_p M.$$

18.2 Scheme-theoretic version

Now let $X = \operatorname{Spec}(R)$.

- A *regular function* on X is a section of the structure sheaf

$$\mathcal{O}_X.$$

Globally,

$$\Gamma(X, \mathcal{O}_X) = R.$$

Thus an element $a \in R$ is a global regular function.

- A *vector field* on X is a section of the tangent sheaf

$$\mathcal{T}_X.$$

Globally,

$$\Gamma(X, \mathcal{T}_X)$$

is generally a different R -module.

So the earlier pointwise interpretation

$$a \in R \quad \rightsquigarrow \quad (x \mapsto a(x) \in \kappa(x))$$

should be understood as the pointwise picture for a section of $\mathcal{O}_X$, not of $\mathcal{T}_X$.

Remark 18.1. Thus the correct analogy is

regular function $\leftrightarrow$ section of $\mathcal{O}_X$,	vector field $\leftrightarrow$ section of $\mathcal{T}_X$.
---	---

18.3 Tangent spaces at points

Let $x \in X = \operatorname{Spec}(R)$ correspond to a prime ideal $\mathfrak{p}_x$. The Zariski tangent space at x is

$$T_x X = \operatorname{Hom}_{\kappa(x)}(\mathfrak{m}_x / \mathfrak{m}_x^2, \kappa(x)) ,$$

where $\mathfrak{m}_x$ is the maximal ideal of the local ring $\mathcal{O}_{X,x}$.

A vector field is therefore something that assigns, at each point x , an element of $T_x X$, just as a regular function assigns an element of $\kappa(x)$.

So pointwise, one has the following comparison:

type of section	fiber over x
regular function	$\kappa(x)$
vector field	$T_x X$

18.4 Example: $X = \operatorname{Spec}(\mathbb{C}[x])$

Let

$$X = \operatorname{Spec}(\mathbb{C}[x]).$$

This is the affine line $\mathbb{A}_{\mathbb{C}}^1$.

Regular functions. A polynomial

$$f(x) \in \mathbb{C}[x]$$

defines a global section of $\mathcal{O}_X$. At a closed point $(x - \alpha)$, its value is

$$f(\alpha) \in \mathbb{C} \cong \kappa((x - \alpha)) .$$

At the generic point (0) , its value is

$$f(x) \in \mathbb{C}(x) = \kappa((0)) .$$

Tangent vectors. At a closed point $(x - \alpha)$, the tangent space is

$$T_{(x-\alpha)} X \cong \operatorname{Hom}_{\mathbb{C}}((x - \alpha) / (x - \alpha)^2, \mathbb{C}) .$$

Since $(x - \alpha) / (x - \alpha)^2$ is one-dimensional over $\mathbb{C}$, we obtain

$$T_{(x-\alpha)} X \cong \mathbb{C}.$$

Vector fields. The tangent sheaf on the affine line is generated by $\frac{d}{dx}$, so

$$\Gamma(X, \mathcal{T}_X) \cong \mathbb{C}[x] \frac{d}{dx}.$$

Hence a global vector field has the form

$$g(x) \frac{d}{dx}, \quad g(x) \in \mathbb{C}[x].$$

Thus:

- $f(x) \in \mathbb{C}[x]$ is a regular function;
- $g(x) \frac{d}{dx}$ is a tangent-bundle section.

Remark 18.2. This is the most useful concrete comparison:

$$f(x) \in \Gamma(X, \mathcal{O}_X), \quad f(x) \frac{d}{dx} \in \Gamma(X, \mathcal{T}_X).$$

So the tangent-bundle analogue of a polynomial $f(x)$ is not $f(x)$ itself, but $f(x) \frac{d}{dx}$.

18.5 Example: $X = \text{Spec}(\mathbb{C}[x]/(g(x)))$

Now let

$$X = \text{Spec}(\mathbb{C}[x]/(g(x))).$$

If $g(x)$ is squarefree, then X is a finite reduced set of points. In that case, each local ring is a field, so the tangent spaces are zero:

$$T_x X = 0 \quad \text{for all closed points } x.$$

Geometrically, this means that a finite reduced set of points has no infinitesimal directions.

If $g(x)$ has repeated roots, then X is nonreduced, and tangent spaces may become nonzero. For example,

$$X = \text{Spec}(\mathbb{C}[x]/((x - \alpha)^2))$$

has only one closed point, but it carries nontrivial infinitesimal structure. In particular, its tangent space at that point is nonzero.

Thus tangent spaces detect nilpotent or infinitesimal thickening that the underlying topological space alone does not see.

18.6 Example: the cyclic-code ring

Now consider

$$A = \mathbb{F}_q[x]/\langle x^n - 1 \rangle, \quad X = \text{Spec}(A).$$

As explained earlier, an element $\overline{a(x)} \in A$ is a regular function on X , that is, a section of $\mathcal{O}_X$. Pointwise it assigns values

$$y \mapsto \overline{a(x)}(y) \in \kappa(y), \quad y \in X.$$

At the same time, every element of A has a unique representative

$$a_0 + a_1x + \cdots + a_{n-1}x^{n-1},$$

so

$$A \cong \mathbb{F}_q^n$$

as an $\mathbb{F}_q$ -vector space.

Thus an element of the cyclic-code ring has two interpretations:

- as a regular function on the closed subscheme

$$V(x^n - 1) \subseteq \operatorname{Spec}(\mathbb{F}_q[x]);$$

- as a word

$$(a_0, a_1, \dots, a_{n-1}) \in \mathbb{F}_q^n.$$

If $x^n - 1$ is squarefree, then X is a finite reduced scheme, so its tangent spaces at closed points vanish. If $x^n - 1$ has repeated irreducible factors, then X is nonreduced, and tangent spaces may be nonzero. Thus tangent spaces again detect the nilpotent structure that is invisible at the level of the underlying set of points.

18.7 Summary dictionary

For a scheme $X = \operatorname{Spec}(R)$, one should compare:

object	fiber over x
regular function	$\kappa(x)$
vector field	$T_x X$

Thus:

sections discussed earlier are sections of $\mathcal{O}_X$, not of the tangent bundle.

The tangent-bundle analogue is obtained by replacing the residue-field value $a(x) \in \kappa(x)$ with a tangent vector $v(x) \in T_x X$.

Table E: regular functions versus tangent-bundle sections

Theme	Regular function	Vector field / tangent-bundle section
Bundle/sheaf	$\mathcal{O}_X$	$\mathcal{T}_X$
Global sections on $X = \text{Spec}(R)$	$\Gamma(X, \mathcal{O}_X) = R$	$\Gamma(X, \mathcal{T}_X)$
Fiber over a point x	$\kappa(x)$	$T_x X$
Pointwise value	$a(x) \in \kappa(x)$	$v(x) \in T_x X$
For $X = \text{Spec}(\mathbb{C}[x])$	$f(x) \in \mathbb{C}[x]$	$g(x) \frac{d}{dx} \in \mathbb{C}[x] \frac{d}{dx}$
For $X = \text{Spec}(\mathbb{C}[x]/(g))$, g squarefree	function on finitely many reduced points	zero tangent spaces at closed points
For $X = \text{Spec}(\mathbb{F}_q[x]/(x^n - 1))$	word in $\mathbb{F}_q^n$ / regular function on $V(x^n - 1)$	tangent directions detect repeated irreducible factors of $x^n - 1$