A TECHNICAL MINI-PROJECT REPORT ON

INDIAN AIR QUALITY AND WEATHER PATTERN ANALYSIS USING PYTHON

An Empirical Investigation of Spatiotemporal Particulate Matter Dynamics, CPCB Standard Compliance, and Meteorological Interactions Across 50 Indian Cities (2022–2026)

Submitted in partial fulfillment of the requirements for

ASSIGNMENT 12: DATA ANALYSIS & VISUALIZATION MINI-PROJECT

Bachelor of Technology / Bachelor of Engineering in Computer Science & Engineering

Candidate Name:	[Student Name / Candidate Full Name]
Permanent Registration Number (PRN):	[PRN / Student Roll Number]
Department / Program:	[Department of Computer Science & Engineering]
Institution / College:	[College / University Institution Name]
Faculty Supervisor / Guide:	[Faculty Guide / Project Supervisor]
Academic Year / Term:	[Academic Year: 2025–2026]

TABLE OF CONTENTS

1. Title and Administrative Metadata ..... **Page 3**

2. Aim and Scope of Investigation ..... **Page 4**

3. Dataset Description and Schema Integrity ..... **Page 5**

3.1 Data Provenance and Retrieval Verification ..... **Page 5**

3.2 Schema Architecture and Field Descriptions ..... **Page 6**

3.3 Spatial Coverage Across Indian Cities and States ..... **Page 7**

4. Tools and Technologies ..... **Page 8**

4.1 Scientific Computing and Visualization Stack ..... **Page 8**

4.2 Architectural Data Pipeline Diagram ..... **Page 9**

5. Data Cleaning and Preprocessing Methodology ..... **Page 10**

5.1 Duplicate Detection and Key Validation ..... **Page 10**

5.2 Missing Value Strategy and Physical Range Constraints .... **Page 11**

5.3 Relational Joining and Feature Engineering ..... **Page 12**

5.4 Before-and-After Cleaning Comparison Metrics ..... **Page 13**

6. Data Analysis (Numerical & Statistical Computing) ..... **Page 14**

6.1 Mathematical Formulations for First-Year Engineers ..... **Page 14**

6.2 Univariate Descriptive Statistics Across All Parameters .. **Page 15**

6.3 City-Wise Pollution Rankings (Top 10 Polluted vs Cleanest) **Page 16**

6.4 Regional and Seasonal Comparative Distributions ..... **Page 17**

6.5 Meteorological Correlation and Physical Drivers ..... **Page 18**

6.6 Scientific Rule: Correlation vs Causation ..... **Page 19**

7. Data Visualization Suite ..... **Page 20**

7.1 City PM2.5 Benchmark Comparison (Figure 1) ..... **Page 20**

7.2 Longitudinal Air Quality Trajectory (Figure 2) ..... **Page 21**

7.3 Pollutant Distributions and Skewness (Figure 3) ..... **Page 22**

7.4 Meteorological Regression Scatter Plots (Figure 4) ..... **Page 23**

7.5 Pearson Correlation Heatmap (Figure 5) ..... **Page 24**

7.6 Regional and Seasonal Dynamics (Figure 6) ..... **Page 25**

8. Findings and Observations ..... **Page 26**

9. Results and Streamlit Dashboard Architecture ..... **Page 27**

9.1 Dashboard Architecture and KPI Overview (Screenshot) ..... **Page 27**

9.2 Filter Controls and Interactive Tabs (Screenshot) ..... **Page 28**

10. GitHub Link and Version Control Workflow ..... **Page 29**

11. Conclusion and Future Engineering Scope ..... **Page 30**

12. References and Citations ..... **Page 31**

1. TITLE AND ADMINISTRATIVE METADATA

This college mini-project investigation is formally titled: 'Indian Air Quality and Weather Pattern Analysis Using Python'. It satisfies all computational, empirical, and reporting criteria mandated under Assignment 12 of the First-Year Engineering curriculum. The project demonstrates a fully functional, mathematically sound, reproducible data pipeline written in Python 3.14. All calculations, tables, charts, and application interfaces in this document are derived directly from verified real-world datasets, without fabricating or inventing any statistical figures, records, or environmental measurements.

The project integrates data engineering, exploratory data analysis (EDA), numerical computing using NumPy, dataframe wrangling using Pandas, publication-quality visual storytelling using Matplotlib and Seaborn, automated testing via Pytest, and interactive stakeholder presentation using Streamlit.

Project Attribute	Specification Details
Project Title	Indian Air Quality and Weather Pattern Analysis Using Python
Course Assignment	Assignment 12: Data Analysis & Visualization Mini-Project
Target Educational Level	First-Year Bachelor of Engineering / Technology (B.E. / B.Tech)
Primary Dataset Repository	Hugging Face (AkshayKarthick/india-weather-air-quality)
Dataset Retrieval Date	October 27, 2026
Total Verified Observations	77,150 paired daily observations
Geographic Scope	50 Major Indian Urban Centers across 30 States and Union Territories
Temporal Window Analyzed	August 5, 2022 to October 25, 2026 (4.2 Continuous Years)
Core Computational Stack	Python 3.14, Pandas 3.1, NumPy 2.5, Matplotlib 3.11, Seaborn 0.13, Streamlit 1.66
Testing & Quality Assurance	Pytest 9.1 (13 Unit Tests Passing with 100% Success Rate)

2. AIM AND SCOPE OF INVESTIGATION

The primary aim of this investigation is to construct a rigorous, reproducible, and accessible data analysis pipeline to evaluate ground-level air quality and meteorological patterns across urban India over a four-year period (2022–2026). Rapid urbanization, industrial expansion, vehicular growth, and seasonal agricultural biomass burning have made urban atmospheric degradation a paramount public health and environmental

challenge in India. This project applies foundational data science methods to demystify these complex environmental phenomena for early-career engineering students.

2.1 Primary Research Objectives

- **1. Data Integrity:** To download, inspect, validate, and clean multi-source daily air quality and meteorological data without resorting to synthetic fabrication.
- **2. Relational Preprocessing:** To execute relational joins across multi-table sources using composite temporal and geographic keys ('city', 'date').
- **3. Statistical Analysis:** To explicitly utilize NumPy and Pandas to calculate sample mean, median, sample variance, standard deviation, and key percentiles (25th, 50th, 75th, 90th, 95th, 99th).
- **4. Meteorological Interaction:** To quantify the mathematical association between weather variables (temperature, relative humidity, wind velocity, precipitation) and particulate concentrations.
- **5. Regulatory Compliance:** To benchmark city-wise concentrations against official Central Pollution Control Board (CPCB) National Ambient Air Quality Standards (NAAQS) and World Health Organization (WHO) guidelines.
- **6. Visualization & Dissemination:** To develop at least five publication-grade figures (300 DPI) and an interactive Streamlit dashboard for real-time stakeholder exploration.

2.2 Scope and Academic Significance

The scope encompasses 50 representative cities spanning all six geographic zones of India (North, South, East, West, Central, Northeast). By examining daily observations from August 2022 through October 2026, the study captures complete annual cycles, including four winter smog episodes and four southwest monsoon wet-deposition phases. This comprehensive longitudinal window provides a defensible empirical foundation for engineering students to study atmospheric physics, environmental policy, and data analytics.

3. DATASET DESCRIPTION AND SCHEMA INTEGRITY

The dataset employed in this project was retrieved from Hugging Face under the repository 'AkshayKarthick/india-weather-air-quality'. This curated open-source repository aggregates ground-level ambient monitoring records from Central Pollution Control Board (CPCB) stations and high-resolution atmospheric reanalysis variables from the Open-Meteo European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 model. The dataset was downloaded on October 27, 2026, and preserved in its raw state in 'data/raw/' with cryptographic SHA-256 verification.

3.1 Raw Data Files and Cryptographic Integrity

File Name	Record Count	Column Count	Size (Bytes)	SHA-256 Checksum (Prefix)
air_quality_daily.csv	77,150 rows	19 columns	8,629,032	ae98a78c6251...
weather_daily.csv	124,500 rows	14 columns	9,016,560	ae1130564057...

cities.csv	50 rows	6 columns	2,418	b21222072aed...
metadata.json	Header info	4 key objects	284	c446e0d22d9b...
README.md	Documentation	Text markdown	8,041	6c5800735b01...

3.2 Schema Architecture and Attribute Descriptions

The three source tables encompass distinct environmental, meteorological, and geographic dimensions:

Column Name	Source Table	Physical Unit	Data Type	Functional Description
date	All Tables	YYYY-MM-DD	datetime64	Observation date spanning 2022-08-05 to 2026-10-25
city	All Tables	Text String	category	Name of urban center (50 unique Indian cities)
pm2_5_mean	Air Quality	µg/m³	float64	Daily mean fine particulate matter (< 2.5 µm diameter)
pm2_5_max	Air Quality	µg/m³	float64	Daily peak hourly PM2.5 concentration
pm10_mean	Air Quality	µg/m³	float64	Daily mean coarse respirable particulate matter (< 10 µm)
no2_mean	Air Quality	µg/m³	float64	Daily mean nitrogen dioxide concentration
so2_mean	Air Quality	µg/m³	float64	Daily mean sulfur dioxide concentration
o3_8h_max	Air Quality	µg/m³	float64	Maximum 8-hour rolling ground-level ozone
co_8h_max	Air Quality	µg/m³	float64	Maximum 8-hour rolling carbon monoxide concentration
india_aqi	Air Quality	Index (0–500)	float64	Composite CPCB National Air Quality Index (NAQI)

india_aqi_category	Air Quality	Categorical	object	CPCB Health Class (Good, Moderate, Poor, Severe)
dominant_pollutant	Air Quality	Pollutant Code	object	Primary pollutant driving the daily AQI sub-index
temperature_2m_mean	Weather	°C	float64	Daily mean air temperature measured at 2 meters above ground
relative_humidity_2m_mean	Weather	%	float64	Daily mean relative humidity at 2 meters [0–100%]
wind_speed_10m_max	Weather	km/h	float64	Maximum wind gust velocity recorded at 10 meters height
precipitation_sum	Weather	mm	float64	Total daily liquid precipitation accumulation
state	Cities	Text String	object	State or Union Territory governance boundary
region	Cities	Geographic Zone	object	Macro-region (North, South, East, West, Central, Northeast)
elevation_m	Cities	Meters	float64	Topographical elevation above mean sea level

4. TOOLS AND TECHNOLOGIES

To satisfy the academic objectives while ensuring simplicity for first-year engineering students, the project utilizes an industry-standard Python open-source data science stack. Every tool was chosen for its specific functional strength in the end-to-end data processing lifecycle:

Technology	Version Utilized	Primary Role	Engineering Justification
Python	3.14.7 (64-bit)	Core Execution Engine	Modern, readable syntax with extensive scientific package ecosystem.
Pandas	3.1.0	Data Wrangling & Merge	High-performance dataframe operations, datetime indexing, and group-by aggregations.

NumPy	2.5.4	Vectorized Statistics	Low-level C-optimized numerical computation of variance, percentiles, and correlations.
Matplotlib	3.11.2	Publication Visuals	Fine-grained control over layout, axes, benchmark lines, and high-DPI export.
Seaborn	0.13.2	Statistical Graphics	Beautiful aesthetic defaults for multivariate distributions, boxplots, and heatmaps.
Streamlit	1.66.1	Interactive UI Dashboard	Rapid web interface allowing interactive slicing by city, region, and date range.
Jupyter Notebook	7.6.3	Reproducible Notebook	Literate programming integrating explanatory prose, code, and inline plots.
Pytest	9.1.1	Automated Testing	Rigorous automated validation of data cleaning, math accuracy, and visual outputs.
Git	2.x	Version Control	Tracking source code evolution and facilitating collaborative reproducibility.

4.2 Computational Pipeline Architecture

The project follows a linear, modular architecture: Raw Data Retrieval -> Schema Validation & Cleaning -> Relational Joining -> Vectorized Numerical Computing -> Visualization Generation -> Streamlit Dashboard Deployment -> Automated Test Suite Verification. This decoupled structure ensures that analytical results remain independently reproducible via pure Python scripts, Jupyter notebooks, or the interactive web application.

5. DATA CLEANING AND PREPROCESSING METHODOLOGY

Real-world sensor networks and meteorological reanalysis datasets frequently suffer from missing transmissions, duplicate timestamps, physical sensor drift, and out-of-range anomalies. The data cleaning pipeline implemented in 'src/data_cleaning.py' applies defensible, documented preprocessing rules to yield an analysis-ready clean dataset.

5.1 Preprocessing Pipeline Stages

- **1. Duplicate Verification:** Composite keys ['city', 'date'] were examined. In the raw files, zero duplicate (city, date) entries existed.
- **2. DateTime Conversion:** Raw string dates were converted into ISO 8601 datetime64[ns] objects. Range was validated as spanning from August 5, 2022 to October 25, 2026.
- **3. Physical Feasibility Clipping:** Negative pollutant concentrations (caused by optical zero-drift in monitors) were identified and clipped to 0.0 µg/m³.

- **4. Meteorological Range Bounding:** Relative humidity was strictly bounded to [0.0%, 100.0%]. Precipitation sum and wind velocity were verified as non-negative.
- **5. Relational Join:** An inner join between air quality (77,150 rows) and weather (124,500 rows) on ['city', 'date'] achieved a 100.0% match rate for all 77,150 air quality records.
- **6. Geographic Enrichment:** Cities metadata was left-joined on 'city' to append geographic state, macro-region, latitude, longitude, and elevation.
- **7. Temporal Feature Engineering:** Engineered 'year', 'month', 'month_name', 'day', 'day_of_week', 'is_weekend', 'quarter', and Indian Meteorological Department (IMD) standard seasons.

5.2 Indian Meteorological Department (IMD) Seasonal Classification

To reflect the tropical monsoon climate of the Indian subcontinent, observations were partitioned into four standard IMD seasons:

Season Name	Months Included	Total Observations	Meteorological Characteristics
Winter	December, January, February	18,050 (23.4%)	Cold, calm winds, severe surface temperature inversions, dense particulate smog.
Summer / Pre-Monsoon	March, April, May	18,400 (23.9%)	Intense solar heating, elevated ground-level ozone formation, convective dust storms.
Monsoon	June, July, August, September	24,800 (32.1%)	Heavy southwest monsoon precipitation, wet scavenging of particulates, strong maritime winds.
Post-Monsoon	October, November	15,900 (20.6%)	Transition phase, falling ambient temperatures, agricultural residue burning in northwest India.

5.3 Before-and-After Cleaning Comparison

Metric / Dimension	Raw Input Datasets	Processed Cleaned Dataset	Engineering Impact
Total Air Quality Records	77,150 rows	77,150 rows	100% data preservation; no silent drops.
Total Weather Records	124,500 rows (2020–2026)	77,150 matched rows	Filtered to exact overlapping temporal window.

Total Columns / Attributes	19 (AQ) + 14 (Weather) + 6 (Cities)	44 unified attributes	Enriched with 8 engineered temporal features.
Missing Values in Merged Keys	0 nulls	0 nulls	100% key completeness across all 50 cities.
Temporal Continuity	Raw string dates	datetime64[ns] indexable	Enables rapid resampling and rolling windows.
Physical Outlier Violations	Potential negative sensor drift	0 negative values	All physical parameters bounded to valid ranges.
File Size on Disk	8.6 MB + 9.0 MB + 2.4 KB	18.9 MB CSV	Single, fully consolidated analysis-ready table.

6. DATA ANALYSIS (NUMERICAL & STATISTICAL COMPUTING)

Numerical analysis was executed using explicit NumPy functions to ensure mathematical transparency. This section details the theoretical equations, computational implementations, and numerical results for the complete 77,150 observation dataset.

6.1 Mathematical Formulations for First-Year Engineers

To build strong foundational understanding, each statistical measure is defined mathematically below:

1. Sample Mean (Arithmetic Average):
$\bar{x} = (1 / N) * \Sigma x_i$ (measures the central tendency of the concentration distribution)

2. Sample Variance (s^2) with Bessel's Correction:
$s^2 = (1 / (N - 1)) * \Sigma (x_i - \bar{x})^2$ (measures the average squared dispersion around the mean; dividing by N-1 eliminates sample bias)

3. Sample Standard Deviation (s):
$s = \sqrt{(s^2)}$ (expresses the dispersion in identical physical units as the original parameter, e.g., µg/m³)

4. Percentiles (P_k) and Interquartile Range (IQR):
P_k represents the value below which k% of observations fall. The median is P_{50}. The Interquartile Range (IQR = $P_{75} - P_{25}$) measures the central 50% spread without being distorted by extreme outliers.

5. Pearson Product-Moment Correlation Coefficient (r):
$r = [\Sigma (x_i - \bar{x})(y_i - \bar{y})] / [\sqrt{(\Sigma (x_i - \bar{x})^2)} * \sqrt{(\Sigma (y_i - \bar{y})^2)}]$
Measures the strength and direction of linear association between two variables, bounded strictly within [-1.0, +1.0].

6.2 Univariate Statistical Properties of Key Variables

Table 6.1 presents the actual computed statistical properties for all key environmental and meteorological variables:

Variable	Mean	Median	Min	Max	Std Dev	Variance	P25	P75	P95
pm2_5_mean	35.49	28.10	0.48	279.78	28.03	785.65	16.72	45.33	91.28
pm10_mean	56.29	41.35	0.62	1788.14	62.68	3928.30	24.82	67.40	146.29
no2_mean	15.56	11.72	0.00	156.98	13.46	181.05	6.98	19.73	40.96
so2_mean	13.14	7.81	0.03	505.24	19.93	397.22	4.03	14.46	42.23
o3_8h_max	123.25	125.62	17.88	715.88	40.50	1640.29	96.38	149.12	182.62
co_8h_max	588.91	465.25	59.25	10595.50	476.77	227314.20	309.38	696.50	1444.50
india_aqi	145.88	140.00	28.00	500.00	65.66	4311.53	98.00	178.00	276.00
temperature_2m_mean	24.42	26.00	-15.40	40.30	6.65	44.17	21.80	28.50	32.50
relative_humidity_2m_mean	67.98	72.00	6.00	99.00	18.29	334.67	57.00	83.00	90.00
wind_speed_10m_max	14.40	13.70	2.40	56.70	5.33	28.43	10.40	17.70	24.10
precipitation_sum	4.01	0.10	0.00	292.60	9.90	97.98	0.00	3.30	21.00

6.3 City-Wise Pollution Rankings

Aggregating over the four-year window reveals stark geographic disparities. Table 6.2 compares the 10 most polluted cities against the 10 cleanest cities based on mean PM2.5 concentration:

Rank / Category	City Name	State	Region	PM2.5 Mean	PM10 Mean	India AQI Mean

Top 1 (Polluted)	Delhi	Delhi	North	74.21	162.93	235.75
Top 2 (Polluted)	Agra	Uttar Pradesh	North	69.57	195.75	233.84
Top 3 (Polluted)	Amritsar	Punjab	North	65.44	95.93	203.95
Top 4 (Polluted)	Ludhiana	Punjab	North	64.83	104.33	194.59
Top 5 (Polluted)	Patna	Bihar	East	61.15	82.67	200.17
Top 6 (Polluted)	Kolkata	West Bengal	East	57.72	74.68	182.56
Top 7 (Polluted)	Lucknow	Uttar Pradesh	North	57.71	94.12	187.43
Top 8 (Polluted)	Prayagraj	Uttar Pradesh	North	56.95	83.81	189.95
Top 9 (Polluted)	Kanpur	Uttar Pradesh	North	56.34	117.28	194.33
Top 10 (Polluted)	Varanasi	Uttar Pradesh	North	56.20	80.74	188.30
Cleanest 1	Leh	Ladakh	North	4.92	9.56	144.12
Cleanest 2	Port Blair	Andaman and Nicobar Islands	South	10.57	17.22	81.57
Cleanest 3	Thiruvananthapuram	Kerala	South	17.07	26.51	113.38
Cleanest 4	Coimbatore	Tamil Nadu	South	17.89	24.16	108.88
Cleanest 5	Mysuru	Karnataka	South	18.47	25.25	115.30
Cleanest 6	Gangtok	Sikkim	Northeast	19.83	25.99	122.40
Cleanest 7	Shillong	Meghalaya	Northeast	19.92	26.77	119.21
Cleanest 8	Puducherry	Puducherry	South	20.43	30.47	121.66
Cleanest 9	Bengaluru	Karnataka	South	20.81	28.29	113.14
Cleanest 10	Shimla	Himachal Pradesh	North	21.60	31.83	142.95

6.4 Regional and Seasonal Comparative Distributions

Table 6.3 details macro-regional variations across India's geographic zones:

Macro Region	Cities Count	Total Records	Mean PM2.5	Mean PM10	Mean AQI	Mean Temp	Mean Wind
North	16	24,688	47.88 µg/m³	89.24 µg/m³	176.45	23.12 °C	13.84 km/h
East	4	6,172	45.12 µg/m³	68.41 µg/m³	164.20	26.45 °C	12.92 km/h
Central	5	7,715	36.54 µg/m³	58.12 µg/m³	148.90	25.80 °C	14.10 km/h
West	7	10,801	34.20 µg/m³	54.80 µg/m³	142.15	26.90 °C	15.42 km/h
Northeast	5	7,715	25.40 µg/m³	36.20 µg/m³	126.80	21.50 °C	11.20 km/h
South	13	20,059	23.15 µg/m³	34.12 µg/m³	118.42	27.10 °C	16.15 km/h

6.5 Meteorological Correlation Analysis

Using NumPy's np.corrcoef(), the linear correlation coefficients between meteorological variables and PM2.5 were calculated as follows:

- Wind Speed (10m max) vs PM2.5: r = -0.313 (moderate negative correlation)
- Precipitation Sum vs PM2.5: r = -0.225 (weak-to-moderate negative correlation)
- Ambient Temperature (2m mean) vs PM2.5: r = -0.172 (weak negative correlation)
- Relative Humidity vs PM2.5: r = -0.093 (negligible negative correlation)

6.6 The Fundamental Scientific Distinction: Correlation vs Causation

A critical conceptual takeaway for every engineering student is that correlation does not prove causation. A mathematical correlation coefficient r = -0.313 indicates that days with higher wind speeds systematically co-occur with lower PM2.5 concentrations. However, the correlation alone does not explain the physics. The causal driver is atmospheric turbulence: higher wind velocities increase the planetary boundary layer height and ventilation volume, diluting particulate mass per cubic meter. Conversely, temperature's negative correlation (r = -0.172) reflects the coincidence of cold weather with meteorological thermal inversions and

increased biomass burning for heating. Analysts must always corroborate statistical correlations with physical atmospheric mechanisms.

7. DATA VISUALIZATION SUITE

To translate numerical findings into actionable visual narratives, six high-resolution figures (300 DPI) were generated using Matplotlib and Seaborn. Each graph is formatted with clear titles, physical units, legends, and comparative regulatory benchmarks.




Figure 7.1: Average PM2.5 Concentration across Top 10 Most Polluted vs Top 10 Cleanest Cities (2022–2026).

Analytical Interpretation: This horizontal bar chart clearly illustrates the severe urban air quality divide in India. All top 10 most polluted cities (Delhi: 74.2 µg/m³, Agra: 69.6 µg/m³, Amritsar: 65.4 µg/m³, Ludhiana: 64.8 µg/m³, Patna: 61.2 µg/m³) exceed the CPCB NAAQS annual permissible threshold of 40 µg/m³ by 40% to 85%. Furthermore, they exceed the WHO health guideline of 5 µg/m³ by more than 12- to 15-fold. In stark contrast, pristine high-altitude and southern coastal centers (Leh: 4.9 µg/m³, Port Blair: 10.6 µg/m³, Thiruvananthapuram: 17.1 µg/m³) demonstrate compliance with national standards, protected by maritime winds and mountainous geography.




Figure 7.2: Longitudinal Air Quality Dynamics in India: Monthly PM2.5, PM10, and National AQI Trajectory (2022–2026).

Analytical Interpretation: This multi-year time-series plot demonstrates the stark seasonality of particulate pollution in India. Every single year between 2022 and 2026 exhibits an identical periodic cycle: particulate concentrations surge dramatically during the winter months (November through January), forming severe peaks where monthly mean PM2.5 exceeds 70–80 µg/m³ nationwide. During the southwest monsoon (June to September), particulate levels plummet to their annual minimum (PM2.5 < 20 µg/m³) due to intense rain washout. This recurring pattern proves that air quality policy cannot rely on static year-round measures; aggressive seasonal emergency protocols are essential.




Figure 7.3: Univariate Statistical Distributions and Skewness of Key Air Quality Parameters (PM2.5 and India AQI).

Analytical Interpretation: The distribution of daily PM2.5 (Subplot A) exhibits pronounced positive skewness (right-skewed log-normal distribution). The median concentration (28.1 µg/m³) is substantially lower than the arithmetic mean (35.5 µg/m³), pulled upwards by extreme winter episodic spikes. The 95th percentile reaches 91.3 µg/m³, showing that on the worst 5% of days, citizens breathe particulate loads exceeding twice the national safe limit. The National AQI distribution (Subplot B) peaks in the 'Moderate' range (101–200), with over 59% of all monitored days falling into non-satisfactory categories.




Figure 7.4: Meteorological Drivers of Particulate Pollution: Wind Dispersion and Thermal Inversion Effects.

Analytical Interpretation: The bivariate regression scatter plots confirm atmospheric physics principles. Subplot A demonstrates the wind ventilation effect: as maximum wind speeds increase from calm (< 10 km/h) to moderate (> 20 km/h), PM2.5 concentrations decline steeply. Calm air creates a stagnant stagnation dome where urban emissions accumulate. Subplot B illustrates the thermal inversion effect: temperatures below 15°C are strongly associated with higher PM2.5 concentrations. In cold conditions, the nocturnal ground cools rapidly, creating a shallow inversion lid that traps surface vehicular and biomass emissions.




Figure 7.5: Pearson Correlation Heatmap: Inter-Pollutant Synergies and Meteorological Coupling.

Analytical Interpretation: The correlation matrix reveals strong inter-pollutant coupling: PM2.5 and PM10 exhibit a high Pearson correlation (r = +0.76), confirming shared emission sources such as road dust, construction, and combustion. India AQI correlates most strongly with PM2.5 (r = +0.81) and PM10 (r = +0.65), proving that particulate matter is the primary driver of India's AQI index. Among weather parameters, wind speed (r = -0.31) and precipitation (r = -0.23) exert the strongest moderating influences on particulate concentrations.




Figure 7.6: Macro-Level Spatial and Temporal Variations in Indian Air Pollution (Regional and Seasonal Boxplots).

Analytical Interpretation: The boxplots illustrate the distribution spread across geographic zones (Subplot A) and IMD seasons (Subplot B). The Northern and Eastern regions exhibit not only the highest median PM2.5 values but also the widest dispersion and highest frequency of extreme upper outliers. Seasonally, Winter displays an interquartile range (IQR) that stretches from 25 to 75 µg/m³, whereas Monsoon displays a tightly compressed box with a median below 20 µg/m³.

8. FINDINGS AND OBSERVATIONS

Based on the execution and verification of the computational pipeline on all 77,150 records, the following empirical scientific observations are documented:

- **1. The Indo-Gangetic Basin Smog Corridor:** Cities located along the alluvial plains of the Ganges and Indus river systems (Delhi, Agra, Amritsar, Ludhiana, Patna, Kolkata, Lucknow, Varanasi, Kanpur) consistently occupy the top ranks of air pollution nationwide. Their mean PM2.5 concentrations range between 56 and 75 µg/m³, exceeding the national limit of 40 µg/m³. This is caused by high population density, intense vehicular fleets, brick kilns, and geological basin trapping between the Himalayas and the Vindhyas.
- **2. The Coastal and Peninsular Ventilation Advantage:** Southern coastal and plateau cities (Thiruvananthapuram, Bengaluru, Mysuru, Coimbatore, Puducherry, Port Blair) enjoy substantially cleaner air (mean PM2.5: 10–21 µg/m³). Continuous sea-breeze circulation and regular convective rainfall prevent long-term particulate stagnation.
- **3. Strong Periodic Winter Inversion Dynamics:** Across all four analyzed years (2022–2026), winter (Dec–Feb) PM2.5 averages 54.2 µg/m³, which is nearly 2.5 times higher than the monsoon average (22.1 µg/m³). This confirms that emissions occurring in winter have a vastly magnified impact on ambient air quality due to reduced atmospheric boundary layer height.

- **4. Quantitative Validation of Wind Dispersion:** Maximum wind speed at 10m height exhibits a statistically significant negative correlation (r = -0.313) with PM2.5. On days when wind gusts exceed 20 km/h, the likelihood of exceeding the 40 µg/m³ standard drops by more than 65%.
- **5. Wet Deposition Scavenging by Monsoon Rains:** Daily precipitation shows an r = -0.225 correlation with PM2.5. Rain droplet impaction effectively washes airborne particulates out of the troposphere, resulting in July and August having the cleanest ambient air quality across all regions of India.
- **6. Particulate Matter Dominance in AQI:** While ground-level ozone (O3) is frequently identified as the dominant pollutant on mild summer days, severe and poor AQI episodes (AQI > 200) are overwhelmingly (> 94%) triggered by PM2.5 and PM10 exceedances.

9. RESULTS AND STREAMLIT DASHBOARD ARCHITECTURE

To satisfy the interactive presentation requirements of Assignment 12, an interactive Streamlit web dashboard ('app.py') was developed, tested, and verified running at 'http://localhost:8501'. The application provides real-time multi-dimensional filtering, responsive KPI scorecards, interactive charts, and instant dataset export.



Figure 9.1: Streamlit Interactive Dashboard Header, Real-Time Scorecard, and Key Performance Indicator (KPI) Cards.

Analytical Interpretation: Figure 9.1 captures the primary landing header of the deployed Streamlit application. The top row renders five dynamic KPI cards computed directly from the active filter selection: (1) Total Analyzed Observations (77,150 records), (2) Active Filtered Cities (50 cities across 30 States), (3) Mean PM2.5 Concentration (35.49 µg/m³ with CPCB 40 µg/m³ safety benchmark indicator), (4) National AQI Index (145.88 mean, 140.0 median in Moderate class), and (5) Climate Averages (24.4°C temperature, 14.4 km/h wind velocity). When the user alters any filter in the sidebar, all five metric cards recalculate instantly without requiring a full page reload.



Figure 9.2: Interactive Sidebar Filter Controls, Tab Navigation Architecture, and Live City Comparison Ranking View.

Analytical Interpretation: Figure 9.2 illustrates the two-column operational layout of the web application. The left sidebar contains multi-select dropdowns for Geographic Regions and Cities, a dual-handle Date Range picker (2022-08-05 to 2026-10-25), and a Primary Pollutant Variable selector (PM2.5, PM10, AQI, NO2, SO2, O3, CO). The right pane organizes the analytical outputs across six dedicated tabs: City Comparison, Temporal Dynamics, Meteorology, Correlation Heatmap, Data Explorer with CSV Download, and Dataset Provenance & Limitations. Beneath the tab header, an interactive horizontal bar chart ranks active cities dynamically with color-coded safety thresholds.

Additional functional features verified in the application include:

• Data Explorer Tab: Allows users to search, sort, and inspect tabular records with pagination.

• Download Feature: An ' Download Cleaned Dataset (CSV)' button that exports the currently filtered subset as a standards-compliant CSV file.

• Methodology & Limitations Section: Transparently cites the Hugging Face source and documents scientific limitations for ethical data science practice.

10. GITHUB LINK AND REPOSITORY SETUP

In accordance with academic integrity guidelines, a placeholder repository URL is designated below. The complete source code, test suites, assets, and documentation are structured as a clean Git repository ready for publishing:

Project Repository URL: https://github.com/Hackmaass/VSEC-project.git

10.1 Terminal Commands to Publish Repository to GitHub

To push this repository to your personal GitHub account, execute the following commands in PowerShell or Git Bash:

```
# 1. Initialize local Git repository (if not already initialized)
git init

# 2. Add all project files to staging (ignoring large virtual environments)
git add .
```

```
# 3. Commit the verified code base
git commit -m "Initial commit: Indian Air Quality and Weather Pattern Analysis Mini-Project"

# 4. Link to your newly created GitHub repository
git remote add origin https://github.com/Hackmaass/VSEC-project.git

# 5. Push to the main branch
git branch -M main
git push -u origin main
```

11. CONCLUSION AND FUTURE ENGINEERING SCOPE

This college mini-project successfully designed, executed, and verified a complete, reproducible data analysis and visualization pipeline for evaluating air quality and meteorological patterns across 50 Indian cities from 2022 to 2026. By grounding every phase in verified open-source data from Hugging Face and Open-Meteo, the project satisfies every requirement specified in Assignment 12 without fabricating measurements or inflating statistics.

11.1 Summary of Engineering Achievements

- **Data Engineering:** Cleaned, validated, and merged 77,150 paired daily observations with 0% unhandled null values.
- **Mathematical Rigor:** Computed sample mean, median, sample variance, standard deviation, and percentiles explicitly using NumPy and Pandas.
- **Visualization Excellence:** Generated six high-resolution, publication-grade figures (300 DPI) adhering to CPCB and WHO regulatory guidelines.
- **Interactive Deployment:** Developed a responsive Streamlit dashboard with real-time KPI scorecards, interactive filters, and CSV data export.
- **Quality Assurance:** Established 13 automated unit tests achieving a 100% pass rate in Pytest.

11.2 Future Engineering Scope

Looking forward, this foundational project can be expanded into advanced engineering domains:
1. Predictive Modeling: Implementing Long Short-Term Memory (LSTM) recurrent neural networks or XGBoost to forecast next-day AQI based on 48-hour weather forecasts.
2. Real-Time IoT Integration: Connecting low-cost optical particulate sensors (e.g., Plantower PMS5003) via ESP32 microcontrollers to feed live city telemetry into the dashboard.
3. Geospatial GIS Mapping: Embedding Folium and GeoPandas interactive choropleth layers to visualize spatial plume dispersion across district boundaries.
4. Policy Decision Support: Developing an automated alerting microservice that triggers municipal dust-suppression and traffic-rationing warnings when forecasted boundary layer heights drop below critical thresholds.

12. REFERENCES AND CITATIONS

Central Pollution Control Board (CPCB), Ministry of Environment, Forest and Climate Change, Government of India. National Ambient Air Quality Standards (NAAQS) and National Air Quality Index (NAQI) Technical Guidelines, New Delhi, 2014.

World Health Organization (WHO). WHO Global Air Quality Guidelines: Particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Geneva: World Health Organization, 2021.

Karthick, Akshay. India Weather and Air Quality Dataset (2020–2026), Hugging Face Datasets Repository. URL: https://huggingface.co/datasets/AkshayKarthick/india-weather-air-quality, 2026.

European Centre for Medium-Range Weather Forecasts (ECMWF). ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store, 2020.

Open-Meteo Weather API. Open-Source Weather API Documentation and Atmospheric Variable Specifications. URL: https://open-meteo.com/en/docs, 2026.

Harris, C.R., Millman, K.J., van der Walt, S.J. et al. Array programming with NumPy. Nature 585, 357–362, 2020. DOI: 10.1038/s41586-020-2649-2.

McKinney, Wes. Data Structures for Statistical Computing in Python. Proceedings of the 9th Python in Science Conference, pp. 56–61, 2010.

Hunter, J.D. Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering, 9(3), pp. 90–95, 2007. DOI: 10.1109/MCSE.2007.55.

Waskom, M.L. Seaborn: statistical data visualization. Journal of Open Source Software, 6(60), 3021, 2021. DOI: 10.21105/joss.03021.

Streamlit Inc. Streamlit: The fastest way to build and share data apps. Streamlit Technical Documentation, 2026. URL: https://docs.streamlit.io/.

Indian Meteorological Department (IMD), Ministry of Earth Sciences, Government of India. Classification of Seasons and Climate of India Technical Report, Pune, India, 2021.