

TUTORIAL 2

ROOT FINDING USING BISECTION METHOD

Numerical Analysis (ENME 602)

Assoc.Prof.Dr. Hesham H. Ibrahim

Eng. Nouran Adel

Table of Contents

- Algorithms Pseudo Code.
- Root Finding Problem Statement.
- Bisection Method.

Algorithms Pseudo Code

Algorithm: is an approximation procedure that describes a finite sequence of steps to be performed in a specified order.

The object of the algorithm is to implement a procedure to solve a problem or approximate a solution to the problem.

Pseudocode: used to describe an algorithm. It follows the rules of structured program construction.

It specifies the input, output and a stopping technique independent of the numerical technique. It can be easily translated in any programming language: e.g Matlab, Python or Java.

Algorithms Pseudo Code

Example: Write an algorithm to compute:

$$\sum_{i=1}^N x_i = x_1 + x_2 + \dots + x_N,$$

where N and the numbers x_1, x_2, \dots, x_N are given.

Pseudocode:

INPUT N, x_1, x_2, \dots, x_N .

OUTPUT $SUM = \sum_{i=1}^N x_i$.

Step 1 Set $SUM = 0$. (*Initialize accumulator.*)

Step 2 For $i = 1, 2, \dots, N$ do
 set $SUM = SUM + x_i$. (*Add the next term.*)


Step 3 **OUTPUT** (SUM);
STOP.

Root Finding Problem Statement

The root of an equation

The value of x that satisfies $f(x) = 0$ is called the root (solution) of the equation $f(x) = 0$ or the zero of the function f .

Graphically, it is the x-intercept when the graph of f crosses the x-axis.



Root Finding Problem Statement


The root-finding problem

is the process of approximating the root (solution) of the equation $f(x) = 0$ (to within a given tolerance).

Bisection Method

Idea of Bisection Method

When there is a root (red dot here), it occurs either in the right half of the interval $[a,b]$ (like in the graph) or in the left half or at the midpoint. Take the half that contains the root!



Bisection Method


We obtain an approximation of the root by repeatedly halving the subintervals of $[a,b]$, and "trapping" the root in smaller and smaller interval at each step.

Bisection Method

Q: How to know that a root exists in $[a, b]$?

Answer: apply the Intermediate Value Theorem:

If f is continuous on $[a, b]$ and $f(a)$ and $f(b)$ are of opposite signs (which is usually expressed as their product $f(a) \cdot f(b) < 0$) then there exist p such that $f(p) = 0$.



Bisection Method

Q: How to select the half that contains the root?


Construct the sequence of subintervals $[a_k, b_k]$ iteratively:

- ▶ Begin with first interval $[a_1, b_1] = [a, b]$ such that $f(a)f(b) < 0$ (so f has a root in $[a, b]$).
- ▶ Compute the midpoint $p = \frac{a+b}{2}$.
- ▶ To choose left or right subinterval: check the sign of $f(p)$ and compare it to the signs of $f(a)$ and $f(b)$.

If $f(a)f(p) < 0$ then f change signs in $[a, p]$, the new interval $[a_2, b_2] = [a, p]$, otherwise, take $[p, b]$.

- ▶ Set either $[a, p]$ or $[p, b]$ as new interval. At each iteration use p as new approximation of the root.

Bisection Method



Bisection Method

Q : How do we know when to stop iterations?

If we iterate until the n^{th} sub-interval length is at most ϵ :

$$\frac{b-a}{2^n} \leq \epsilon \quad (1)$$

The Theorem guarantees that the absolute error is at most ϵ :

$$|p - p_n| \leq \frac{b-a}{2^n} \leq \epsilon$$

To determine the number of iteration **n** necessary to reach a tolerance ϵ in $[a, b]$, solve (1) for **n**: taking \ln of both sides

$$\ln\left(\frac{b-a}{2^n}\right) < \ln(\epsilon)$$

$$\ln(b-a) - n \ln 2 < \ln(\epsilon) \implies n > \frac{\ln(b-a) - \ln(\epsilon)}{\ln 2} = \frac{\ln\left(\frac{b-a}{\epsilon}\right)}{\ln 2}$$

Bisection Method

Bisection Algorithm

To find a solution to $f(x) = 0$ given the continuous function f on the interval $[a, b]$, where $f(a)$ and $f(b)$ have opposite signs:

Bisection Method

INPUT endpoints a, b ; tolerance TOL ; maximum number of iterations N_0 .

OUTPUT approximate solution p or message of failure.

Step 1 Set $i = 1$;
 $FA = f(a)$.

Step 2 While $i \leq N_0$ do Steps 3–6.

Step 3 Set $p = a + (b - a)/2$; (Compute p_i .)
 $FP = f(p)$.

Step 4 If $FP = 0$ or $(b - a)/2 < TOL$ then
 OUTPUT (p); (Procedure completed successfully.)
 STOP.

Step 5 Set $i = i + 1$.

Step 6 If $FA \cdot FP > 0$ then set $a = p$; (Compute a_i, b_i .)
 $FA = FP$
 else set $b = p$. (FA is unchanged.)

Step 7 **OUTPUT** ('Method failed after N_0 iterations, $N_0 =', N_0$);
 (The procedure was unsuccessful.)
 STOP.

Problem 1

2. Let $f(x) = 3(x + 1) \left(x - \frac{1}{2}\right) (x - 1)$. Use the Bisection method on the following interval to find p_3 :
[-2, 1.5]

Problem 1

3. Use the Bisection method to find solutions to within 10^{-2} for $x^3 - 7x^2 + 14x - 6 = 0$ on the following interval:
[0, 1]

Problem 2

1. Sketch the graphs of $y = x$ and $y = \sin(x)$.

Problem 2

2. Use the Bisection method to find an approximation to within 10^{-5} to the first positive value of x with $x = 2 \sin(x)$.

Problem 2


4. Use Theorem 2.1 to find a bound for the number of iterations needed to achieve an approximation with accuracy 10^{-3} to the solution of $x^3 + x - 4 = 0$ lying in the interval $[1, 4]$. Find an approximation to the root with this degree of accuracy.

Problem 3

A trough of length L has a cross-section in the shape of a semicircle with radius r . (see the accompanying figure). When filled with water to within a distance h of the top, the volume V of the water is:

$$V = L \left[0.5\pi r^2 - r^2 \arcsin\left(\frac{h}{r}\right) - h(r^2 - h^2)^{\frac{1}{2}} \right]$$

Suppose $L = 10$ ft, $r = 1$ ft and $V = 10.4$ ft³. Find the depth of water in the trough to within 0.01 ft.



*Thank
you!*