

Results Summary

A summary of the results obtained from the analysis of MD simulations

Hadi Sedigh

2026-05-18

Table of contents

1	Results Summary	2
1.1	Distance and Radius of Gyration at 20 °C	2
1.2	BSA and Hbond at 20 °C	4
1.3	RMSF at 20 °C	5
1.4	RMSD at 20 °C	6

1 Results Summary

1.1 Distance and Radius of Gyration at 20 °C




Figure 1: (A) Distance between the center of mass (COM) of Twist-20 and Lsal-20 at 20 °C , (B) and the minimum distance between any two atoms of the complexes.




Figure 2: (C) and the minimum distance between the backbone atoms of the complexes. (D) Radius of gyration of subunits of Twist-20 and Lsal-20.

1.2 BSA and Hbond at 20 °C




Figure 3: (A) Buried surface area (BSA) of Twist-20 and Lsal-20 at 20 °C , (B) the number of hydrogen bonds between them.

1.3 RMSF at 20 °C




Figure 4: (A) RMSF of CA for Twest-20 and Lsal-20 at 20 °C aligned to their own reference structure, (B) RMSF of CA for Twest-20 and Lsal-20 at 20 °C aligned to the reference structure of the complex.

1.4 RMSD at 20 °C




Figure 5: RMSD of CA for Twist-20 and Lsal-20 at 20 °C .