



Fast solution of stiff PDEs in 1D, 2D and 3D periodic domains and on the sphere

ICOSAHOM 2016

Hadrien Montanelli & Niall Bootland

Numerical Analysis Group, University of Oxford

June 30, 2016





- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:**



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}
This leads to a system of N ODEs for the coefficients $\hat{u}(t) = \{\hat{u}_k(t)\}$,

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad t \in [0, T], \quad \hat{u}(0) = \hat{u}_0$$



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}
This leads to a system of N ODEs for the coefficients $\hat{u}(t) = \{\hat{u}_k(t)\}$,

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad t \in [0, T], \quad \hat{u}(0) = \hat{u}_0$$

Example (1D KdV): $u_t = -u_{xxx} - \frac{1}{2}(u^2)_x \implies \hat{u}' = ik^3\hat{u} - \frac{ik}{2}\mathcal{F}((\mathcal{F}^{-1}\hat{u})^2)$



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}
This leads to a system of N ODEs for the coefficients $\hat{u}(t) = \{\hat{u}_k(t)\}$,

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad t \in [0, T], \quad \hat{u}(0) = \hat{u}_0$$

Example (1D KdV): $u_t = -u_{xxx} - \frac{1}{2}(u^2)_x \implies \hat{u}' = ik^3\hat{u} - \frac{ik}{2}\mathcal{F}((\mathcal{F}^{-1}\hat{u})^2)$

- **Why is this challenging?**



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}
This leads to a system of N ODEs for the coefficients $\hat{u}(t) = \{\hat{u}_k(t)\}$,

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad t \in [0, T], \quad \hat{u}(0) = \hat{u}_0$$

Example (1D KdV): $u_t = -u_{xxx} - \frac{1}{2}(u^2)_x \implies \hat{u}' = ik^3\hat{u} - \frac{ik}{2}\mathcal{F}((\mathcal{F}^{-1}\hat{u})^2)$

- **Why is this challenging?** Stiffness, i.e., standard methods need very small time-steps (related to \mathbf{L} having large eigenvalues)



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}
This leads to a system of N ODEs for the coefficients $\hat{u}(t) = \{\hat{u}_k(t)\}$,

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad t \in [0, T], \quad \hat{u}(0) = \hat{u}_0$$

Example (1D KdV): $u_t = -u_{xxx} - \frac{1}{2}(u^2)_x \implies \hat{u}' = ik^3\hat{u} - \frac{ik}{2}\mathcal{F}((\mathcal{F}^{-1}\hat{u})^2)$

- **Why is this challenging?** Stiffness, i.e., standard methods need very small time-steps (related to \mathbf{L} having large eigenvalues)
- **Time-stepping scheme:**



- **Problem:** Stiff PDEs of the form (scalar or systems)

$$u_t = \mathcal{L}u + \mathcal{N}(u), \quad u(t=0, X) = u_0(X),$$

in 1D, 2D and 3D periodic domains and on the sphere

- **Space discretization:** Fourier spectral method in space (N equispaced points)

$$u(t, x) \approx \sum_{k=-N/2}^{N/2} \hat{u}_k(t) e^{ikx}, \quad \hat{u}_k(t) = \frac{1}{N} \sum_{j=0}^{N-1} u(t, x_j) e^{-ikx_j}$$

The linear part \mathcal{L} is discretized with Fourier differentiation matrices \Rightarrow matrix \mathbf{L}
This leads to a system of N ODEs for the coefficients $\hat{u}(t) = \{\hat{u}_k(t)\}$,

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad t \in [0, T], \quad \hat{u}(0) = \hat{u}_0$$

Example (1D KdV): $u_t = -u_{xxx} - \frac{1}{2}(u^2)_x \implies \hat{u}' = ik^3\hat{u} - \frac{ik}{2}\mathcal{F}((\mathcal{F}^{-1}\hat{u})^2)$

- **Why is this challenging?** Stiffness, i.e., standard methods need very small time-steps (related to \mathbf{L} having large eigenvalues)
- **Time-stepping scheme:** Exponential integrators





- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$



- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$

- **Two special features:** The matrix \mathbf{L} is diagonal and the stiffness comes from \mathbf{L}



- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$

- **Two special features:** The matrix \mathbf{L} is diagonal and the stiffness comes from \mathbf{L}
- **Key idea of exponential integrators:** Integrate \mathbf{L} exactly with matrix exponential (trivial) and apply a numerical scheme to \mathbf{N}



- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$

- **Two special features:** The matrix \mathbf{L} is diagonal and the stiffness comes from \mathbf{L}
- **Key idea of exponential integrators:** Integrate \mathbf{L} exactly with matrix exponential (trivial) and apply a numerical scheme to \mathbf{N}
- **Formula:** For given value \hat{u}_n at time $t_n = nh$, the numerical approximation \hat{u}_{n+1} at time $t_{n+1} = (n+1)h$ is given by

$$\hat{u}_{n+1} = e^{h\mathbf{L}}\hat{u}_n + h \sum_{i=1}^s B_i(h\mathbf{L})\mathbf{N}(\hat{v}_i)$$



- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$

- **Two special features:** The matrix \mathbf{L} is diagonal and the stiffness comes from \mathbf{L}
- **Key idea of exponential integrators:** Integrate \mathbf{L} exactly with matrix exponential (trivial) and apply a numerical scheme to \mathbf{N}
- **Formula:** For given value \hat{u}_n at time $t_n = nh$, the numerical approximation \hat{u}_{n+1} at time $t_{n+1} = (n+1)h$ is given by

$$\hat{u}_{n+1} = e^{h\mathbf{L}}\hat{u}_n + h \sum_{i=1}^s B_i(h\mathbf{L})\mathbf{N}(\hat{v}_i),$$

where the *stages* \hat{v}_i are defined by

$$\begin{aligned}\hat{v}_1 &= \hat{u}_n, \\ \hat{v}_2 &= e^{C_2 h \mathbf{L}} \hat{u}_n + h A_{2,1}(h \mathbf{L}) \mathbf{N}(\hat{v}_1), \\ \hat{v}_3 &= e^{C_3 h \mathbf{L}} \hat{u}_n + h A_{3,1}(h \mathbf{L}) \mathbf{N}(\hat{v}_1) + h A_{3,2}(h \mathbf{L}) \mathbf{N}(\hat{v}_2), \\ &\vdots\end{aligned}$$



- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$

- **Two special features:** The matrix \mathbf{L} is diagonal and the stiffness comes from \mathbf{L}
- **Key idea of exponential integrators:** Integrate \mathbf{L} exactly with matrix exponential (trivial) and apply a numerical scheme to \mathbf{N}
- **Formula:** For given value \hat{u}_n at time $t_n = nh$, the numerical approximation \hat{u}_{n+1} at time $t_{n+1} = (n+1)h$ is given by

$$\hat{u}_{n+1} = e^{h\mathbf{L}}\hat{u}_n + h \sum_{i=1}^s B_i(h\mathbf{L})\mathbf{N}(\hat{v}_i),$$

where the *stages* \hat{v}_i are defined by

$$\begin{aligned}\hat{v}_1 &= \hat{u}_n, \\ \hat{v}_2 &= e^{C_2 h \mathbf{L}} \hat{u}_n + h A_{2,1}(h \mathbf{L}) \mathbf{N}(\hat{v}_1), \\ \hat{v}_3 &= e^{C_3 h \mathbf{L}} \hat{u}_n + h A_{3,1}(h \mathbf{L}) \mathbf{N}(\hat{v}_1) + h A_{3,2}(h \mathbf{L}) \mathbf{N}(\hat{v}_2), \\ &\vdots\end{aligned}$$

Coefficients $A_{i,j}$, B_i and C_i are functions of \mathbf{L} (computed with contour integrals around each eigenvalue of \mathbf{L})



- **Problem:** System of ODEs of the form

$$\hat{u}'(t) = \mathbf{L}\hat{u} + \mathbf{N}(\hat{u}), \quad \hat{u}(0) = \hat{u}_0$$

- **Two special features:** The matrix \mathbf{L} is diagonal and the stiffness comes from \mathbf{L}
- **Key idea of exponential integrators:** Integrate \mathbf{L} exactly with matrix exponential (trivial) and apply a numerical scheme to \mathbf{N}
- **Formula:** For given value \hat{u}_n at time $t_n = nh$, the numerical approximation \hat{u}_{n+1} at time $t_{n+1} = (n+1)h$ is given by

$$\hat{u}_{n+1} = e^{h\mathbf{L}}\hat{u}_n + h \sum_{i=1}^s B_i(h\mathbf{L})\mathbf{N}(\hat{v}_i),$$

where the *stages* \hat{v}_i are defined by

$$\begin{aligned}\hat{v}_1 &= \hat{u}_n, \\ \hat{v}_2 &= e^{C_2 h \mathbf{L}} \hat{u}_n + h A_{2,1}(h \mathbf{L}) \mathbf{N}(\hat{v}_1), \\ \hat{v}_3 &= e^{C_3 h \mathbf{L}} \hat{u}_n + h A_{3,1}(h \mathbf{L}) \mathbf{N}(\hat{v}_1) + h A_{3,2}(h \mathbf{L}) \mathbf{N}(\hat{v}_2), \\ &\vdots\end{aligned}$$

Coefficients $A_{i,j}$, B_i and C_i are functions of \mathbf{L} (computed with contour integrals around each eigenvalue of \mathbf{L})

- **Cost per time-step:** $O(2sN \log(N))$



Method	Type	Order	Stages s	Steps q
ABNørsett4	ETD Adams-Bashforth	4	1	4
ABNørsett5	ETD Adams-Bashforth	5	1	5
ABNørsett6	ETD Adams-Bashforth	6	1	6
Friedli (VRK4)	ETD Runge-Kutta	4	4	1
Strehmel-Weiner	ETD Runge-Kutta	4	4	1
Cox-Matthews (ETDRK4)	ETD Runge-Kutta	4	4	1
Krogstad (ETDRK4-B)	ETD Runge-Kutta	4	4	1
Minchev	ETD Runge-Kutta	4	4	1
Hochbruck-Ostermann	ETD Runge-Kutta	4	5	1
Luan-Ostermann (EXPRK5S8)	ETD Runge-Kutta	5	8	1
(Mod)GenLawson41	(Mod.) Gen. Lawson	4	4	1
(Mod)GenLawson42	(Mod.) Gen. Lawson	4	4	2
(Mod)GenLawson43	(Mod.) Gen. Lawson	4	4	3
(Mod)GenLawson44	(Mod.) Gen. Lawson	5	4	4
(Mod)GenLawson45	(Mod.) Gen. Lawson	6	4	5
PEC423	Predictor-corrector	4	2	3
PECEC433	Predictor-corrector	4	3	3
PEC524	Predictor-corrector	5	2	4
PECEC534	Predictor-corrector	5	3	4
PEC625	Predictor-corrector	6	2	5
PECEC635	Predictor-corrector	6	3	5
PEC726	Predictor-corrector	7	2	6
PECEC736	Predictor-corrector	7	3	6



Method	Type	Order	Stages s	Steps q
ABNørsett4	ETD Adams-Bashforth	4	1	4
ABNørsett5	ETD Adams-Bashforth	5	1	5
ABNørsett6	ETD Adams-Bashforth	6	1	6
Friedli (VRK4)	ETD Runge-Kutta	4	4	1
Strehmel-Weiner	ETD Runge-Kutta	4	4	1
Cox-Matthews (ETDRK4)	ETD Runge-Kutta	4	4	1
Krogstad (ETDRK4-B)	ETD Runge-Kutta	4	4	1
Minchev	ETD Runge-Kutta	4	4	1
Hochbruck-Ostermann	ETD Runge-Kutta	4	5	1
Luan-Ostermann (EXPRK5S8)	ETD Runge-Kutta	5	8	1
(Mod)GenLawson41	(Mod.) Gen. Lawson	4	4	1
(Mod)GenLawson42	(Mod.) Gen. Lawson	4	4	2
(Mod)GenLawson43	(Mod.) Gen. Lawson	4	4	3
(Mod)GenLawson44	(Mod.) Gen. Lawson	5	4	4
(Mod)GenLawson45	(Mod.) Gen. Lawson	6	4	5
PEC423	Predictor-corrector	4	2	3
PECEC433	Predictor-corrector	4	3	3
PEC524	Predictor-corrector	5	2	4
PECEC534	Predictor-corrector	5	3	4
PEC625	Predictor-corrector	6	2	5
PECEC635	Predictor-corrector	6	3	5
PEC726	Predictor-corrector	7	2	6
PECEC736	Predictor-corrector	7	3	6

- ETD Adams-Bashforth and Lawson methods have bad stability properties



Method	Type	Order	Stages s	Steps q
ABNørsett4	ETD Adams-Bashforth	4	1	4
ABNørsett5	ETD Adams-Bashforth	5	1	5
ABNørsett6	ETD Adams-Bashforth	6	1	6
Friedli (VRK4)	ETD Runge-Kutta	4	4	1
Strehmel-Weiner	ETD Runge-Kutta	4	4	1
Cox-Matthews (ETDRK4)	ETD Runge-Kutta	4	4	1
Krogstad (ETDRK4-B)	ETD Runge-Kutta	4	4	1
Minchev	ETD Runge-Kutta	4	4	1
Hochbruck-Ostermann	ETD Runge-Kutta	4	5	1
Luan-Ostermann (EXPRK5S8)	ETD Runge-Kutta	5	8	1
(Mod)GenLawson41	(Mod.) Gen. Lawson	4	4	1
(Mod)GenLawson42	(Mod.) Gen. Lawson	4	4	2
(Mod)GenLawson43	(Mod.) Gen. Lawson	4	4	3
(Mod)GenLawson44	(Mod.) Gen. Lawson	5	4	4
(Mod)GenLawson45	(Mod.) Gen. Lawson	6	4	5
PEC423	Predictor-corrector	4	2	3
PECEC433	Predictor-corrector	4	3	3
PEC524	Predictor-corrector	5	2	4
PECEC534	Predictor-corrector	5	3	4
PEC625	Predictor-corrector	6	2	5
PECEC635	Predictor-corrector	6	3	5
PEC726	Predictor-corrector	7	2	6
PECEC736	Predictor-corrector	7	3	6

- ETD Adams-Bashforth and Lawson methods have bad stability properties
- ETD Runge-Kutta of order four have similar accuracy and stability properties (hard to beat ETDRK4), **EXPRK5S8** the most accurate but sometimes unstable



Method	Type	Order	Stages s	Steps q
ABNørsett4	ETD Adams-Bashforth	4	1	4
ABNørsett5	ETD Adams-Bashforth	5	1	5
ABNørsett6	ETD Adams-Bashforth	6	1	6
Friedli (VRK4)	ETD Runge-Kutta	4	4	1
Strehmel-Weiner	ETD Runge-Kutta	4	4	1
Cox-Matthews (ETDRK4)	ETD Runge-Kutta	4	4	1
Krogstad (ETDRK4-B)	ETD Runge-Kutta	4	4	1
Minchev	ETD Runge-Kutta	4	4	1
Hochbruck-Ostermann	ETD Runge-Kutta	4	5	1
Luan-Ostermann (EXPRK5S8)	ETD Runge-Kutta	5	8	1
(Mod)GenLawson41	(Mod.) Gen. Lawson	4	4	1
(Mod)GenLawson42	(Mod.) Gen. Lawson	4	4	2
(Mod)GenLawson43	(Mod.) Gen. Lawson	4	4	3
(Mod)GenLawson44	(Mod.) Gen. Lawson	5	4	4
(Mod)GenLawson45	(Mod.) Gen. Lawson	6	4	5
PEC423	Predictor-corrector	4	2	3
PECEC433	Predictor-corrector	4	3	3
PEC524	Predictor-corrector	5	2	4
PECEC534	Predictor-corrector	5	3	4
PEC625	Predictor-corrector	6	2	5
PECEC635	Predictor-corrector	6	3	5
PEC726	Predictor-corrector	7	2	6
PECEC736	Predictor-corrector	7	3	6

- ETD Adams-Bashforth and Lawson methods have bad stability properties
- ETD Runge-Kutta of order four have similar accuracy and stability properties (hard to beat ETDRK4), EXPRK5S8 the most accurate but sometimes unstable
- Predictor-Corrector methods of order ≥ 5 more accurate but often unstable





Example 1: ETD RK schemes for KdV equation from $t = 0$ to $t = 0.005$

We plot relative error = $\frac{\|u_{\text{approx}} - u_{\text{exact}}\|_{\infty}}{\|u_{\text{exact}}\|_{\infty}}$ vs relative time-step = $\frac{dt}{0.005}$ and time



Example 1: ETD RK schemes for KdV equation from $t = 0$ to $t = 0.005$

We plot relative error = $\frac{\|u_{\text{approx}} - u_{\text{exact}}\|_{\infty}}{\|u_{\text{exact}}\|_{\infty}}$ vs relative time-step = $\frac{dt}{0.005}$ and time







Example 2: ETD RK schemes for KdV equation from $t = 0$ to $t = 0.015$



Example 2: ETD RK schemes for KdV equation from $t = 0$ to $t = 0.015$







Example 3: ETD RK schemes for Gray-Scott equations in 2D from $t = 0$ to $t = 30$



Example 3: ETD RK schemes for Gray-Scott equations in 2D from $t = 0$ to $t = 30$







- **Preloaded demos:** Nine demos in 1D, four in 2D, four in 3D

```
u = spin('kdv');
```

```
u = spin2('gs2');
```

```
u = spin3('sh3');
```




- **Preloaded demos:** Nine demos in 1D, four in 2D, four in 3D

```
u = spin('kdv');
```

```
u = spin2('gs2');
```

```
u = spin3('sh3');
```

- **Define your own problem:**

```
dom = [0 32*pi]; tspan = [0 100];
```

```
S = spinop(dom, tspan);
```

```
S.linearPart = @(u) -diff(u,2)-diff(u,4);
```

```
S.nonlinearPart = @(u) -.5*diff(u.^2);
```


```
S.init = chebfun(@(x) cos(x/16).*(1 + sin(x/16)), dom, 'trig');
```

```
u = spin(S);
```






Our approach is based on the Double Fourier Sphere method:






Our approach is based on the Double Fourier Sphere method:



- When using longitude-latitude coordinate (λ, θ) , functions are 2π -periodic in λ but not periodic in θ




Our approach is based on the Double Fourier Sphere method:



- When using longitude-latitude coordinate (λ, θ) , functions are 2π -periodic in λ but not periodic in θ
- Double up the function and flip it




Our approach is based on the Double Fourier Sphere method:



- When using longitude-latitude coordinate (λ, θ) , functions are 2π -periodic in λ but not periodic in θ
- Double up the function and flip it
- Constant values along lines $\theta = 0$ and $\theta = \pm\pi$



Our approach is based on the Double Fourier Sphere method:



- When using longitude-latitude coordinate (λ, θ) , functions are 2π -periodic in λ but not periodic in θ
- Double up the function and flip it
- Constant values along lines $\theta = 0$ and $\theta = \pm\pi$

```
u = spherefun(@(lam,tt) 1./(1 + (cos(lam).*sin(tt)).^2) + cos(tt).^2);
```





- **Problem:** We want to solve

$$u_t = \mathcal{L}u + \mathcal{N}(u) = u_{\theta\theta} + \frac{\cos \theta}{\sin \theta} u_\theta + \frac{1}{\sin^2 \theta} u_{\lambda\lambda} + \mathcal{N}(u)$$

with the DFS method and exponential integrators



- **Problem:** We want to solve

$$u_t = \mathcal{L}u + \mathcal{N}(u) = u_{\theta\theta} + \frac{\cos \theta}{\sin \theta} u_\theta + \frac{1}{\sin^2 \theta} u_{\lambda\lambda} + \mathcal{N}(u)$$

with the DFS method and exponential integrators

- **Space discretization:** DFS in coefficient space with $m \times n$ points, leads to a block diagonal matrix \mathbf{L} (n blocks of size m)



- **Problem:** We want to solve

$$u_t = \mathcal{L}u + \mathcal{N}(u) = u_{\theta\theta} + \frac{\cos \theta}{\sin \theta} u_\theta + \frac{1}{\sin^2 \theta} u_{\lambda\lambda} + \mathcal{N}(u)$$

with the DFS method and exponential integrators

- **Space discretization:** DFS in coefficient space with $m \times n$ points, leads to a block diagonal matrix \mathbf{L} (n blocks of size m)
- **Matrix exponential:** Diagonalize $\mathbf{L} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^{-1}$ by block $\Rightarrow e^{h\mathbf{L}} = \mathbf{V}e^{h\mathbf{\Lambda}}\mathbf{V}^{-1}$, condition number $O(100)$, $O(nm^3)$ cost



- **Problem:** We want to solve

$$u_t = \mathcal{L}u + \mathcal{N}(u) = u_{\theta\theta} + \frac{\cos \theta}{\sin \theta} u_\theta + \frac{1}{\sin^2 \theta} u_{\lambda\lambda} + \mathcal{N}(u)$$

with the DFS method and exponential integrators

- **Space discretization:** DFS in coefficient space with $m \times n$ points, leads to a block diagonal matrix \mathbf{L} (n blocks of size m)
- **Matrix exponential:** Diagonalize $\mathbf{L} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^{-1}$ by block $\Rightarrow e^{h\mathbf{L}} = \mathbf{V} e^{h\mathbf{\Lambda}} \mathbf{V}^{-1}$, condition number $O(100)$, $O(nm^3)$ cost
- **ETDRK4 coefficients:** Same as previously, contour integrals around each eigenvalue of \mathbf{L}



- **Problem:** We want to solve

$$u_t = \mathcal{L}u + \mathcal{N}(u) = u_{\theta\theta} + \frac{\cos \theta}{\sin \theta} u_\theta + \frac{1}{\sin^2 \theta} u_{\lambda\lambda} + \mathcal{N}(u)$$

with the DFS method and exponential integrators

- **Space discretization:** DFS in coefficient space with $m \times n$ points, leads to a block diagonal matrix \mathbf{L} (n blocks of size m)
- **Matrix exponential:** Diagonalize $\mathbf{L} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^{-1}$ by block $\Rightarrow e^{h\mathbf{L}} = \mathbf{V}e^{h\mathbf{\Lambda}}\mathbf{V}^{-1}$, condition number $O(100)$, $O(nm^3)$ cost
- **ETDRK4 coefficients:** Same as previously, contour integrals around each eigenvalue of \mathbf{L}
- **Cost per time-step:** $O(nm^2)$





- In the past 15 years it has been discovered that the highly non-obvious formulas known as exponential integrators are the best tool for the accurate solution of stiff PDEs



- In the past 15 years it has been discovered that the highly non-obvious formulas known as exponential integrators are the best tool for the accurate solution of stiff PDEs
- We have compared 30 of them and found that the ETDRK4 formula of Cox & Matthews (2002) is hard to beat



- In the past 15 years it has been discovered that the highly non-obvious formulas known as exponential integrators are the best tool for the accurate solution of stiff PDEs
- We have compared 30 of them and found that the ETDRK4 formula of Cox & Matthews (2002) is hard to beat
- Chebfun's `spin` implements this, and it is the easiest and most efficient way to solve many stiff PDEs in 1D/2D/3D periodic domains



- In the past 15 years it has been discovered that the highly non-obvious formulas known as exponential integrators are the best tool for the accurate solution of stiff PDEs
- We have compared 30 of them and found that the ETDRK4 formula of Cox & Matthews (2002) is hard to beat
- Chebfun's **spin** implements this, and it is the easiest and most efficient way to solve many stiff PDEs in 1D/2D/3D periodic domains
- If you work with stiff PDEs, you definitely want to try **spin**



- In the past 15 years it has been discovered that the highly non-obvious formulas known as exponential integrators are the best tool for the accurate solution of stiff PDEs
- We have compared 30 of them and found that the ETDRK4 formula of Cox & Matthews (2002) is hard to beat
- Chebfun's **spin** implements this, and it is the easiest and most efficient way to solve many stiff PDEs in 1D/2D/3D periodic domains
- If you work with stiff PDEs, you definitely want to try **spin**
- Fourier spectral methods and exponential integrators can be used for solving PDEs on the sphere with the Double Fourier Sphere method (in coefficient space) and diagonalization by block