



## Numerical computation with periodic functions

8th International Congress on Industrial and Applied Mathematics, Beijing

Hadrien Montanelli (Oxford)

**Collaborators:** N.I. Gushterov, M. Javed, and L.N. Trefethen (Oxford),  
G.B. Wright (Boise)

August 10, 2015



**1** Extension of Chebfun to periodic functions

**2** Application 1: Nonlinear stiff PDEs

**3** Application 2: Choreographies



UNIVERSITY OF  
OXFORD

## Trigonometric Interpolation

---

<sup>1</sup>Wright, Javed, M., and Trefethen, "Extension of Chebfun to periodic functions", SISC, accepted.



# Trigonometric Interpolation

- $f$  Lipschitz continuous and periodic function on  $[0, 2\pi]$

---

<sup>1</sup>Wright, Javed, M., and Trefethen, "Extension of Chebfun to periodic functions", SISC, accepted.



## Trigonometric Interpolation

- $f$  Lipschitz continuous and periodic function on  $[0, 2\pi]$
- **Continuous world (true object):**  $f$  has a unique trigonometric series of the form

$$f(x) = \sum_{k=-\infty}^{\infty} a_k e^{ikx}, \quad a_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx$$

---

<sup>1</sup>Wright, Javed, M., and Trefethen, "Extension of Chebfun to periodic functions", SISC, accepted.



# Trigonometric Interpolation

- $f$  Lipschitz continuous and periodic function on  $[0, 2\pi]$
- **Continuous world (true object):**  $f$  has a unique trigonometric series of the form

$$f(x) = \sum_{k=-\infty}^{\infty} a_k e^{ikx}, \quad a_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx$$

- **Discrete world (approximation):**  $f$  has a unique trigonometric polynomial  $p_N$  that interpolates the data  $f_j = f(x_j)$  at the  $N$  points  $x_j = 2\pi j/N$  of the form

$$p_N(x) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikx}, \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} f_j e^{-ikx_j}$$

<sup>1</sup>Wright, Javed, M., and Trefethen, "Extension of Chebfun to periodic functions", SISC, accepted.



# Trigonometric Interpolation

- $f$  Lipschitz continuous and periodic function on  $[0, 2\pi]$
- **Continuous world (true object):**  $f$  has a unique trigonometric series of the form

$$f(x) = \sum_{k=-\infty}^{\infty} a_k e^{ikx}, \quad a_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx$$

- **Discrete world (approximation):**  $f$  has a unique trigonometric polynomial  $p_N$  that interpolates the data  $f_j = f(x_j)$  at the  $N$  points  $x_j = 2\pi j/N$  of the form

$$p_N(x) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikx}, \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} f_j e^{-ikx_j} = \sum_{j=-\infty}^{\infty} a_{k+jN}$$

<sup>1</sup>Wright, Javed, M., and Trefethen, "Extension of Chebfun to periodic functions", SISC, accepted.



# Trigonometric Interpolation

- $f$  Lipschitz continuous and periodic function on  $[0, 2\pi]$
- **Continuous world (true object):**  $f$  has a unique trigonometric series of the form


$$f(x) = \sum_{k=-\infty}^{\infty} a_k e^{ikx}, \quad a_k = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx$$

- **Discrete world (approximation):**  $f$  has a unique trigonometric polynomial  $p_N$  that interpolates the data  $f_j = f(x_j)$  at the  $N$  points  $x_j = 2\pi j/N$  of the form

$$p_N(x) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikx}, \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} f_j e^{-ikx_j} = \sum_{j=-\infty}^{\infty} a_{k+jN}$$

- Extension of Chebfun to periodic functions<sup>1</sup>:

```
f = @(x) tanh(5*cos(5*x));
p = chebfun(f,[0 2*pi],'trig');
plot(p), plotcoeffs(p)
```



<sup>1</sup>Wright, Javed, M., and Trefethen, "Extension of Chebfun to periodic functions", SISC, accepted.





UNIVERSITY OF  
OXFORD

## Periodic ODEs and e-value problems (1/3)



- **Problem:** We want to solve

$$\mathcal{L}u = f, \quad \mathcal{L}u = \lambda u,$$

with periodic operator  $\mathcal{L}$  on  $[0, 2\pi]$  defined by

$$\mathcal{L} = u'' + a(x)u' + b(x)u = \mathcal{D}^{(2)} + \mathcal{M}[a]\mathcal{D} + \mathcal{M}[b]$$



- **Problem:** We want to solve

$$\mathcal{L}u = f, \quad \mathcal{L}u = \lambda u,$$

with periodic operator  $\mathcal{L}$  on  $[0, 2\pi]$  defined by

$$\mathcal{L} = u'' + a(x)u' + b(x)u = \mathcal{D}^{(2)} + \mathcal{M}[a]\mathcal{D} + \mathcal{M}[b]$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size



## Periodic ODEs and e-value problems (1/3)

- **Problem:** We want to solve

$$\mathcal{L}u = f, \quad \mathcal{L}u = \lambda u,$$

with periodic operator  $\mathcal{L}$  on  $[0, 2\pi]$  defined by

$$\mathcal{L} = u'' + a(x)u' + b(x)u = \mathcal{D}^{(2)} + \mathcal{M}[a]\mathcal{D} + \mathcal{M}[b]$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size
- Discretize  $[0, 2\pi]$  with grid  $x_j = 2\pi j/N$ ,  $f$  with trigonometric polynomial  $f_N$  that interpolates  $f(x_j) = \text{ifft}(f_k)$ , and look for trigonometric polynomial solution  $u_N$

$$u_N(x) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} u_k e^{ikx}, \quad u_k = \text{fft}(u(x_j))$$



## Periodic ODEs and e-value problems (1/3)

- **Problem:** We want to solve

$$\mathcal{L}u = f, \quad \mathcal{L}u = \lambda u,$$

with periodic operator  $\mathcal{L}$  on  $[0, 2\pi]$  defined by

$$\mathcal{L} = u'' + a(x)u' + b(x)u = \mathcal{D}^{(2)} + \mathcal{M}[a]\mathcal{D} + \mathcal{M}[b]$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size
- Discretize  $[0, 2\pi]$  with grid  $x_j = 2\pi j/N$ ,  $f$  with trigonometric polynomial  $f_N$  that interpolates  $f(x_j) = \text{ifft}(f_k)$ , and look for trigonometric polynomial solution  $u_N$

$$u_N(x) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} u_k e^{ikx}, \quad u_k = \text{fft}(u(x_j))$$

- Leads to the following systems

$$\left( D_v^{(2)} + M_v[a]D_v + M_v[b] \right) \mathbf{u} = \mathbf{f} \qquad \left( D_c^{(2)} + M_c[a]D_c + M_c[b] \right) \mathbf{u} = \mathbf{f}$$

$$\mathbf{u} = \{u(x_j)\}, \quad \mathbf{f} = \{f(x_j)\}$$

$$\mathbf{u} = \{u_k\}, \quad \mathbf{f} = \{f_k\}$$



- **Problem:** We want to solve

$$\mathcal{L}u = f, \quad \mathcal{L}u = \lambda u,$$

with periodic operator  $\mathcal{L}$  on  $[0, 2\pi]$  defined by

$$\mathcal{L} = u'' + a(x)u' + b(x)u = \mathcal{D}^{(2)} + \mathcal{M}[a]\mathcal{D} + \mathcal{M}[b]$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size
- Discretize  $[0, 2\pi]$  with grid  $x_j = 2\pi j/N$ ,  $f$  with trigonometric polynomial  $f_N$  that interpolates  $f(x_j) = \text{ifft}(f_k)$ , and look for trigonometric polynomial solution  $u_N$

$$u_N(x) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} u_k e^{ikx}, \quad u_k = \text{fft}(u(x_j))$$

- Leads to the following systems

$$\left( D_v^{(2)} + M_v[a]D_v + M_v[b] \right) \mathbf{u} = \mathbf{f} \qquad \left( D_c^{(2)} + M_c[a]D_c + M_c[b] \right) \mathbf{u} = \mathbf{f}$$

$$\mathbf{u} = \{u(x_j)\}, \quad \mathbf{f} = \{f(x_j)\} \qquad \mathbf{u} = \{u_k\}, \quad \mathbf{f} = \{f_k\}$$

- Automatic Fréchet differentiation, and Newton's method in function space for nonlinear ODEs



UNIVERSITY OF  
OXFORD

## Periodic ODEs and e-value problems (2/3)



$$0.004u'' + uu' - u = \cos(2\pi x), \quad x \in [-1, 1]$$






$$0.004u'' + uu' - u = \cos(2\pi x), \quad x \in [-1, 1]$$

```
N = chebop(-1,1);  
N.op = @(x,u) .004*diff(u,2) + u.*diff(u) - u;  
N.bc = 'periodic';  
f = chebfun('cos(2*pi*x)','trig');  
u = N\f; plot(u)
```



$$0.004u'' + uu' - u = \cos(2\pi x), \quad x \in [-1, 1]$$

```
N = chebop(-1,1);
N.op = @(x,u) .004*diff(u,2) + u.*diff(u) - u;
N.bc = 'periodic';
f = chebfun('cos(2*pi*x)','trig');
u = N\f; plot(u)
```





UNIVERSITY OF  
OXFORD

## Periodic ODEs and e-value problems (3/3)



$$-u'' + 2q \cos(2x)u = \lambda u, \quad x \in [0, 2\pi]$$




$$-u'' + 2q \cos(2x)u = \lambda u, \quad x \in [0, 2\pi]$$

```
q=2;
L = chebop([0 2*pi]);
L.op = @(x,u) -diff(u,2) + 2*q*cos(2*x).*u;
L.bc = 'periodic';
[V,D] = eigs(L,5); plot(V)
```



$$-u'' + 2q \cos(2x)u = \lambda u, \quad x \in [0, 2\pi]$$

```
q=2;
L = chebop([0 2*pi]);
L.op = @(x,u) -diff(u,2) + 2*q*cos(2*x).*u;
L.bc = 'periodic';
[V,D] = eigs(L,5); plot(V)
```





UNIVERSITY OF  
OXFORD

## Linear PDEs (1/2)



- **Problem:** We want to solve

$$u_t(x, t) = \mathcal{L}u(x, t),$$

for some linear periodic operator  $\mathcal{L}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$





- **Problem:** We want to solve

$$u_t(x, t) = \mathcal{L}u(x, t),$$

for some linear periodic operator  $\mathcal{L}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- When the problem is well-posed, solution given by the operator exponential

$$u(x, t) = e^{\mathcal{L}t}u(x, 0)$$



## Linear PDEs (1/2)

- **Problem:** We want to solve

$$u_t(x, t) = \mathcal{L}u(x, t),$$

for some linear periodic operator  $\mathcal{L}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- When the problem is well-posed, solution given by the operator exponential

$$u(x, t) = e^{\mathcal{L}t}u(x, 0)$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size



## Linear PDEs (1/2)

- **Problem:** We want to solve

$$u_t(x, t) = \mathcal{L}u(x, t),$$

for some linear periodic operator  $\mathcal{L}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- When the problem is well-posed, solution given by the operator exponential

$$u(x, t) = e^{\mathcal{L}t} u(x, 0)$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size
- E.g., advection equation

$$u_t = \mathcal{L}u, \quad \mathcal{L} = a(x) \frac{\partial}{\partial x} = \mathcal{M}[a] \mathcal{D}$$



## Linear PDEs (1/2)

- **Problem:** We want to solve

$$u_t(x, t) = \mathcal{L}u(x, t),$$

for some linear periodic operator  $\mathcal{L}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- When the problem is well-posed, solution given by the operator exponential

$$u(x, t) = e^{\mathcal{L}t}u(x, 0)$$

- **Method:** Spectral method, based on values or coefficients, with adaptive grid size
- E.g., advection equation

$$u_t = \mathcal{L}u, \quad \mathcal{L} = a(x) \frac{\partial}{\partial x} = \mathcal{M}[a]\mathcal{D}$$

$$\mathbf{u}(\mathbf{t}) = e^{M_v[a]D_v t} \mathbf{u}(\mathbf{0})$$

$$\mathbf{u}(\mathbf{t}) = e^{M_c[a]D_c t} \mathbf{u}(\mathbf{0})$$

$$\mathbf{u}(\mathbf{t}) = \{u(x_j, t)\}, \quad \mathbf{u}(\mathbf{0}) = \{u(x_j, 0)\}$$

$$\mathbf{u}(\mathbf{t}) = \{u_k(t)\}, \quad \mathbf{u}(\mathbf{0}) = \{u_k(0)\}$$



UNIVERSITY OF  
OXFORD

## Linear PDEs (2/2)



## Linear PDEs (2/2)

$$u_t = -\left(\frac{1}{5} + \sin^2(x-1)\right)u_x, \quad x \in [0, 2\pi], \quad t \in [0, 20]$$



$$u_t = -\left(\frac{1}{5} + \sin^2(x-1)\right)u_x, \quad x \in [0, 2\pi], \quad t \in [0, 20]$$


```
c = chebfun(@(x) -(1/5 + sin(x-1).^2), [0 2*pi]);
L = chebop(@(x,u) c.*diff(u), [0 2*pi]);
L.bc = 'periodic';
u0 = chebfun(@(x) exp(-100*(x-1).^2), [0 2*pi]);
u = expm(L, 0:.5:20, u0), waterfall(u, 0:.5:20)
```



## Linear PDEs (2/2)

$$u_t = -\left(\frac{1}{5} + \sin^2(x-1)\right)u_x, \quad x \in [0, 2\pi], \quad t \in [0, 20]$$

```
c = chebfun(@(x) -(1/5 + sin(x-1).^2), [0 2*pi]);
L = chebop(@(x,u) c.*diff(u), [0 2*pi]);
L.bc = 'periodic';
u0 = chebfun(@(x) exp(-100*(x-1).^2), [0 2*pi]);
u = expm(L, 0:.5:20, u0), waterfall(u, 0:.5:20)
```







1 Extension of Chebfun to periodic functions

2 Application 1: Nonlinear stiff PDEs

3 Application 2: Choreographies



UNIVERSITY OF  
OXFORD

## Problem of interest



## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$



## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- In Chebfun: `pde15s`, based on the methods of lines, works for nonlinear PDEs but slow for some stiff PDEs, need another approach



## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- In Chebfun: `pde15s`, based on the methods of lines, works for nonlinear PDEs but slow for some stiff PDEs, need another approach
- **Method:** Discretization with a spectral method in space with  $M$  points leads to

$$u'(t) = Lu(t) + N(t, u(t)),$$

with  $u(t) = \{u_k(t)\}$ , i.e., a system of  $M$  ODEs



## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- In Chebfun: `pde15s`, based on the methods of lines, works for nonlinear PDEs but slow for some stiff PDEs, need another approach
- **Method:** Discretization with a spectral method in space with  $M$  points leads to

$$u'(t) = Lu(t) + N(t, u(t)),$$

with  $u(t) = \{u_k(t)\}$ , i.e., a system of  $M$  ODEs

- If  $N = 0$ ,

$$u(t) = e^{tL}u(x, 0)$$



## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- In Chebfun: `pde15s`, based on the methods of lines, works for nonlinear PDEs but slow for some stiff PDEs, need another approach
- **Method:** Discretization with a spectral method in space with  $M$  points leads to

$$u'(t) = Lu(t) + N(t, u(t)),$$

with  $u(t) = \{u_k(t)\}$ , i.e., a system of  $M$  ODEs

- If  $N = 0$ ,

$$u(t) = e^{tL}u(x, 0)$$

- If  $N$  is a constant,

$$u(t) = e^{tL}u_0 + L^{-1}[e^{tL} - I]N$$



## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- In Chebfun: `pde15s`, based on the methods of lines, works for nonlinear PDEs but slow for some stiff PDEs, need another approach
- **Method:** Discretization with a spectral method in space with  $M$  points leads to

$$u'(t) = Lu(t) + N(t, u(t)),$$

with  $u(t) = \{u_k(t)\}$ , i.e., a system of  $M$  ODEs

- If  $N = 0$ ,

$$u(t) = e^{tL}u(x, 0)$$

- If  $N$  is a constant,

$$u(t) = e^{tL}u_0 + L^{-1}[e^{tL} - I]N$$

- If  $N = N(t)$ ,

$$u(t) = e^{tL}u_0 + \int_0^t e^{(t-s)L}N(s)ds$$





## Problem of interest

- **Problem:** We want to solve nonlinear stiff PDEs of the form

$$u_t = \mathcal{L}u + \mathcal{N}(t, u(x, t)),$$

for some periodic op.  $\mathcal{L}$  and  $\mathcal{N}$  on  $[0, 2\pi] \times [0, T]$  and initial condition  $u(x, 0)$

- In Chebfun: `pde15s`, based on the methods of lines, works for nonlinear PDEs but slow for some stiff PDEs, need another approach
- **Method:** Discretization with a spectral method in space with  $M$  points leads to

$$u'(t) = Lu(t) + N(t, u(t)),$$

with  $u(t) = \{u_k(t)\}$ , i.e., a system of  $M$  ODEs

- If  $N = 0$ ,

$$u(t) = e^{tL}u(x, 0)$$

- If  $N$  is a constant,

$$u(t) = e^{tL}u_0 + L^{-1}[e^{tL} - I]N$$

- If  $N = N(t)$ ,

$$u(t) = e^{tL}u_0 + \int_0^t e^{(t-s)L}N(s)ds$$

- For general  $N$ , one need to find a way to capture the nonlinearity in a time-stepping method



UNIVERSITY OF  
OXFORD

## Exponential integrators



## Exponential integrators

- **Problem:** We want to solve a system of  $M$  ODEs

$$u'(t) = Lu(t) + N(t, u(t))$$



## Exponential integrators

- **Problem:** We want to solve a system of  $M$  ODEs

$$u'(t) = Lu(t) + N(t, u(t))$$

- **Method:** From the value  $u_n = u(t_n)$  at time  $t_n$ , the numerical approximation  $u_{n+1}$  at time  $t_{n+1} = t_n + h$  is given by the formula

$$u_{n+1} = e^{hL}u_n + h \sum_{i=1}^s B_i(hL)N(t_n + c_i h, v_i),$$

with internal stages  $v_1 = u_n$  and, for  $2 \leq i \leq s$ ,

$$v_i = e^{c_i hL}u_n + h \sum_{j=1}^{i-1} A_{i,j}(hL)N(t_n + c_j h, v_j)$$



## Exponential integrators

- **Problem:** We want to solve a system of  $M$  ODEs

$$u'(t) = Lu(t) + N(t, u(t))$$

- **Method:** From the value  $u_n = u(t_n)$  at time  $t_n$ , the numerical approximation  $u_{n+1}$  at time  $t_{n+1} = t_n + h$  is given by the formula

$$u_{n+1} = e^{hL}u_n + h \sum_{i=1}^s B_i(hL)N(t_n + c_i h, v_i),$$

with internal stages  $v_1 = u_n$  and, for  $2 \leq i \leq s$ ,

$$v_i = e^{c_i hL}u_n + h \sum_{j=1}^{i-1} A_{i,j}(hL)N(t_n + c_j h, v_j)$$

- Coefficients  $A_{i,j}$ ,  $B_i$  and  $c_i$  given via Butcher tableau



## Exponential integrators

- **Problem:** We want to solve a system of  $M$  ODEs

$$u'(t) = Lu(t) + N(t, u(t))$$

- **Method:** From the value  $u_n = u(t_n)$  at time  $t_n$ , the numerical approximation  $u_{n+1}$  at time  $t_{n+1} = t_n + h$  is given by the formula

$$u_{n+1} = e^{hL}u_n + h \sum_{i=1}^s B_i(hL)N(t_n + c_i h, v_i),$$

with internal stages  $v_1 = u_n$  and, for  $2 \leq i \leq s$ ,

$$v_i = e^{c_i hL}u_n + h \sum_{j=1}^{i-1} A_{i,j}(hL)N(t_n + c_j h, v_j)$$

- Coefficients  $A_{i,j}$ ,  $B_i$  and  $c_i$  given via Butcher tableau
- In Chebfun: available soon, use 4th- and 5th-order time-stepping schemes, easy to add another scheme, essentially just give the Butcher tableau



UNIVERSITY OF  
OXFORD

## Example 1: Kuramoto–Sivashinsky



## Example 1: Kuramoto–Sivashinsky

$$u_t = -u_{xx} - u_{xxxx} - uu_x, \quad x \in [0, 32\pi], \quad t \in [0, 200]$$

$$u(x, 0) = \cos(x/16)(1 + \sin(x/16))$$





## Example 1: Kuramoto–Sivashinsky

$$u_t = -u_{xx} - u_{xxxx} - uu_x, \quad x \in [0, 32\pi], \quad t \in [0, 200]$$

$$u(x, 0) = \cos(x/16)(1 + \sin(x/16))$$

```
S = spin('ks',{[0 32*pi], [0 200]});
u0 = chebfun('cos(x/16).*(1 + sin(x/16))',[0 32*pi],'trig');
u = solvepde(S,u0);
```




## Example 1: Kuramoto–Sivashinsky

$$u_t = -u_{xx} - u_{xxxx} - uu_x, \quad x \in [0, 32\pi], \quad t \in [0, 200]$$

$$u(x, 0) = \cos(x/16)(1 + \sin(x/16))$$

```
S = spin('ks',{[0 32*pi], [0 200]});
u0 = chebfun('cos(x/16).*(1 + sin(x/16))',[0 32*pi],'trig');
u = solvepde(S,u0);
```





UNIVERSITY OF  
OXFORD

## Example 2: Korteweg–de Vries



## Example 2: Korteweg–de Vries

$$u_t = -u_{xxx} - uu_x, \quad x \in [-\pi, \pi], \quad t \in [0, .01]$$

$$u(x, 0) = 3A^2 \operatorname{sech}^2\left(\frac{A(x+2)}{2}\right) + 3B^2 \operatorname{sech}^2\left(\frac{B(x+1)}{2}\right)$$



## Example 2: Korteweg–de Vries

$$u_t = -u_{xxx} - uu_x, \quad x \in [-\pi, \pi], \quad t \in [0, .01]$$

$$u(x, 0) = 3A^2 \operatorname{sech}^2\left(\frac{A(x+2)}{2}\right) + 3B^2 \operatorname{sech}^2\left(\frac{B(x+1)}{2}\right)$$

```
S = spin('kdv',{[-pi pi], [0 .01]}); A = 25; B = 16;
u0 = @(x) 3*A^2*sech(.5*A*(x+2)).^2+3*B^2*sech(.5*B*(x+1)).^2;
u0 = chebfun(u0,[-pi pi],'trig'); u = solvepde(S,u0);
```




## Example 2: Korteweg–de Vries

$$u_t = -u_{xxx} - uu_x, \quad x \in [-\pi, \pi], \quad t \in [0, .01]$$

$$u(x, 0) = 3A^2 \operatorname{sech}^2\left(\frac{A(x+2)}{2}\right) + 3B^2 \operatorname{sech}^2\left(\frac{B(x+1)}{2}\right)$$

```
S = spin('kdv',{[-pi pi], [0 .01]}); A = 25; B = 16;
u0 = @(x) 3*A^2*sech(.5*A*(x+2)).^2+3*B^2*sech(.5*B*(x+1)).^2;
u0 = chebfun(u0,[-pi pi],'trig'); u = solvepde(S,u0);
```





1 Extension of Chebfun to periodic functions

2 Application 1: Nonlinear stiff PDEs

**3 Application 2: Choreographies**



UNIVERSITY OF  
OXFORD

## Minimization of the action (Simó)



UNIVERSITY OF  
OXFORD

## Minimization of the action (Simó)

- Based on the principle of least action applied to the  $n$ -body problem



## Minimization of the action (Simó)

- Based on the principle of least action applied to the  $n$ -body problem

Choreographies of the  $n$ -body problem

$$z_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{z_i(t) - z_j(t)}{|z_i(t) - z_j(t)|^3} = 0, \quad 0 \leq j \leq n-1,$$

with  $z_j(t) = q\left(t + \frac{2\pi j}{n}\right)$  for some  $2\pi$ -periodic function  $q(t) : [0, 2\pi] \rightarrow \mathbb{C}$



## Minimization of the action (Simó)

- Based on the principle of least action applied to the  $n$ -body problem

Choreographies of the  $n$ -body problem

$$z_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{z_i(t) - z_j(t)}{|z_i(t) - z_j(t)|^3} = 0, \quad 0 \leq j \leq n-1,$$

with  $z_j(t) = q\left(t + \frac{2\pi j}{n}\right)$  for some  $2\pi$ -periodic function  $q(t) : [0, 2\pi] \rightarrow \mathbb{C}$

## Minima of the action

$$A = \int_0^{2\pi} (K(t) - U(t)) dt,$$

$$K(t) = \frac{n}{2} |q'(t)|^2, \quad U(t) = -n \sum_{j=1}^{n-1} \left| q(t) - q\left(t + \frac{2\pi j}{n}\right) \right|^{-1}$$



UNIVERSITY OF  
OXFORD

## Method

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.



- *Trigonometric interpolation:*  $q(t) \rightarrow p_N(t) \rightarrow A_N$

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.



- *Trigonometric interpolation:*  $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$  instead of  $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.



- *Trigonometric interpolation:*  $q(t) \rightarrow p_N(t) \rightarrow A_N$

$$q(t) \approx p_N(t) \text{ instead of } q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$$

- *A closed-form expression for the gradient:*  $p_N(t) \rightarrow \nabla A_N$

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.



- *Trigonometric interpolation:*  $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$  instead of  $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- *A closed-form expression for the gradient:*  $p_N(t) \rightarrow \nabla A_N$

- *Optimization algorithm:*  $p_N(t) \rightarrow p_N^*(t)$

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.





- *Trigonometric interpolation:*  $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$  instead of  $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- *A closed-form expression for the gradient:*  $p_N(t) \rightarrow \nabla A_N$

- *Optimization algorithm:*  $p_N(t) \rightarrow p_N^*(t)$

Quasi-Newton methods (BFGS) instead of Gradient methods

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.



- *Trigonometric interpolation:*  $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$  instead of  $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- *A closed-form expression for the gradient:*  $p_N(t) \rightarrow \nabla A_N$

- *Optimization algorithm:*  $p_N(t) \rightarrow p_N^*(t)$

Quasi-Newton methods (BFGS) instead of Gradient methods

- Can be generalized to the sphere using stereographic projection <sup>2</sup>

---

<sup>2</sup>M. and Gushterov, "Computing planar and spherical choreographies", SIADS, submitted.



UNIVERSITY OF  
OXFORD

## Numerical results




## Numerical results

- At convergence check  $\|\nabla A_N\|_2$ ,  $|c_k|$ , and  $\|\text{residual}\|_\infty$



## Numerical results

- At convergence check  $\|\nabla A_N\|_2$ ,  $|c_k|$ , and  $\|\text{residual}\|_\infty$



Action	68.8516	71.3312	77.1588	88.4397	109.6366	119.3191
Computer time (s)	0.79	0.49	0.44	0.70	0.98	0.86
2-norm of the gradient	1.26e-02	1.39e-02	8.87e-03	9.68e-03	1.18e-02	1.28e-02
Smallest coefficient	4.71e-06	6.45e-08	2.26e-06	3.33e-06	2.08e-05	2.75e-05
$\infty$ -norm of the residual	9.31e-02	1.09e-03	1.30e-02	4.43e-02	2.83e-01	6.56e-01



UNIVERSITY OF  
OXFORD

## Conclusion and future work



## Conclusion and future work

- Chebfun now uses trigonometric interpolation for periodic problems



## Conclusion and future work

- Chebfun now uses trigonometric interpolation for periodic problems
- It includes the solution of linear and nonlinear ODEs, e-value problems, and linear and nonlinear stiff PDEs





## Conclusion and future work

- Chebfun now uses trigonometric interpolation for periodic problems
- It includes the solution of linear and nonlinear ODEs, e-value problems, and linear and nonlinear stiff PDEs
- Trigonometric interpolation can also be used when looking for periodic solutions of dynamical systems



## Conclusion and future work

- Chebfun now uses trigonometric interpolation for periodic problems
- It includes the solution of linear and nonlinear ODEs, e-value problems, and linear and nonlinear stiff PDEs
- Trigonometric interpolation can also be used when looking for periodic solutions of dynamical systems
- Future work includes the automatic computation of adjoint for optimal control



## Conclusion and future work

- Chebfun now uses trigonometric interpolation for periodic problems
- It includes the solution of linear and nonlinear ODEs, e-value problems, and linear and nonlinear stiff PDEs
- Trigonometric interpolation can also be used when looking for periodic solutions of dynamical systems
- Future work includes the automatic computation of adjoint for optimal control

$$Lu = \sum_{k=0}^n a_k u^{(k)},$$

$$L^* v = \sum_{k=0}^n (-1)^k (\overline{a_k} v)^{(k)}$$



## Conclusion and future work

- Chebfun now uses trigonometric interpolation for periodic problems
- It includes the solution of linear and nonlinear ODEs, e-value problems, and linear and nonlinear stiff PDEs
- Trigonometric interpolation can also be used when looking for periodic solutions of dynamical systems
- Future work includes the automatic computation of adjoint for optimal control

$$Lu = \sum_{k=0}^n a_k u^{(k)},$$

$$L^* v = \sum_{k=0}^n (-1)^k (\overline{a_k} v)^{(k)}$$

```
L = chebop([0 2*pi]);
L.op = @(t,u) diff(u) + cos(2*t).*u;
L.bc = 'periodic';
L'
```