



Computing planar and spherical choreographies

17th Biennial Oxford/Cambridge Applied Mathematics Meeting

Hadrien Montanelli (Oxford)

Collaborator: Nikola I. Gushterov (Oxford)

June 10, 2015



1 Introduction

2 Computing planar choreographies

3 Computing spherical choreographies



UNIVERSITY OF
OXFORD

Minimization of the action (Simó)



Minimization of the action (Simó)

- Based on the principle of least action applied to the n -body problem



Minimization of the action (Simó)

- Based on the principle of least action applied to the n -body problem

Choreographies of the n -body problem

$$z_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{z_i(t) - z_j(t)}{|z_i(t) - z_j(t)|^3} = 0, \quad 0 \leq j \leq n-1,$$

with $z_j(t) = q\left(t + \frac{2\pi j}{n}\right)$ for some 2π -periodic function $q(t) : [0, 2\pi] \rightarrow \mathbb{C}$



Minimization of the action (Simó)

- Based on the principle of least action applied to the n -body problem

Choreographies of the n -body problem

$$z_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{z_i(t) - z_j(t)}{|z_i(t) - z_j(t)|^3} = 0, \quad 0 \leq j \leq n-1,$$

with $z_j(t) = q\left(t + \frac{2\pi j}{n}\right)$ for some 2π -periodic function $q(t) : [0, 2\pi] \rightarrow \mathbb{C}$

Minima of the action

$$A = \int_0^{2\pi} (K(t) - U(t)) dt,$$

$$K(t) = \frac{n}{2} |q'(t)|^2, \quad U(t) = -n \sum_{j=1}^{n-1} \left| q(t) - q\left(t + \frac{2\pi j}{n}\right) \right|^{-1}$$



1 Introduction

2 Computing planar choreographies

3 Computing spherical choreographies



UNIVERSITY OF
OXFORD

Method



- *Trigonometric interpolation:* $q(t) \rightarrow p_N(t) \rightarrow A_N$



- **Trigonometric interpolation:** $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$ instead of $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$



- *Trigonometric interpolation:* $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$ instead of $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- *A closed-form expression for the gradient:* $p_N(t) \rightarrow \nabla A_N$



- *Trigonometric interpolation:* $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$ instead of $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- *A closed-form expression for the gradient:* $p_N(t) \rightarrow \nabla A_N$
- *Optimization algorithm:* $p_N(t) \rightarrow p_N^*(t)$



- **Trigonometric interpolation:** $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$ instead of $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- **A closed-form expression for the gradient:** $p_N(t) \rightarrow \nabla A_N$

- **Optimization algorithm:** $p_N(t) \rightarrow p_N^*(t)$

Quasi-Newton methods (BFGS) instead of Gradient methods



Method


- **Trigonometric interpolation:** $q(t) \rightarrow p_N(t) \rightarrow A_N$

$q(t) \approx p_N(t)$ instead of $q(t) = x(t) + iy(t) \approx x_N(t) + iy_N(t)$

- **A closed-form expression for the gradient:** $p_N(t) \rightarrow \nabla A_N$

- **Optimization algorithm:** $p_N(t) \rightarrow p_N^*(t)$

Quasi-Newton methods (BFGS) instead of Gradient methods





Trigonometric Interpolation: $q(t) \rightarrow p_N(t) \rightarrow A_N$

¹Wright, Javed, Montanelli, and Trefethen, "Extension of Chebfun to periodic functions", 2015.

Trigonometric Interpolation: $q(t) \rightarrow p_N(t) \rightarrow A_N$

- Trigonometric interpolant $p_N(t)$ of $q(t)$ at N points:

$$p_N(t) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikt}, \quad t \in [0, 2\pi], \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} q_j e^{-ikt_j}, \quad |k| \leq \frac{N-1}{2}$$

¹Wright, Javed, Montanelli, and Trefethen, "Extension of Chebfun to periodic functions", 2015.

Trigonometric Interpolation: $q(t) \rightarrow p_N(t) \rightarrow A_N$

- Trigonometric interpolant $p_N(t)$ of $q(t)$ at N points:

$$p_N(t) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikt}, \quad t \in [0, 2\pi], \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} q_j e^{-ikt_j}, \quad |k| \leq \frac{N-1}{2}$$

- Recent extension of Chebfun to periodic functions ¹

¹Wright, Javed, Montanelli, and Trefethen, "Extension of Chebfun to periodic functions", 2015.


Trigonometric Interpolation: $q(t) \rightarrow p_N(t) \rightarrow A_N$

- Trigonometric interpolant $p_N(t)$ of $q(t)$ at N points:

$$p_N(t) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikt}, \quad t \in [0, 2\pi], \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} q_j e^{-ikt_j}, \quad |k| \leq \frac{N-1}{2}$$

- Recent extension of Chebfun to periodic functions ¹

```
>> q = chebfun(@(t)cos(3*pi*t)+1i*sin(2*pi*t),'trig')
q =
chebfun column (1 smooth piece)
      interval      length  endpoint values trig
[      -1,         1]         7      complex values
```



¹Wright, Javed, Montanelli, and Trefethen, "Extension of Chebfun to periodic functions", 2015.


Trigonometric Interpolation: $q(t) \rightarrow p_N(t) \rightarrow A_N$

- Trigonometric interpolant $p_N(t)$ of $q(t)$ at N points:

$$p_N(t) = \sum_{k=-\frac{N-1}{2}}^{\frac{N-1}{2}} c_k e^{ikt}, \quad t \in [0, 2\pi], \quad c_k = \frac{1}{N} \sum_{j=0}^{N-1} q_j e^{-ikt_j}, \quad |k| \leq \frac{N-1}{2}$$

- Recent extension of Chebfun to periodic functions ¹

```
>> q = chebfun(@(t)cos(3*pi*t)+1i*sin(2*pi*t),'trig')
q =
chebfun column (1 smooth piece)
      interval      length  endpoint values trig
[      -1,         1]         7      complex values
```



- Action computed with the exponentially accurate trapezoidal rule

$$A_N = \frac{n}{2} \int_0^{2\pi} |p'_N(t)|^2 dt + n \sum_{j=1}^{n-1} \int_0^{2\pi} \left| p_N(t) - p_N\left(t + \frac{2\pi j}{n}\right) \right|^{-1} dt$$

¹Wright, Javed, Montanelli, and Trefethen, "Extension of Chebfun to periodic functions", 2015.




UNIVERSITY OF
OXFORD

Numerical results



Numerical results



Action	68.8516	71.3312	77.1588	88.4397	109.6366	119.3191
Computer time (s)	0.79	0.49	0.44	0.70	0.98	0.86
2-norm of the gradient	1.26e-02	1.39e-02	8.87e-03	9.68e-03	1.18e-02	1.28e-02
Smallest coefficient	4.71e-06	6.45e-08	2.26e-06	3.33e-06	2.08e-05	2.75e-05
∞ -norm of the residual	9.31e-02	1.09e-03	1.30e-02	4.43e-02	2.83e-01	6.56e-01



- 1 Introduction
- 2 Computing planar choreographies
- 3 Computing spherical choreographies**



UNIVERSITY OF
OXFORD

Spherical n -body problem

Spherical n -body problem

Equations of the motion

$$X_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{R^3 X_i(t) - R(X_i(t) \cdot X_j(t))X_j(t)}{[R^4 - (X_i(t) \cdot X_j(t))^2]^{3/2}} + R^{-2} \|X_j'(t)\|^2 X_j(t) = 0,$$

with $X_j(t) \in \mathbb{S}_R^2 = \{X \in \mathbb{R}^3, \|X\| = R\} \subset \mathbb{R}^3, 0 \leq j \leq n-1$

Spherical n -body problem

Equations of the motion

$$X_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{R^3 X_i(t) - R(X_i(t) \cdot X_j(t))X_j(t)}{[R^4 - (X_i(t) \cdot X_j(t))^2]^{3/2}} + R^{-2} \|X_j'(t)\|^2 X_j(t) = 0,$$

with $X_j(t) \in \mathbb{S}_R^2 = \{X \in \mathbb{R}^3, \|X\| = R\} \subset \mathbb{R}^3$, $0 \leq j \leq n-1$

Potential

$$U(t) = -\frac{1}{R} \sum_{j=0}^{n-1} \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \cot \frac{d_{GC}(X_i(t), X_j(t))}{R},$$

Spherical n -body problem

Equations of the motion

$$X_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{R^3 X_i(t) - R(X_i(t) \cdot X_j(t))X_j(t)}{[R^4 - (X_i(t) \cdot X_j(t))^2]^{3/2}} + R^{-2} \|X_j'(t)\|^2 X_j(t) = 0,$$

with $X_j(t) \in \mathbb{S}_R^2 = \{X \in \mathbb{R}^3, \|X\| = R\} \subset \mathbb{R}^3, 0 \leq j \leq n-1$

Potential

$$U(t) = -\frac{1}{R} \sum_{j=0}^{n-1} \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \cot \frac{d_{GC}(X_i(t), X_j(t))}{R},$$

with

$$d_{GC}(X_i(t), X_j(t)) = R \arccos \frac{X_i(t) \cdot X_j(t)}{R^2}$$



UNIVERSITY OF
OXFORD

Spherical choreographies

Choreographies of the spherical n -body problem

$$X_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{R^3 X_i(t) - R(X_i(t) \cdot X_j(t)) X_j(t)}{[R^4 - (X_i(t) \cdot X_j(t))^2]^{3/2}} + R^{-2} \|X_j'(t)\|^2 X_j(t) = 0,$$

with $X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$ for some 2π -periodic function $Q(t) : [0, 2\pi] \rightarrow \mathbb{S}_R^2 \subset \mathbb{R}^3$

Choreographies of the spherical n -body problem

$$X_j''(t) - \sum_{\substack{i=0 \\ i \neq j}}^{n-1} \frac{R^3 X_i(t) - R(X_i(t) \cdot X_j(t)) X_j(t)}{[R^4 - (X_i(t) \cdot X_j(t))^2]^{3/2}} + R^{-2} \|X_j'(t)\|^2 X_j(t) = 0,$$

with $X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$ for some 2π -periodic function $Q(t) : [0, 2\pi] \rightarrow \mathbb{S}_R^2 \subset \mathbb{R}^3$

Minima of the action

$$A = \int_0^{2\pi} (K(t) - U(t)) dt,$$

$$K(t) = \frac{n}{2} \|Q'(t)\|^2, \quad U(t) = -\frac{n}{R} \sum_{j=1}^{n-1} \frac{Q(t) \cdot Q\left(t + \frac{2\pi j}{n}\right)}{\sqrt{R^4 - (Q(t) \cdot Q\left(t + \frac{2\pi j}{n}\right))^2}}$$



Computing spherical choreographies

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



Computing spherical choreographies

- Our method ² is based on stereographic projection

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



- Our method ² is based on stereographic projection

\mathbb{S}_R^2	\mathbb{C}
$X = (x_1, x_2, x_3)^T$	$z = P_R(X) = \frac{Rx_1 + iRx_2}{R - x_3}$

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



- Our method ² is based on stereographic projection

\mathbb{S}_R^2	\mathbb{C}
$X = (x_1, x_2, x_3)^T$	$z = P_R(X) = \frac{Rx_1 + iRx_2}{R - x_3}$
$Q(t), X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$	$q(t) = P_R(Q(t)), z_j(t) = P_R(X_j(t))$

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



- Our method ² is based on stereographic projection

\mathbb{S}_R^2	\mathbb{C}
$X = (x_1, x_2, x_3)^T$	$z = P_R(X) = \frac{Rx_1 + iRx_2}{R - x_3}$
$Q(t), X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$	$q(t) = P_R(Q(t)), z_j(t) = P_R(X_j(t))$
$d(X, Y) = \ X - Y\ $	$d(z, \xi) = \frac{2R^2 z - \xi }{\sqrt{(R^2 + z ^2)(R^2 + \xi ^2)}}$

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



- Our method ² is based on stereographic projection

\mathbb{S}_R^2	\mathbb{C}
$X = (x_1, x_2, x_3)^T$	$z = P_R(X) = \frac{Rx_1 + iRx_2}{R - x_3}$
$Q(t), X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$	$q(t) = P_R(Q(t)), z_j(t) = P_R(X_j(t))$
$d(X, Y) = \ X - Y\ $	$d(z, \xi) = \frac{2R^2 z - \xi }{\sqrt{(R^2 + z ^2)(R^2 + \xi ^2)}}$
$d_{GC}(X, Y) = R \arccos\left(\frac{X \cdot Y}{R^2}\right)$	$d_{GC}(z, \xi) = 2R \arcsin \frac{d(z, \xi)}{2R}$

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



- Our method ² is based on stereographic projection

\mathbb{S}_R^2	\mathbb{C}
$X = (x_1, x_2, x_3)^T$	$z = P_R(X) = \frac{Rx_1 + iRx_2}{R - x_3}$
$Q(t), X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$	$q(t) = P_R(Q(t)), z_j(t) = P_R(X_j(t))$
$d(X, Y) = \ X - Y\ $	$d(z, \xi) = \frac{2R^2 z - \xi }{\sqrt{(R^2 + z ^2)(R^2 + \xi ^2)}}$
$d_{GC}(X, Y) = R \arccos\left(\frac{X \cdot Y}{R^2}\right)$	$d_{GC}(z, \xi) = 2R \arcsin \frac{d(z, \xi)}{2R}$
$A = \frac{n}{2} \int_0^{2\pi} \left(\frac{2R^2 q'(t) }{R^2 + q(t) ^2} \right)^2 dt + \frac{n}{R} \sum_{j=1}^{n-1} \int_0^{2\pi} \frac{2R^2 - d(q(t), z_j(t))^2}{d(q(t), z_j(t))\sqrt{4R^2 - d(q(t), z_j(t))^2}} dt$	

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.



- Our method ² is based on stereographic projection

\mathbb{S}_R^2	\mathbb{C}
$X = (x_1, x_2, x_3)^T$	$z = P_R(X) = \frac{Rx_1 + iRx_2}{R - x_3}$
$Q(t), X_j(t) = Q\left(t + \frac{2\pi j}{n}\right)$	$q(t) = P_R(Q(t)), z_j(t) = P_R(X_j(t))$
$d(X, Y) = \ X - Y\ $	$d(z, \xi) = \frac{2R^2 z - \xi }{\sqrt{(R^2 + z ^2)(R^2 + \xi ^2)}}$
$d_{GC}(X, Y) = R \arccos\left(\frac{X \cdot Y}{R^2}\right)$	$d_{GC}(z, \xi) = 2R \arcsin \frac{d(z, \xi)}{2R}$
$A = \frac{n}{2} \int_0^{2\pi} \left(\frac{2R^2 q'(t) }{R^2 + q(t) ^2} \right)^2 dt + \frac{n}{R} \sum_{j=1}^{n-1} \int_0^{2\pi} \frac{2R^2 - d(q(t), z_j(t))^2}{d(q(t), z_j(t))\sqrt{4R^2 - d(q(t), z_j(t))^2}} dt$	

- Trigonometric interpolation, closed-form of the gradient, quasi-Newton methods

²Montanelli and Gushterov, "Computing planar and spherical choreographies", 2015.





UNIVERSITY OF
OXFORD

Numerical results (1/2)





Numerical results (1/2)





Numerical results (1/2)



Action	22.036690
Number of coefficients	45
Computer time (s)	0.64
2-norm of the gradient	1.14e-02
Smallest coefficient	8.57e-07
∞ -norm of the residual	1.07e-03






UNIVERSITY OF
OXFORD

Numerical results (2/2)




Numerical results (2/2)











Numerical results (2/2)





Numerical results (2/2)



	$R = 2$	3	5	10	20
Left	3.23e-01	1.05e-01	3.38e-02	8.08e-03	1.99e-03
Middle	3.06e-01	1.04e-01	3.39e-02	8.16e-03	2.02e-03
Right	3.35e-01	1.12e-01	3.64e-02	8.77e-03	1.97e-03



UNIVERSITY OF
OXFORD

Conclusion and future work



Conclusion and future work

- Choreographies are very special periodic solutions of the n -body problem characterized by a single orbit



Conclusion and future work

- Choreographies are very special periodic solutions of the n -body problem characterized by a single orbit
- They also exist on a sphere in a cotangent potential, and as in the plane, they can be computed using trigonometric interpolation and minimization of the action



Conclusion and future work

- Choreographies are very special periodic solutions of the n -body problem characterized by a single orbit
- They also exist on a sphere in a cotangent potential, and as in the plane, they can be computed using trigonometric interpolation and minimization of the action
- Stability properties of spherical choreographies have not been discussed



Conclusion and future work

- Choreographies are very special periodic solutions of the n -body problem characterized by a single orbit
- They also exist on a sphere in a cotangent potential, and as in the plane, they can be computed using trigonometric interpolation and minimization of the action
- Stability properties of spherical choreographies have not been discussed
- Apply these ideas to the n -vortex problem

$$z_j'(t) = \frac{i}{2\pi} \sum_{\substack{k=0 \\ k \neq j}}^{n-1} \Gamma_k \frac{z_j(t) - z_k(t)}{|z_j(t) - z_k(t)|^2}, \quad 0 \leq j \leq n-1$$