

# Agenda

- Concept of Incidence
- Cumulative Incidence
- Incidence Rate
- Relationship between Cumulative Incidence and Incidence Rate
- Relationship between Incidence and Prevalence
- Problems



# Definition of Epidemiology

*“ The study of the distribution and determinants of disease frequency in man”*

MacMahon B, Pugh TF. Epidemiology: principles and methods. Little Brown. 1970



# Implication: Measurement



- ***“study of the distribution ... of disease frequency in human populations”***
  - Measure Outcomes (Disease)
  - Example: **prevalence of CHD, incidence of CHD, risk, odds, rates**
- ***“study of the ... determinants of disease frequency in human populations”***
  - Measure Associations between Risk Factors and Outcomes
  - Example: Risk Ratios, Odds Ratios, Rate Ratios, Regression Coefficients





HARVARD  
School of Public Health

1



ic H

# Why is Prevalence of CHD higher among Non-Smokers?

- Non-Smokers have higher risk of developing CHD (**higher incidence – unlikely**)
- Non-Smokers live longer after developing CHD (**longer duration – unlikely**)
- CHD causes people to stop smoking (**reversed causation - likely**)
- **Bias**
- **Confounding**
- **Chance**



# Conclusion

- Prevalence data are not good for identify risk factors for developing disease
- Alternative: Measure **Incidence** – development of disease in a population followed over time



# Incidence

- Measures **development** of disease in a population over a time
- **Population-at-Risk:** Free of disease at the start of the follow-up period and at risk for developing the disease
- **Cumulative Incidence (Estimated Risk, Average Risk)**
  - $(\# \text{ Cases})/(\# \text{ Subjects at Risk})$
- **Incidence Rate:**
  - $(\# \text{ Cases})/(\text{Person-Time})$



# Incidence

- Measures **development** of disease in a population over a time
- **Population-at-Risk:** Free of disease at the start of the follow-up period and at risk for developing the disease
- **Cumulative Incidence (Estimated Risk, Average Risk)**
  - $(\# \text{ Cases})/(\# \text{ Subjects at Risk})$
- **Incidence Rate:**
  - $(\# \text{ Cases})/(\text{Person-Time})$



# Incidence

- Measures development of disease in a population over a time
- Population-at-Risk: Free of disease at the start of the follow-up period and at risk for developing the disease
- **Cumulative Incidence (Estimated Risk, Average Risk)**
  - $(\# \text{ Cases})/(\# \text{ Subjects at Risk})$
- **Incidence Rate:**
  - $(\# \text{ Cases})/(\text{Person-Time})$



**Example: Incidence of First Headaches (D) among 100 students over 7 days**

ID#	Days of Follow-up						
	1	2	3	4	5	6	7
1.	D						
2.	D						
3.	D						
4.	D						
5.	X	D					
6.	X	X	D				
7.	X	X	D				
8.	X	X	D				
9.	X	X	D				
10.	X	X	X	D			
11.	X	X	X	D			
12.	X	X	X	X	D		
13.	X	X	X	X	X	D	
14.	X	X	X	X	X	D	
15.	X	X	X	X	X	X	D
16.	X	X	X	X	X	X	X
...							
100.	X	X	X	X	X	X	X
Day	1	2	3	4	5	6	7



**Example: Incidence of First Headaches (D) among 100 students over 7 days**

ID#

1.

→

D

2.

→

D

3.

→

D

4.

→

D

5.

→

D

6.

→

D

7.

→

D

8.

→

D

9.

→

D

10.

→

D

11.

→

D

12.

→

D

13.

→

D

14.

→

D

15.

→

D

16.

→

→

100.

→

D

Day

1

2

3

4

5

6

7



# Options for Describing the Incidence of Headaches

- **15 students developed headaches**
  - **Problem:** doesn't account for the population at risk
  - **Problem:** doesn't account for length of follow-up
  - Need a point of reference




# **Option #1 for Describing the Incidence of Headaches**


- 0.15 developed Headaches within 7 days
  - Proportion
- **Cumulative incidence (Estimated Risk)**
  - $(\# \text{new cases of disease}) / (\# \text{ in Pop.-at-Risk})$
- Problem: Value depends on chosen length of follow-up
  - 0.11 developed Headaches within 4 days
  - 0.04 developed Headaches within 1 day



# Data Needed for Calculating Cumulative Incidence of CHD



# Problem: Competing Risks and Losses-to-Follow-up



# Problem

- Don't know if people lost-to-follow-up developed CHD
- Don't know if people who died from other causes (competing risks) would have developed CHD if they had not died.
- Don't know who should be in the numerator of the Cumulative Incidence calculation



Press Esc to exit full screen

## Problem

- Don't know if people lost-to-follow-up developed CHD
- Don't know if people who died from other causes (competing risks) would have developed CHD if they had not died.
- Don't know who should be in the numerator of the Cumulative Incidence calculation



HARVARD  
School of Public Health

Activate Windows  
Go to Settings to activate Windows.  
The Windows logo, consisting of the word "Windows" in a blue, blocky font with a red "W" and a blue "n".

# Solution

- If reason for incomplete follow-up is unrelated to risk of CHD, then survival analysis methods provide valid estimates of the risk for developing CHD during a fixed time period
- Special Case: **Incidence Rates**



## Option # 2 for Describing the Incidence of Headaches

- 15 students developed Headaches from 645 person-days of observation
- **Incidence Rate** =  $15/(645 \text{ person-days})$   
–  $15/645 \text{ pd} = .0233/1 \text{ pd} = 2.33/100 \text{ pd}$
- **Problem:** Summary measure and may not reflect high or low incidence rates on certain days



# Incidence Rate Calculation

- $IR(t) = (\# \text{ cases that develop during the interval around } t) / (\text{amount of person-time observed during the interval})$
- Length of the time interval should reflect constancy of  $IR(t)$



# Incidence Rate Calculation

Incidence Rate at time “t” =  $IR(t)$

= (# of new cases occurring during interval around ‘t’)  
/(amount of person-time observed during the interval)

**Question:** Why divide by person-time?



# Incidence Rate Calculation

Suppose you have 100 people who are “at-risk” during the one-week period around “t” and 5 develop the disease

$$IR(t) \approx$$



# Incidence Rate Calculation

Suppose you have 100 people who are “at-risk” during the one-week period around “t” and 5 develop the disease

$$IR(t) \approx \frac{\text{5 CASES}}{\text{700 P-days}}$$



# Incidence Rate Calculation

Suppose you have 100 people who are “at-risk” during the one-week period around “t” and 5 develop the disease

$$IR(t) \approx 5/(700 \text{ person-days})$$



If incidence rate was constant and 100 people followed for a two-week period then we expect 10 cases to develop disease

$$IR(t) \approx \frac{10}{1400P}$$





HARVARD  
School of Public Health



# Incidence Rate Calculation

- Special case: constant  $IR(t)$  during the entire follow-up period  
$$IR = (\text{total number of cases}) / (\text{total amount of follow-up})$$
- General Case:  $IR = \text{Average } IR(t)$



# Incidence Rate Calculation

- Special case: constant  $IR(t)$  during the entire follow-up period

$$IR = \frac{\text{total number of cases}}{\text{total amount of follow-up}}$$

- General Case:  $IR = \text{Average } IR(t)$

## Mortality Experience for Three Populations Followed for Four Years

	Population		
	A	B	C
Number-at-Risk	100	100	100
# Deaths in first year	10	2	10
# Deaths in second year	10	2	9
# Deaths in third year	10	2	8
# Deaths in fourth year	10	34	7
Total # Deaths	40	40	34
4-yr Cum. Incidence			



## Mortality Experience for Three Populations Followed for Four Years

	Population		
	A	B	C
Number-at-Risk	100	100	100
# Deaths in first year	10	2	10
# Deaths in second year	10	2	9
# Deaths in third year	10	2	8
# Deaths in fourth year	10	34	7
Total # Deaths	40	40	34
4-yr Cum. Incidence	0.40	0.40	0.34

Activate Windows  
Go to Settings to activate Windows



## Incidence Rates: Population A

Year	Population-at-Risk at start of year	# Deaths	Person-Years	IR (deaths/py)
1	100	10	95	.11
2	90	10	85	.12
3	80	10	75	.13
4	70	10	65	.15
Total		40	320	

Activate Windows  
Go to Settings to activate Windows



## Incidence Rates: Population(A)

Year	Population-at-Risk at start of year	# Deaths	Person-Years	IR (deaths/py)
1	100	10	95	$10/95 = .11$
2	90	10	85	$10/85 = .12$
3	80	10	75	.13
4	70	10	65	.15
Total		40	320	.12


## Incidence Rates: Population B

Year	Population-at-Risk at start of year	# Deaths	Person-Years	IR (deaths/py)
1	100	2	99	$2/99 = .02$
2	98	2	97	$2/97 = .02$
3	96	2	95	$2/95 = .02$
4	94	34	77	$34/77 = .44$
Total		40	368	



## Incidence Rates: Population C

Year	Population-at-Risk at start of year	# Deaths	Person-Years	IR (deaths/py)
1	100	10	95	.11
2	90	9	85.5	.11
3	81	8	77	.10
4	73	7	69.5	.10
Total		34	327	10



## Incidence Rates: Population C

Year	Population-at-Risk at start of year	# Deaths	Person-Years	IR (deaths/py)
1	100	10	95	.11
2	90	9	85.5	.11
3	81	8	77	.10
4	73	7	69.5	.10
Total		34	327	10.41



Activate Windows  
Go to settings to activate Windows.



## Comments:

For **Population C**,  $IR(t)$  appears to be constant over time,

$IR = (\text{total no. of new cases}) /$   
 $(\text{total amount of person-time of follow-up})$

total no. of new cases =  $10 + 9 + 8 + 7 = 34$

total amount of follow-up =  $95 + 85.5 + 77 + 69.5$   
= 327

$IR = (34 \text{ cases}) / (327 \text{ person-years})$   
= (.10 case) / (1 person-year)  
= (10 cases) / (100 person years)



If the Incidence Rate function,  $IR(t)$ , is not constant over time then  $IR$  is a **weighted** average of  $IR(t)$  for different time periods.

Example: Average Incidence Rate for **Population B**

$$\text{Average IR} = 40 / 368\text{py}$$

$$\begin{aligned} &= [(99\text{py})(2/99\text{py}) + (97\text{py})(2/97\text{py}) \\ &\quad + (95\text{py})(2/95\text{py}) + (77\text{py})(34/77\text{py})] \\ &/ [99\text{py} + 97\text{py} + 95\text{py} + 77\text{py}] \end{aligned}$$



## Example

Suppose you followed 1000 people for 1 year  
and the **Incidence Rate (IR)** was  $10/(100y)$

How much person-time would you observe?

How many cases of disease would you observe?

What is one-year **Cumulative Incidence**?



## Example

Suppose you followed 1000 people for 1 year and the **Incidence Rate (IR)** was  $10/(100y)$

- How much person-time would you observe?  
 $\approx 1000y$
- How many cases of disease would you observe?  
 $100$

What is one-year **Cumulative Incidence**?



## Example

Suppose you followed 1000 people for 1 year and the **Incidence Rate (IR)** was  $10/(100y)$

- How much person-time would you observe?  
 $\approx 1000y$
- How many cases of disease would you observe?  
 $100$
- What is one-year **Cumulative Incidence**?  
 $100/1000 \approx .10$

# Relationship Between Cumulative Incidence (Risk) and Incidence Rate

- Approximation (for time period  $\Delta t$ ) :  
$$\text{Risk} \approx (\text{IR})(\Delta t)$$
- Example:
  - $\text{IR} = 10/100\text{y}$
  - One-year Risk =  $(10/100\text{y})(1\text{y}) = .10$
- Problem: Formula doesn't take into account correct amount of person-time
- Approximation works well for small IR and short  $\Delta t$



Example (Population C):  $IR = .10/y = 1/10y$

Implications from formula:

→ One-year Risk  $= (1/10y)(1y) = 0.10$   
→ Four-year Risk  $= (1/10y)(4y) = 0.40$   
Ten-year Risk  $= (1/10y)(10y) = 1.0$   
Twelve-year Risk  $= (1/10y)(12y) = 1.2 ???$

**Problem:** Formula does not account for decreasing population-at-risk

**Solution:** Survival Analysis Methods



Example (Population C):  $IR = .10/y = 1/10y$

Implications from formula:

→ One-year Risk	$= (1/10y)(1y)$	$= 0.10$
→ Four-year Risk	$= (1/10y)(4y)$	$= 0.40$
→ Ten-year Risk	$= (1/10y)(10y)$	$= 1.0$
→ Twelve-year Risk	$= (1/10y)(12y)$	$= 1.2$ ???

**Problem:** Formula does not account for decreasing population-at-risk

**Solution:** Survival Analysis Methods



## Incidence Rates: Population C

Year	Population-at-Risk at start of year	# Deaths	Person-Years	IR (deaths/py)
1	100	10	95	.11
2	90	9	85.5	.11
3	81	8	77	.10
4	73	7	69.5	.10
Total		34	327	.10

$$4\text{-Year Risk} = 34/100 = .34 \neq (.10/1y)(4y) = .40$$



## Relationship Between Cumulative Incidence (Risk) and Incidence Rate

- General Formula:  $R = 1 - \exp(-\int IR(x)dx)$
- Special Case:  $R = 1 - \exp(-(IR)(\Delta t))$ 
  - Assumes Constant Incidence Rate
- Example: **Population C**
  - $IR = .10/1y$
  - Four-year Risk =  $1 - \exp(-(.10/1y)(4y)) = 0.33$





# Relationship Between Cumulative Incidence (Risk) and Incidence Rate


- General Formula:  $R = 1 - \exp(-\int IR(x)dx)$
- Special Case:  $R = 1 - \exp(-(IR)(\Delta t))$ 
  - Assumes Constant Incidence Rate
- Example: Population C
  - $IR = .10/1y$
  - Four-year Risk =  $1 - \exp(-(.10/1y)(4y)) = 0.33$


$$1 - \exp(-(.10/1y)(4y)) = 0.33$$



HARVARD  
School of Public Health







# Steady State

Disease:  $N \times (\text{Prevalence})$

↓ Cure Rate

Non-Disease:  $N \times (1 - \text{Prevalence})$

↑

Incidence Rate

# Steady State

Disease :  $N \times (\text{Prevalence})$

↓ Cure Rate

Non-Disease:  $N \times (1-\text{Prevalence})$

↑

Incidence Rate

$[N \times \text{Prevalence}] \times \text{Cure Rate}$

$= [N \times (1-\text{Prevalence})] \times \text{Incidence Rate}$



# Steady State

$$[N \times \text{Prevalence}] \times \text{Cure Rate} = [N \times (1-\text{Prevalence})] \times \text{Incidence Rate}$$


4



# Steady State

$$[N \times \text{Prevalence}] \times \text{Cure Rate} = [N \times (1-\text{Prevalence})] \times \text{Incidence Rate}$$

$$\frac{\text{Prevalence}}{(1-\text{Prevalence})} = \frac{\text{Incidence Rate}}{\text{Cure Rate}}$$

# Steady State

Prevalence/(1-Prevalence)

$$= (\text{Incidence Rate})/(\text{Cure Rate})$$


Cure Rate = (#cures)/(# disease-days)

= 1/(Average Duration of Disease)

$$= 1/D$$

Prevalence/(1-Prevalence) = (Incidence Rate)/(1/D)

$$= (\text{Incidence Rate}) \times D$$



# Steady State

Prevalence/(1-Prevalence)

$$= (\text{Incidence Rate})/(\text{Cure Rate})$$

Cure Rate = (#cures)/(# disease-days)

= 1/(Average Duration of Disease)

$$= 1/D$$

Prevalence/(1-Prevalence) = (Incidence Rate)/(1/D)

$$= (\text{Incidence Rate}) \times D$$



# Steady State

$\text{Prevalence}/(1-\text{Prevalence}) = (\text{Incidence Rate}) \times D$

**Prevalence Odds = (Incidence Rate)  $\times D$**

Special Case: If Prevalence is small


Prevalence Odds  $\approx$  Prevalence

**Prevalence  $\approx$  (Incidence Rate)  $\times D$**





HARVARD  
School of Public Health



## 24-Year Cumulative Incidence of Death for Males and Females

Sex	Died	Survived	Total	Cumulative Incidence
Males	843	1101	1944	
Females	707	1783	2490	
Total	1550	2884	4434	



## Questions

- What is the 24-year Cumulative Incidence of Death among all participants?
- What is the 24-year Cumulative Incidence among Males?
- What is the 24-year Cumulative Incidence among Females?



HARVARD  
School of Public Health



# Questions

- What is the 24-year Cumulative Incidence of Death among **all participants**?
- What is the 24-year Cumulative Incidence among **Males**?
- What is the 24-year Cumulative Incidence among Females?

## 24-Year Cumulative Incidence of Death for Males and Females

Sex	Died	Survived	Total	Cumulative Incidence
Males	843	1101	1944	
Females	707	1783	2490	
Total	1550	2884	4434	1550/4434



## 24-Year Cumulative Incidence of Death for Males and Females

Sex	Died	Survived	Total	Cumulative Incidence
Males	843	1101	1944	8
Females	707	1783	2490	
Total	1550	2884	4434	1550/4434



## 24-Year Cumulative Incidence of Death for Males and Females

Sex	Died	Survived	Total	Cumulative Incidence
Males	843	1101	1944	843/194
Females	707	1783	2490	
Total	1550	2884	4434	1550/4434



### 24-Year Cumulative Incidence of Death for Males and Females

Sex	Died	Survived	Total	Cumulative Incidence
Males	843	1101	1944	843/1944
Females	707	1783	2490	707/2490
Total	1550	2884	4434	1550/4434



HARVARD  
School of Public Health



## 24-Year Cumulative Incidence of Death for Males and Females

Sex	Died	Survived	Total	Cumulative Incidence
Males	843	1101	1944	$843/1944 = 0.43$
Females	707	1783	2490	$707/2490 = 0.28$
Total	1550	2884	4434	$1550/4343 = 0.35$



## Incidence Rate of Death for Males and Females

Sex	Died	Person-Years of Follow-up	Incidence Rate
Males	843	38287.33	
Females	707	52828.26	
Total	1550	91115.59	



# Questions

- What is the Incidence Rate of Death among all participants?
- What is the Incidence Rate of Death among Males?
- What is the Incidence Rate of Death among Females?



4



## Incidence Rate of Death for Males and Females

Sex	Died	Person-Years of Follow-up	Incidence Rate
Males	843	38287.33	
Females	707	52828.26	
Total	1550	91115.59	$1550/91115$



Activate Windows  
Go to Settings to activate Windows

## Incidence Rate of Death for Males and Females

Sex	Died	Person-Years of Follow-up	Incidence Rate
Males	843	38287.33	$843/38287.33\text{py} = 2.20/100\text{py}$
Females	707	52828.26	$707/52828.26\text{py} = 1.34/100\text{py}$
Total	1550	91115.59	$1550/91115.59\text{py} = 1.70/100\text{py}$

# Incidence of CHD for Males and Females

Sex	CHD	Population-at-Risk	Person-Years of Follow-up
Males	710	1944	32763.95
Females	530	2490	48161.20
Total	1240	4434	80925.15



# Question

- How would you measure the incidence of CHD among men?
- How would you measure the incidence of CHD among women?

# Problem

- Cannot measure Cumulative Incidence directly
- Not all participants are “at-risk” for developing (and recording) CHD during 24 years of follow-up
- Reasons
  - Deaths from other causes
  - Losses-to-follow-up
- Solution: Report Incidence Rate




# Questions

- What is the Incidence Rate of CHD among all participants?
- What is the Incidence Rate of CHD among Males?
- What is the Incidence Rate of CHD among Females?

## Incidence Rate of CHD for Males and Females

Sex	CHD	Person-Years of Follow-up	Incidence Rate
Males	710	32763.95	
Females	530	48161.20	
Total	1240	80925.15	$1240 / (80,925.15)$



Activate Windows  
Go to Settings to activate Windows



## Incidence Rate of CHD for Males and Females

Sex	CHD	Person-Years of Follow-up	Incidence Rate
Males	710	32763.95	$710/32763.95 =$ $2.17/100\text{py}$
Females	530	48161.20	$530/48161.20 =$ $1.10/100\text{py}$
Total	1240	80925.15	$1240/80925.15 =$ $1.53/100\text{py}$

