

CE 311K: Linear System of Equations

Krishna Kumar


University of Texas at Austin

krishnak@utexas.edu

December 2, 2019

1 Linear System of Equations


Which one will fill first?






Solving Linear System of Equations

$$\begin{aligned}3x_1 + 2x_2 &= 18 \\ -x_1 + 2x_2 &= 2\end{aligned}$$

Solving Linear System of Equations



Singularity and Ill-conditioned



Solving Linear System of Equations

① Direct Methods

- ① Gauss Elimination
- ② Gauss-Jordan Elimination
- ③ LU decomposition

② Iterative Methods

- ① Jacobi iterative
- ② Gauss-Seidel

Direct methods

Consider a system of 3 linear equations for simplicity:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3$$

Matrix form is:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

Concise form: $Ax = b$

Systems that can be solved easily

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix} \quad \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \quad \begin{bmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{bmatrix}$$

Solve by “back substitution” Upper triangle system (U)

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

Start with the last equation 3: $x_3 = b_3/a_{33}$

Equation 2: $a_{22}x_2 + a_{23}x_3 = b_2$ so

$$x_2 = \frac{b_2 - a_{23}x_3}{a_{22}}$$

Equation 1: $a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1$ so

$$x_1 = \frac{b_1 - (a_{12}x_2 + a_{13}x_3)}{a_{11}}$$

General for ‘n’ systems: $x_n = b_n/a_{nn}$ $x_i = \frac{b_i - \sum_{j=i+1}^n a_{ij}x_j}{a_{ii}}$

Gauss Elimination

Consider a system of 3 linear equations:

$$\begin{bmatrix} 2 & 4 & 6 \\ 4 & 11 & 21 \\ 6 & 21 & 52 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 24 \\ 72 \\ 158 \end{bmatrix}$$

Divide row 2 by -2 and row 3 by -3

$$\begin{bmatrix} 2 & 4 & 6 \\ 0 & 3 & 9 \\ 0 & 9 & 34 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 24 \\ 24 \\ 86 \end{bmatrix}$$

Second reduction:

$$\begin{bmatrix} 2 & 4 & 6 \\ 0 & 3 & 9 \\ 0 & 0 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 24 \\ 24 \\ 14 \end{bmatrix}$$

$$x_3 = 2, x_2 = 2, x_1 = 2$$

CE 311K: Linear Systems

└ Linear System of Equations

└ Gauss Elimination

Consider a system of 3 linear equations:

$$\begin{bmatrix} 2 & 4 & 6 \\ 4 & 11 & 21 \\ 6 & 21 & 52 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 24 \\ 72 \\ 158 \end{bmatrix}$$

Divide row 2 by -2 and row 3 by -3

$$\begin{bmatrix} 2 & 4 & 6 \\ 0 & 3 & 9 \\ 0 & 9 & 34 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 24 \\ 24 \\ 86 \end{bmatrix}$$

Second reduction:

$$\begin{bmatrix} 2 & 4 & 6 \\ 0 & 3 & 9 \\ 0 & 0 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 24 \\ 24 \\ 14 \end{bmatrix}$$

$$x_3 = 2, x_2 = 2, x_1 = 2$$

```
a = np.array([[2,4,6], [4,11,21], [6, 21, 52]])
b = np.array([24, 72, 158])
x = np.linalg.solve(a, b)
```