

Interpolation of DCMs

Bonnie Jonkman

November 13, 2015

1 Logarithmic maps for DCMs

For any direction cosine matrix (DCM), Λ , the logarithmic map for the matrix is a skew-symmetric matrix, λ :

$$\lambda = \log(\Lambda) = \begin{bmatrix} 0 & \lambda_3 & -\lambda_2 \\ -\lambda_3 & 0 & \lambda_1 \\ \lambda_2 & -\lambda_1 & 0 \end{bmatrix} \quad (1)$$

2 Matrix exponentials

The angle of rotation for the skew-symmetric matrix, λ is

$$\theta = \|\lambda\| = \sqrt{\lambda_1^2 + \lambda_2^2 + \lambda_3^2} \quad (2)$$

The matrix exponential is

$$\Lambda = \exp(\lambda) = \begin{cases} I & \theta = 0 \\ I + \frac{\sin \theta}{\theta} \lambda + \frac{1 - \cos \theta}{\theta^2} \lambda^2 & \theta > 0 \end{cases} \quad (3)$$

3 Solving for λ

If the logarithmic map and matrix exponential are truly inverses, we need

$$\exp(\log(\Lambda)) = \Lambda. \quad (4)$$

Using the expression for λ from Equation 1, we get

$$\exp \left(\begin{bmatrix} 0 & \lambda_3 & -\lambda_2 \\ -\lambda_3 & 0 & \lambda_1 \\ \lambda_2 & -\lambda_1 & 0 \end{bmatrix} \right) = \Lambda = \begin{bmatrix} \Lambda_{11} & \Lambda_{12} & \Lambda_{13} \\ \Lambda_{21} & \Lambda_{22} & \Lambda_{23} \\ \Lambda_{31} & \Lambda_{32} & \Lambda_{33} \end{bmatrix} \quad (5)$$

Doing a little algebra for $\theta > 0$, Equation 5 becomes

$$\Lambda = \begin{bmatrix} 1 - \frac{1 - \cos \theta}{\theta^2} (\lambda_3^2 + \lambda_2^2) & \frac{\sin \theta}{\theta} \lambda_3 + \frac{1 - \cos \theta}{\theta^2} \lambda_1 \lambda_2 & -\frac{\sin \theta}{\theta} \lambda_2 + \frac{1 - \cos \theta}{\theta^2} \lambda_1 \lambda_3 \\ -\frac{\sin \theta}{\theta} \lambda_3 + \frac{1 - \cos \theta}{\theta^2} \lambda_1 \lambda_2 & 1 - \frac{1 - \cos \theta}{\theta^2} (\lambda_3^2 + \lambda_1^2) & \frac{\sin \theta}{\theta} \lambda_1 + \frac{1 - \cos \theta}{\theta^2} \lambda_2 \lambda_3 \\ \frac{\sin \theta}{\theta} \lambda_2 + \frac{1 - \cos \theta}{\theta^2} \lambda_1 \lambda_3 & -\frac{\sin \theta}{\theta} \lambda_1 + \frac{1 - \cos \theta}{\theta^2} \lambda_2 \lambda_3 & 1 - \frac{1 - \cos \theta}{\theta^2} (\lambda_2^2 + \lambda_1^2) \end{bmatrix} \quad (6)$$

It follows that the trace is

$$\begin{aligned}\text{Tr}(\Lambda) &= 3 - 2 \frac{1 - \cos \theta}{\theta^2} (\lambda_1^2 + \lambda_2^2 + \lambda_3^2) \\ &= 3 - 2 (1 - \cos \theta) \\ &= 1 + 2 \cos \theta\end{aligned}$$

or

$$\theta = \cos^{-1} \left(\frac{1}{2} (\text{Tr}(\Lambda) - 1) \right) \quad \theta \in [0, \pi] \quad (7)$$

It also follows that

$$\Lambda - \Lambda^T = \frac{2 \sin \theta}{\theta} \begin{bmatrix} 0 & \lambda_3 & -\lambda_2 \\ -\lambda_3 & 0 & \lambda_1 \\ \lambda_2 & -\lambda_1 & 0 \end{bmatrix} \quad (8)$$

or, when $\sin \theta \neq 0$

$$\lambda = \frac{\theta}{2 \sin \theta} (\Lambda - \Lambda^T) \quad (9)$$

We need an equation that works when $\sin \theta$ approaches 0, that is, when θ approaches 0 or θ approaches π . When θ approaches 0, Equation 9 actually behaves well:

$$\lim_{\theta \rightarrow 0} \frac{\theta}{2 \sin \theta} = \frac{1}{2} \quad (10)$$

and since θ is the l_2 norm of the individual components of λ , it follows that they approach zero, and we get

$$\lambda = 0 \quad (11)$$

However, when θ approaches π , $\Lambda - \Lambda^T$ approaches 0, and

$$\lim_{\theta \rightarrow \pi} \frac{\theta}{2 \sin \theta} = \infty \quad (12)$$

We need a different method to find λ . Going back to Equations 5 and 6, we can compute the following:

$$\Lambda_{11} + \Lambda_{22} - \Lambda_{33} = 1 - \frac{2\lambda_3^2(1 - \cos \theta)}{\theta^2} \quad (13)$$

or

$$\lambda_3 = \pm \theta \sqrt{\frac{(1 + \Lambda_{33} - \Lambda_{11} - \Lambda_{22})}{2(1 - \cos \theta)}} \quad (14)$$

Equations for the other two components of λ are similar:

$$\lambda_1 = \pm \theta \sqrt{\frac{(1 + \Lambda_{11} - \Lambda_{22} - \Lambda_{33})}{2(1 - \cos \theta)}} \quad (15)$$

$$\lambda_2 = \pm \theta \sqrt{\frac{(1 + \Lambda_{22} - \Lambda_{11} - \Lambda_{33})}{2(1 - \cos \theta)}} \quad (16)$$

Equations 14-16 give us the magnitude, not the sign of the vector we are looking for. Here is one possibility for choosing the sign: If $(\lambda_1) \neq 0$, choose $\text{sign}(\lambda_1)$ positive.

$$\Lambda_{12} + \Lambda_{21} = \frac{2(1 - \cos \theta)}{\theta^2} \lambda_1 \lambda_2 \quad (17)$$

so

$$\text{sign}(\lambda_2) = \text{sign}(\Lambda_{12} + \Lambda_{21}) \quad (18)$$

and similarly,

$$\text{sign}(\lambda_3) = \text{sign}(\Lambda_{13} + \Lambda_{31}) \quad (19)$$

If $(\lambda_1) = 0$, similar arguments can be used to choose $\text{sign}(\lambda_2)$ positive, and

$$\text{sign}(\lambda_3) = \text{sign}(\Lambda_{23} + \Lambda_{32}) \quad (20)$$

At this point, the relationships between the components of λ are set, so we have computed $\pm\lambda$. If $\theta = \pi$, both values are a solution, so this good enough.

If θ is close to π , we will need to determine if we have the negative of the solution. This is required for numerical stability of the solution. In this case, $\Lambda - \Lambda^T$ is not exactly zero, so we can look at the sign of the solution we would have computed if we had used Equation 9:

$$\Lambda_{23} - \Lambda_{32} = \left| \frac{\sin \theta}{\theta} \right| \lambda_1 \quad (21)$$

$$\Lambda_{31} - \Lambda_{13} = \left| \frac{\sin \theta}{\theta} \right| \lambda_2 \quad (22)$$

$$\Lambda_{12} - \Lambda_{21} = \left| \frac{\sin \theta}{\theta} \right| \lambda_3 \quad (23)$$

For numerical reasons, we don't want to use these equations to get the magnitude of the components of λ , but we can look at the sign of the element with largest magnitude and ensure our λ has the same sign.

4 Interpolation

4.1 Periodicity of solutions

Given $\lambda_k = \lambda \left(1 + \frac{2k\pi}{\|\lambda\|} \right)$ for any integer k , it follows that

$$\theta_k = \left| 1 + \frac{2k\pi}{\|\lambda\|} \right| \theta \quad (24)$$

or

$$\theta_k = |\theta + 2k\pi| \quad (25)$$

$$\begin{aligned}
\Lambda_k &= \exp(\lambda_k) \\
&= I + \frac{\sin \theta_k}{\theta_k} \lambda_k + \frac{1 - \cos \theta_k}{\theta_k^2} \lambda_k^2 \\
&= I + \frac{\sin |\theta + 2k\pi|}{|\theta + 2k\pi|} \left(\frac{\theta + 2k\pi}{\theta} \right) \lambda + \frac{1 - \cos |\theta + 2k\pi|}{|\theta + 2k\pi|^2} \left(\frac{\theta + 2k\pi}{\theta} \right)^2 \lambda^2 \\
&= I + \frac{\sin |\theta + 2k\pi|}{\theta} \frac{\theta + 2k\pi}{|\theta + 2k\pi|} \lambda + \frac{1 - \cos |\theta + 2k\pi|}{\theta^2} \lambda^2 \\
&= I + \frac{\sin \theta}{\theta} \lambda + \frac{1 - \cos \theta}{\theta^2} \lambda^2 \\
&= \exp(\lambda) \\
&= \Lambda
\end{aligned}$$

Thus, if λ is one solution to $\log(\Lambda)$, then so is $\lambda_k = \lambda \left(1 + \frac{2k\pi}{\|\lambda\|} \right)$ for any integer k .

4.2 Finding values of λ for interpolation

Given a set of λ^j to be interpolated, find equivalent $\tilde{\lambda}^j$ for integers $j = 1, 2, \dots, n$: Set $\tilde{\lambda}^1 = \lambda^1$. For each $j \in [2, n]$, check to see if $\tilde{\lambda}^{j-1}$ is closer (in the l_2 -norm sense) to λ^j or $\lambda^j \left(1 + \frac{2\pi}{\|\lambda^j\|} \right)$. If the latter, set $\tilde{\lambda}^j = \lambda^j \left(1 + \frac{2\pi}{\|\lambda^j\|} \right)$ and continue checking if we need to add more 2π periods. Otherwise, check to see if $\tilde{\lambda}^{j-1}$ is closer to λ^j or $\lambda^j \left(1 - \frac{2\pi}{\|\lambda^j\|} \right)$. If the latter, set $\tilde{\lambda}^j = \lambda^j \left(1 - \frac{2\pi}{\|\lambda^j\|} \right)$ and continue checking if we need to subtract more 2π periods. Otherwise set $\tilde{\lambda}^j = \lambda^j$.

Interpolation must occur on the $\tilde{\lambda}^j$ and not the λ^j .