

Your Paper

You

March 4, 2025

Denote $G_5 := G_5(\mathbb{Z}_p)$, and $G_5^+ := G_5^+(\mathbb{Q}_p)$.

$\zeta_{L_{5,p}}^\wedge(s) = \int_{G_5^+} |\det g|_p^s d\mu(G_5) = \int_{G_5^+} |\det uh|_p^s d\mu(G_5)$, where $h \in H$ and $u \in N_h$.

Each u is unipotent, hence $\zeta_{L_{5,p}}^\wedge(s) = \int_{G_5^+} |\det h|_p^s d\mu(G_5) = \int_{G_5^+} |\lambda_1^4 \lambda_2^6 \lambda_3^6 \lambda_4^4|_p^s d\mu(G_5) =$
 $= \int_{G_5^+} \left[|\lambda_1^4|_p |\lambda_2^6|_p |\lambda_3^6|_p |\lambda_4^4|_p \right]^s d\mu(G_5)$, by the inductive formula we have found
for every $|h|$.

We denote $v_i := v_p(\lambda_i)$,

and so $\zeta_{L_{5,p}}^\wedge(s) = \int_{G_5^+} \left[p^{-4v_1} p^{-6v_2} p^{-6v_3} p^{-4v_4} \right]^s d\mu(G_5) = \int_{G_5^+} p^{-(4v_1+6v_2+6v_3+4v_4)s} d\mu(G_5)$.

We denote $I(\underline{\lambda}) := p^{-(4v_1+6v_2+6v_3+4v_4)s}$. Now we use the natural matrix decomposition of the N_h matrix of Berman's, which means that

$\zeta_{L_{5,p}}^\wedge(s) = \int_{G_5^+} I(\underline{\lambda}) d\mu(G_5) = \int_{\underline{\lambda}} \int_{\underline{a}} \int_{\underline{b}} \int_{\underline{c}} I(\underline{\lambda}) d\mu(\underline{c}) d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda})$. Since $I(\underline{\lambda})$ depends only on $\lambda_1, \lambda_2, \lambda_3, \lambda_4$, which appear only in the computation of the outermost integral, we consider them as constants for all the inner integrals, which means that we have $\zeta_{L_{5,p}}^\wedge(s) = \int_{\underline{\lambda}} I(\underline{\lambda}) \int_{\underline{a}} \int_{\underline{b}} \int_{\underline{c}} 1 d\mu(\underline{c}) d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda})$,

hence all the inner integrals evaluate to the measure of their domains of integration. now we compute the innermost integral by considering \underline{a} , \underline{b} and $\underline{\lambda}$ as constants, and integrating only over \underline{c} . Considering the multiplication uh , we observe that for each element c_j , we must have that $\rho_j = c_j \lambda_1 \lambda_2 \lambda_3 \lambda_4 \in \mathbb{Z}_p$, which means that $v(\rho_i) = v(c_i \lambda_1 \lambda_2 \lambda_3 \lambda_4) \geq 0 \Rightarrow v(c_i) + v_1 + v_2 + v_3 + v_4 \geq 0 \Rightarrow v(c_i) \geq -(v_1 + v_2 + v_3 + v_4)$. But this means that $c_i \in p^{-(v_1+v_2+v_3+v_4)} \mathbb{Z}_p$, and since the domain of integration for this integral is $\underline{c} = \{c_1, c_2, c_3, c_4\}$, then $\mu(\underline{c}) = |c_j|_p^4 = p^{4(v_1+v_2+v_3+v_4)}$. Denote $I(\underline{\lambda}, \underline{c}) := I(\underline{\lambda}) p^{4(v_1+v_2+v_3+v_4)}$, we now

have that $\zeta_{L_{5,p}}^\wedge(s) = \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}) \int_{\underline{a}} \int_{\underline{b}} 1 d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda})$.

Denote $\lambda_{13} := \lambda_1 \lambda_2 \lambda_3$, $\lambda_{24} := \lambda_2 \lambda_3 \lambda_4$, and $\lambda_{14} := \lambda_1 \lambda_2 \lambda_3 \lambda_4$. We now consider the constraints on \underline{b} .

$b_{11} \lambda_{13}, b_{31} \lambda_{13}, b_{41} \lambda_{13} \in \mathbb{Z}_p$, and $b_{12} \lambda_{24}, b_{22} \lambda_{24} \in \mathbb{Z}_p$. These constraints are obtained by multiplying elements in block M_{13} with elements in h , but one

observes that we have b_{22} also in location $(5, 10)$ of the matrix, and b_{31} in location $(7, 10)$, which means that $b_{22}\lambda_{14}, b_{31}\lambda_{14} \in \mathbb{Z}_p$. But since we already have $b_{22}\lambda_{24}, b_{31}\lambda_{13} \in \mathbb{Z}_p$, the constraints $b_{22}\lambda_{14}$ and $b_{31}\lambda_{14}$ do not contribute any new information. In addition, we have one of the elements of \underline{b} that forms a constraint together with elements from \underline{a} , namely $(a_{11}a_{22} - b_{11})\lambda_{24} \in \mathbb{Z}_p$. The constraints $b_{31}\lambda_{13}, b_{41}\lambda_{13}, b_{12}\lambda_{24}, b_{22}\lambda_{24} \in \mathbb{Z}_p$ from above translate to $p^{-2(v_1+v_2+v_3)}p^{-2(v_2+v_3+v_4)} = p^{-2(v_1+2v_2+2v_3+v_4)}$. On the other hand, b_{11} is a part of two constraints, hence we must have both $b_{11} \in p^{-(v_1+v_2+v_3)}\mathbb{Z}_p$ and $a_{11}a_{22} - b_{11} \in p^{-(v_2+v_3+v_4)}\mathbb{Z}_p \Rightarrow b_{11} \in a_{11}a_{22} + p^{-(v_2+v_3+v_4)}\mathbb{Z}_p$, which means that we need to compute the measure $\mu(A)$, where $A = p^{-(v_1+v_2+v_3)}\mathbb{Z}_p \cap a_{11}a_{22} + p^{-(v_2+v_3+v_4)}\mathbb{Z}_p$. Denote $\alpha := v_1 + v_2 + v_3$, $\beta := v_2 + v_3 + v_4$ and $x := a_{11}a_{22}$, and we need to find a formula for a generic intersection of the form $A = p^{-\alpha}\mathbb{Z}_p \cap x + p^{-\beta}\mathbb{Z}_p$. We need to find a formula for this generic form. Since b_{11} is in the intersection, we have that $b_{11} = z = x + y$ where $y \in p^{-\beta}$ and $z \in p^{-\alpha}\mathbb{Z}_p \Rightarrow z - x \in p^{-\beta}\mathbb{Z}_p$. Assume $\beta \geq \alpha \Rightarrow -\beta \leq -\alpha$, and since $v_p(b_{11}) = v_p(z - x) \geq \min\{v_p(z), v_p(x)\}$, and $v_p(z) \geq -\alpha \geq -\beta$, then we have two cases. If $v_p(x) \geq -\beta$, then $v_p(z - x) \geq \beta \Rightarrow z - x \in p^{-\beta}\mathbb{Z}_p$. But $-\alpha \geq -\beta \Rightarrow p^{-\alpha}\mathbb{Z}_p \subseteq p^{-\beta}\mathbb{Z}_p \Rightarrow A = p^{-\alpha}\mathbb{Z}_p$. If $v_p(x) < -\beta$, then $v_p(z - x) = v_p(x) < -\beta \Rightarrow z - x \notin p^{-\beta}\mathbb{Z}_p$, which means that $A = \emptyset$. One checks that if we assume $\alpha \geq \beta$, then we obtain that $A = p^{-\beta}\mathbb{Z}_p$ if $v_p(x) \geq -\alpha$, and $A = \emptyset$ if $v_p(x) < -\alpha$. Therefore, $\mu(A) = p^{\min\{\alpha, \beta\}}$ for every x such that $v_p(x) \geq \min\{-\alpha, -\beta\} = -\max\{\alpha, \beta\}$, which means, in our case, that $v_p(x) = v_p(a_{11}a_{22}) \geq -\max\{v_1 + v_2 + v_3, v_2 + v_3 + v_4\} = -v_2 - v_3 - \max\{v_1, v_4\}$. Thus, denoting $I(\underline{\lambda}, \underline{c}, \underline{b}) := I(\underline{\lambda}, \underline{c})p^{-(v_2+v_3)-\max\{v_1, v_4\}}$, we have that $\zeta_{L_{5,p}}^\wedge(s) =$

$\int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}) \int_{\underline{a}} 1 d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda})$. Denote $v_{ij} := v_p(a_{ij})$. For the constraints on \underline{a} , we have

$$\begin{aligned} a_{11}\lambda_1\lambda_2, -a_{11}\lambda_2\lambda_3, -a_{11}\lambda_2\lambda_3\lambda_4 \in \mathbb{Z}_p &\Rightarrow v_{11} \geq -(v_1 + v_2), v_{11} \geq -(v_2 + v_3) \Rightarrow \\ v_{11} &\geq -v_2 - \min\{v_1, v_3\}. \\ a_{21}\lambda_1\lambda_2, a_{21}\lambda_1\lambda_2\lambda_3, a_{21}\lambda_1\lambda_2\lambda_3\lambda_4 \in \mathbb{Z}_p &\Rightarrow v_{21} \geq -(v_1 + v_2). \\ a_{22}\lambda_2\lambda_3, -a_{22}\lambda_3\lambda_4, a_{22}\lambda_1\lambda_2\lambda_3 \in \mathbb{Z}_p &\Rightarrow \\ \Rightarrow v_{22} &\geq -(v_2 + v_3), v_{22} \geq -(v_3 + v_4), v_{22} \geq -(v_1 + v_2 + v_3) \Rightarrow v_{22} \geq -v_3 - \min\{v_2, v_4\}. \\ a_{33}\lambda_3\lambda_4, a_{33}\lambda_2\lambda_3\lambda_4, a_{33}\lambda_1\lambda_2\lambda_3\lambda_4 \in \mathbb{Z}_p &\Rightarrow v_{33} \geq -(v_3 + v_4). \\ a_{21}a_{22}\lambda_1\lambda_2\lambda_3 \in \mathbb{Z}_p &\Rightarrow v_{21} + v_{22} \geq -(v_1 + v_2 + v_3). \\ -a_{11}a_{33}\lambda_2\lambda_3\lambda_4 \in \mathbb{Z}_p &\Rightarrow v_{11} + v_{33} \geq -(v_2 + v_3 + v_4). \\ a_{21}a_{33}\lambda_1\lambda_2\lambda_3\lambda_4 \in \mathbb{Z}_p &\Rightarrow v_{21} + v_{33} \geq -(v_1 + v_2 + v_3 + v_4). \end{aligned}$$

And we also have the constraint found earlier,

$$v_{11} + v_{22} \geq -(v_2 + v_3 + \max\{v_1, v_4\}).$$

We have three constraints on a_{21}

1. $v_{21} \geq -(v_1 + v_2)$
2. $v_{21} \geq -(v_1 + v_2 + v_3 + v_{22})$
3. $v_{21} \geq -(v_1 + v_2 + v_3 + v_4 + v_{33})$

But the third constraint does not add new information, because we already have the two separate constraints $v_{21}, v_{33} \geq -(v_1 + v_2 + v_3 + v_4)$.

The two valid constraints translate to

$$v_{21} \geq \min\{-(v_1 + v_2), -(v_1 + v_2 + v_3 + v_{22})\} = -(v_1 + v_2) - \{0, v_3 + v_{22}\}.$$

In the same way, we obtain the constraint $v_{33} \geq -(v_3 + v_4) - \min\{0, v_2 + v_{11}\}$.

Thus, we decompose the inner integral for \underline{a} into separate integrals, to obtain

$$\begin{aligned} \zeta_{L_{5,p}}^\wedge(s) &= \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}) \int_{\underline{a}} 1 d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda}) = \\ &= \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}) \int_{a_{11}} \int_{a_{22}} \int_{a_{33}} \int_{a_{21}} 1 d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda}). \end{aligned}$$

Hence, we have the measures $\mu(a_{21}) = p^{v_1 + v_2 + \min\{0, v_3 + v_{22}\}}$ and $\mu(a_{33}) = p^{v_3 + v_4 + \min\{0, v_2 + v_{11}\}}$. Denote $I(\underline{\lambda}, \underline{c}, \underline{b}, a_{21}, a_{33}) := I(\underline{\lambda}, \underline{c}, \underline{b}) p^{v_1 + v_2} p^{v_3 + v_4}$. We

$$\begin{aligned} \text{have } \zeta_{L_{5,p}}^\wedge(s) &= \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}) \int_{\underline{a}} 1 d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda}) = \\ &= \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}) \int_{a_{11}} \int_{a_{22}} \int_{a_{33}} \int_{a_{21}} 1 d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda}) = \\ &= \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}, a_{21}, a_{33}) \int_{a_{11}} p^{\min\{0, v_2 + v_{11}\}} \int_{a_{22}} p^{\min\{0, v_3 + v_{22}\}} d\mu(\underline{b}) d\mu(\underline{a}) d\mu(\underline{\lambda}). \end{aligned}$$

By the constraints we found earlier on a_{22} , we have the following.

1. $v_{22} \geq -v_3 - \min\{v_2, v_4\}$
2. $v_{22} \geq -(v_2 + v_3) - \max\{v_1, v_4\} - v_{11}$

which translates into $v_{22} \geq -v_3 - \min\{\min\{v_2, v_4\}, v_2 + \max\{v_1, v_4\} + v_{11}\} = -v_3 - \min\{v_2, v_4, v_2 + \max\{v_1, v_4\} + v_{11}\}$.

Denote $\alpha := v_2 + \max\{v_1, v_4\} + v_{11}$ and $\beta := \min\{v_2, v_4, \beta\}$. We already have the constraint $v_{11} \geq -(v_2 + \min\{v_1, v_3\})$, which means that, in either case, $v_{11} \geq -(v_1 + v_2) \geq -\min\{v_1, v_4\} - v_2$

$$\Rightarrow \alpha = v_2 + \max\{v_1, v_4\} + v_{11} \geq \max\{v_1, v_4\} - \min\{v_1, v_4\} \geq 0$$

$$\Rightarrow \beta = \min\{v_2, v_4, \alpha\} \geq 0 \Rightarrow v_3 + \alpha > 0 \Rightarrow v_{22} \geq -(v_3 + \beta).$$

For the inner integral $\int_{a_{22}} p^{\min\{0, v_3 + v_{22}\}} d\mu(a_{22})$, we have two cases. If $v_3 +$

$$v_{22} \geq 0, \text{ then } \min\{0, v_3 + v_{22}\} = 0 \Rightarrow \int_{a_{22}} p^{\min\{0, v_3 + v_{22}\}} d\mu(a_{22}) = \int_{v_{22} \geq -v_3} 1 d\mu(a_{22}) = p^{v_3}.$$

$$\text{If } v_3 + v_{22} < 0, \text{ then } \int_{a_{22}} p^{\min\{0, v_3 + v_{22}\}} d\mu(a_{22}) = \int_{v_{22} < -v_3} p^{v_3 + v_{22}} d\mu(a_{22}).$$

But we said earlier that $v_{22} \geq -(v_3 + \beta)$, hence $-v_3 - \beta \leq v_{22} \leq -v_3 - 1 \Rightarrow -\beta \leq v_3 + v_{22} \leq -1$, which means that we can compute the integral over a_{22} as

$$\text{a sum of } \beta \text{ integrals, } \int_{-v_3 - \beta \leq v_{22} \leq -v_3 - 1} p^{v_3 + v_{22}} d\mu(a_{22}) = \sum_{\tau=1}^{\beta} \int_{v_{22} = -v_3 - \tau} p^{-\tau} d\mu(a_{22}).$$

To evaluate each integral in the sum, we need to calculate the measure of its domain, namely $\mu(\{v_{22} = -(v_3 + \tau)\}) = \mu(\{v_{22} \geq -(v_3 + \tau + 1)\} \setminus \{v_{22} \geq -(v_3 + \tau)\}) = \mu(p^{-(v_3 + \tau + 1)} \mathbb{Z}_p \setminus p^{-(v_3 + \tau)} \mathbb{Z}_p) = p^{v_3 + \tau + 1} - p^{v_3 + \tau} = p^{v_3 + \tau} (p - 1)$, which means that each integral evaluates as $p^{v_3 + \tau} (p - 1) p^{-\tau} = p^{v_3} (p - 1)$, and the sum is over

β such integrals, so we have that $\int_{a_{22}} p^{\min\{0, v_3+v_{22}\}} d\mu(a_{22}) = p^{v_3} + \beta p^{v_3}(p-1)$,

where β depends also on v_{11} .

Hence, we need to compute the integral

$$\begin{aligned} & \int_{a_{11}} p^{\min\{0, v_2+v_{11}\}} p^{v_3} [1+\beta(p-1)] d\mu(a_{11}). \text{ We denote } I(\underline{\lambda}, \underline{c}, \underline{b}, a_{21}, a_{33}, a_{22}) := \\ & I(\underline{\lambda}, \underline{c}, \underline{b}, a_{21}, a_{33}, a_{22}) p^{v_3}, \text{ so } \zeta_{L^5, p}^\wedge(s) = \\ & = \int_{\underline{\lambda}} I(\underline{\lambda}, \underline{c}, \underline{b}, a_{21}, a_{33}, a_{22}) \int_{a_{11}} p^{\min\{0, v_2+v_{11}\}} [1+\beta(p-1)] d\mu(a_{11}). \text{ But as we} \\ & \text{saw earlier, } p^{\min\{0, v_2+v_{11}\}}. \end{aligned}$$