UVa Online Judge Page 1 of 2



Home < *URL*: http://uva.onlinejudge.org/index.php> ▶ Browse Problems

Root < *URL*: *index.php*?

option=com onlinejudge&Itemid=8&category=0>



option=com_onlinejudge&Itemid=8&page=submit_problem&prob

option=com_onlinejudge&Itemid=8&page=problem stats&prob

834 - Continued Fractions

Time limit: 3.000 seconds

Continued Fractions

Let b_0 , b_1 , b_2 ,..., b_n be integers with $b_k > 0$ for k > 0. The *continued fraction* of order n with coefficients b_1 , b_2 ,..., b_n and the initial term b_0 is defined by the following expression

$$b_0 + rac{1}{b_1 + rac{1}{b_{2+...+rac{1}{b_{m}}}}}$$

which can be abbreviated as $[\boldsymbol{b}_0; \boldsymbol{b}_1, ..., \boldsymbol{b}_n]$.

An example of a continued fraction of order n = 3 is [2;3, 1, 4]. This is equivalent to

$$2 + \frac{1}{3 + \frac{1}{1 + \frac{1}{4}}} = \frac{43}{19}$$

Write a program that determines the expansion of a given rational number as a continued fraction. To ensure uniqueness, make $b_n > 1$.

Input

UVa Online Judge Page 2 of 2