## MVE136 Random Signals Analysis

## Written home exam Monday 17 August 2020 2 PM-6 PM

TEACHER AND EXAMINER: Patrik Albin 0317723512 palbin@chalmers.se.

AIDS: All aids are permitted. (See the Canvas course "MVE136 Re-Exam MVE136" with instructions for this reexam for clarifications.)

GRADES: 12 (40%), 18 (60%) and 24 (80%) points for grade 3, 4 and 5, respectively. MOTIVATIONS: All answers/solutions must be motivated. Good Luck!

Task 1. Calculate  $\Pr(X(1)Y(2) > 0)$  when X(t) and Y(t),  $t \in \mathbb{R}$ , are independent zeromean WSS Gaussian processes with PSD's  $S_{XX}(f) = S_{YY}(f) = e^{-|f|}$ . (5 points)

**Task 2.** Let X(t) and Y(t),  $t \ge 0$ , be independent Poission processes with intensity (/rate) 1. Find Pr(X(1) = Y(2)). (5 points)

**Task 3.** Let X(n),  $n = 0, 1, 2, \ldots$ , be a Markov chain with possible values 0 and 1, transition matrix  $P = \binom{1/2}{1/2} \binom{1/2}{1/2}$  and starting distribution  $\pi(0) = (1 \ 0)$ . Find the expected value E(T) of the random time

$$T = \min\{n > 0 : X(m) = 1 \text{ for some } 0 < m < n \text{ and } X(n) = 0\}$$

it takes for the chain to make a journey from 0 to 1 and back to 0. [HINT: A geometric discrete random variable  $\xi$  with PMF  $\Pr(\xi = k) = p(1-p)^{k-1}$  for k = 1, 2, ... has expected value  $E(\xi) = 1/p$ .] (5 points)

**Task 4.** Show by an example that two WSS random processes X(t) and Y(t) need not be jointly WSS, i.e., that  $R_{XY}(t, t+\tau)$  may depend on t. (5 points)

Task 5. A WSS process X(t) with PSD  $S_{XX}(f)$  is sent on a noisy channel where an independent zero-mean WSS noise process N(t) with PSD  $S_{NN}(f)$  is added. The received signal Y(t) = X(t) + N(t) is processed through a Wiener filter LTI system with transfer function  $H(f) = S_{XX}(f)/(S_{XX}(f) + S_{NN}(f))$  and outsignal Z(t) that minimizes the mean-square distance  $D = E((X(t) - Z(t))^2)$  between the sent signal and the processed received signal. It is possible to have D = 0: What exactly is required for this to happen? (5 points)

**Task 6.** How can one do parametric spectral estimation for an AR(1) process?

(5 points)