

MAT367 pset2 Problem 2

Problem 2: (Tangent vectors to a sphere) The unit sphere S^n in \mathbb{R}^{n+1} is defined by the equation:

$$\sum_{i=1}^{n+1} (x^i)^2 = 1.$$

For $p = (p^1, \dots, p^{n+1}) \in S^n$, show that a necessary and sufficient condition for

$$X_p = \sum a^i \frac{\partial}{\partial x^i} \Big|_p \in T_p \mathbb{R}^{n+1}$$

to be tangent to S^n at p is:

$$\sum a^i p^i = 0.$$

Solution:

Consider $X_p = \sum a^i \frac{\partial}{\partial x^i} \Big|_p \in T_p(\mathbb{R}^{n+1})$, where $T_p(\mathbb{R}^{n+1})$ is an $(n+1)$ -dimensional vector space centered at $p = (p^1, \dots, p^{n+1})$.

Since $x^i(X_p) = \sum a^i \frac{\partial x^i}{\partial x^i} = a^i$ for a coordinate function x^i , we see that $X_p = (a^1, a^2, \dots, a^{n+1})$.

Now, suppose $(a^1, \dots, a^{n+1}) \in X_p$ is tangent to S^n at p . Then, by Proposition 8.16 in the textbook by Loring Tu, there exists a curve $c : (-\epsilon, \epsilon) \rightarrow S^n$ such that

$$c(0) = p, \quad c'(0) = X_p.$$

Write $c(t) = (x^1(t), x^2(t), \dots, x^{n+1}(t))$. Since $c(t) \in S^n$ for all t , we have:

$$\sum_{i=1}^{n+1} (x^i(t))^2 = 1.$$

Differentiating both sides with respect to t and evaluating at $t = 0$ gives:

$$\sum_{i=1}^{n+1} 2x^i(t)x^{i'}(t) \Big|_{t=0} = 0.$$

Since $x^i(0) = p^i$ and $x^{i'}(0) = a^i$, this simplifies to:

$$2 \sum_{i=1}^{n+1} p^i a^i = 0 \quad \Rightarrow \quad \sum_{i=1}^{n+1} p^i a^i = 0.$$

Now, consider the subspace of \mathbb{R}^{n+1} :

$$N = \{(a^1, \dots, a^{n+1}) \in \mathbb{R}^{n+1} \mid \sum a^i p^i = 0\}.$$

To confirm that N is a subspace, we check the conditions:

- The zero vector $\mathbf{0} = (0, 0, \dots, 0)$ is in N since $\sum 0 \cdot p^i = 0$.
- Let $a = (a^1, \dots, a^{n+1}) \in N$ and $b = (b^1, \dots, b^{n+1}) \in N$. Then,

$$\sum (a^i + cb^i)p^i = \sum a^i p^i + c \sum b^i p^i = 0 + c \cdot 0 = 0.$$

This shows that $ca + b \in N$ for any scalar $c \in \mathbb{R}$.

Thus, N is a subspace of \mathbb{R}^{n+1} .

From the previous derivation, we have $T_p(S^n) \subseteq N$.

We also observe that $p \notin N$ since $\sum (p^i)^2 = 1 \neq 0$. Since S^n is an n -dimensional manifold, $T_p(S^n)$ is an n -dimensional subspace of \mathbb{R}^{n+1} . Moreover, since $T_p(S^n) \subseteq N$ and both are subspaces of \mathbb{R}^{n+1} , their dimensions satisfy:

$$\dim(T_p(S^n)) = n \leq \dim(N) \leq n + 1 = \dim(\mathbb{R}^{n+1}).$$

Since $p \notin N$ and $\dim(\mathbb{R}^{n+1}) = n + 1$, we obtain $\dim(N) = n$. Since $T_p(S^n)$ and N have the same dimension, we conclude:

$$T_p(S^n) = N.$$

Thus, any X_p tangent to S^n at p is in N , and any $(a^1, \dots, a^{n+1}) \in N$ is also tangent to S^n at p , as required.