

# Assessing lead sources in fishes of the northeast Pacific Ocean



Miling Li<sup>a,b,\*</sup>, Dominique Weis<sup>a,b</sup>, Kate E. Smith<sup>a,b</sup>, Alyssa E. Shiel<sup>a,c</sup>, Wade D. Smith<sup>d,e</sup>, Brian P.V. Hunt<sup>a,d,e</sup>, Abe Torchinsky<sup>a</sup>, Evgeny A. Pakhomov<sup>a,d,e</sup>

<sup>a</sup> Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia, Canada

<sup>b</sup> Pacific Centre for Isotopic and Geochemical Research, University of British Columbia, Canada

<sup>c</sup> College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, USA

<sup>d</sup> Institute for the Oceans and Fisheries, University of British Columbia, Canada

<sup>e</sup> Hakai Institute, Heriot Bay, BC, Canada

## ARTICLE INFO

### Article history:

Received 2 August 2019

Received in revised form 14 December 2019

Accepted 17 December 2019

Available online 19 December 2019

### Keywords:

Lead isotopes

Heavy metal

Fish

Marine ecosystem

Bioindicators

## ABSTRACT

Industrial lead (Pb) emissions have changed oceanic Pb concentrations and isotopic compositions significantly over the last century. Asian industrial emissions are currently the dominant Pb sources in the northwest and central Pacific Ocean. This study investigated major Pb sources in the northeast Pacific Ocean (inland, coastal, and open ocean), where no comprehensive data exist currently. We measured Pb concentrations and isotopic compositions of a variety of shellfish and fish collected in British Columbia (BC) and used them for Pb source apportionment applications in the marine environment. We identified a clear trend in Pb isotopic compositions, from higher  $^{206}\text{Pb}/^{207}\text{Pb}$  in inland fish to lower  $^{206}\text{Pb}/^{207}\text{Pb}$  in open ocean fish, with coastal fish possessing mid-range values. These results indicate that natural sources of Pb dominate freshwater ecosystems in the central BC coast, whereas anthropogenic Pb sources prevail in western North America and the northeast Pacific. Pacific herring that forage along BC coast exhibit Pb isotopic composition that deviates toward the Chinese Pb regression line. We estimate an Asian origin for about one third of their total Pb. This study shows that Pb isotopes in aquatic organisms are valuable tools for assessing Pb sources across coastal and offshore regions of the northeast Pacific, with great potential for identifying primary foraging areas of marine organisms.

© 2019 Elsevier Ltd. All rights reserved.

## 1. Introduction

Changes in the types and magnitudes of global industrial lead (Pb) emissions have altered Pb concentrations and isotopic compositions in the environment over the last century. Emissions from leaded gasoline used to be the dominant source of the total global Pb in the environment but have drastically decreased due to its nearly complete phase-out over the past four decades (Eisenreich et al., 1986; Wu and Boyle, 1997; Boyle et al., 2014). In parallel, rapid industrialization in Asia, accompanied by intense coal combustion and metal smelting, has released a large amount of Pb into the atmosphere (Li et al., 2012; Flegal et al., 2013; Zurbick et al., 2017). Diminishing impacts from leaded gasoline combined with growing Pb emissions of Asian industrial activities have been recorded in a variety of environmental samples, including seawater (Wu and Boyle, 1997; Boyle et al., 2014), ice and snow (Gross et al., 2012; Bory

et al., 2014), corals (Inoue and Tanimizu, 2008), and sediments (Chen et al., 2016; Hosono et al., 2016).

A number of studies have observed marked Asian Pb contributions in various regions of the Northern Hemisphere (Osterberg et al., 2008; Díaz-Somoano et al., 2009; Ewing et al., 2010; Gallon et al., 2011; Gross et al., 2012; Li et al., 2012; Zurbick et al., 2017; Asher et al., 2018). Bory et al. (2014) reported growing impacts of anthropogenic Pb from Chinese aerosols and dust deposited on Greenland snow since the 1990s. Intergovernmental Oceanographic Commission cruise data and global Pb emission estimates indicate that present-day anthropogenic Pb in surface and near-surface waters of the western and central Pacific Ocean is likely dominated by emissions from East Asia; mostly from China, Russia, and Japan (Gallon et al., 2011; Zurbick et al., 2017). The Pb concentrations and isotopic compositions of ice cores and airborne particulate samples collected in western North America suggest that Chinese industrial aerosols have undergone long-range transport via mid-latitude westerlies and contributed 29–46 % of total atmospheric Pb deposition during the peak seasons of Asian impacts, i.e., spring and summer (Osterberg et al., 2008; Ewing et al., 2010; Gross et al., 2012; Asher et al., 2018).

\* Corresponding author at: 306 EOS Main, 2020-2207 Main Mall, Vancouver, BC V6T 1Z4, Canada.

E-mail address: [mli@eoas.ubc.ca](mailto:mli@eoas.ubc.ca) (M. Li).

Lead isotopes are a powerful tool for tracking sources of contamination in the environment (Patterson and Settle, 1987; Komárek et al., 2008; Cheng and Hu, 2010). The relative abundance of Pb isotopes ( $^{204}\text{Pb}$ ,  $^{206}\text{Pb}$ ,  $^{207}\text{Pb}$ , and  $^{208}\text{Pb}$ ) varies systematically with geological setting and the time since geologic formation (e.g., age of the bedrock, ore body) (Faure and Mensing, 2005; Weis, 2017). Different environmental Pb sources are frequently distinguished by their characteristic isotope ratios (Komárek et al., 2008; Cheng and Hu, 2010). Lead enters marine food webs through primary producers and undergoes negligible isotopic fractionation within the biosphere. Therefore, the environmental source(s) of Pb are the determinant of isotopic composition in marine species, rather than physical, chemical, and biological processes (Flegal et al., 1987; Michaels and Flegal, 1990; Smith et al., 1990). Consequently, Pb isotope ratios in shellfish and fish are expected to reflect the Pb isotopic composition of the ambient water, sediment, and diet (Prosi, 1989). Previous studies have shown that the Pb isotopic composition of biological samples can provide a fingerprint for sources of Pb (Spencer et al., 2000; Ip et al., 2005).

Scarce Pb data exist for seawater and marine biota from the northeast Pacific Ocean. Substantial inputs of Pb from Asian dusts and aerosols have been observed in western North America (Osterberg et al., 2008; Ewing et al., 2010; Gross et al., 2012; Asher et al., 2018). In this study, we address the following research questions: What are the dominant sources of present-day Pb in the northeast Pacific Ocean? To what extent do Pb emissions from Asian industries, principally China, account for the Pb found in this region? We investigate the relative importance of various Pb sources across inland, coastal, and the open ocean waters of the northeast Pacific. We use a variety of fish and bivalves from British Columbia (BC), Canada and report Pb isotope and concentration data for these species. Juvenile and returning adult Pacific salmon reflect the Pb isotopic composition of their freshwater and open ocean habitats, respectively, and coastal fish (herring) and shellfish (mussels, and clams) act as bioindicators for Pb exposure in coastal waters.

## 2. Materials and methods

### 2.1. Sample collection and processing

Fish and bivalve species were collected from diverse habitats across >800 km of coastline in British Columbia, Canada (Table 1, Fig. 1). All samples were stored frozen after collection. Four species of returning adult Pacific salmon (chinook, sockeye, chum, pink) were collected in 2014 and 2018 from Rivers Inlet on the central coast of BC. Juvenile sockeye salmon (fry) were

captured with beach seines in 2015 in Owikeno Lake, a large glacial lake that is surrounded by coastal volcanic mountains and empties into Rivers Inlet via the Wannock River. Pacific herring were collected from five sites spanning Strait of Georgia and central and north BC coast. Spring pre-spawning and spawning herring aggregations were sampled in 2016 from purse seine surveys (Haida Bay; Denman Island) conducted by Fisheries and Oceans Canada and with cast nets by independent research surveys (other sites). Mussel samples were handpicked from Third Beach on Calvert Island on the central coast of BC in 2018. Clams were collected from four sites near Quadra Island in the northern Strait of Georgia in 2007. The Strait of Georgia is a semi-enclosed estuarine sea between the BC mainland and Vancouver Island and has different oceanographic conditions from other BC coastal environments that connect directly to open ocean (Ianson et al., 2016). Hereafter, we call other coastal regions including the west coast of Vancouver Island and the central and north BC coast as coastal ocean to distinguish them from Strait of Georgia (Fig. 1).

The Pb concentrations of fish muscle tissue have direct implications for human health and trophic transfer along the food chain but are generally much lower than other biological tissues in fish (e.g., gill, liver, intestine, hard tissues) (Jeziarska and Witeska, 2006). Limited data have been reported for Pb levels in the muscle of marine fish, which is partly due to analytical challenges with low concentration levels. At the Department of Earth, Ocean and Atmospheric Sciences at the University of British Columbia (UBC), we subsampled fish muscle tissue (salmon and herring) with an acid-cleaned ceramic knife and freeze-dried the tissue prior to digestion and chemical analysis. Juvenile salmon samples were too small to fillet muscle tissue so were freeze-dried and digested as whole fish samples. Bivalve samples were shelled and prepared for analysis as individual samples, rather than pooled. After freeze-drying, all subsequent sample preparation and Pb concentration and isotopic analyses were performed in Class 1000 (or better) clean laboratories (with Class 100 laminar flow hoods) in the Pacific Centre for Isotopic and Geochemical Research at UBC.

### 2.2. Lead concentration and isotope analysis


Approximately 0.2 g of freeze-dried fish tissue was digested in 10 mL 14 M  $\text{HNO}_3$  using a laboratory microwave system (CEM Corporation, Matthews, NC, USA). The digests were completely evaporated and then reconstituted in a mixed solution of 0.5 % HCl, 1.5 %  $\text{HNO}_3$  and 10 ng/g indium (added as an internal standard). Following established sample preparation and analytical procedures (Smith et al., 2019), the Pb concentration and isotopic composition of the solution for all samples but clams were

**Table 1**  
Sample collection locations and predominant feeding habitats of fish and shellfish species used in this study.

Common name (Scientific name)	Sample Size	Capture Locations	Feeding Habitats	Mean Pb (1 SD) (ng/g dry weight)	
Sockeye Salmon Fry <sup>a</sup> ( <i>Oncorhynchus nerka</i> )	10	Owikeno lake	Freshwater Inland	230.8	(162.1)
Manila clams ( <i>Venerupis philippinarum</i> )	8	Quadra Island	Strait of Georgia	33.7	(24.9)
Pacific Oyster <sup>b</sup> ( <i>Crassostrea gigas</i> )	3	Desolation Sound	Strait of Georgia	89.3	(41.0)
Pacific Oyster <sup>b</sup> ( <i>Crassostrea gigas</i> )	8	Barkley Sound	Coastal ocean	126.8	(48.9)
Blue Mussel ( <i>Mytilus edulis</i> )	10	Calvert Island	Coastal ocean	208.6	(61.5)
Pacific Herring ( <i>Clupea pallasii</i> )	3	Denman Island	Coastal ocean	10.6	(2.8)
Pacific Herring ( <i>Clupea pallasii</i> )	4	Haida Bay	Coastal ocean	38.8	(10.8)
Pacific Herring ( <i>Clupea pallasii</i> )	7	Kitasu Bay	Coastal ocean	34.5	(22.0)
Pacific Herring ( <i>Clupea pallasii</i> )	2	Kwakshua Channel	Coastal ocean	23.9	(17.0)
Pacific Herring ( <i>Clupea pallasii</i> )	4	Smith Inlet	Coastal ocean	31.2	(15.4)
Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> )	3	Rivers Inlet	Open ocean	17.7	(3.3)
Chum Salmon ( <i>Oncorhynchus keta</i> )	2	Rivers Inlet	Open ocean	21.1	(13.3)
Pink Salmon ( <i>Oncorhynchus gorbuscha</i> )	3	Rivers Inlet	Open ocean	9.3	(4.4)
Sockeye Salmon ( <i>Oncorhynchus nerka</i> )	9	Rivers Inlet	Open ocean	8.5	(4.3)

<sup>a</sup> Whole-body digestion.

<sup>b</sup> Data from Shiel et al. (2012).



**Fig. 1.** Sample locations of fish and shellfish collected along the BC coast. All Pacific salmon are adult salmon that returned to Rivers Inlet to spawn. Sockeye fry are the juvenile salmon that lived in Owikeno Lake, upstream of Rivers Inlet. The oyster samples were collected in southern BC (1) Mainland and (2) Vancouver Island and data were reported by Shiel et al. (2012). Pacific herring samples were collected from five sites along the BC coast from north to south (1) Haida Bay, (2) Kitasu Bay, (3) Kwakshua channel, (4) Smith Inlet, (5) Denman Island. The striped areas and transitional domain signify the major feeding areas of Pacific herring and Rivers Inlet sockeye salmon, respectively (Espinasse et al., 2018; Fisheries and Oceans Canada, 2019). The grey star signifies the approximate location of Trail Teck Operations, a large Canadian metal and mining company.

analyzed using an Agilent 7700x quadrupole inductively coupled plasma mass spectrometer (ICP-MS, Agilent Technologies, Santa Clara, CA, USA) and a Nu AttoM high-resolution (sector-field) inductively coupled plasma mass spectrometer (HR-ICP-MS, Nu Instruments Ltd., UK), respectively. Clam samples were prepared and measured in the same way as oyster samples in a previous study and details regarding sample preparation and analytical methods can be found in Shiel et al. (2012). Briefly, ELEMENT2 HR-ICP-MS (Thermo Finnigan, Germany) and Nu Plasma multi-collector inductively coupled plasma mass spectrometer (MC-ICP-MS, Nu Instruments Ltd., UK) analyzed Pb concentration and isotopic composition of clam samples, respectively (Shiel et al., 2012).

There are no marine biological standard reference materials (SRMs) currently commercially available that are certified for Pb isotopic composition. We used TORT-3 (Lobster Hepatopancreas Reference Material, National Research Council Canada) as our in-house reference material because it is certified for Pb concentrations and is typically used in studies that focus on trace metals in marine fauna. We found that TORT-3, among other SRMs (e.g., DORM-4 fish protein; SRM 1566b oyster tissue), behaved most similarly to our fish and shellfish samples during sample digestion. TORT-3 was prepared and analyzed in the same manner as samples. Precision of Pb concentrations, estimated by replicate analysis of TORT-3 and fish samples, was better than 9 % ( $n = 6$ ) and 10 % ( $n = 4$ ), respectively. For Pb isotope analysis by HR-ICP-MS, each sample was bracketed by NIST SRM 981 for mass discrimination correction, using the standard-sample bracketing (SSB) correction method. For a given batch, the samples and bracketing standards were all prepared so the final concentration of Pb matched within 15 % for all analytes. We are currently building a Pb isotope database for SRM TORT-3 and use it as an in-house reference. Characterization of TORT-3 by both HR-ICP-MS and MC-ICP-MS yields consistent Pb isotope results

( $^{206}\text{Pb}/^{207}\text{Pb} = 1.165 \pm 0.005$  (2SD),  $^{208}\text{Pb}/^{206}\text{Pb} = 2.090 \pm 0.008$  from HR-ICP-MS ( $n = 9$ );  $^{206}\text{Pb}/^{207}\text{Pb} = 1.164 \pm 0.001$ ,  $^{208}\text{Pb}/^{206}\text{Pb} = 2.090 \pm 0.002$  from MC-ICP-MS ( $n = 3$ )).

### 3. Calculations


We estimated the fraction of Pb contributed by Asian sources in each sample using methods modified from previous studies (Ewing et al., 2010; Asher et al., 2018). As Chinese source ores and coals have characteristic higher  $^{208}\text{Pb}/^{206}\text{Pb}$  due to higher Th/U in source rocks (Bollhöfer and Rosman, 2001; Bi et al., 2017), Asian Pb influence can be quantified using the vertical divergence of  $^{208}\text{Pb}/^{206}\text{Pb}$  ratios from the local BC Pb isotopic array without Asian Pb influence. Here we use the leaded gasoline line to represent historical BC Pb isotope characteristics given its dominant impacts on the Pb isotopic signatures found in North America (Bollhöfer and Rosman, 2001). The fraction of Asian influence ( $F_{\text{Asia}}$ ) is calculated as the ratio of  $\Delta^{208}\text{Pb}$  Obs/ $\Delta^{208}\text{Pb}$  Asia (Eq. (3)), where  $\Delta^{208}\text{Pb}$  signifies the offset between the expected BC  $^{208}\text{Pb}/^{206}\text{Pb}$  ratio and either the observed or the expected Asian  $^{208}\text{Pb}/^{206}\text{Pb}$  ratio for a given  $^{206}\text{Pb}/^{207}\text{Pb}$  ratio (Eqs. (1) and (2) and Fig. 2) (Ewing et al., 2010).

$$\Delta^{208}\text{Pb Obs} = ^{208}\text{Pb}/^{206}\text{Pb}_{\text{Observed}} - ^{208}\text{Pb}/^{206}\text{Pb}_{\text{BC}} \quad (1)$$

$$\Delta^{208}\text{Pb Asia} = ^{208}\text{Pb}/^{206}\text{Pb}_{\text{Asia}} - ^{208}\text{Pb}/^{206}\text{Pb}_{\text{BC}} \quad (2)$$

$$F_{\text{Asia}} (\%) = \Delta^{208}\text{Pb Obs} / \Delta^{208}\text{Pb Asia} * 100 \quad (3)$$

For local and regional context, we compare the Pb data from this study with environmental samples previously collected in the North Pacific Ocean and west coast of North America (California and northward). The Pb isotopic signatures in the ocean have



**Fig. 2.** Illustration of the calculation method for Asian Pb as a fraction of total Pb. The leaded gasoline line connects two major ore deposits (Australian Broken Hill and US Mississippi Valley) previously used for producing tetraethyllead used worldwide (Sangster et al., 2000; Soto-Jimenez et al., 2006).

changed substantially with time due to bans on leaded gasoline and rapidly growing industries in Asia. As the residence time of Pb in North Pacific surface water is relatively short (6–20 months) (Tsunogai and Nozaki, 1971; Gallon et al., 2006; Boyle et al., 2014), we limited our comparison to recent data in the literature (2000–present day). Differences in isotopic compositions and the extent of Asian Pb influence among fish from different feeding habitats were examined using ANOVA and Tukey's Honest Significant Difference (HSD) test. All statistical analyses were performed using R version 3.1.1 (R Core Team, 2014).

## 4. Results

### 4.1. Lead concentrations in fish and shellfish

Lead concentrations in muscle tissue of adult Pacific salmon ( $11.7 \pm 7.0$  ng/g dry weight) are higher than fresh albacore tuna muscle (1.5 ng/g, assuming 80 % moisture content) measured by Settle and Patterson (1980), which could be explained by a variety of factors including different Pb contamination levels in seawater between present day and four decades ago and distinct life histories and habitats across species. Lead concentrations in Pacific salmon are nearly 3 and 11 times lower than those of Pacific herring ( $30.1 \pm 17.7$  ng/g dry weight) and bivalves ( $125.1 \pm 84.7$  ng/g dry weight). This is consistent with prior findings that biomagnification of Pb generally does not occur during trophic transfer and inverse relationship between trophic levels and Pb concentrations in muscle tissue has been observed in laboratory and field research (Michaels and Flegal, 1990; Suedel et al., 1994; Cardwell et al., 2013). The average species-specific Pb concentrations by capture locations can be found in Table 1 and detailed Pb concentrations and isotopic compositions of each sample can be viewed in Table S1. Due to the whole-body digestion involving calcareous skeletal materials where Pb is concentrated relative to muscle tissue, the Pb concentrations of sockeye fry should not be compared directly with the muscle tissue of other fish.

The differences among Pb concentrations of salmon, herring, and bivalves largely result from distinct exposure pathways and physiological characteristics of individual species. For example, the primary exposure route of Pb for finfish appears to be the direct

uptake of aqueous  $Pb^{2+}$  across the gills and dietary Pb intake can be readily excreted (Spry and Wiener, 1991; Jezierska and Witeska, 2006). In contrast, besides aqueous Pb intake, suspended particulate matters adsorbing much higher levels of Pb than ambient water concentrations also contributes considerably to the body burden of Pb in filter-feeding organisms such as bivalves (Prosi, 1989). This explains the much higher Pb tissue concentrations found in bivalves than finfish.

### 4.2. Lead isotopic compositions of marine biota vary with feeding habitat


Lead isotopic compositions of fish and shellfish that feed in lake, coastal, and open ocean waters are notably different, exhibiting a clear trend of Pb isotope ratios varying from highest  $^{206}Pb/^{207}Pb$  in Owikeno Lake to lowest  $^{206}Pb/^{207}Pb$  in the open ocean northeast Pacific (Fig. 3; see Fig. 1 for capture locations). Along with previously published local geological and environmental samples, the Pb isotopic composition of samples collected in this study form the BC Pb regression line ( $R^2 = 0.95$ , Fig. 3). Some of our Pacific herring results deviate from the BC Pb regression line and we address this anomaly in the discussion section. Sockeye salmon fry have not left Owikeno Lake since hatching and their Pb isotopic compositions mostly overlap those of BC volcanic rocks and Cascadia basin sediments (Mullen and Weis, 2013; Carpentier et al., 2014). This indicates that the sockeye salmon fry were primarily exposed to natural sources of Pb and that anthropogenic sources had minimal contribution to their Pb burden.

Bivalves (clams, mussels, and oysters) are filter feeders living on or near the seafloor and they obtain metals from both seawater and suspended solids including suspended sedimentary particles (Prosi, 1989; Gagnon and Fisher, 1997). Prior studies have found that Pb concentration and isotopic compositions in bivalves are more likely determined by sediment than seawater (Flegal et al., 1987; Wang et al., 1996; Shulkin et al., 2003; Dang et al., 2015). Bivalve samples collected in south, central BC coast, and Strait of Georgia display a wide range of Pb isotope ratios ( $^{206}Pb/^{207}Pb$ : 1.15–1.18), which overlap those of other environmental samples (road dust, lichen, river particulate matter) collected from southern BC where most of the BC human population and industries are located (Fig. 3). We infer from our data that diverse Pb sources, probably from southern BC, have contributed to the Pb content of suspended particles in southern and central BC coastal waters and Strait of Georgia, and ultimately lead to the variability of the Pb compositions in bivalves.

In contrast to the observed variation of Pb isotopic compositions in bivalves, Pacific herring exhibit smaller variation in  $^{206}Pb/^{207}Pb$  (range: 1.16–1.17) and are not correlated with capture locations (Fig. S1). Herring is a coastal pelagic fish species that forage along BC coast and spawn in bays and estuaries (Fig. 1) (Hay et al., 2009; Therriault et al., 2009), and therefore their Pb isotopic compositions are expected to reflect those of seawater and be minimally influenced by sediment. The observed limited Pb isotopic variation in herring indicates a relatively homogenous Pb source in BC coastal ocean seawater and/or their migratory behaviors have likely helped integrate Pb exposures from various coastal locations (Fisheries and Oceans Canada, 2019).

Adult Pacific salmon exhibit distinct Pb isotopic compositions from those of inland and coastal fish (Fig. 3). Before returning to freshwater to spawn, the Rivers Inlet sockeye salmon typically spend 2–3 years foraging and obtain 90 % of their final adult body weight in offshore waters of the northeast Pacific Ocean, primarily in the easterly flowing subarctic current of the transitional domain (Fig. 1) (Groot and Margolis, 1991; Ishida et al., 1998; Espinasse et al., 2018). Therefore, these adult sockeye






**Fig. 3.** Lead isotopic composition of all analyzed fish and shellfish from this study, with local geologic and environmental context. The solid line illustrates the Pb regression lines between local lithology, including the coast plutonic complex (Cui and Russell, 1995), Garibaldi volcanic belt (Mullen and Weis, 2013), Pacific sediment (Carpentier et al., 2014), Sullivan mine ore (Sangster et al., 2000), and a variety of environmental samples collected in BC, including fish and shellfish (this study), lichens (Simonetti et al., 2003), road dust (Preciado et al., 2007), BC oysters (Shiel et al., 2012), seawater (McAlister, 2015), and Fraser river particulate matter (Snauffer et al., 2010). The dashed line represents the leaded gasoline curve connecting two major ore deposits previously used for producing tetraethyllead used worldwide (Sangster et al., 2000; Soto-Jimenez et al., 2006). Error bars denote analytical uncertainty (2SE). Clams and oysters were analyzed by MC-ICP-MS and their error bars are smaller than the symbol size.

salmon would have accumulated most of their Pb from open ocean habitats and their Pb isotope ratios indicate that the open ocean seawater has a lower  $^{206}\text{Pb}/^{207}\text{Pb}$  ratio compared to that of inland and coastal waters (Fig. 3). Adult chum and pink salmon from Rivers Inlet also mature in the offshore northeast Pacific and they show similar Pb isotopic compositions to those of sockeye

salmon (Groot and Margolis, 1991; Ishida et al., 1998), confirming that lower  $^{206}\text{Pb}/^{207}\text{Pb}$  compositions exist in the northeast Pacific Ocean. Unlike the other salmon species, the Pb isotopic compositions of Chinook salmon overlap with both oceanic and coastal fish (Fig. 3). This can be explained by Chinook salmon's behavioral differences in migration — some complete



**Fig. 4.** Comparison of Pb isotopic composition between fish and shellfish collected from the BC coast and relevant environmental data from western North America and the northern Pacific Ocean. Data for seawater in the northwest Pacific (Zurbrick et al., 2017), northeast Pacific (McAlister, 2015), air samples from central California (Ewing et al., 2010), and honey samples from the Greater Vancouver Regional District (Smith et al., 2019) are also illustrated. Figure legend is the same as Fig. 3. The regression line of Chinese coal and ore deposit is from data compiled by Bi et al. (2017) and Cheng and Hu (2010). The selected sampling locations within the Line P transect in the northeast Pacific Ocean are depicted in the bottom left (Schuback, 2016).

extensive offshore oceanic migration while others spend most of their ocean life in coastal waters (Groot and Margolis, 1991). Our results therefore imply the great potential of Pb isotopes to distinguish different habitat use among fish stocks.

## 5. Discussion

### 5.1. Input of anthropogenic Pb to the northeast Pacific Ocean

Given that natural Pb sources in southern BC (Garibaldi volcanic belt; Cascadia basin sediments) and Alaska (Aleutians) exhibit high  $^{206}\text{Pb}/^{207}\text{Pb}$  and low  $^{208}\text{Pb}/^{206}\text{Pb}$  ratios (Fig. 3) (Kay et al., 1978; Mullen and Weis, 2013; Carpentier et al., 2014), the Pb isotopic compositions of Pacific salmon imply that there must be some anthropogenic Pb sources with low  $^{206}\text{Pb}/^{207}\text{Pb}$  and high  $^{208}\text{Pb}/^{206}\text{Pb}$  values dominating in the northeast Pacific Ocean. Recent studies reported Pb isotopic composition of honey samples from the Greater Vancouver Regional District, BC and air samples from cities in California's Central Valley also resemble those of Pacific salmon (Fig. 4) (Ewing et al., 2010; Smith et al., 2019). We therefore suggest that similar anthropogenic Pb sources prevail in western North America and the northeast Pacific Ocean. We further explore this anthropogenic Pb source by examining the relationship between  $^{208}\text{Pb}/^{206}\text{Pb}$  values and the inverse of Pb concentration ( $1/[\text{Pb}]$ ) in Pacific salmon (Fig. 5a). We find a two-end member mixing model can explain the variability in the Pb content and isotopic compositions in all salmon samples except for two Chinook samples. The higher Pb concentrations in these salmon are potentially a result of more influence from Pb source(s) with  $^{208}\text{Pb}/^{206}\text{Pb}$  near 2.14 in the open ocean (black star in Fig. 5a).

The BC Pb regression line based on local geological and environmental samples ( $y = -1.05x + 3.31$ ) is very similar to the world legacy gasoline Pb line resulting from the addition of tetraethyl Pb (TEL) as an antiknock agent in gasoline in the late 20th century ( $y = -1.13 + 3.40$ ) in Fig. 3. The majority of fish samples in this study align well along the legacy gasoline Pb line, suggesting that they are likely impacted by Pb sources resembling legacy gasoline Pb (Australian type and US Mississippi Valley type ores). The Pb end-member in the northeast Pacific Ocean reflected by salmon samples points to Australian-type Pb ore compositions (Mt. Isa and Broken Hill).

Some regional emissions of Pb ultimately sourced from Mt. Isa ore deposit may explain the observed Pb isotopic composition in salmon from the northeast Pacific. Prior studies attributed anthropogenic Pb in BC to the smelting and refining Trail facility of Teck Metals Ltd in southern BC (Fig. 1) and the use of aviation gasoline by small, propeller aircraft, which contains high levels of TEL (Simonetti et al., 2003; Shiel et al., 2012). The Trail facility has the largest on-site release of Pb to the atmosphere in BC (>700 kg/year during 2010–2014 and >1500 kg/year pre-2010) (Government of Canada, 2019). The Pb used for TEL production and for smelting at Trail is largely from Mt. Isa ore deposit after BC Sullivan mine was closed in 2001 (Shiel et al., 2012), characterized by a distinctly low  $^{206}\text{Pb}/^{207}\text{Pb}$  ratio ( $\approx 1.04$ ) (Sangster et al., 2000). Additional Pb isotopic measurements that assess major regional industrial sectors (e.g., fuel combustion, smelting, wastewater) are needed to further identify specific emission sources contributing Pb to the northeast Pacific Ocean and their evolution over time.


### 5.2. Distinct Pb sources in coastal waters from open ocean

The differences of Pb isotopic compositions observed in coastal samples (bivalves and herring) and adult salmon imply that the dominant Pb sources in coastal waters have lower  $^{208}\text{Pb}/^{206}\text{Pb}$  than that of the open ocean and Australian-type Pb inputs (Fig. 3). Consistently, the predicted dominant Pb source for coastal fish (herring) has a lower  $^{208}\text{Pb}/^{206}\text{Pb}$  value (2.11) than that derived from the majority of salmon samples (2.14) (black stars in Fig. 5). The two Chinook samples that do not follow the general pattern of Pb source attribution for other salmon have Pb isotopic compositions overlapping those of coastal fish and shellfish in Fig. 3. We infer that the same coastal Pb sources likely dominate Pb exposure in these Chinook and result in their lower  $^{208}\text{Pb}/^{206}\text{Pb}$  values and higher Pb concentrations relative to those of other salmon samples (Fig. 5a).

Based on the literature values of Pb isotope ratios in other environmental samples, we postulate that the Pb in coastal waters are from urban runoff and river flows into coastal ocean and/or the coastal upwelling that brings deeper waters with legacy Asian and North American industrial lead ( $^{206}\text{Pb}/^{207}\text{Pb} = 1.16\text{--}1.21$ ;  $^{208}\text{Pb}/^{206}\text{Pb} = 2.02\text{--}2.12$ ) to surface waters (Flegal et al., 1986; McAlister, 2015). It is worth noting that five herring samples collected from the southern and central BC coast have the lowest Pb concentrations among all herring samples and fall off the mixing line derived from more contaminated herring (Fig. 5b). Although aqueous Pb concentration is the determining factor for Pb uptake, Pb concentrations in fish are also affected by many environmental, ecological, and physiological characteristics of individual species (Prosi, 1989). Our data reveal that it is challenging to use Pb isotopes in biological materials for tracking contamination sources when the Pb exposure is very low.

### 5.3. Influences of Chinese Pb in coastal and open ocean waters

Chinese coal, ore deposits, and aerosols have a distinct Pb isotopic fingerprint characterized by enrichment in  $^{208}\text{Pb}$ , a



**Fig. 5.** Lead isotope ratio  $^{208}\text{Pb}/^{206}\text{Pb}$  vs.  $1/[\text{Pb}]$  concentration ( $\text{ng/g dry weight}^{-1}$ ) for (a) Pacific salmon and (b) Pacific herring. The stars represent the Pb pollution sources predicted based on linear regression. Samples in parentheses are not considered for linear regression (see explanation in text). The numbers in square symbols in (b) illustrate the capture location of each herring sample, as described in Fig. 1.

radioactive decay product of  $^{232}\text{Th}$  (Bollhöfer and Rosman, 2001; Bi et al., 2017). A number of North American studies have shown that environmental samples influenced by Chinese emissions tend to deviate from the local Pb isotopic mixing line toward the Chinese Pb regression line (Ewing et al., 2010; Christensen et al., 2015; Asher et al., 2018). Strong influence of Chinese Pb emissions in other East Asian countries such as Japan and South Korea is well known and previous studies concluded that Chinese emissions are the dominant source of Pb input to surface waters of the northwest and central Pacific Ocean (Wu et al., 2010; Gallon et al., 2011; Zurbrick et al., 2017). This is confirmed by Pb isotopic compositions of the surface waters of the Kuroshio current and the western subarctic gyre located offshore of Japan, plotting nearer to the Chinese Pb regression line (purple shade in Fig. 4) (Zurbrick et al., 2017). In contrast, Pb isotopic ratios of surface waters collected along line P in the northeast Pacific Ocean align well with the BC Pb mixing line and show little deviation toward the Chinese Pb line (green and blue shades in Fig. 4) (McAlister, 2015). The difference in seawater Pb isotope ratios between the northwest and northeast Pacific Ocean highlights the difference in predominant Pb sources between these geographic regions and indicates lower contribution of Pb from Chinese emissions in the northeast Pacific relative to the northwest Pacific.

Fig. 6 shows the species-specific fractions of Asian Pb influence derived from the vertical divergence of their  $^{208}\text{Pb}/^{206}\text{Pb}$  ratios from the legacy gasoline line (see method in Fig. 2). We find that Asian Pb contributes significantly higher input (31 % of total Pb) in BC coastal ocean as reflected in the herring and central coast shellfish than other ecological zones (4 % in the open ocean, 5 % in northern Strait of Georgia, and minimal in inland freshwater ecosystems) (Figs. 6 and S2). Similar elevated Asian Pb impact was found in the sediment of lower Alberni Inlet and Barkley Sound on the west coast of Vancouver Island relative to the upper Alberni Inlet directly connected and influenced by upstream terrestrial/freshwater ecosystems (Ikehata, 2013). Determining the exact cause for this geographic discrepancy in Asian Pb impact will require further research. Here we provide two scenarios that may lead to the elevated Asian Pb fractions in coastal ocean observed in this study.

In the first scenario, heavier precipitation along the BC coast as an effect of the coastal mountain range may result in transporting higher amounts of Pb from the atmosphere to coastal ocean


through wet deposition, compared to the open ocean or inland BC. Most Pb in the ocean originates from atmospheric deposition and surface water Pb concentrations tend to follow atmospheric deposition fluxes of the previous few years (Boyle et al., 2014). Castello and Shelton (2004) estimated that the spatial distribution of total winter precipitation on the Pacific coast of BC and Washington rapidly decreases from 60 cm on the coast to 10 cm in open ocean within ~500 km. Several studies estimated trans-Pacific influence of Asian aerosols and dust account for an average of ~46 % of Pb along the coast of California during peak seasons (spring-summer) of Asian dust influence on North America (Ewing et al., 2010; Asher et al., 2018). Pacific herring in this study would reflect an accumulation of Pb from all seasons over 3–5 years (Ware, 1985), and suggest annual average Asian Pb input (39 % of total Pb) along the BC coast.

An alternative explanation involves anthropogenic source(s) that occur predominantly in the coastal ocean but not in the open ocean or inland. We consider emissions from heavy ship traffic in BC coastal ocean as the most plausible source (Johannessen and Macdonald, 2009). Central and northern BC have little industrial activities and small human populations (<0.5 people per sq. kilometer), therefore, the direct release of Pb to the oceans from BC inland activities is likely minimal (Statistics Canada, 2019). The BC coast serves 25 routes for frequent ferries within BC and additional routes from Washington, USA. It is also the inevitable cruise route to Alaska, which hosts about 500 voyages per year (Cruiselines International Association, 2019). Oceangoing traffic could release a considerable amount of Pb into the air and seawater given the high Pb content of these fuels (e.g., diesel and heavy fuel oil) (Akpoveta and Osakwe, 2014; Bi et al., 2017). The lower Asian Pb fractions observed in bivalves from the northern Strait of Georgia, relative to those from coastal ocean, could be a result of its restricted seawater and sediment exchange with coastal and open ocean (Ianson et al., 2016) (Fig. 6). Future research that investigates the Pb isotope compositions of fuels used for various kinds of marine transport along BC coast is required to elucidate the Pb contribution of marine traffic in the northeast Pacific coastal waters.

## 6. Conclusions

This study takes the initial step to establish the Pb isotope landscape of the northeast Pacific Ocean and shows a clear trend in Pb isotopic compositions, from highest  $^{206}\text{Pb}/^{207}\text{Pb}$  in inland fish to lowest  $^{206}\text{Pb}/^{207}\text{Pb}$  in open ocean fish. We conclude that some Australian-type Pb end-member source(s) with low  $^{206}\text{Pb}/^{207}\text{Pb}$  and high  $^{208}\text{Pb}/^{206}\text{Pb}$  values largely contribute to the Pb in the northeast Pacific whereas natural sources of Pb dominate freshwater ecosystems in the central BC coast. We estimate that Chinese-sourced Pb contributes to about 31 % of total Pb in coastal species but it has minimal input to the Pb in the freshwater lakes of central BC and open ocean. We propose heavy marine traffic and precipitation in coastal ocean as the most plausible explanations and thus themes for future research.

The distinct Pb isotopic compositions across species primarily foraging in inland, coastal, and offshore regions indicate great potential for using Pb isotopes to distinguish among habitats (e.g., coastal vs. open ocean) of migratory species in the northeast Pacific Ocean. To date, very few studies have explored the use of Pb isotopes for distinguishing stocks of marine organisms (fish, cephalopods, mammals) and identifying their nursery or feeding grounds (Outridge and Stewart, 1999; Spencer et al., 2000; Raimundo et al., 2009). Our results show the promise of Pb isotopes to discern two different behavioral patterns (offshore oceanic vs. coastal migration) within Chinook stocks. With further efforts to develop the Pb isotope landscape in the northeast Pacific and BC coast (e.g., GEOTRACES cruise GP15 in 2018, seasonal line



**Fig. 6.** The fractions of Asian Pb influence in each species used in this study, estimated by the deviation toward the Chinese Pb line from the leaded gasoline Pb line (Fig. 2). Box and whiskers show interquartile and 1.5 times the interquartile, respectively. Estimated negative values of Asian Pb contribution results from the uncertainty associated with the regression slopes and analytical uncertainty of fish measurements and they are not significant from zero (see Fig. 3).



P cruises by UBC), we expect that Pb isotopes will become an effective tool for answering critical ecological and environmental questions in the region such as identifying pollution sources and foraging hotspots for species that have tremendous ecological and economic importance.

## Supporting information

Additional two figures and all data produced in this study are available in supporting information.

## Declaration of Competing Interest

The authors declare no competing financial interest.

## Acknowledgements

M. Li's postdoctoral fellowship was provided by Multidisciplinary Applied Geochemistry Network. All chemical analyses were covered by D. Weis's Discovery Grant from Natural Sciences and Engineering Research Council of Canada. Collection of salmon samples in Rivers Inlet was supported by a grant from the Monell and Vetlesen foundations, and logistical support from the Wuikinuxv First Nation and the Hakai Institute. The authors thank Maureen Soon, Vivian Lai, Kathy Gordon, and Marghalery Amini (Pacific Centre for Isotopic and Geochemical Research, Dept. of Earth, Ocean and Atmospheric Sciences, UBC) for their technical and analytical assistance. We also thank Alyssa Gehman (UBC, Hakai Institute) for collecting mussel samples and Patricia Schulte (Dept. of Zoology at UBC) for insightful discussions.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.ancene.2019.100234>.

## References

- Akpoveta, O., Osakwe, S., 2014. Determination of heavy metal contents in refined petroleum products. *IOSR J. Appl. Chem.* 7, 01–02.
- Asher, E.C., Christensen, J.N., Post, A., Perry, K., Cliff, S.S., Zhao, Y., Trousdell, J., Faloon, I., 2018. The transport of asian dust and combustion aerosols and associated ozone to North America as observed from a mountaintop monitoring site in the California coast Range. *J. Geophys. Res. Atmos.* 123, 5667–5680.
- Bi, X.Y., Li, Z.G., Wang, S.X., Zhang, L., Xu, R., Liu, J.L., Yang, H.M., Guo, M.Z., 2017. Lead isotopic compositions of selected coals, Pb/Zn ores and fuels in China and the application for source tracing. *Environ. Sci. Technol.* 51, 13502–13508.
- Bollhöfer, A., Rosman, K.J.R., 2001. Isotopic source signatures for atmospheric lead: the Northern Hemisphere. *Geochim. Cosmochim. Acta* 65, 1727–1740.
- Bory, A.J.M., Abouchami, W., Galer, S.J.G., Svensson, A., Christensen, J.N., Biscaye, P.E., 2014. A Chinese imprint in insoluble pollutants recently deposited in central Greenland as indicated by lead isotopes. *Environ. Sci. Technol.* 48, 1451–1457.
- Boyle, E., Lee, J.-M., Echegoyen, Y., Noble, A., Moos, S., Carrasco, G., Zhao, N., Kayser, R., Zhang, J., Gamo, T., Obata, H., Norisuye, K., 2014. Anthropogenic lead emissions in the ocean: the evolving global experiment. *Oceanography* 27, 69–75.
- Cardwell, R.D., DeForest, D.K., Brix, K.V., Adams, W.J., 2013. Do Cd, Cu, Ni, Pb, and Zn biomagnify in aquatic ecosystems? Reviews of Environmental Contamination and Toxicology, Vol. 226. Springer, pp. 101–122.
- Carpentier, M., Weis, D., Chauvel, C., 2014. Fractionation of Sr and Hf isotopes by mineral sorting in Cascadia basin terrigenous sediments. *Chem. Geol.* 382, 67–82.
- Castello, A.F., Shelton, M., 2004. Winter precipitation on the US Pacific Coast and El Niño–Southern oscillation events. *Int. J. Clim.* 24, 481–497.
- Chen, M., Boyle, E.A., Switzer, A.D., Gouramanis, C., 2016. A century long sedimentary record of anthropogenic lead (Pb), Pb isotopes and other trace metals in Singapore. *Environ. Pollut.* 213, 446–459.
- Cheng, H., Hu, Y., 2010. Lead (Pb) isotopic fingerprinting and its applications in lead pollution studies in China: a review. *Environ. Pollut.* 158, 1134–1146.
- Christensen, J.N., Weiss-Penzias, P., Fine, R., McDade, C.E., Trzepla, K., Brown, S.T., Gustin, M.S., 2015. Unraveling the sources of ground level ozone in the Intermountain Western United States using Pb isotopes. *Sci. Total Environ.* 530–531, 519–525.
- Cruiselines International Association, 2019. History of the Alaska Cruise Industry. . Accessed on Feb.8th, 2019 <http://www.cliaalaska.org/economy/alaska-cruise-history/>.
- Cui, Y., Russell, J., 1995. Nd-Sr-Pb isotopic studies of the southern Coast plutonic complex, southwestern British Columbia. *Geol. Soc. Am. Bull.* 107, 127–138.
- Dang, D.H., Schafer, J.R., Brach-Papa, C., Lenoble, V.R., Durrieu, G.L., Dutruich, L., Chiffolleau, J.-F., Gonzalez, J.-L., Blanc, G.R., Mullot, J.-U., 2015. Evidencing the impact of coastal contaminated sediments on mussels through Pb stable isotopes composition. *Environ. Sci. Technol.* 49, 11438–11448.
- Díaz-Somoano, M., Kylander, M.E., López-Antón, M.A., Suárez - Ruiz, I., Martínez-Tarazona, M.R., Ferrat, M., Kober, B., Weiss, D.J., 2009. Stable lead isotope compositions in selected coals from around the world and implications for present day aerosol source tracing. *Environ. Sci. Technol.* 43, 1078–1085.
- Eisenreich, S.J., Metzger, N.A., Urban, N.R., Robbins, J.A., 1986. Response of atmospheric lead to decreased use of lead in gasoline. *Environ. Sci. Technol.* 20, 171–174.
- Espinasse, B., Hunt, B.P., Doson Coll, Y., Pakhomov, E.A., 2018. Investigating high seas foraging conditions for salmon in the North Pacific: insights from a 100 year scale archive for Rivers Inlet sockeye salmon. *Can. J. Fish. Aquat. Sci.* 76, 918–927.
- Ewing, S.A., Christensen, J.N., Brown, S.T., Vancuren, R.A., Cliff, S.S., Depaolo, D.J., 2010. Pb isotopes as an indicator of the Asian contribution to particulate air pollution in urban California. *Environ. Sci. Technol.* 44, 8911–8916.
- Faure, G., Mensing, T.M., 2005. *Isotopes: Principles and Applications*. John Wiley & Sons.
- Fisheries and Oceans Canada, 2019. Herring Migratory Behaviour. . Accessed on January 15, 2019 <http://www.pac.dfo-mpo.gc.ca/science/species-especes/pelagic-pelagique/herring-hareng/hertags/pages/default4-eng.htm>.
- Flegal, A., Itoh, K., Patterson, C., Wong, C., 1986. Vertical profile of lead isotopic compositions in the north-east Pacific. *Nature* 321, 689.
- Flegal, A.R., Gallon, C., Ganguli, P.M., Conaway, C.H., 2013. All the lead in China. *Crit. Rev. Environ. Sci. Technol.* 43, 1869–1944.
- Flegal, A.R., Rosman, K.J., Stephenson, M.D., 1987. Isotope systematics of contaminant leads in Monterey Bay. *Environ. Sci. Technol.* 21, 1075–1079.
- Gagnon, C., Fisher, N.S., 1997. The bioavailability of sediment-bound Cd, Co, and Ag to the mussel *Mytilus edulis*. *Can. J. Fish. Aquat. Sci.* 54, 147–156.
- Gallon, C., Ranville, M.A., Conaway, C.H., Landing, W.M., Buck, C.S., Morton, P.L., Flegal, A.R., 2011. Asian industrial lead inputs to the north pacific evidenced by lead concentrations and isotopic compositions in surface waters and aerosols. *Environ. Sci. Technol.* 45, 9874–9882.
- Gallon, C., Tessier, A., Gobeil, C., Carignan, R., 2006. Historical perspective of industrial lead emissions to the atmosphere from a Canadian smelter. *Environ. Sci. Technol.* 40, 741–747.
- Government of Canada, 2019. National Pollutant Release Inventory. . Accessed on January 14, 2019 <https://pollution-waste.canada.ca/national-release-inventory/archives/index.cfm?do=query&lang=en>.
- Groot, C., Margolis, L., 1991. *Pacific Salmon Life Histories*. UBC press.
- Gross, B.H., Kreutz, K.J., Osterberg, E.C., McConnell, J.R., Handley, M., Wake, C.P., Yalcin, K., 2012. Constraining recent lead pollution sources in the North Pacific using ice core stable lead isotopes. *J. Geophys. Res.* 117, 1–15.
- Hay, D.E., McCarter, P.B., Daniel, K.S., Schweigert, J.F., 2009. Spatial diversity of Pacific herring (*Clupea pallasii*) spawning areas. *ICES J. Mar. Sci.* 66, 1662–1666.
- Hosono, T., Alvarez, K., Kuwae, M., 2016. Lead isotope ratios in six lake sediment cores from Japan Archipelago: historical record of trans-boundary pollution sources. *Sci. Total Environ.* 559, 24–37.
- Ianson, D., Allen, S.E., Moore-Maley, B.L., Johannessen, S.C., Macdonald, R.W., 2016. Vulnerability of a semienclosed estuarine sea to ocean acidification in contrast with hypoxia. *Geophys. Res. Lett.* 43, 5793–5801.
- Ikehata, M., 2013. Evaluation of Trace Metal Distributions (arsenic, Cadmium, Lead) and Lead Sources in Sediments From a Sound and an Inlet on the West Coast of Vancouver Island, British Columbia MSc thesis. University of British, Columbia. <https://open.library.ubc.ca/cIRcle/collections/ubctheses/24/items/1.0166846>.
- Inoue, M., Tanimizu, M., 2008. Anthropogenic lead inputs to the western Pacific during the 20th century. *Sci. Total Environ.* 406, 123–130.
- Ip, C.C.M., Li, X.D., Zhang, G., Wong, C.S.C., Zhang, W.L., 2005. Heavy metal and Pb isotopic compositions of aquatic organisms in the Pearl River Estuary, South China. *Environ. Pollut.* 138, 495–505.
- Ishida, Y., Ito, S.-O., Ueno, Y., Sakai, J., 1998. Seasonal growth patterns of Pacific salmon (*Oncorhynchus* spp.) in offshore waters of the North Pacific Ocean. *N. Pac. Anadr. Fish Comm. Bull.* 1, 66–80.
- Jeziarska, B., Witeska, M., 2006. The Metal Uptake and Accumulation in Fish Living in Polluted Waters. Pages 107–114 Soil and Water Pollution Monitoring, Protection and Remediation. Springer.
- Johannessen, S., Macdonald, R., 2009. Effects of local and global change on an inland sea: the Strait of Georgia, British Columbia. *Canada. Climate Research* 40, 1–21.
- Kay, R.W., Sun, S.S., Lee-Hu, C.N., 1978. Pb and Sr isotopes in volcanic rocks from the Aleutian Islands and Pribilof Islands. *Alaska. Geochimica et Cosmochimica Acta* 42, 263–273.
- Komárek, M., Ettler, V., Chrastný, V., Mihaljevič, M., 2008. Lead isotopes in environmental sciences: a review. *Environ. Int.* 34, 562–577.
- Li, Q., Cheng, H., Zhou, T., Lin, C., Guo, S., 2012. The estimated atmospheric lead emissions in China, 1990–2009. *Atmos. Environ.* 60, 1–8.
- McAlister, J., 2015. Biogeochemistry of Dissolved Gallium and Lead Isotopes in the Northeast Pacific and Western Arctic Oceans PhD thesis. University of British Columbia. <https://open.library.ubc.ca/cIRcle/collections/ubctheses/24/items/1.0166278>.
- Michaels, A.F., Flegal, A.R., 1990. Lead in marine planktonic organisms and pelagic food webs. *Limnol. Oceanogr.* 35, 287–295.
- Mullen, E.K., Weis, D., 2013. Sr-Nd-Hf-Pb isotope and trace element evidence for the origin of alkalic basalts in the Garibaldi Belt, northern Cascade arc. *Geochim. Geophys. Geosystems* 14, 3126–3155.



- Osterberg, E., Mayewski, P., Kreutz, K., Fisher, D., Handley, M., Sneed, S., Zdanowicz, C., Zheng, J., Demuth, M., Waskiewicz, M., 2008. Ice core record of rising lead pollution in the North Pacific atmosphere. *Geophys. Res. Lett.* 35.
- Outridge, P., Stewart, R.E., 1999. Stock discrimination of Atlantic walrus (*Odobenus rosmarus rosmarus*) in the eastern Canadian Arctic using lead isotope and element signatures in teeth. *Can. J. Fish. Aquat. Sci.* 56, 105–112.
- Patterson, C.C., Settle, D.M., 1987. Review of data on eolian fluxes of industrial and natural lead to the lands and seas in remote regions on a global scale. *Mar. Chem.* 22, 137–162.
- Preciado, H.F., Li, L.Y., Weis, D., 2007. Investigation of past and present multi-metal input along two highways of British Columbia, Canada, using lead isotopic signatures. *Water Air Soil Pollut.* 184, 127–139.
- Prosi, F., 1989. Factors controlling biological availability and toxic effects of lead in aquatic organisms. *Sci. Total Environ.* 79, 157–169.
- R Core Team, 2014. R: a Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Raimundo, J., Vale, C., Caetano, M., Cesario, R., Moura, I., 2009. Total lead and its stable isotopes in the digestive gland of *Octopus vulgaris* as a fingerprint. *Aquat. Biol.* 6, 25–30.
- Sangster, D.F., Outridge, P.M., Davis, W.J., 2000. Stable lead isotope characteristics of lead ore deposits of environmental significance. *Environ. Rev.* 8, 115–147.
- Schuback, N., 2016. On the Light and Iron Dependent Coupling of Carbon Fixation and Photosynthetic Electron Transport in Arctic and Subarctic Marine Phytoplankton MSc thesis. University of British Columbia. <https://open.library.ubc.ca/cIRcle/collections/ubctheses/24/items/1.0314322>.
- Settle, D.M., Patterson, C.C., 1980. Lead in albacore : guide to lead pollution in americans. *Science* 207, 1167–1176.
- Shiel, A.E., Weis, D., Orians, K.J., 2012. Tracing cadmium, zinc and lead sources in bivalves from the coasts of western Canada and the USA using isotopes. *Geochim. Cosmochim. Acta* 76, 175–190.
- Shulkin, V., Presley, B., Kavun, V.I., 2003. Metal concentrations in mussel *Crenomytilus grayanus* and oyster *Crassostrea gigas* in relation to contamination of ambient sediments. *Environ. Int.* 29, 493–502.
- Simonetti, A., Clement, G., Carignan, J., 2003. Tracing sources of atmospheric pollution in Western Canada using the Pb isotopic composition and heavy metal abundances of epiphytic lichens. *Atmos. Environ.* 37, 2853–2865.
- Smith, D.R., Flegal, A.R., Niemeyer, S., Estes, J.A., 1990. Stable lead isotopes evidence anthropogenic contamination in Alaskan sea otters. *Environ. Sci. Technol.* 24, 1517–1521.
- Smith, K.E., Weis, D., Amini, M., Shiel, A.E., Lai, V.W.-M., Gordon, K., 2019. Honey as a biomonitor for a changing world. *Nat. Sustain.* 2, 223–232.
- Snauffer, A., Menard, O., Kieffer, B., Francois, R., Weis, D., 2010. Trace Element Signatures of Particles in the Fraser River Estuary. AGU Fall Meeting Abstracts.
- Soto-Jimenez, M.F., Hibdon, S.A., Rankin, C.W., Aggarawl, J., Ruiz-Fernandez, A.C., Paez-Osuna, F., Flegal, A.R., 2006. Chronicling a century of lead pollution in Mexico: stable lead isotopic composition analyses of dated sediment cores. *Environ. Sci. Technol.* 40, 764–770.
- Spencer, K., Shafer, D.J., Gaudie, R.W., DeCarlo, E.H., 2000. Stable lead isotope ratios from distinct anthropogenic sources in fish otoliths: a potential nursery ground stock marker. *Comp. Biochem. Physiol., Part A Mol. Integr. Physiol.* 127, 273–284.
- Spry, D.J., Wiener, J.G., 1991. Metal bioavailability and toxicity to fish in low-alkalinity lakes: a critical review. *Environ. Pollut.* 71, 243–304.
- Statistics Canada, 2019. Census Profile 2016. . Accessed on Feb.8th, 2019 <https://www12.statcan.gc.ca/census-recensement/2016/dp-pd/prof/index.cfm?Lang=E>.
- Suedel, B., Boraczek, J., Peddicord, R., Clifford, P., Dillon, T., 1994. Trophic transfer and biomagnification potential of contaminants in aquatic ecosystems. *Reviews of Environmental Contamination and Toxicology*. Springer, pp. 21–89.
- Therriault, T.W., Hay, D.E., Schweigert, J.F., 2009. Biological overview and trends in pelagic forage fish abundance in the Salish Sea (Strait of Georgia, British Columbia). *Mar. Ornithol.* 37, 3–8.
- Tsunogai, S., Nozaki, Y., 1971. Lead-210 and polonium-210 in the surface water of the Pacific. *Geochem. J.* 5, 165–173.
- Wang, W.-X., Fisher, N.S., Luoma, S.N., 1996. Kinetic determinations of trace element bioaccumulation in the mussel *Mytilus edulis*. *Mar. Ecol. Prog. Ser.* 140, 91–113.
- Ware, D., 1985. Life history characteristics, reproductive value, and resilience of Pacific herring (*Clupea harengus pallasii*). *Can. J. Fish. Aquat. Sci.* 42, s127–s139.
- Weis, D., 2017. Lead isotopes. In: White, W.M. (Ed.), *Encyclopedia of Geochemistry: A Comprehensive Reference Source on the Chemistry of the Earth*. Springer International Publishing, Cham, pp. 1–5.
- Wu, J., Boyle, E.A., 1997. Lead in the western North Atlantic Ocean: completed response to leaded gasoline phaseout. *Geochim. Cosmochim. Acta* 61, 3279–3283.
- Wu, J., Rember, R., Jin, M., Boyle, E.A., Flegal, A.R., 2010. Isotopic evidence for the source of lead in the North Pacific abyssal water. *Geochim. Cosmochim. Acta* 74, 4629–4638.
- Zurbrick, C.M., Gallon, C., Flegal, A.R., 2017. Historic and industrial lead within the Northwest Pacific Ocean evidenced by lead isotopes in seawater. *Environ. Sci. Technol.* 51, 1203–1212.