

Article Type: Primary Research Articles

Sound the Alarm: A Meta-Analysis on the Effect of Aquatic Noise on Fish Behavior and Physiology

Running Title: Meta-Analysis on the Effects of Noise on Fish

Kieran Cox^{1,2,3*}, Lawrence P. Brennan¹, Travis G. Gerwing^{1,2,4}, Sarah E. Dudas^{1,2,3}, Francis Juanes¹

¹ Department of Biology, University of Victoria, Victoria, British Columbia, Canada

² Hakai Institute, Calvert Island, British Columbia, Canada

³ Department of Biology, Vancouver Island University, Nanaimo, British Columbia, Canada

⁴ Ecosystem Science and Management Program, University of Northern British Columbia, Prince George, British Columbia, Canada

*Corresponding Author. Kieran Cox. Email: kcox@uvic.ca Phone: +1 778 977-0142

Keywords: Aquatic Noise, Soundscape, Fish Behavior, Fish Physiology, Fitness

Consequences, Global Change, Noise Pollution, Systematic Review

Paper type: Primary Research Article

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.14106

This article is protected by copyright. All rights reserved.

Abstract

The aquatic environment is increasingly bombarded by a wide variety of noise pollutants, whose range and intensity are increasing with each passing decade. Yet little is known about how aquatic noise affects marine communities. To determine the implications that changes to the soundscape may have on fishes, a meta-analysis was conducted focusing on the ramifications of noise on fish behavior and physiology. Our meta-analysis identified 42 studies that produced 2,354 data points, which in turn indicated that anthropogenic noise negatively affects fish behavior and physiology. The most predominate responses occurred within foraging ability, predation risk, and reproductive success. Additionally, anthropogenic noise was shown to increase the hearing thresholds and cortisol levels of numerous species while tones, biological, and environmental noise were most likely to affect complex movements and swimming abilities. These findings suggest that the majority of fish species are sensitive to changes in the aquatic soundscape, and depending on the noise source, species responses may have extreme and negative fitness consequences. As such this global synthesis should serve as a warning of the potentially dire consequences facing marine ecosystems if alterations to aquatic soundscapes continue on their current trajectory.

Introduction

The range and intensity of anthropogenic noise in aquatic environments has increased considerably in recent decades, yet little is known about its effects on marine communities (Popper and Hastings, 2009; Popper and Fay, 2011; Simmonds *et al.*, 2014). Specifically, low-frequency ambient noise levels in the open ocean have increased by 3.3 dB per decade since the 1950's, representing a doubling of the noise budget every decade (McDonald *et al.*, 2006; Frisk 2012). Recent research suggests that these increases in noise occur regardless of an area's protection designation. Anthropogenic noise now doubles background sound levels

in over half of all protected units within the continental United States, and 14% of habitats critical to endangered species are exposed to a tenfold increase in sound levels relative to the natural soundscape (Buxton *et al.*, 2017). Although derived from terrestrial parks, this phenomenon is likely more severe in marine protected areas as sound travels five times faster in water than air, therefore, moving greater distances in less time, with less attenuation (Hawkins and Myrberg; 1983; Finfer *et al.*, 2008). Additionally, coastal regions are three times more populated than the global average, resulting in 40% of the global population living within 100 km of the coastline (Cohen *et al.*, 1997; Small & Nicholls 2003). This influx of anthropogenic activity indicates that noise production is centered around coastlines. In all cases, increased noise levels are positively correlated with transportation, development, and resource extraction; all of which are increasing to keep pace with the ever-expanding global economy (Frisk 2012; Buxton *et al.*, 2017).

The majority of sound produced by anthropogenic activities is considered to be noise pollution as it contains little to no intentional information (Popper and Hastings, 2009; Pijanowski *et al.*, 2011). Many fish species have adaptations that have increased their abilities to detect and produce sound, including the roughly 800 fish species from 109 families known to be soniferous, however, these adaptations have also made them susceptible to noise pollution (Kaatz 2002; Popper and Hastings, 2009; Picciulin *et al.*, 2010; Shannon *et al.*, 2016). Investigations into the potential implications of noise on marine life have determined that these disturbances would likely first lead to physiological followed by behavioral changes (Kight and Swaddle, 2011; Gedamke *et al.*, 2016). Depending on the intensity and duration of exposure, noise pollution has the potential to temporarily or permanently alter auditory thresholds, mask the detection of important environmental cues, and lead to increased mortality due to predation (Bass and Clark, 2003; Ladich, 2008; Popper and Hastings, 2009; Picciulin *et al.*, 2010; Simpson *et al.*, 2016). Despite the concerns regarding

increasing anthropogenic noise, the monitoring and regulation of this pollutant has been limited (Simmonds *et al.*, 2014). This is in part due to a lack of understanding the effects of noise on aquatic organisms (Bass and Clark, 2003; Popper and Hastings, 2009).

To determine the implications that disturbances in the aquatic soundscape may have on marine and freshwater fishes, we conducted a meta-analysis addressing the effect of noise on fish behavior and physiology. Meta-analyses are an effective method for assessing ecological trends (Mann, 1990; Côté & Sutherland 1997; Harrison 2011), as they allow for generalizable conclusions to be reached through the utilization of data from multiple sources (Cadotte *et al.*, 2012). Although high-quality syntheses of noise pollution in both the terrestrial and aquatic environments have been conducted (see Popper and Hastings, 2009; Radford *et al.*, 2014; Shannon *et al.*, 2016), these studies have been limited to thorough reviews or vote counting methods that lack statistical power and were therefore unable to quantify the impact to species (Harrison 2011). Through this analysis we reveal the main sources of noise that fishes are exposed to both experimentally and in the wild: anthropogenic noise, biological noise, environmental noise, tones, and music. We also examined how species responses, including foraging ability, movement, responses to predation and reproductive ability, were affected by the various noise sources. This study constitutes the first global quantitative synthesis addressing the effect of noise pollution on fish behavior and physiology.

Methods

Systematic Literature Search

Thompson's Web of Science was used to conduct a systematic literature search. Search results were limited to peer-reviewed articles published between 1950 and 2015. The specific search terms were 'fish' and 'noise or sound or acoustic or ecoacoustics or

bioacoustics’ and ‘behav* or physiol* or response or morphology’, which returned 2,817 potentially relevant peer-reviewed articles. An additional 526 potentially relevant peer-reviewed articles were retrieved through other search engines including ScienceDirect and JSTOR, and by thoroughly reviewing bibliographies of relevant reviews. The titles and abstracts of the 3,343 studies were reviewed to determine which papers addressed the effects of anthropogenic noise on fish behavior or physiology (Fig. 1; Moher *et al.*, 2009). Articles that met these criteria (452) were then further evaluated to identify those that met the criteria of: original research, behavior or physiology focus, listed sound source, experimental control, included mean value, listed standard error or standard deviation, and the sample size stated.

Forty-two studies from 11 countries met the search criteria (Fig. S1). Thirty-six of the studies were conducted within a range of laboratory settings, and six of the studies were conducted in situ. We then extracted the sample size, mean, and standard deviation of the treatment and control groups from each study. All data were obtained from tables and text when possible; if necessary, the reliable and accurate extraction software GraphClick was used to retrieve data from figures (Boyle *et al.*, 2013). A total of 2,354 data points were collected from the 42 studies (Fig. S2).

Effect Size Calculation

The Metafor package was used in R-studio to calculate the effect sizes and variances for each study (Viechtbauer, 2010; R Core Team, 2013). Mean difference (md) was calculated using Eq. (1), where \bar{Y}_1 and \bar{Y}_2 are the mean values of the treatment and control group, respectively.

$$md = \bar{Y}_1 - \bar{Y}_2 \quad (1)$$

The standardized mean difference (Hedge's d), which is an indication of the overall effect and weights of studies based on their sample sizes and standard deviations, was determined using Eq. (2). Sample sizes are indicated by n_1 and n_2 with standard deviations s_1 and s_2 .

$$d = \frac{\bar{Y}_1 - \bar{Y}_2}{\sqrt{\frac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{n_1+n_2-2}}} \quad (2)$$

The variance for Hedge's d was determined via Eq. (3) (Hedges and Olkin, 1985).

$$V_d = \frac{n_1+n_2}{n_1n_2} + \frac{d^2}{2(n_1+n_2)} \quad (3)$$

To account for the large amount of dissimilarity present within the response variables, the directionality of each study was determined to ensure that negative and positive effect sizes represented negative and positive responses, respectively. For example, for a response variable such as growth rate, an increase would result in a positive effect size, while an increase in a response variable such as cortisol (a common stress hormone) would also lead to a positive effect size, despite being an undesirable response. Accounting for the directionality of each response variable is thus a critical step for a meta-analysis of this nature.

Statistical Analyses

Separate analyses were conducted on studies addressing behavioral and physiological responses, which were further split according to sound source: anthropogenic noise, environmental noise, tones, music, and biological noise. We considered biological noise both a positive and negative source, *e.g.* mating calls vs. aggressive conspecifics calls, therefore studies examining positive and negative biological noises were split accordingly and analyzed separately. Furthermore, in rare cases when studies evaluated the effect of both

behavior and physiology or multiple noise sources, each case was treated independently as effect sizes for each sound source were never pooled during the analysis.

To determine which aspects of species behavior and physiology would be affected by the various sound sources, the response variables for each study were grouped into the following behavioral categories: foraging, movement, predation avoidance, reproduction, social interactions, and vocalization; and the following physiological categories: auditory system, blood chemistry, body condition, body size and growth, fatty acid, immune system, neurotransmitters, organ health, reproduction, and stress (Table S3 & S4).

The analysis was conducted in R-studio (R Core Team, 2013; R Studio, 2015). Forest and funnel plots were generated for each of the sound sources within the behavioral and physiological categories using the ‘metafor’ and ‘MAd’ packages to determine the summary effect and confidence intervals of each model, and establish if any potential publication bias was present (Viechtbauer, 2010; Del Re and Hoyt 2014). Separate linear mixed effect models were used to determine which of the behavioral and physiological categories effect sizes differed from the ‘zero’ (no effect) line. A separate linear mixed effect model was run for each of the sound sources within the behavioral and physiological categories, which used response category as a fixed factor and study as a random effect. The resulting model value and standard deviations were then plotted as a visual indication of how each response variable was affected by the various sound sources.

Results

Effect of Noise on Behavior and Physiology

Forest plots indicated that behavioral and physiological responses varied strongly based on sound source. Anthropogenic noise had a significantly negative effect on fish behavior and physiology, as studies addressing behavioral responses yielded an overall effect

size (ES) of -4.73 with 95% confidence intervals (CI) of -8.75 and -0.70, while studies addressing physiological responses had an ES of -1.35 with 95% CI of -2.07 and -0.63 (Fig. 2a; Fig. 3a). These results indicate that there is a negative impact of 4.73 and 1.35 standard deviations to behavior and physiology when fish are exposed to anthropogenic noise, compared to control groups. Tones also had a negative (but not significant) impact on behavior and physiology with overall ES's of -10.5 (CI -23.43, 2.42) and -6.75 (CI -15.62, 2.12), respectively (Fig. 2b; Fig. 3b). Biological noise that was deemed to be adverse, had a negative, but non-significant effect on behavior (ES -0.61, CI -1.16, 0.07; Fig. 2c), while positive biological noise had no effect on fish behavior (ES 0.08, CI -0.41, 0.57; Fig. 2d). A single study addressed the effect of biological noise on fish physiology, and indicated that noises from positive biological sources do not affect fish physiology (ES 0.61, CI -0.17, 1.40; Fig. 3c). Fish behavior was not affected by environmental noise (ES 0.01, CI -0.35, 0.37; Fig. 2e), while fish physiology was negatively affected (ES -0.67, CI -1.29, 0.06; Fig. 3d). No studies addressing how fish respond behaviorally to music were conducted, however, studies addressing the effect of music on fish physiology indicated that music has a positive, although not significant, effect on fish physiology (ES 41.36, CI -39.60, 122.32; Fig. 3e). Funnel plots indicated that there was a minor but acceptable amount of publication bias within studies focusing on fish behavior and physiology (data not shown).

Behavioral Responses to Aquatic Noises

Anthropogenic noise increased movement (mixed-effects model: Estimate=18.52, DF= 155, t-value= 3.53, and P= 0.00) and reproduction (mixed-effects model: Estimate= 17.78, DF= 155, t-value= 3.09, and P= 0.002) related behaviors, which included swimming depth, directional changes, schooling adjustments, swimming speed, and the time parents spent caring for their nests (Fig. 4a; Table S5). In contrast, anthropogenic noise decreased predation responses, including starting responses, time until caught and responses to predatory strikes (mixed-effects model: Estimate= -9.02,

DF= 155, t-value= -1.69, and P= 0.092), as well as social interactions *e.g.*, encounters won and number of social interactions (mixed-effects model: Estimate= -7.34, DF= 150, t-value= -1.23, and P= 0.220), although both decreases were not significant. Foraging behaviors, however, significantly decreased due to anthropogenic noise (mixed-effects model: Estimate= -12.17, DF= 155, t-value= -2.24, and P= 0.027), indicating that the proportion and number of food items consumed, foraging efficiency, and discrimination error, were all negatively affected by anthropogenic noise. Substantial variation existed in the vocalization response to anthropogenic noise, with both positive and negative responses with no unanimous direction observed (mixed-effects model: Estimate= 0.81, DF= 10, t-value= 0.05, and P= 0.960).

Tones significantly increased foraging (mixed-effects model: Estimate= 15.98, DF= 51, t-value= 2.27, and P= 0.03), movements (mixed-effects model: Estimate= 17.94, DF= 51, t-value= 2.78, and P= 0.01), and predation responses, (mixed-effects model: Estimate= 21.01, DF= 51, t-value= 2.69, and P= 0.01). These increases were associated with activities that included food handling error, discrimination error, swimming speed, and startle responses (Fig. 4b; Table S5). As with anthropogenic noise, species vocalization responses to tones showed a large amount of variation, with no clear directionality (mixed-effects model: Estimate= 0.23, DF= 4, t-value= 0.03, and P= 0.98).

Negative biological noise induces significant increases in behaviors related to foraging (mixed-effects model: Estimate= 1.33, DF= 319, t-value= 3.55, and P= 0.001), movement (mixed-effects model: Estimate= 0.28, DF= 319, t-value= 5.4, and P= 0.000), and social interactions (mixed-effects model: Estimate= 1.21, DF= 2, t-value= 1.86, and P= 0.204), while positive biological noise had no impact on species vocalizations (mixed-effects model: Estimate= -0.1, DF= 4, t-value= -0.23, and P= 0.833; Fig. 4c,d; Table S5). Specifically, exposure to negative biological noise increased species bite rate, the amount of time they spent isolated, and their aggression.

Neither movement nor vocalization-related behaviors increased or decreased when exposed to environmental noise (mixed-effects model: Estimate= -0.07, DF= 18, t-value= -0.53, and P= 0.600; mixed-effects model: Estimate= 0.22, DF= 18, t-value= 0.85, and P= 0.408; Fig. 4e; Table S5)

Physiological Responses to Aquatic Noises

Anthropogenic noise caused significant increases in the auditory system (mixed-effects model: Estimate= 2.63, DF= 18, t-value= 4.11, and P= 0.001), and stress levels (mixed-effects model: Estimate= 1.72, DF= 991, t-value= 2.11, and P= 0.035) indicating that hearing threshold (including the inability to hear conspecifics) and cortisol levels increase when exposed to anthropogenic noise (Fig. 5a; Table S6).

Neither reproduction nor neurotransmitters showed a significant response to positive biological noise (mixed-effects model: Estimate= -2.43, DF= 2, t-value= -1.83, and P= 0.209; mixed-effects model: Estimate= -0.01, DF= 2, t-value= -0.01, and P= 0.995; Fig. 5b).

Environmental noise led to positive response within the auditory system, which was predominately due to increases in species hearing thresholds when exposed to environmental noise (mixed-effects model: Estimate= 1.02, DF= 2, t-value= 5.16, and P= 0.036). However, none of the other areas tested, which included blood chemistry, body condition, size and growth, immune system, organ health, reproduction, and stress exhibited significant positive or negative responses to environmental noise (Fig. 5c; Table S6).

Exposure to music caused decreases in response variables associated with blood chemistry, body condition, size and growth, fat stores, and organ health, and increased neurotransmitters, although all of these alterations were subject to large amounts of variation and none of them were significant (Fig. 5d; Table S6).

Tones significantly increased body condition, measured as carcass moisture and condition factor (mixed-effects model: Estimate= 95.63, DF= 214, t-value= 8.29, and P= 0.000). No responses to tones were observed within any of the other physiological responses tested (Fig. 5e; Table S6).

Discussion

Given growing concerns surrounding species' responses to the ever-increasing bombardment of noise in aquatic environments, there is a pressing need for innovative ways to evaluate this issue (Popper and Hastings, 2009; Popper and Fay, 2011). Unfortunately, understanding how species respond to noise pollution is greatly obstructed by a lack of concrete knowledge (Hawkins *et al.*, 2015). To further our understanding of fish responses to aquatic noise we analyzed 42 peer-reviewed studies from 11 countries, focusing on the impact of noise on fish behavior or physiology. The studies included freshwater, estuarine and marine species and were conducted in variety of field and laboratory settings. The resulting 2,354 data points were used to develop models summarizing trends across studies as well as specific responses.

Recent research suggests that all stages of species' life histories may be negatively affected by anthropogenic noise (see Popper and Hastings, 2009; Radford et al 2014; Shannon *et al.*, 2016). Our findings support this trend as anthropogenic noise was shown to increase movement, nest care, hearing thresholds and stress levels, while decreasing foraging-related behaviors. Increases in movement were attributed to directional changes and alterations to swimming behaviors, the majority of which appeared to be responses to perceived predation and were energetically costly (Fraser and Gilliam 1992). Increased parental care also carries additional costs with implications for both parents and offspring, as nest care is a strenuous and time-consuming activity that can exhaust certain species, especially those with complex mating strategies, to the point of death (Williams, 1966; Alonso-Alvarez and Velando 2012; Bose et al., 2015). Increased hearing thresholds and cortisol levels were associated with an increase in stress-related hormones, and suggest that anthropogenic noise has the potential to cause both short and long-term physiological effects. As the potential consequences of hearing threshold increases are well documented within mammals, if fishes respond in a similar fashion then it is likely that the development of the auditory cortical will be severely impaired (Fay and Popper, 2000; Chang and Merzenich 2003). Recovery from threshold shifts will vary according to the stimulus as well as the auditory sensitivity of the affected species (Clark 1991). However, alterations to a species ability to forage and move through its environment have the potential to

impede many species at all life history stages (Slabbekoorn *et al.*, 2010). Our results indicate that across a wide range of experimental and natural exposure conditions, anthropogenic noise can have adverse effects on fish behavior and physiology, which are not limited to specific responses or species. These findings quantify Popper and Hastings' (2009) concerns surrounding the potential implications of anthropogenic noise on fishes, and advance the notion that this issue is in urgent need of further attention and regulation.

In general, similar to anthropogenic noise, exposure to pure tones appeared to have a negative, but weaker, effect on fish behavior and physiology. The lack of statistical significance in these results is potentially a function of a low sample size in the studies summarized here, as tones have a long history of eliciting adverse physiological responses in fishes (see Popper and Clarke 1976), especially in relation to species' hearing thresholds (Slabbekoorn *et al.*, 2010; Kight and Swaddle 2011). In unique cases, tones increase auditory thresholds at a wider range of frequencies than anthropogenic noise (Scholik and Yan 2001), indicating that, much like anthropogenic noise, tones have the potential to negatively affect fish behavior and physiology across a wide range of species and conditions. Results of mixed effect models indicated that tones caused increased foraging, movement, and predation related responses. Furthermore, tones caused an increase in body condition-related response variables; the majority of which were attributed to increases in carcass moisture. Alterations to movement, foraging, and predation responses suggest that species exposed to tones will be more likely to startle, and experience discrimination or handling errors when feeding. Species exposed to tones will likely alter their swimming depth and speed, increase time spent in isolation, and potentially modify schooling shape or direction. These alterations in activity levels support and possibly explain the mechanisms behind fishes fleeing from seismic exploration. Fishes within these areas become more susceptible to predation, and school less effectively, resulting in decreased fitness and potentially size, which in turn, drastically reduces trawler and longline catch rates within surrounding waters (Hirst and Rodhouse 2000; Wardle *et al.*, 2001). Although untested to date, if decreasing catch rates are any indication, aquatic soundscapes may play more of a role in shaping fisheries and recent declines than currently assumed.

Unfortunately, investigations into how biological noise affects fishes are extremely rare and require extensive further inquiry. Additionally, the necessity to split these studies based on the nature of the sounds further reduces the likelihood of detecting overarching trends. However, despite this current lack of existing research, results generally supported our predictions. As expected, negative biological noises were determined to have an adverse effect on fish behavior, whereas studies investigating positive biological noise all resulted in positive responses, but these responses failed to elicit a significant effect size. This is likely due to the low number of studies considered, as it has been suggested and well supported that species, especially those with anatomical modifications for sound projection or reception, can recognize and potentially evaluate sounds related to biological interactions (Fay and Popper 2000; Horne 2008). The lack of research into this topic is understandable given the difficulty associated with replicating natural soundscapes under laboratory conditions; mostly due to issues surrounding the balance between sound pressure and particle velocity within a closed environment (Akamatsu *et al.*, 2002). As such, there is a dire need to address this issue and conduct more in situ research to increase the acoustic validity of testing environments (Slabbekoorn, 2016).

Our results unanimously showed that fish physiology is negatively affected by environmental noise. However, as only a single study addressed how environmental noise affects fish behavior, results surrounding behavioral responses were inconclusive and not considered indicative of a trend. Much like anthropogenic noise, environmental noise caused an increase in species' hearing thresholds. Although hearing impairment is an obvious response to extreme environmental noises (Kight and Swaddle 2011), the effect that hearing loss will have on species' life history is not as evident since studies investigating the long-term consequences of hearing loss are rare, and inferring potential consequences is essential. Over 800 species from 109 families are known to produce sounds (Kaatz 2002; Rountree *et*

al., 2006), with distinct variation between sounds existing at the species, population and even gender level (Kihlsinger and Klimley 2002; Ueng *et al.*, 2007). The inability to hear and thus communicate with conspecifics could impact critical interactions such as agonistic communication, courtship, and spawning (Myrberg, 1997; McKibben and Bass 1998; Aalbers, 2008) and have dire fitness consequences (Brumm and Slabbekoorn, 2005; Wysocki and Ladich 2005). Given that anthropogenic noise elicited similar but more predominant hearing shifts, species living within areas subjected to intense anthropogenic and environmental noise, like exposed coastlines along shipping routes, are likely bombarded with acoustic stimuli to the point that the previously listed consequences become extremely probable.

The finding that music may positively affect fish physiology, although potentially comical to some, does represent a potential point of interest. These studies occurred within aquatic facilities, where mechanical noise can produce a stressful environment (Papoutsoglou *et al.*, 2008). One of the observed effects was that classical music increased dopamine levels in fish. Dopamine neuropathways are essential for positive reinforcement learning and occur across all animal phyla, including vertebrates (Barron *et al.*, 2010). This result suggests that classical music may be interpreted as a positive stimulation and could be used as an environmental enrichment tool to improve animal welfare, which in turn can result in improved growth rates (Newberry, 1995) for fish hatcheries.

It is worth noting that under these experimental conditions, a lack of significance does not necessarily indicate that these areas of species behavior or physiology will not be affected by noise pollution. As meta-analyses draw from the existing literature, unbalanced samples sizes and insufficient data are inevitable and represent areas in need of further study. This is evident in the area of study concerning environmental noise, music, and biological noise, which are currently data insufficient and require further research. However, as these findings indicate that multiple fish species respond similarly to noise across a wide range of experimental conditions, our study aligns with

current reviews on the topic (see Popper and Hastings, 2009; Radford et al 2014; Shannon *et al.*, 2016), but is the first to do so in a robust quantitative fashion through the use of meta-analysis.

These findings also illuminate an ongoing debate within soundscape research: the study of species responses to aquatic noise in laboratory conditions. Some have concluded that sound fields within tanks will never match those of natural aquatic environments (Akamatsu *et al.*, 2002). Species, however, may be adapted to detect sound under a wider variety of conditions than previously considered. For example, as natural habitats are far more structurally complex than the experimental designs used in these studies, it is reasonable to assume that species whose hearing capabilities and sound production evolved within these habitats are well suited to detect noise despite distortion. Although currently untested, noise distortion under experimental conditions may not be magnitudes more severe than that of noise propagating through a kelp forest, coral reef or other complex habitats. Undoubtedly, a large portion of the information embedded within the noise has the potential to be lost in an experimental setting, however, this loss may not differ drastically from natural conditions and thus the extent to which studies conducted in imperfect sound fields should be discredited remains a topic in need of further inquiry.

It is now well-recognized that to regulate, monitor and predict the effect that aquatic noise has on fishes, increasing our knowledge of marine soundscapes is essential (Hawkins *et al.*, 2015), especially as aquatic soundscapes continue to be subjected to anthropogenic activity. A major limitation of this effort is the lack of understanding of how sound affects marine organisms (Bass and Clark, 2003; Popper and Hastings, 2009; Simmonds *et al.*, 2014). Our findings suggest that aquatic noise, depending largely on the source, has the potential to disturb species' ability to interact with conspecifics, forage efficiently, produce viable offspring, and school effectively, while inducing potentially reversible hearing loss. These findings are likely not limited to fishes; although a thorough synthesis is still lacking, recent research suggests that invertebrates may be just as susceptible to noise pollution. Solan and colleagues (2016) determined that invertebrate species, even those that do not

communicate via acoustics, can exhibit a suite of behavioral responses when exposed to shipping and construction-related noise. These behavioral responses can ultimately alter how species mediate key ecosystem processes, possibly undermining the ecosystem services provided by benthic invertebrates.

Our study illustrates how the ever increasing, potentially unstoppable, bombardment of aquatic noise may negatively affect the majority of marine species. However, it is important to remember that fish and other marine organisms have not evolved in a quiet environment (Slabbekoorn *et al.*, 2008). Although most of the responses observed in this synthesis are associated with an inability to function properly, not all responses resulted in permanent damage (Bass and Clark, 2003; Ladich, 2008; Popper and Hastings, 2009; Picciulin *et al.*, 2010). This result suggests that despite unregulated noise pollution in marine soundscapes having the potential to drastically affect organisms, there is also the option of regulation and recovery within these systems. As anthropogenic noise is easily the most negative of the sound sources that exists within any body of water, and noise pollution is increasing exponentially, this global synthesis should serve as a warning and echo the growing concerns regarding the potentially dire consequences for marine ecosystems if noise continues to climb in an unregulated manner.

Acknowledgments

This research was supported by the Canada Research Chairs Program, Natural Sciences and Engineering Research Council of Canada, the Liber Ero Foundation, Canadian Healthy Oceans Network, the Canada Foundation for Innovation and the British Columbia Knowledge Development Fund. We would like to thank the Hakai Institute, the Juanes Lab, and the University of Victoria for their continued support. We would also like to thank Hailey Davies for assistance in revising this manuscript. Furthermore, we would like to thank Arthur Popper and Anthony Hawkins for providing valuable feedback on the oral version of this work, and their lifelong commitment to the study of soundscape ecology.

References

- Aalbers SA (2008) Seasonal, diel, and lunar spawning periodicities and associated sound production of white seabass (*Atractoscion nobilis*). *Fishery Bulletin*, **106**, 143–151.
- Akamatsu T, Okumura T, Novarini N, Yan HY (2002) Empirical refinements applicable to the recording of fish sounds in small tanks. *The Journal of the Acoustical Society of America*, **112**, 3073–3082.
- Alonso-Alvarez C, Velando A. (2012). Benefits and costs of parental care. *The evolution of parental care*, 40-61.
- Amoser S, Ladich F (2003) Diversity in noise-induced temporary hearing loss in otophysine fishes. *The Journal of the Acoustical Society of America*, **113**, 2170–2179.
- Anderson PA, Berzins IK, Fogarty F, Hamlin HJ, Guillette LJ (2011) Sound, stress, and seahorses: The consequences of a noisy environment to animal health. *Aquaculture*, **311**, 129–138.
- Andersson MH, Dock-Akerman E, Ubral-Hedenberg R, Ohman MC (2007) Swimming behavior of roach (*Rutilus rutilus*) and three-spined stickleback (*Gasterosteus aculeatus*) in response to wind power noise and single-tone frequencies. *Ambio*, **36**, 636–638.
- Barron AB, Søvik E, Cornish JL (2010) The Roles of Dopamine and Related Compounds in Reward-Seeking Behavior Across Animal Phyla. *Frontiers in Behavioral Neuroscience*, **4**, 1–9.
- Bass AH, Clark CW (2003) The Physical Acoustics of Underwater Sound Communication. In: *Acoustic Communication* (eds Simmons AM, Fay RR, Popper AN), pp. 15–64. Springer New York, New York, NY.
- Bertucci F, Beauchaud M, Attia J, Mathevon N (2010) Sounds modulate males' aggressiveness in a cichlid fish. *Ethology*, **116**, 1179–1188.

- Bose APH, McClelland GB, Balshine S (2015) Cannibalism, competition, and costly care in the plainfin midshipman fish, *Porichthys notatus*. *Behavioral Ecology*, **27**, 628–636.
- Boyle MA, Samaha AL, Rodewald AM, Hoffmann AN (2013) Evaluation of the reliability and validity of GraphClick as a data extraction program. *Computers in Human Behavior*, **29**, 1023–1027.
- Brumm H, Slabbekoorn H (2005) Acoustic Communication in Noise. *Advances in the Study of Behavior*, **35**, 151–209.
- Buscaino G, Filiciotto F, Buffa G *et al.* (2010) Impact of an acoustic stimulus on the motility and blood parameters of European sea bass (*Dicentrarchus labrax L.*) and gilthead sea bream (*Sparus aurata L.*). *Marine Environmental Research*, **69**, 136–142.
- Buxton RT, McKenna MF, Mennitt D, Fristrup K, Crooks K, Angeloni L, Wittemyer G (2017) Noise pollution is pervasive in U.S. protected areas. *Science*, **356**, 531–533.
- Cadotte MW, Mehrkens LR, Menge DNL (2012) Gauging the impact of meta-analysis on ecology. *Evolutionary Ecology*, **26**, 1153–1167.
- Catli T, Yildirim O, Turker A (2015) The Effect of Different Tempos of Music During Feeding, on Growth Performance, Chemical Body Composition, and Feed Utilization of Turbot (*Psetta maeotica*, Pallas 1814). *The Israeli Journal of Aquaculture-Bamidgeh*, **67**.
- Chang EF, Merzenich MM (2003) Environmental noise retards auditory cortical development. *Science*, **300**, 498–502.
- Clark WW (1991) Recent studies of temporary threshold shift (TTS) and permanent threshold shift (PTS) in animals. *Journal of the Acoustical Society of America*, **90**, 155–163.
- Codarin A, Wysocki LE, Ladich F, Picciulin M (2009) Effects of ambient and boat noise on hearing and communication in three fish species living in a marine protected area (Miramare, Italy). *Marine Pollution Bulletin*, **58**, 1880–1887.

Cohen JE, Small C, Mellinger A, Gallup J, Sachs J, Vitousek PM, Mooney HA (1997)

Estimates of Coastal Populations. *Science*, **278**, 1211–1212.

Côté IM, Sutherland WJ (1997) The Effectiveness of Removing Predators on Bird

Populations. *Conservation Biology*, **11**, 395–405.

Crovo JA, Mendonça MT, Holt DE, Johnston CE (2015) Stress and auditory responses of the

otophysan fish, *Cyprinella venusta*, to road traffic noise. *PLoS ONE*, **10**, e0137290.

Del Re AC, Hoyt WT. (2014) MAd: Meta-analysis with mean differences. R package version 0.8-2.

Finfer DC, Leighton TG, White PR (2008) Issues relating to the use of a 61.5 dB conversion

factor when comparing airborne and underwater anthropogenic noise levels. *Applied*

Acoustics, **69**, 464–471.

Flammang MK, Weber MJ, Thul MD (2014) Laboratory Evaluation of a Bioacoustic Bubble

Strobe Light Barrier for Reducing Walleye Escapement. *North American Journal of*

Fisheries Management, **34**, 1047–1054.

Fraser DF, Gilliam J. (1992) Nonlethal impacts of predation invasion: facultative suppression

of growth and reproduction. *Ecology*, **73**, 959–970.

Frisk G V. (2012) Noiseconomics: The relationship between ambient noise levels in the sea

and global economic trends. *Scientific Reports*, **2**, 437.

Gedamke J, Ferguson M, Harrison J *et al.* (2016) Predicting anthropogenic noise

contributions to US waters. In: *Advances in Experimental Medicine and Biology*, Vol.

875 (eds Popper AN, Hawkins A), pp. 341–347. Springer New York, New York, NY.

Halvorsen MB, Wysocki LE, Stehr CM, Baldwin DH, Chicoine DR, Scholz NL, Popper AN

(2009) Barging Effects on Sensory Systems of Chinook Salmon Smolts. *Transactions of the American Fisheries Society*, **138**, 777–789.

- Halvorsen MB, Zeddies DG, Ellison WT, Chicoine DR, Popper AN (2012) Effects of mid-frequency active sonar on hearing in fish. *The Journal of the Acoustical Society of America*, **131**, 599–607.
- Harrison F (2011) Getting started with meta-analysis. *Methods in Ecology and Evolution*, **2**, 1–10.
- Hawkins, AD, Myrberg AA. (1983). Hearing and sound communication under water. *Bioacoustics, a comparative approach*. Academic Press, London, 347-405.
- Hawkins AD, Pembroke AE, Popper AN (2015) Information gaps in understanding the effects of noise on fishes and invertebrates. *Reviews in Fish Biology and Fisheries*, **25**, 39–64.
- Hedges L V., Olkin I (1988) Statistical Methods for Meta-Analysis. *Journal of Educational Statistics*, **13**, 75.
- Hirst AG, Rodhouse PG (2000) Impacts of geophysical seismic surveying on fishing success. *Reviews in Fish Biology and Fisheries*, **10**, 113–118.
- Holt DE, Johnston CE (2014) Evidence of the Lombard effect in fishes. *Behavioral Ecology*, **25**, 819–826.
- Horne JK (2008) Acoustic ontogeny of a teleost. *Journal of Fish Biology*, **73**, 1444–1463.
- Isabella-Valenzi L, Higgs DM (2013) Sex- and state-dependent attraction of round gobies, *Neogobius melanostomus*, to conspecific calls. *Behaviour*, **150**, 1509–1530.
- Jordão JM, Fonseca PJ, Amorim MCP (2012) Chorusing Behaviour in the Lusitanian Toadfish: Should I Match My Neighbours' Calling Rate? *Ethology*, **118**, 885–895.
- Kaatz IM (2002) Multiple sound-producing mechanisms in teleost fishes and hypotheses regarding their behavioural significance. *Bioacoustics*, **12**, 230–233.
- Kight CR, Swaddle JP (2011) How and why environmental noise impacts animals: An integrative, mechanistic review. *Ecology Letters*, **14**, 1052–1061.

- Kihlsinger RL, Klimley AP (2002) Species identity and the temporal characteristics of fish acoustic signals. *Journal of Comparative Psychology*, **116**, 210–214.
- Ladich F (2008) Sound communication in fishes and the influence of ambient and anthropogenic noise. *Bioacoustics*, **17**, 34–38.
- Leong H, Ros AFH, Oliveira RF (2009) Effects of putative stressors in public aquaria on locomotor activity, metabolic rate and cortisol levels in the Mozambique tilapia *Oreochromis mossambicus*. *Journal of Fish Biology*, **74**, 1549–1561.
- Liu M, Wei QW, Du H, Fu ZY, Chen QC (2013) Ship noise-induced temporary hearing threshold shift in the Chinese sucker *Myxocyprinus asiaticus* (Bleeker, 1864). *Journal of Applied Ichthyology*, **29**, 1416–1422.
- Mann C. (1990). Meta-analysis in the breech. *Science*, 249(4968), 476–481.
- McCauley RD, Fewtrell J, Popper AN (2003) High intensity anthropogenic sound damages fish ears. *The Journal of the Acoustical Society of America*, **113**, 638–642.
- McDonald MA, Hildebrand JA, Wiggins SM (2006) Increases in deep ocean ambient noise in the Northeast Pacific west of San Nicolas Island, California. *The Journal of the Acoustical Society of America*, **120**, 711–718.
- McKibben JR, Bass AH (1998) Behavioral assessment of acoustic parameters relevant to signal recognition and preference in a vocal fish. *The Journal of the Acoustical Society of America*, **104**, 3520–3533.
- Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009) Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine*, **6**.
- Moore A, Waring CP (1999) Reproductive priming in mature male Atlantic salmon parr exposed to the sound of redd cutting. *Journal of Fish Biology*, **55**, 884–887.

- Myrberg JR AA (1997) Sound production by a coral reef fish (*Pomacentrus partitus*): evidence for a vocal, territorial “keep-out” signal. *Bulletin of Marine Sciences*, **60**(3), 1017–1025.
- Nedelec SL, Simpson SD, Morley EL, Nedelec B, Radford AN (2015) Impacts of regular and random noise on the behaviour, growth and development of larval Atlantic cod (*Gadus morhua*). *Proceedings of the Royal Society B*, **282**, 20151943.
- Newberry RC (1995) Environmental enrichment: Increasing the biological relevance of captive environments. *Applied Animal Behaviour Science*, **44**, 229–243.
- Nichols TA, Anderson TW, Širović A (2015) Intermittent noise induces physiological stress in a coastal marine fish. *PLoS ONE*, **10**, 1–13.
- Papoutsoglou SE, Karakatsouli N, Batzina A, Papoutsoglou ES, Tsopelakos A (2008) Effect of music stimulus on gilthead seabream *Sparus aurata* physiology under different light intensity in a re-circulating water system. *Journal of Fish Biology*, **73**, 980–1004.
- Papoutsoglou SE, Karakatsouli N, Louizos E *et al.* (2007) Effect of Mozart’s music (Romanze-Andante of “Eine Kleine Nacht Musik”, sol major, K525) stimulus on common carp (*Cyprinus carpio* L.) physiology under different light conditions. *Aquacultural Engineering*, **36**, 61–72.
- Papoutsoglou SE, Karakatsouli N, Papoutsoglou ES, Vasilikos G (2010) Common carp (*Cyprinus carpio*) response to two pieces of music (“Eine Kleine Nachtmusik” and “Romanza”) combined with light intensity, using recirculating water system. *Fish Physiology and Biochemistry*, **36**, 539–554.
- Papoutsoglou SE, Karakatsouli N, Psarrou A *et al.* (2015) Gilthead seabream (*Sparus aurata*) response to three music stimuli (Mozart—“Eine Kleine Nachtmusik,” Anonymous—“Romanza,” Bach—“Violin Concerto No. 1”) and white noise under recirculating water conditions. *Fish Physiology and Biochemistry*, **41**, 219–232.

Papoutsoglou SE, Karakatsouli N, Skouradakis C, Papoutsoglou ES, Batzina A, Leondaritis

G, Sakellaridis N (2013) Effect of musical stimuli and white noise on rainbow trout

(*Oncorhynchus mykiss*) growth and physiology in recirculating water conditions.

Aquacultural Engineering, **55**, 16–22.

Pereira R, Rismondo S, Caiano M, Pedroso SS, Fonseca PJ, Amorim MCP (2014) The role of

agonistic sounds in male nest defence in the painted goby *Pomatoschistus pictus*.

Ethology, **120**, 53–63.

Picciulin M, Sebastianutto L, Codarin A, Calcagno G, Ferrero EA (2012) Brown meagre

vocalization rate increases during repetitive boat noise exposures: A possible case of

vocal compensation. *The Journal of the Acoustical Society of America*, **132**, 3118–

3124.

Picciulin M, Sebastianutto L, Codarin A, Farina A, Ferrero EA (2010) In situ behavioural

responses to boat noise exposure of *Gobius cruentatus* (Gmelin, 1789; fam. Gobiidae)

and *Chromis chromis* (Linnaeus, 1758; fam. Pomacentridae) living in a Marine

Protected Area. *Journal of Experimental Marine Biology and Ecology*, **386**, 125–132.

Pijanowski BC, Villanueva-Rivera LJ, Dumyahn SL *et al.* (2011) Soundscape Ecology: The

Science of Sound in the Landscape. *BioScience*, **61**, 203–216.

Popper AN, Clarke NL (1976) The auditory system of the goldfish (*Carassius auratus*):

Effects of intense acoustic stimulation. *Comparative Biochemistry and Physiology*

Part A: Physiology, **53**, 11–18.

Popper AN, Hastings MC (2009) The effects of anthropogenic sources of sound on fishes.

Journal of Fish Biology, **75**, 455–489.

Popper AN, Fay RR (2011) Rethinking sound detection by fishes. *Hearing Research*, **273**,

25–36.

Popper AN, Smith ME, Cott PA, Hanna BW, MacGillivray AO, Austin ME, Mann DA

(2005) Effects of exposure to seismic airgun use on hearing of three fish species. *The Journal of the Acoustical Society of America*, **117**, 3958–3971.

Purser J, Radford AN (2011) Acoustic noise induces attention shifts and reduces foraging performance in three-spined sticklebacks (*Gasterosteus aculeatus*). *PLoS ONE*, **6**.

R Core Team (2013) R: A Language and Environment for Statistical Computing. Version 3.0.1. R Foundation for Statistical Computing, Vienna, Austria. Available at: www.r-project.org (accessed 10 April 2015).

R Studio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA. Available at: <http://www.rstudio.com/> (accessed 10 April 2015).

Radford AN, Kerridge E, Simpson SD (2014) Acoustic communication in a noisy world: Can fish compete with anthropogenic noise? *Behavioral Ecology*, **25**, 1022–1030.

Rountree RA, Gilmore G, Goudey CA, Hawkins AD, Luczkovich JJ, David A. Mann (2006) Listening to Fish. *Fisheries*, **31**, 433–446.

Sarà G, Dean JM, D'Amato D *et al.* (2007) Effect of boat noise on the behaviour of bluefin tuna *Thunnus thynnus* in the Mediterranean Sea. *Marine Ecology Progress Series*, **331**, 243–253.

Scholik AR, Yan HY (2001) Effects of underwater noise on auditory sensitivity of a cyprinid fish. *Hearing Research*, **152**, 17–24.

Scholik AR, Yan HY (2002) Effects of noise on auditory sensitivity of fishes. *Bioacoustics*, **12**, 186–188.

Sebastianutto L, Picciulin M, Costantini M, Ferrero EA (2011) How boat noise affects an ecologically crucial behaviour: The case of territoriality in *Gobius cruentatus* (Gobiidae). *Environmental Biology of Fishes*, **92**, 207–215.

- Shafiei Sabet S, Neo YY, Slabbekoorn H (2015) The effect of temporal variation in sound exposure on swimming and foraging behaviour of captive zebrafish. *Animal Behaviour*, **107**, 49–60.
- Shannon G, McKenna MF, Angeloni LM *et al.* (2016) A synthesis of two decades of research documenting the effects of noise on wildlife. *Biological Reviews*, **91**, 982–1005.
- Sierra-Flores R, Attack T, Migaud H, Davie A (2015) Stress response to anthropogenic noise in Atlantic cod *Gadus morhua* L. *Aquacultural Engineering*, **67**, 67–76.
- Simmonds MP, Dolman SJ, Jasny M, Parsons EC., Weilgart L, Wright AJ, Leaper R (2014) Marine noise pollution-increasing recognition but need for more practical action. *Journal of Ocean Technology*, **9**, 71–90.
- Simpson SD, Purser J, Radford AN (2015) Anthropogenic noise compromises antipredator behaviour in European eels. *Global Change Biology*, **21**, 586–593.
- Simpson SD, Radford AN, Nedelec SL, Ferrari MCO, Chivers DP, McCormick MI, Meekan MG (2016) Anthropogenic noise increases fish mortality by predation. *Nature Communications*, **7**, 10544.
- Slabbekoorn H. (2016). Aiming for progress in understanding underwater noise impact on fish: complementary need for indoor and outdoor studies. In *The effects of noise on aquatic life II* (pp. 1057-1065). Springer New York.
- Slabbekoorn H, Bouton N, van Opzeeland I, Coers A, ten Cate C, Popper AN (2010) A noisy spring: The impact of globally rising underwater sound levels on fish. *Trends in Ecology and Evolution*, **25**, 419–427.
- Slabbekoorn H, Ripmeester EAP (2008) Birdsong and anthropogenic noise: Implications and applications for conservation. *Molecular Ecology*, **17**, 72–83.
- Small C, Robert J. Nichols (2003) A Global Analysis of Human Settlement in Coastal Zones. *Journal of Coastal Research*, **19**, 584–599.

- Smith ME, Kane AS, Popper AN (2004) Noise-induced stress response and hearing loss in goldfish (*Carassius auratus*). *Journal of Experimental Biology*, **207**, 427–435.
- Solan M, Hauton C, Godbold JA, Wood CL, Leighton TG, White P (2016) Anthropogenic sources of underwater sound can modify how sediment-dwelling invertebrates mediate ecosystem properties. *Scientific Reports*, **6**, 20540.
- Ueng JP, Huang B-Q, Mok HK (2007) Sexual differences in the spawning sounds of the Japanese croaker, *Argyrosomus japonicus* (Sciaenidae). *Zoological Studies*, **46**, 103–110.
- Vasconcelos RO, Amorim MCP, Ladich F (2007) Effects of ship noise on the detectability of communication signals in the Lusitanian toadfish. *Journal of Experimental Biology*, **210**, 2104–2112.
- Viechtbauer W (2010) Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software*, **36**, 1–48.
- Voellmy IK, Purser J, Flynn D, Kennedy P, Simpson SD, Radford AN (2014) Acoustic noise reduces foraging success in two sympatric fish species via different mechanisms. *Animal Behaviour*, **89**, 191–198.
- Wardle CS, Carter TJ, Urquhart GG, Johnstone ADF, Ziolkowski AM, Hampson G, Mackie D (2001) Effects of seismic air guns on marine fish. *Continental Shelf Research*, **21**, 1005–1027.
- Williams GC (1966) Natural Selection, the costs of reproduction, and a refinement of lack's principle. *The American Naturalist*, **100**, 687–690.
- Wilson B, Dill LM (2002) Pacific herring respond to simulated odontocete echolocation sounds. *Canadian Journal of Fisheries and Aquatic Sciences*, **59**, 542–553.
- Wysocki LE, Dittami JP, Ladrich F (2006) Ship noise and cortisol secretion in European freshwater fishes. *Biological Conservation*, **128**, 501–508.

Wysocki LE, Ladich F (2005) Hearing in fishes under noise conditions. *Journal of the Association for Research in Otolaryngology*, **6**, 28–36.

Figure Captions


Figure 1. A PRISMA diagram outlining the selection processes of papers for the meta-analysis on the effect of aquatic noise on fish behavior and physiology.


Figure 2. Forest plots illustrating how various aquatic noises affect fish behavior. Studies were divided into the following categories based on noise source: a) Anthropogenic Noise b) Tones c) Negative Biological Noise d) Positive Biological Noise e) Environmental Noise. Author(s) and publication year are listed within each plot.

Figure 3. Forest plots illustrating how various aquatic noises affect fish physiology. Studies were divided into the following categories base on noise source: a) Anthropogenic Noise b) Tones c) Positive Biological Noise d) Environmental e) Music. Author(s) and publication year are listed within each plot.


Figure 4: Fish behavioral responses to aquatic noise, derived via separate mixed effect models evaluating fish responses to each noise source: a) Anthropogenic Noise b) Tone c) Positive Biological Noise d) Negative Biological Noise e) Environmental Noise. Horizontal lines indicate the 0 effect size or ‘no change’ line.

Figure 5: Fish physiology responses to aquatic noise derived via separate mixed effect models evaluating fish responses to each noise source: a) Anthropogenic Noise b) Positive Biological Noise c) Environmental Noise d) Music e) Tone. Horizontal lines indicate the 0 effect size or ‘no change’ line.







(a)




(b))



(c)



(d)



(e)

