

## Short communication

## Species-specific wet-dry mass calibrations for dominant Northeastern Pacific Ocean macroalgae and seagrass

Sara B. Wickham<sup>a,b,\*</sup>, Chris T. Darimont<sup>b,c,d</sup>, John D. Reynolds<sup>b,e</sup>, Brian M. Starzomski<sup>a,b</sup><sup>a</sup> School of Environmental Studies, University of Victoria, P.O. Box 1700, STN CSC, Victoria, British Columbia V8W 2Y2 Canada<sup>b</sup> Hakai Institute, P.O. Box 309, Heriot Bay, British Columbia, V0P 1H0, Canada<sup>c</sup> Department of Geography, University of Victoria, P.O. Box 3060 STN CSC, Victoria, British Columbia, V8W 3R4, Canada<sup>d</sup> Raincoast Conservation Foundation, P.O. Box 2429, Sidney, British Columbia, V8L 3Y3, Canada<sup>e</sup> Earth to Ocean Research Group, Department of Biological Sciences, 8888 University Drive, Simon Fraser University, Burnaby, BC V5A 1S6, Canada

## ARTICLE INFO

## Keywords:

Biomass  
Mass calibration  
Wet weight  
Dry weight  
Ratio  
Macroalgae  
Macrophyte  
Sea wrack  
Seagrass  
Kelp  
Northeast Pacific Ocean

## ABSTRACT

Macroalgae and seagrasses form the base of productive ecosystems in the Northeastern Pacific Ocean. Often, ecological research on macroalgae, seagrasses, and sea wrack requires the conversion of biomass from wet to dry to create consistency across investigations. This process, however, can be destructive, impractical, time consuming, and labour intensive. We collected samples of twelve common Northeastern Pacific Ocean seaweed species (*Alaria marginata*, *Codium fragile*, *Egregia menziesii*, *Fucus distichus*, *Macrocystis pyrifera*, *Mazzaella* spp., *Nereocystis luetkeana*, *Pterygophora californica*, *Pyropia* spp., *Ulva* spp., and the seagrasses *Zostera marina* and *Phyllospadix* spp.) in two states: wet and fresh, or aged and partially desiccated. We weighed, dried, and compared samples, finding that many species displayed a strong ( $R^2 > 0.8$ ) and predictable linear relationship between wet and dried conditions. In contrast, half the aged samples did not have a significant relationship between partially desiccated and dried conditions. Our results provide practical wet weight to dry weight ratios for many species, and with further research, a reliable set of species-specific wet to dry weight ratios for all species could be established. Wet to dry weight ratios are useful for macroalgae, seagrass, and sea wrack research or commercial applications and would reduce the need to conduct extensive wet-dry calibrations in each study.

## 1. Introduction

Ecological research on macroalgae, seagrass, and sea wrack (detached, shore-cast macroalgae) has been fundamental to our understanding of ecosystem functioning (Filbee-Dexter and Scheibling, 2014), habitat connectivity (Dugan et al., 2003), community structure (Smale et al., 2016) and species distribution (Smith et al., 2018). Macroalgal research also continues to offer a key understanding of anthropogenic stressors to oceanic communities (Johnson et al., 2011; Brodie et al., 2014; Filbee-Dexter and Wernberg, 2018) and the ecosystem services provided by macroalgae (Krause-Jensen and Duarte, 2016). Additionally, the demand for macroalgae in commercial products (such as nutraceuticals and foods) has intensified and is predicted to rise (Dhargalkar and Pereira, 2005; Jiménez-Escrig et al., 2012). Obtaining accurate estimates of the biomass of macroalgae, seagrasses, and sea wrack is central to their study. However, some biomass estimation methods are destructive and can have long term ecological

consequences (Watt and Scrosati, 2013). Therefore, non-destructive methods of accurate and efficient biomass estimation of macroalgae, seagrass, and sea wrack can provide a valuable contribution to future research.

In the Northeastern Pacific Ocean, sub-tidal and intertidal biomass is dominated by various species of macroalgae (hereafter referred to as seaweeds) and seagrass. Kelps (Order Laminariales) are among the fastest growing algae in the world, and form vast and ecologically important underwater forests (Duggins et al., 1989; Steneck et al., 2002). Others, like eelgrass (*Zostera* spp.) comprise extensive mats across the shallow subtidal that provide key nursery habitats for species of ecological, cultural, and economic value, such as salmon (*Oncorhynchus* spp.) (Simenstad et al., 1982) and rockfish (*Sebastes* spp.) (Olson, 2017). Although work on seaweeds and sea wrack has well-established methods for sampling and surveying techniques and equipment in other parts of the world (e.g., Stagnol et al., 2016), a gap in knowledge remains in understanding the relationship between wet and dry biomass

\* Corresponding author. Present address: School of Environment, Resources and Sustainability, University of Waterloo, 200 University Avenue West, Waterloo, Ontario, N2L 3G1, Canada.

E-mail address: [swickham@uvic.ca](mailto:swickham@uvic.ca) (S.B. Wickham).

<https://doi.org/10.1016/j.aquabot.2018.09.006>

Received 11 October 2017; Received in revised form 11 September 2018; Accepted 16 September 2018

Available online 20 September 2018

0304-3770/ © 2018 Elsevier B.V. All rights reserved.

for seaweed species commonly found in the Northeast Pacific Ocean. Establishing wet-dry calibrations for dominant seaweeds of this large region would facilitate simplified future research on these ecologically and economically important species.

Dry mass measurement is considered the most reliable method for precise biomass estimates of seaweeds (Stagnol et al., 2016). This method enables consistent results across investigations (Bickel and Perrett, 2015). Dry mass can also account for variation in water content of different seaweed species. Additionally, it has the potential to account for the variation in desiccation levels of sea wrack deposits along shorelines. When feasible, seaweeds are sorted by species, functional group, or a subsample of a mixture of several seaweed species is oven dried to a constant weight at which no further mass loss is recorded (Madsen, 1993). After dry mass determination, the most commonly used approach is to regress (e.g. Orr et al., 2005; Fox et al., 2014) or use standard major axis testing (e.g. Gevaert et al., 2001, 2008; Stagnol et al., 2016) to determine the relationship between wet and dry mass. The slope is then used to create a wet weight to dry weight ratio (DW:WW) (e.g. Gevaert et al., 2001; Orr et al., 2005).

Obtaining dry mass measurements is labour intensive, time consuming, and sometimes problematic. For instance, in field studies with large amounts of macrophyte biomass, dry mass measurements may not be feasible. Performing wet-dry calibration work in remote field locations without electrical power can be impractical. Additionally, one common genus of marine macroalgae (*Desmarestia* spp.) contains sulphuric acid in its tissue that will rapidly disintegrate itself and any co-occurring specimens after collection (Druehl and Clarkston, 2016). A potential problem in beach wrack dry mass measurements is that many studies determine their dry mass estimates based on a subsample of a mixture of species, not a species-specific subsample (e.g. Mews et al., 2006; Orr et al., 2005). Lastly, dry mass determination is destructive and will render the organism unsuitable for further research and may remove large amounts of seaweeds from natural settings, which can disturb ecological processes (Dugan et al., 2003).

Despite the substantial amount of information available regarding seaweed survey and sampling techniques available from locations across the globe, we found no information on standardized regression coefficients obtained from dry mass determination for the twelve common seaweed species we sampled specific to the Northeastern Pacific Ocean. Therefore, each study that requires dry seaweed biomass measurements for these species must perform the time-consuming and labour-intensive step of dry mass determination, or resort to destructive biomass measurements. In this paper, we present the results of wet-dry calibrations for twelve common macroalgae and seagrass species collected from the Northeastern Pacific. By sharing these results, and with further research, we can provide a suite of standardized wet to dry weight ratios to use in estimating the dry biomass of macroalgae, seagrass, and wrack in western North America.

## 2. Methods

We collected 252 macroalgae and seagrass specimens from three sandy beaches and two rocky intertidal outcrops located along the northwest end of Calvert Island, British Columbia, Canada (51.6536°N, 128.1301°W) in July (n = 223) and September (n = 29) 2016. Samples contained twelve species: *Alaria marginata*, *Codium fragile*, *Egregia menziesii*, *Fucus distichus*, *Macrocystis pyrifera*, *Mazzaella* spp., *Nereocystis luetkeana*, *Pterygophora californica*, *Pyropia* spp., *Ulva* spp. (bladed form), and the seagrasses *Zostera marina* and *Phyllospadix* spp. Samples of various weights (27–265 g) were collected indiscriminately from shoreline wrack accumulations (n = 134) and harvested live from rocky intertidal shorelines (n = 60). Collected samples were sorted into two categories based on their moisture saturation levels: aged or wet (see Supplementary Table 1 for details of the number of samples in the aged and wet categories collected during each period). Aged samples were collected from wrack deposited in lines parallel to the water's

edge that had been stagnant on the beach for over 24 h, confirmed visually during beach surveys over a six-day period. Aged samples were collected opportunistically from these wrack lines and contained a subset of the studied species: *A. marginata*, *C. fragile*, *E. menziesii*, *F. distichus*, *M. pyrifera*, *N. luetkeana*, *Phyllospadix* spp., and *Ulva* spp., all of which were partially desiccated. Wet samples were collected opportunistically from samples that had been deposited recently on the shoreline (confirmed visually during surveys) and harvested live from rocky intertidal outcrops at low tide. All wet samples showed no signs of desiccation.

Blades, stipes, and (when applicable) pneumatocysts were collected. Samples were sorted and separated into species and for three kelp species (*M. pyrifera*, *N. luetkeana*, and *P. californica*), samples were processed as “blade only” or “stipe only” categories. However, because most researchers may not separate kelp blades from stipes, we also processed samples that were portions of entire kelp individuals: stipes with attached blades and pneumatocysts (“mixed” category). Samples were rinsed, cleaned free of sand and epiphytes, and shaken to remove surface water. Wet weights of all samples were measured on a Denver Instrument MXX-612 balance and transferred into a Fisher Scientific Isotemp drying oven within one hour of collection. Samples were dried at 80 °C until their dry weight reached a constant mass ( $\pm 0.005$  g) for three consecutive measurements (typically this took about 72 h).

All data were analyzed in R package version 3.3.3 (R Core Team, 2017). We performed standardized major axis tests (as per Warton et al., 2006) and used the slope of the linear equation to generate wet to dry weight ratios (Table 1). Using package ‘smatr’ (Warton et al. 2012) we compared the slopes of wet samples that were harvested live against wet samples that were collected from wrack lines as well as samples that were collected in July against samples that were collected in September. Additionally, we compared slopes for the three dried combinations of samples (blade only, stipe only, and mixed) of the kelps *M. pyrifera*, *N. luetkeana*, and *P. californica*. The ‘smatr’ package fits bivariate lines in allometry using standardized major axis testing to make inferences about the lines.

Using the package ‘lmodel2’ (Legendre, 2014) we plotted the confidence intervals of the wet samples for four species that had potential outliers, low sample size, and heterogeneous data.

## 3. Results

Fifteen of eighteen categories from twelve species of macroalgae and seagrass exhibited strong ( $R^2 > 0.8$ ) and significant linear relationships ( $p$ -values  $< 0.05$ ) between wet and dry mass when dried from wet samples (*C. fragile*, *E. menziesii*, *F. distichus*, *M. pyrifera* mixed, *M. pyrifera* blade, *M. pyrifera* stipe, *Mazzaella* spp., *N. luetkeana* blade,

**Table 1**

Wet to dry weight ratios (WW:DW) for the nine species across twelve categories of seaweeds that displayed strong ( $R^2 > 0.8$ ) and significant ( $p$ -value  $< 0.05$ ) relationships when dried from wet. DW stands for dry weight and WW stands for wet weight. Species with outlier driven relationships were not included.

Species	WW:DW
<i>Codium fragile</i>	DW = 0.069WW
<i>Egregia menziesii</i>	DW = 0.189WW
<i>Fucus distichus</i>	DW = 0.195WW
<i>Macrocystis pyrifera</i> mixed	DW = 0.115WW
<i>Macrocystis pyrifera</i> blade	DW = 0.168WW
<i>Macrocystis pyrifera</i> stipe	DW = 0.121WW
<i>Mazzaella</i> spp.	DW = 0.192WW
<i>Nereocystis luetkeana</i> stipe	DW = 0.045WW
<i>Pterygophora californica</i> mixed	DW = 0.151WW
<i>Pterygophora californica</i> stipe	DW = 0.245WW
<i>Pyropia</i> spp.	DW = 0.307WW
<i>Zostera marina</i>	DW = 0.224WW




Fig. 1. Relationships between wet and dry mass for the eighteen categories of twelve species of marine macroalgae and seagrass commonly found in the Northeastern Pacific Ocean. All samples were collected fresh from recently deposited wrack lines ( $n = 134$ ) or harvested live ( $n = 60$ ). The species-specific wet to dry weight ratio is the slope of the linear equation, displayed in the top right corner of each plot.

*N. leutkeana* stipe, *P. californica* mixed, *P. californica* blade, *P. californica* stipe, *Pyropia* spp., *Ulva* spp., *Z. marina*; Fig. 1). Three species categories displayed weak ( $R^2 < 0.8$ ) but significant relationships ( $p$ -value  $< 0.05$ ) between wet and dry mass when dried from wet samples (*A. marginata*, *N. leutkeana* mixed, *Phyllospadix* spp.; Fig. 1). Four species (*C. fragile*, *E. menziesii*, *N. leutkeana* mixed, and *Phyllospadix* spp.) out of eight also exhibited significant linear relationships ( $p$ -value  $< 0.05$ ) between aged and dry mass (Fig. 2).

We found no significant difference between wet samples that were harvested live from rocky outcrops and wet samples that were collected from freshly deposited wrack lines (ANCOVA; all  $p$ -values  $> 0.1$ ). We found no significant differences between samples harvested in July and samples harvested in September (ANCOVA; all  $p$ -values  $> 0.1$ ). For the kelps *M. pyrifera*, *N. leutkeana*, and *P. californica*, we found no significant difference between the slopes of wet and dry weight for blade, stipe, and mixed samples (ANCOVA; all  $p$ -values  $> 0.1$ ).

The majority of collected samples had a wet weight of under 200 g. For some species, there was a good distribution across the weight spectrum. For other species the weight distribution could be characterized by few samples of high weight and aggregations of samples with low weight (Supplementary Material Figs. 1 and 2).

Of the wet samples, for three of the four species (*N. leutkeana* blade, *P. californica* blade, and *Ulva* spp.), that had outliers, low sample size, and heterogeneous data, the slope of the data with the outlier removed was different than the slope of the data with the outlier (Supplementary

Material Fig. 3). Low replication size may be affecting these samples and with further analysis these outliers may fall within the predicted range of these species' wet-to-dry weight ratios. Therefore, we did not remove these data point from the standardized major axis tests used to generate dry to wet weight ratios.

#### 4. Discussion

For several species of marine seaweeds and seagrasses common to the Northeast Pacific, we found that water content lost from the tissues of live collected and shore-cast marine seaweed species generally displayed strong and predictable relationships. We are not aware of any data that display the differences in water content of seaweed species native to the Northeast Pacific Ocean. Therefore, our results may offer a first estimation for wet to dry weight ratios across this range until more local-scale data with a larger sampling effort are available. Using these calibrations, researchers can estimate the dry biomass for six of the twelve macroalgae and seagrasses tested, and for three different parts of two different kelp species (Table 1). Three species (*A. marginata*, *N. leutkeana* mixed, and *Phyllospadix* spp.) display weak relationships when dried from wet samples. Three collections with low samples sizes (*N. leutkeana* blade, *P. californica* blade, and *Ulva* spp.) had relationships impacted by outliers and more data are needed to draw a conclusion on the wet-dry mass relationship of these species. Further analysis should be undertaken to improve our understanding of the wet-dry




Fig. 2. Relationships between aged and dry mass for eight species of marine macroalgae commonly found in the Northeastern Pacific Ocean. All samples were collected from aged wrack lines and had signs of desiccation. The species-specific wet to dry weight ratio is the slope of the linear equation, displayed in the top right corner of each plot. Statistically significant results (*Codium fragile*, *Egregia menziesii*, *Nereocystis leutkeana*, and *Phyllospadix* spp.), are marked by \*.

relationships for these latter six seaweed categories.

The reliable water content loss relationships only existed for the wet seaweeds. Establishing a relationship between aged (partially desiccated) to dry ratios was less dependable. From the aged samples collected along wrack lines, *A. marginata*, *F. distichus*, *M. pyrifera*, and *Ulva* spp. did not display significant water content loss relationships. This could be due to variation in how desiccated individual samples were at time of collection and a lack of precision in recording moisture saturation categories. In the future, it would be helpful to develop several more specific categories of desiccation in sea wrack. These categories could be based on the time the seaweeds were deposited along a shoreline, inferred from tidal information.

For the kelps we dried three categories of wet samples: blade only, stipe only, and blade and stipe together (mixed). For all kelp species tested the difference between blade, stipe, and mixed categories had no significant effect on the relationship between wet and dry weight. This indicates that the blades and stipes of these kelps have similar potential for water content loss and the wet to dry weight ratios for 'mixed' samples can be applied. However, if possible, separating the blades from the stipes would yield more precise estimates (Supplementary Table 2) as the relationships displayed by the blade only and stipe only categories were stronger than those of the mixed category (Fig. 1). Additionally, other research has displayed significantly different water content values for differing parts of kelp (Smale et al., 2016).

Our sampling occurred during two periods of the year, once in July and once in early September. Research on several species of brown seaweeds (Phaeophyceae) in the North Atlantic ocean found water

content differences between summer and winter (Black, 1948; Gevaert et al., 2001, 2008; Wernberg et al., 2001), which could be the result of differences in carbohydrate amounts throughout different seasons (Manns et al., 2017). Results from our study indicated no significant water content differences in samples harvested at these two different time periods. As both collection dates fell within the summer growing season for the Northeastern Pacific Ocean (Druehl and Clarkston, 2016), perhaps the seaweed species we collected had not undergone significant changes in carbohydrate levels. However, to achieve the most accurate wet to dry weight ratios it would be important to perform further tests on seasonal changes of tissue water content for all common Northeastern Pacific Ocean seaweed species, as many industries harvest seaweeds and sea wrack outside of the summer growing season.

Each species demonstrated its own particular wet-dry mass relationship. This variability could be due to variation among species in chemical composition. Research worldwide has investigated the chemical compositions of marketable substances within seaweeds (e.g., mannitol, ash, alginate, and carrageenan) (Schiener et al., 2014; Ismail et al., 2017). Overall, each seaweed species has a different chemical profile and, as discussed above, the composition of these chemical profiles can differ between seasons (Marinho-Soriano et al., 2006; Manns et al., 2017). Kelps such as *M. pyrifera*, *N. leutkeana*, and *P. californica* contain high amounts of water soluble alginate, which leaches quickly when then kelp is out of water and increases mass loss (Mews et al., 2006). The seagrasses *Z. marina* and *Phyllospadix* spp. contain higher amounts of water insoluble lignin and cellulose, which do not leach upon drying and may contribute to relatively decreased amounts



of mass loss in these species (Mews et al., 2006). These well documented chemical differences between seaweed species should encourage investigators to apply species-specific wet to dry weight ratios to their research.

A potential weakness of this study is the low sample size of seaweeds collected per species. In other studies it is not uncommon to collect upwards of 100 samples per species over the course of one year (Gevaert et al., 2001; Wernberg et al., 2001). The relationships obtained with low sample sizes may be affected by factors such as the difference between seaweed categories (i.e. mixed versus blade versus stipe) and time of year (as discussed above). To confirm the utility of the wet to dry weight ratios derived from this research, and to further investigate seasonal, morphological, and wrack water content differences, more research into the biometrics of Northeastern Pacific Ocean seaweed species should be pursued and shared.

Undoubtedly future research will continue to investigate the many aspects of marine, aquatic, and shore-cast macroalgae and seagrass ecology. Whereas considerable effort has gone into developing effective sampling techniques, prior to this research little information existed on wet-dry mass relationships for individual seaweed species of the Northeast Pacific Ocean. Our research shows promising evidence of strong wet-dry biomass relationships across six common seaweeds and two species of kelp within three different sampling categories. These calculated relationships enable estimations of tissue water content loss within seaweeds and may be useful to the seaweed industry and ecological researchers, reducing the destructive, time consuming, and potentially unnecessary step of collecting and drying seaweed samples.

## Acknowledgements

We thank colleagues from the Heiltsuk and Wuikinuxv Nations for the opportunity to perform this research. We deeply appreciate the field, laboratory, and funding support from the Hakai Institute and the Tula Foundation, which made this paper possible. Thank you also to Beatrice Proudfoot for field and laboratory work and to Natalie Ban, Katy Hind, and three anonymous reviewers for helpful edits. BMS thanks the Wallace Stegner House and the Eastend (SK) Arts Council for writing support.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi: <https://doi.org/10.1016/j.aquabot.2018.09.006>.

## References

- Bickel, T.O., Perrett, C., 2015. Precise determination of aquatic plant wet mass using a salad spinner. *Can. J. Fish. Aquat. Sci.* 73, 1–4.
- Black, W.A.P., 1948. The seasonal variation in chemical composition of some of the littoral seaweeds common to Scotland. Part II. *Laminaria digitata*. *J. Soc. Chem. Ind.* 67, 169–172.
- Brodie, J., Williamson, C.J., Smale, D.A., Kamenos, N.A., Mieszkowska, N., Santos, R., Cunliffe, M., Steinke, M., Yesson, C., Anderson, K.M., Asnaghi, V., Brownlee, C., Burdett, H.L., Burrows, M.T., Collins, S., Donohue, P.J.C., Harvey, B., Foggo, A., Noisette, F., Nunes, J., Ragazzola, F., Raven, J.A., Schmidt, D.N., Suggett, D., Teichberg, M., Hall-Spencer, J.M., 2014. The future of the northeast Atlantic benthic flora in a high CO<sub>2</sub> world. *Ecol. Evol.* 4, 2787–2798.
- Dhargalkar, V.K., Pereira, N., 2005. Seaweed: promising plant of the millennium. *Sci. Cult.* 71, 60–66.
- Druehl, L.D., Clarkston, B.E., 2016. *Pacific Seaweeds: a Guide to Common Seaweeds of the West Coast*. Harbour Publishing, Madeira Park, BC.
- Dugan, J.E., Hubbard, D.M., McCrary, M.D., Pierson, M.O., 2003. The response of macrofauna communities and shorebirds to macrophyte wrack subsidies on exposed sandy beaches of southern California. *Estuarine Coast. Shelf Sci.* 58S, 25–40.
- Duggins, D.O., Simeanstad, C.A., Estes, J.A., 1989. Magnification of secondary production by kelp detritus in coastal marine ecosystems. *Science (New York N.Y.)* 245, 170–173.
- Filbee-Dexter, K., Scheibling, R.E., 2014. Sea urchin barrens as alternative stable states of collapsed kelp ecosystems. *Mar. Ecol. Prog. Ser.* 495, 1–25.
- Filbee-Dexter, K., Wernberg, T., 2018. Rise of turfs: a new battlefield for globally declining kelp forests. *BioScience* 1–13.
- Fox, C.H., El-Sabaawi, R., Paquet, P.C., Reimchen, T.E., 2014. Pacific herring *Clupea pallasii* and wrack macrophytes subsidize semi-terrestrial detritivores. *Mar. Ecol. Prog. Ser.* 495, 49–64.
- Gevaert, F., Davoult, D., Creach, a., Kling, R., Janquin, M.-a., Seuront, L., Lemoine, Y., 2001. Carbon and nitrogen content of *Laminaria saccharina* in the eastern english channel: biometrics and seasonal variations. *J. Mar. Biol. Assoc. U. K.* 81, 727.
- Gevaert, F., Janquin, M.A., Davoult, D., 2008. Biometrics in *Laminaria digitata*: a useful tool to assess biomass, carbon and nitrogen contents. *J. Sea Res.* 60, 215–219.
- Ismail, M.M., El Zokm, G.M., El-Sayed, A.A.M., 2017. Variation in biochemical constituents and master elements in common seaweeds from Alexandria Coast, Egypt, with special reference to their antioxidant activity and potential food uses: prospective equations. *Environ. Monit. Assess.* 189, 175–648.
- Jiménez-Escrig, A., Gómez-Ordóñez, E., Rupérez, P., 2012. Brown and red seaweeds as potential sources of antioxidant nutraceuticals. *J. Appl. Phycol.* 24, 1123–1132.
- Johnson, C.R., Banks, S.C., Barrett, N.S., Cazassus, F., Dunstan, P.K., Edgar, G.J., Frusher, S.D., Gardner, C., Haddon, M., Helidoniotis, F., Hill, K.L., Holbrook, N.J., Hosie, G.W., Last, P.R., Ling, S.D., Melbourne-Thomas, J., Miller, K., Pecl, G.T., Richardson, A.J., Ridgway, K.R., Rintoul, S.R., Ritz, D.A., Ross, D.J., Sanderson, J.C., Shepherd, S.A., Slotwinski, A., Swadling, K.M., Taw, N., 2011. Climate change cascades: shifts in oceanography, species' ranges and subtidal marine community dynamics in eastern Tasmania. *J. Exp. Mar. Biol. Ecol.* 400, 17–32.
- Krause-Jensen, D., Duarte, C.M., 2016. Substantial role of macroalgae in marine carbon sequestration. *Nat. Geosci.* 9, 737–742.
- Legendre, P., 2014. *lmodel2: Model II Regression*.
- Madsen, J.D., 1993. Biomass techniques for monitoring and assessing control of aquatic vegetation. *Lake Reserv. Manag.* 7, 141–154.
- Manns, D., Nielsen, M.M., Bruhn, A., Saake, B., Meyer, A.S., 2017. Compositional variations of brown seaweeds *Laminaria digitata* and *Saccharina latissima* in Danish waters. *J. Appl. Phycol.* 29, 1493–1506.
- Marinho-Soriano, E., Fonseca, P.C., Carneiro, M.A.A., Moreira, W.S.C., 2006. Seasonal variation in the chemical composition of two tropical seaweeds. *Bioresour. Technol.* 97, 2402–2406.
- Mews, M., Zimmer, M., Jelinski, D.E., 2006. Species-specific decomposition rates of beach-cast wrack in barkley sound, British Columbia, Canada. *Mar. Ecol. Prog. Ser.* 328, 155–160.
- Olson, A., 2017. *Seagrass Meadows as Seascape Nurseries for Rockfish (Sebastes Spp.)*. MSc Thesis. University of Victoria, Victoria, BC.
- Orr, M., Zimmer, M., Jelinski, D.E., Mews, M., 2005. Wrack deposition on different beach types: spatial and temporal variation in the pattern of subsidy. *Ecology* 86, 1496–1507.
- Schiener, P., Black, K.D., Stanley, M.S., Green, D.H., 2014. The seasonal variation in the chemical composition of the kelp species *Laminaria digitata*, *Laminaria hyperborea*, *Saccharina latissima* and *Alaria esculenta*. *J. Appl. Phycol.* 27, 363–373.
- Simenstad, C.A., Fresh, K.L., Salo, E.O., 1982. The role of puget Sound and Washington coastal estuaries in the life history of pacific salmon: an unappreciated function. Page In: Kennedy, V.S. (Ed.), *Estuarine Comparisons*, first edition. Academic Press, New York.
- Smale, D.A., Burrows, M.T., Evans, A.J., King, N., Sayer, M.D.J., Yunnice, A.L.E., Moore, P.J., 2016. Linking environmental variables with regionalscale variability in ecological structure and standing stock of carbon within UK kelp forests. *Mar. Ecol. Prog. Ser.* 542, 79–95.
- Smith, T.M., York, P.H., Broitman, B.R., Thiel, M., Hays, G.C., van Sebille, E., Putman, N.F., Macreadie, P.I., Sherman, C.D.H., 2018. Rare long-distance dispersal of a marine angiosperm across the Pacific Ocean. *Glob. Ecol. Biogeogr.* 28, 487–496.
- Stagnol, D., Macé, M., Destombe, C., Davoult, D., 2016. Allometric relationships for intertidal macroalgae species of commercial interest. *J. Appl. Phycol.* 28, 3407–3411.
- Steneck, R.S., Graham, M.H., Bourque, B.J., Corbett, D., Erlandson, J.M., Estes, J.A., Tegner, M.J., 2002. Kelp forest ecosystems: biodiversity, stability, resilience and future. *Environ. Conserv.* 29, 436–459.
- R.Core Team, 2017. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Warton, D.I., Wright, I.J., Falster, D.S., Westoby, M., 2006. Bivariate line-fitting methods for allometry. *Biol. Rev. Camb. Philos. Soc.* 81, 259–291.
- Warton, D. I., R. A. Duursma, D. S. Falster, and S. Taskinen. 2012. *smatr 3- an R package for estimation and inference about allometric lines*. *Methods in Ecology and Evolution* 3:257–259.
- Watt, C.A., Scrosati, R.A., 2013. Regional consistency of intertidal elevation as a mediator of seaweed canopy effects on benthic species richness, diversity, and composition. *Mar. Ecol. Prog. Ser.* 491, 91–99.
- Wernberg, T., Thomsen, M.S., Stæhr, P.A., Pedersen, M.F., 2001. Comparative phenology of *Sargassum muticum* and *Halidrys siliquosa* (Phaeophyceae: fucales) in Limfjorden, Denmark. *Bot. Marina* 44, 31–39.