

Fundamentals of Information Theory

Homework 2

Problem 1 (10 points) Suppose you're on a game show, and you've given the choice of three doors: Behind one door is a car; behind the others, goat. You pick a door, say No. 1, and the host, who knows what's behind the doors, opens another door, say No. 3, which has a goat. He then says to you, "Do you want to pick door No. 2?" Is it to your advantage to switch your choice so that you have a higher chance to win the car instead of the goat?

Problem 2 (15 points) One is given 24 coins. It is known that precisely one coin is fake, which weights differently compared with genuine coins. It is not clear whether this fake coin is heavier or lighter, though. Now we use a balance to identify the fake coin, but we do not have the weights for this balance.

- (a) What is the minimum number of the weighting operations in order to identify this fake coin.
- (b) Explain briefly about your weighting procedure to identify this fake coin.

Problem 3 (10 points)

Let $p(x, y)$ be given in Table 1, i.e., $p(X = 0, Y = 1) = 0$.

Table 1: $p(x, y)$

$X \backslash Y$	0	1
0	$1/3$	0
1	$1/3$	$1/3$

Find.

- (a) $H(X)$, $H(Y)$.
- (b) $H(X|Y)$, $H(Y|X)$.
- (c) $H(X, Y)$.
- (d) $H(Y) - H(Y|X)$.
- (e) $I(X; Y)$.
- (f) Draw a Venn diagram for the quantities in (a) through (e).

Problem 4 (10 points) Let the random variable X have three possible outcomes $\{a, b, c\}$. Consider two distributions on this random variable:

Symbol	$p(x)$	$q(x)$
a	$1/2$	$1/3$
b	$1/4$	$1/3$
c	$1/4$	$1/3$

Calculate $H(p(x))$, $H(q(x))$, $D(p(x)||q(x))$, and $D(q(x)||p(x))$. Verify that in this case, $D(p(x)||q(x)) \neq D(q(x)||p(x))$.

Problem 5 (10 points)

Consider a non-uniform random variable X with $M = 8$ possible outcomes and probabilities $(1/2, 1/4, 1/8, 1/16, 1/64, 1/64, 1/64, 1/64)$. We use one symbol to represent one outcome, so we have 8 types of symbols (A, B, C, D, E, F, G, H) . Then we transmit 1024 symbols from point s to point d . In this 1024-symbol sequence, it turns out that the numbers of these 8 type symbols are $(516, 255, 126, 68, 17, 18, 10, 14)$, respectively

- (a) Compute the entropy $H(X)$ in bits.
- (b) Consider a coding scheme c_1 , $A \rightarrow 000$, $B \rightarrow 001$, $C \rightarrow 010$, $D \rightarrow 011$, $E \rightarrow 100$, $F \rightarrow 101$, $G \rightarrow 110$, $H \rightarrow 111$, to transmit these 1024 symbols. Compute the total bits using scheme c_1 .
- (c) Similarly, consider another coding scheme c_2 , $A \rightarrow 0$, $B \rightarrow 10$, $C \rightarrow 110$, $D \rightarrow 1110$, $E \rightarrow 11110$, $F \rightarrow 111110$, $G \rightarrow 1111110$, $H \rightarrow 11111110$, to transmit these 1024 symbols. Compute the total bits using scheme c_2 . On average, how many bits are used to transmit one symbol in this transmitted sequence?

Problem 6 (10 points) Let X_1 and X_2 be identically distributed but not necessarily independent. Let

$$\rho = 1 - \frac{H(X_2|X_1)}{H(X_1)}.$$

- (a) Show that $\rho = \frac{I(X_1; X_2)}{H(X_1)}$.
- (b) Show that $0 \leq \rho \leq 1$.
- (c) When is $\rho = 0$?
- (d) When is $\rho = 1$?

Problem 7 (10 points)

Let X , Y and Z be joint random variables. Prove the following inequalities and find conditions for equality.

- (a) $H(X, Y|Z) \geq H(X|Z)$.
- (b) $I(X, Y; Z) \geq I(X; Z)$.
- (c) $H(X, Y, Z) - H(X, Y) \leq H(X, Z) - H(X)$.
- (d) $I(X; Z|Y) \geq I(Z; Y|X) - I(Z; Y) + I(X; Z)$.