

Chap 4 Spectral Analysis of Stochastic Processes

Dong Yan
EI. HUST.

Chapter 4: Spectral Analysis



- References:

1. Stochastic Processes and Their Applications,
Frank E. Beichelt, O211. W138
Chap 8, Spectral analysis of stationary
processes
2. Probability, Random Variables and Random
Signal Principles, 2nd Edition, Peyton Z. Peebles,
O211 W58/2
Chap 7, Spectral characteristic of random
processes

Chapter 4: Spectral Analysis



Content:

4.1 Spectral Density Functions

4.2 Spectral Analysis of Linear Systems

4.3 Spectrum of Amplitude-modulated Signals

4.4 Narrow-band Gaussian Processes

4.1 Spectral Density Functions



Content:

4.1.1 Autospectral Density Functions

4.1.2 Wiener-Khintchine Theorem

4.1.3 Crossspectral Density Functions

4.1.4 S.D.F. of Derived Random Processes

Average of the power



- If $X(t)$ is ergodic, then

$$R_{XX}(0) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T x^2(t) dt$$

- $R(0)$ represents the time average of the power of a random process $X(t)$.
- **Def. 1** average of the power

$$\bar{P}_X = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T x^2(t) dt$$

- **Parseval theorem**

$$\int_{-\infty}^{\infty} x^2(t) dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} X^2(\omega) d\omega$$

- The representation of the average energy in terms of time and frequency of a signal.
- Thus,

$$\bar{P}_X = \lim_{T \rightarrow \infty} \frac{1}{4\pi T} \int_{-T}^T X^2(\omega) d\omega$$

4.1.1 Autospectral Density Functions



Def.2 Spectral Density Function (S.D.F.)

- The S.D.F. of a random process $X(t)$ is

$$S_{XX}(\omega) = \lim_{T \rightarrow \infty} \frac{1}{2T} |X_T(\omega)|^2 \quad S_{XX}(f) = \lim_{T \rightarrow \infty} \frac{1}{2T} |X_T(f)|^2$$

- (autospectral density function) (**Power Spectrum**)
- From Def.1 and Def. 2,

$$\begin{aligned} \overline{P}_X &= \lim_{T \rightarrow \infty} \frac{1}{4\pi T} \int_{-T}^T X^2(\omega) d\omega \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} S_{XX}(\omega) d\omega = \frac{1}{\pi} \int_0^{\infty} S_{XX}(\omega) d\omega \\ &= \int_{-\infty}^{\infty} S_{XX}(f) df = 2 \int_0^{\infty} S_{XX}(f) df \end{aligned}$$

Definitions associated with the S.D.F.



1. The k th moment of the S.D.F.

$$m_k = \int_0^{\infty} \omega^k S_{XX}(\omega) d\omega$$

2. Modal frequency ω_m

the frequency where the S.D.F. peaks.

3. Mean frequency

$$\bar{\omega}_X = \frac{\int_0^{\infty} \omega S_{XX}(\omega) d\omega}{\int_0^{\infty} S_{XX}(\omega) d\omega} = \frac{m_1}{m_0}$$

the mean frequency is not equal to the average value of the various frequencies involved in a random process; it is the mathematical mean of the S.D.F.



4. The k th moment about the mean of the S.D.F.

$$\mu_k = \int_0^\infty (\omega - \bar{\omega})^k S_{XX}(\omega) d\omega$$

Thus,

$$\mu_0 = m_0 = \int_0^\infty S_{XX}(\omega) d\omega = \pi \bar{P}_X$$

$$\mu_1 = 0$$

$$\mu_2 = m_2 - \frac{m_1^2}{m_0}$$



5. The bandwidth parameter of the S.D.F.

$$\varepsilon = \sqrt{1 - \frac{m_2^2}{m_0 m_4}}$$

$\varepsilon = 0$, the random process has a narrow-band spectrum;

$\varepsilon = 1$, the random process has a wide-band spectrum;



6. The spectral width parameter of the S.D.F.

$$\nu = \sqrt{\frac{\mu_2}{\mu_0} \frac{1}{\bar{\omega}}}$$

7. The spectral peakedness parameter

$$Q_P = \frac{\int_0^{\infty} \omega S_{XX}^2(\omega) d\omega}{\left(\int_0^{\infty} S_{XX}(\omega) d\omega \right)^2}$$

The value increases with increasing sharpness of the S.D.F.

4.1.1 Autospectral Density Functions



- Properties of the S.D.F.

- i) Nonnegative $S_{XX}(\omega) = \lim_{T \rightarrow \infty} \frac{1}{2T} |X_T(\omega)|^2 \geq 0$

- ii) Real

- iii) Even, if $X(t)$ is real, $S_{XX}(\omega) = S_{XX}(-\omega)$

- iv) If $\int_{-\infty}^{\infty} |R_{XX}(\tau)| d\tau < \infty$, then $S_{XX}(\omega)$ is a continuous function of ω

4.1 Spectral Density Functions



Content:

4.1.1 Autospectral Density Functions

4.1.2 Wiener-Khintchine Theorem

4.1.3 Crossspectral Density Functions

4.1.4 S.D.F. of Derived Random
Processes

4.1.2 Wiener-Khintchine Theorem



- The Fourier transform of the correlation function of an ergodic process:

$$\int_{-\infty}^{\infty} R_{XX}(\tau) e^{-i\omega\tau} d\tau = \lim_{T \rightarrow \infty} \frac{1}{2T} |X_T(\omega)|^2$$

- And the definition of S.D.F.,

$$S_{XX}(\omega) = \lim_{T \rightarrow \infty} \frac{1}{2T} |X_T(\omega)|^2$$

- Obtain,

$$S_{XX}(\omega) = \int_{-\infty}^{\infty} R_{XX}(\tau) e^{-i\omega\tau} d\tau$$

$$R_{XX}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{\infty} S_{XX}(\omega) e^{i\omega\tau} d\omega$$

4.1.2 Wiener-Khintchine Theorem



Theorem: Wiener-Khintchine Theorem

For a **weakly stationary random process** $X(t)$, its correlation function and the power spectrum are Fourier transform pair.

$$S_{XX}(\omega) = \int_{-\infty}^{\infty} R_{XX}(\tau) e^{-i\omega\tau} d\tau$$

$$R_{XX}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{\infty} S_{XX}(\omega) e^{i\omega\tau} d\omega$$

Standard Fourier Transform

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$$

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{i\omega t} d\omega$$

4.1.2 Wiener-Khintchine Theorem



- Both correlation function and power spectrum are **real** and **even** functions.

$$S_{xx}(\omega) = \int_{-\infty}^{\infty} R_{xx}(\tau) e^{-i\omega\tau} d\tau$$

$$= 2 \int_0^{\infty} R_{xx}(\tau) \cos \omega\tau d\tau$$

$$R_{xx}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{\infty} S_{xx}(\omega) e^{i\omega\tau} d\omega$$

$$= \frac{1}{\pi} \int_0^{\infty} S_{xx}(\omega) \cos \omega\tau d\omega$$

4.1.2 Wiener-Khintchine Theorem



- The average power and correlation function

$$\frac{1}{\pi} \int_0^{\infty} S_{XX}(\omega) d\omega = \bar{P}_X = R_{XX}(0) = E[X^2(t)]$$

- If $E[X(t)] = 0$,

$$\text{Var}[X(t)] = \frac{1}{\pi} \int_0^{\infty} S_{XX}(\omega) d\omega = \bar{P}_X = R_{XX}(0)$$

4.1.2 Wiener-Khintchine Theorem



Example: White Noise

The process with autocorrelation function,

$$R(\tau) = A\delta(\tau)$$

and power spectrum,

$$S(\omega) = A = \text{constant}$$

The **continuous white noise** is a real, stationary, continuous-time stochastic process with constant spectral density.

the “most random” stochastic process





4.1.2 Wiener-Khintchine Theorem

Example 2. Band-limited White Noise

$$S(\omega) = A, \text{ where } -\frac{W}{2} < \omega < \frac{W}{2}$$

Thus,

$$R_{XX}(\tau) = \frac{1}{2\pi} \int_{-W/2}^{W/2} A e^{i\omega\tau} d\omega = \frac{A}{\pi\tau} \sin \frac{1}{2} W\tau$$



Review 2.2.1 Stationary Processes



- weakly stationary

Example 1. Random phase processes

$X(t) = A \cos(\omega t + \varepsilon)$, $t > 0$, whereas A and ω are constants and ε is random variable uniformly distributed between $-\pi$ and π .

$$E[x(t)] = 0$$

$$\begin{aligned} C_{XX}(t_1, t_2) &= \frac{A^2}{2} \cos \omega_0 (t_2 - t_1) \\ &= \frac{A^2}{2} \cos \omega_0 \tau = C_{XX}(\tau) \end{aligned}$$



4.1.2 Wiener-Khintchine Theorem



Example 3. Random Phase Processes

$$R_{XX}(\tau) = \frac{A^2}{2} \cos \omega_0 \tau$$


Obtain: $S(\omega) = \frac{A^2 \pi}{2} [\delta(\omega - \omega_0) + \delta(\omega + \omega_0)]$



4.1.2 Wiener-Khintchine Theorem




- Summary



$$\text{Ave. Energy } P_x = \frac{1}{2T} \int_{-T}^T \{x(t)\}^2 dt$$

$$R_{xx}(\tau) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T x(t) x(t + \tau) dt$$

Auto-Correlation Function




(Fourier Transform)

4.1.2 Wiener-Khintchine Theorem



- Summary



4.1.2 Wiener-Khintchine Theorem



- Summary

$R_X(\tau)$	$S_X(\omega)$
$\begin{cases} (T - \tau)/T, & \tau < T, \\ 0, & \text{otherwise} \end{cases}$	$T \{\sin(\omega T/2)/(\omega T/2)\}^2 = T \text{sinc}^2(fT)$
1	$2\pi\delta(\omega)$
$\delta(\tau)$	1
$\exp(-\alpha \tau)$	$2\alpha/(\alpha^2 + \omega^2)$
$\cos(\omega_0\tau)$	$\pi\delta(\omega - \omega_0) + \pi\delta(\omega + \omega_0)$
$\exp(-\alpha \tau) \cos(\omega_0\tau)$	$\alpha/[\alpha^2 + (\omega - \omega_0)^2] + \{\alpha/[\alpha^2 + (\omega + \omega_0)^2]\}$
$2W \text{sinc}(2W\tau) = \sin(2\pi W\tau)/\pi\tau$	$\begin{cases} 1, & \omega \leq 2\pi W \\ 0, & \text{otherwise} \end{cases}$
$\text{sinc}(u) = \sin(\pi u)/\pi u, -\infty < u < \infty$	

4.1.2 Wiener-Khintchine Theorem



e.g. Suppose the continuous-time random process $X(t)$ has autocorrelation function

$$R_X(\tau) = 1 + e^{-\alpha|\tau|}, \quad -\infty < \tau < \infty, \quad \alpha > 0$$

Find the spectral density function for $X(t)$.

$$S_X(\omega) = 2\pi\delta(\omega) + \frac{2\alpha}{\alpha^2 + \omega^2}$$

4.1 Spectral Density Functions



Content:

4.1.1 Autospectral Density Functions

4.1.2 Wiener-Khintchine Theorem

4.1.3 Crossspectral Density
Functions

4.1.4 S.D.F. of Derived Random
Processes

4.1.3 Cross-Spectral Density Functions



1. Definition of Cross-Spectral Density Function (CS.D.F)

Cross-Spectral Density Function of two random processes $X(t)$ and $Y(t)$ is

$$\tau = t_1 - t_2 \quad S_{XY}(\omega) = \lim_{T \rightarrow \infty} \frac{1}{2T} X(\omega) Y^*(\omega)$$

$$S_{YX}(\omega) = \lim_{T \rightarrow \infty} \frac{1}{2T} X^*(\omega) Y(\omega)$$

4.1.3 Cross-Spectral Density Functions



Theorem: Wiener-Khintchine Theorem

For two **jointly stationary random processes** $X(t)$ and $Y(t)$, their cross-correlation function and cross-power density function are Fourier transform pair.

$$S_{XY}(\omega) = \int_{-\infty}^{\infty} R_{XY}(\tau) e^{-i\omega\tau} d\tau \quad S_{YX}(\omega) = \int_{-\infty}^{\infty} R_{YX}(\tau) e^{-i\omega\tau} d\tau$$

$$R_{XY}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{\infty} S_{XY}(\omega) e^{i\omega\tau} d\omega \quad R_{YX}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{\infty} S_{YX}(\omega) e^{i\omega\tau} d\omega$$

Standard Fourier Transform

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$$
$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{i\omega t} d\omega$$

4.1.3 Cross-Spectral Density Functions



2. Real part and imaginary part

$$\begin{aligned} S_{XY}(\omega) &= \int_{-\infty}^{\infty} R_{XY}(\tau) e^{-i\omega\tau} d\tau \\ &= \int_{-\infty}^{\infty} R_{XY}(\tau) \cos \omega\tau d\tau - i \int_{-\infty}^{\infty} R_{XY}(\tau) \sin \omega\tau d\tau \\ &= C_{XY}(\omega) + iQ_{XY}(\omega) \end{aligned}$$

$$\begin{aligned} C_{XY}(\omega) &= \int_{-\infty}^{\infty} R_{XY}(\tau) \cos \omega\tau d\tau \\ &= \int_{-\infty}^0 R_{XY}(\tau) \cos \omega\tau d\tau + \int_0^{\infty} R_{XY}(\tau) \cos \omega\tau d\tau \\ &= \int_0^{\infty} R_{XY}(-\tau) \cos \omega\tau d\tau + \int_0^{\infty} R_{XY}(\tau) \cos \omega\tau d\tau \\ &= \int_0^{\infty} [R_{YX}(\tau) + R_{XY}(\tau)] \cos \omega\tau d\tau \end{aligned}$$

4.1.3 Cross-Spectral Density Functions



Imaginary part

$$\begin{aligned} Q_{YX}(\omega) &= -\int_{-\infty}^{\infty} R_{XY}(\tau) \sin \omega \tau d\tau \\ &= \int_0^{\infty} [-R_{XY}(\tau) + R_{YX}(\tau)] \sin \omega \tau d\tau \end{aligned}$$

Amplitude Spectrum:

$$|S_{XY}(\omega)| = \sqrt{C_{XY}^2(\omega) + Q_{XY}^2(\omega)}$$

Phase Spectrum:

$$\varepsilon_{XY}(\omega) = \tan^{-1} \frac{Q_{XY}(\omega)}{C_{XY}(\omega)}$$

4.1.3 Cross-Spectral Density Functions



3. Properties of Cross-Spectral Density Function

i) $C_{XY}(\omega)$ and $C_{YX}(\omega)$ are even functions

ii) $Q_{XY}(\omega)$ and $Q_{YX}(\omega)$ are odd functions

iii) $S_{XY}(\omega) = S_{YX}(-\omega) = S_{YX}^*(\omega)$

iv) If $X(t)$ and $Y(t)$ are jointly stationary random processes and uncorrelated,

$$S_{XY}(\omega) = S_{YX}(\omega) = 2\pi m_X m_Y \delta(\omega)$$

4.1.3 Cross-Spectral Density Functions



4. Coherency function

$$\gamma_{XY}(\omega) = \frac{|S_{XY}(\omega)|^2}{S_{XX}(\omega)S_{YY}(\omega)} = \frac{C_{XY}^2(\omega) + Q_{XY}^2(\omega)}{S_{XX}(\omega)S_{YY}(\omega)}$$

Mutual correlation coefficients

$$\rho_{XY}[t_1, t_2] = \frac{C_{XY}(t_1, t_2)}{\sqrt{\text{Var}[X(t_1)]\text{Var}[Y(t_2)]}}$$

$$0 \leq \gamma_{XY}(\omega) \leq 1$$

$\gamma_{XY}(\omega) \equiv 1$ if and only if $X(t)$ and $Y(t)$ are exactly linearly related.

4.1.3 Cross-Spectral Density Functions



Example 1.

Given: a cross-spectral density function

$$S_{XY}(\omega) = \begin{cases} a + i \frac{b\omega}{\omega_0} & -\omega_0 < \omega < \omega_0 \\ 0 & \text{otherwise} \end{cases}$$

where $\omega_0 > 0$, a and b are real constants.

Obtain: the cross-correlation function.

4.1 Spectral Density Functions



Content:

4.1.1 Autospectral Density Functions

4.1.2 Wiener-Khintchine Theorem

4.1.3 Crossspectral Density Functions

4.1.4 S.D.F. of Derived Random Processes

4.1.4 S.D.F. of Derived Random Processes



1. Autocorrelation function of derived random processes

$$R_{\dot{x}\dot{x}}(t_1, t_2) = \frac{\partial}{\partial t_2} R_{xx}(t_1, t_2) = -\frac{d}{d\tau} R_{xx}(\tau)$$

$$R_{\ddot{x}\ddot{x}}(\tau) = \frac{\partial^2}{\partial t_1 \partial t_2} R_{xx}(t_1, t_2) = -\frac{d^2}{d\tau^2} R_{xx}(\tau)$$

$$R_{\ddot{x}\ddot{x}}(\tau) = \frac{d^4}{d\tau^4} R_{xx}(\tau)$$

4.1.4 S.D.F. of Derived Random Processes



2. Spectral density function of derived random processes

$$\mathcal{F}\{x(t)\} = X(\omega)$$

$$\mathcal{F}\{\dot{x}(t)\} = \dot{X}(\omega) = i\omega X(\omega)$$

$$\mathcal{F}\{\ddot{x}(t)\} = \ddot{X}(\omega) = -\omega^2 X(\omega)$$

$$S_{\dot{x}\dot{x}}(\omega) = \lim_{T \rightarrow \infty} \frac{1}{2T} |\dot{X}(\omega)|^2$$

$$= \omega^2 \lim_{T \rightarrow \infty} \frac{1}{2T} |X(\omega)|^2 = \omega^2 S_{xx}(\omega)$$

4.1.4 S.D.F. of Derived Random Processes



2. Spectral density function of derived random processes

$$\begin{aligned} S_{\ddot{x}\ddot{x}}(\omega) &= \lim_{T \rightarrow \infty} \frac{1}{2T} |\ddot{X}(\omega)|^2 \\ &= \omega^4 \lim_{T \rightarrow \infty} \frac{1}{2T} |X(\omega)|^2 = \omega^4 S_{xx}(\omega) \end{aligned}$$

Homework

4.1

4.2

4.7