

Extensions of Abelian Automaton Groups

Chris Grossack
(Advisor: Klaus Sutner)

May 8, 2019

Let's Unpack That

Extensions of Abelian Automaton Groups

Let's Unpack That

Extensions of Abelian Automaton Groups

Let's Unpack That

Extensions of **Abelian** Automaton Groups

Let's Unpack That

Extensions of Abelian Automaton Groups

Let's Unpack That

Extensions of Abelian Automaton Groups

1 Automata


2 Abelian

3 Groups

4 Extensions

Finite State Automata


- Combinatorial Objects
- Encode length preserving functions on binary strings
 - ▶ states
 - ▶ transitions



- One function per state
- Evaluate by following edges


Evaluating Functions

- \mathcal{A}_2^3 . This automaton will be our friend for the rest of this talk
- Defines three functions:
 - ▶ f
 - ▶ f_0
 - ▶ f_1




Evaluating Functions

- \mathcal{A}_2^3 . This automaton will be our friend for the rest of this talk
- Defines three functions:
 - ▶ f
 - ▶ f_0
 - ▶ f_1




- How do we compute, say, f of a string?

Evaluating Functions




- $f(011010)$

Evaluating Functions




• $f(011010)$

Evaluating Functions




• $1f_0(11010)$

Evaluating Functions




- $1f_0(11010)$

Evaluating Functions




- $11f(1010)$

Evaluating Functions




- $110f_1(010)$

Evaluating Functions




- $1100f_0(10)$

Evaluating Functions




- $11001f(0)$

Evaluating Functions



- $110011f_0(\varepsilon)$

Evaluating Functions



- 110011

Definition

$\partial_0 f$ (resp. $\partial_1 f$) is the unique function such that for every $w \in 2^*$,
 $f(0w) = (f(0))(\partial_0 f)(w)$

Definition

$\partial_0 f$ (resp. $\partial_1 f$) is the unique function such that for every $w \in 2^*$,
 $f(0w) = (f(0))(\partial_0 f)(w)$

Definition


A state f is called *Odd* if it flips its input bit, and *Even* otherwise.

Definition

$\partial_0 f$ (resp. $\partial_1 f$) is the unique function such that for every $w \in 2^*$,
 $f(0w) = (f(0))(\partial_0 f)(w)$

Definition

A state f is called *Odd* if it flips its input bit, and *Even* otherwise.



- $\partial_0 f = f_0$ and $\partial_1 f = f_1$
- f is odd, f_0 and f_1 are even

Definition

For f and g in an automaton \mathcal{A} , write $f + g$ for the function
$$(f + g)(x) = f(g(x))$$

Definition

For f and g in an automaton \mathcal{A} , write $f + g$ for the function $(f + g)(x) = f(g(x))$

- We are interested in the **Abelian** case.

Definition

For f and g in an automaton \mathcal{A} , write $f + g$ for the function
$$(f + g)(x) = f(g(x))$$

- We are interested in the **Abelian** case.
- For all of our machines, $f + g = g + f$

Definition

For f and g in an automaton \mathcal{A} , write $f + g$ for the function
 $(f + g)(x) = f(g(x))$

- We are interested in the **Abelian** case.
- For all of our machines, $f + g = g + f$
- Given a machine \mathcal{A} , this condition is checkable in polynomial time

Definition

Recall a *Group* is a set \mathcal{G} equipped with

- $0 \in \mathcal{G}$
- $+: \mathcal{G} \rightarrow \mathcal{G} \rightarrow \mathcal{G}$ (associative)
- $-: \mathcal{G} \rightarrow \mathcal{G}$
- satisfying $0 + x = x + 0 = x$ and $x + (-x) = (-x) + x = 0$

Definition

Recall a *Group* is a set \mathcal{G} equipped with

- $0 \in \mathcal{G}$
 - $+: \mathcal{G} \rightarrow \mathcal{G} \rightarrow \mathcal{G}$ (associative)
 - $-: \mathcal{G} \rightarrow \mathcal{G}$
 - satisfying $0 + x = x + 0 = x$ and $x + (-x) = (-x) + x = 0$
-
- If S is the state set of an automaton \mathcal{A} , consider \mathcal{G} to be the closure of S under $+$
 - take $0 = id : 2^* \rightarrow 2^*$ to be the empty sum

Definition

Recall a *Group* is a set \mathcal{G} equipped with

- $0 \in \mathcal{G}$
 - $+: \mathcal{G} \rightarrow \mathcal{G} \rightarrow \mathcal{G}$ (associative)
 - $-: \mathcal{G} \rightarrow \mathcal{G}$
 - satisfying $0 + x = x + 0 = x$ and $x + (-x) = (-x) + x = 0$
-
- If S is the state set of an automaton \mathcal{A} , consider \mathcal{G} to be the closure of S under $+$
 - take $0 = id : 2^* \rightarrow 2^*$ to be the empty sum
 - This does not have $-$ in general
 - Our functions don't even need to be *invertible*

Definition

Recall a *Group* is a set \mathcal{G} equipped with

- $0 \in \mathcal{G}$
 - $+: \mathcal{G} \rightarrow \mathcal{G} \rightarrow \mathcal{G}$ (*associative*)
 - $-: \mathcal{G} \rightarrow \mathcal{G}$
 - *satisfying* $0 + x = x + 0 = x$ and $x + (-x) = (-x) + x = 0$
-
- If S is the state set of an automaton \mathcal{A} , consider \mathcal{G} to be the closure of S under $+$
 - take $0 = id : 2^* \rightarrow 2^*$ to be the empty sum
 - This does not have $-$ in general
 - Our functions don't even need to be *invertible*
 - Each function is invertible iff each state is invertible in one step


The Inverse Automaton

- Since each state sees an invertible function. . . invert it.


The Inverse Automaton

- Since each state sees an invertible function. . . invert it.

\mathcal{A}_2^3




$-\mathcal{A}_2^3$




The Inverse Automaton

- Since each state sees an invertible function. . . invert it.

\mathcal{A}_2^3



$-\mathcal{A}_2^3$




- Take Care: $\partial_0(-f) = -\partial_1 f$


The Inverse Automaton

- Since each state sees an invertible function. . . invert it.

\mathcal{A}_2^3



$-\mathcal{A}_2^3$




- Take Care: $\partial_0(-f) = -\partial_1 f$
- But: $(f + (-f))(01) = f((-f)(01)) = f(11) = 01$


The Inverse Automaton

- Since each state sees an invertible function. . . invert it.

\mathcal{A}_2^3



$-\mathcal{A}_2^3$



- Take Care: $\partial_0(-f) = -\partial_1 f$
- But: $(f + (-f))(01) = f((-f)(01)) = f(11) = 01$
- An easy induction shows these are actually inverses.

- So $\mathcal{G}(\mathcal{A})$ is a group whenever each state sees an invertible function

- So $\mathcal{G}(\mathcal{A})$ is a group whenever each state sees an invertible function
- If \mathcal{A} is abelian, so is $\mathcal{G}(\mathcal{A})$.

- So $\mathcal{G}(\mathcal{A})$ is a group whenever each state sees an invertible function
- If \mathcal{A} is abelian, so is $\mathcal{G}(\mathcal{A})$.
- What groups can we get?

- So $\mathcal{G}(\mathcal{A})$ is a group whenever each state sees an invertible function
- If \mathcal{A} is abelian, so is $\mathcal{G}(\mathcal{A})$.
- What groups can we get?

Theorem (Nekrashevych and Sidki)

The only abelian automaton groups are $(\mathbb{Z}/2\mathbb{Z})^m$ and \mathbb{Z}^m .

- So $\mathcal{G}(\mathcal{A})$ is a group whenever each state sees an invertible function
- If \mathcal{A} is abelian, so is $\mathcal{G}(\mathcal{A})$.
- What groups can we get?

Theorem (Nekrashevych and Sidki)

The only abelian automaton groups are $(\mathbb{Z}/2\mathbb{Z})^m$ and \mathbb{Z}^m .

- Given an abelian automaton \mathcal{A} , one can check in polynomial time which group it generates.

- So $\mathcal{G}(\mathcal{A})$ is a group whenever each state sees an invertible function
- If \mathcal{A} is abelian, so is $\mathcal{G}(\mathcal{A})$.
- What groups can we get?

Theorem (Nekrashevych and Sidki)

The only abelian automaton groups are $(\mathbb{Z}/2\mathbb{Z})^m$ and \mathbb{Z}^m .

- Given an abelian automaton \mathcal{A} , one can check in polynomial time which group it generates.
- We will focus on the \mathbb{Z}^m case here

Theorem

\mathbb{Z}^m equipped with a matrix \mathbf{A} and \bar{e} forms a (infinite state) abelian automaton (called $\mathfrak{C}(\mathbf{A}, \bar{e})$) with residuation as shown below. Further, for every abelian automaton \mathcal{A} whose group is \mathbb{Z}^m , there exists an \mathbf{A} and \bar{e} such that \mathcal{A} is a finite subautomaton. The odd states are exactly the states with odd first component.

$$\mathbf{A} = \begin{pmatrix} \frac{a_1}{2} & 1 & 0 & \cdots & 0 \\ \frac{a_2}{2} & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{a_{m-1}}{2} & 0 & 0 & \cdots & 1 \\ \frac{a_m}{2} & 0 & 0 & \cdots & 0 \end{pmatrix}$$


$$\partial_0 \bar{v} = \begin{cases} A(\bar{v}) & \bar{v} \text{ is even} \\ A(\bar{v} - \bar{e}) & \bar{v} \text{ is odd} \end{cases}$$

$$\partial_1 \bar{v} = \begin{cases} A(\bar{v}) & \bar{v} \text{ is even} \\ A(\bar{v} + \bar{e}) & \bar{v} \text{ is odd} \end{cases}$$

- $a_i \in \mathbb{Z}$
- \mathbf{A} has irreducible characteristic polynomial
- \bar{e} (the **Residuation Vector**) is odd

Example:

Take $\mathbf{A} = \begin{pmatrix} -1 & 1 \\ -\frac{1}{2} & 0 \end{pmatrix}$, and $\bar{e} = (3, 2)$. Then:



- It is natural to ask for what matrices \mathbf{A} and vectors \bar{e} can we find a given \mathcal{A} in $\mathfrak{C}(\mathbf{A}, \bar{e})$, and at what vectors \bar{v} are its states?

- It is natural to ask for what matrices \mathbf{A} and vectors \bar{e} can we find a given \mathcal{A} in $\mathfrak{C}(\mathbf{A}, \bar{e})$, and at what vectors \bar{v} are its states?
- It can be shown that only one \mathbf{A} works

- It is natural to ask for what matrices \mathbf{A} and vectors \bar{e} can we find a given \mathcal{A} in $\mathfrak{C}(\mathbf{A}, \bar{e})$, and at what vectors \bar{v} are its states?
- It can be shown that only one \mathbf{A} works
- Becker even found a way of computing \mathbf{A} given the automaton

- It is natural to ask for what matrices \mathbf{A} and vectors \bar{e} can we find a given \mathcal{A} in $\mathfrak{C}(\mathbf{A}, \bar{e})$, and at what vectors \bar{v} are its states?
- It can be shown that only one \mathbf{A} works
- Becker even found a way of computing \mathbf{A} given the automaton
- It can also be shown that for any \bar{e} , if \mathcal{A} is a subautomaton, its location in the structure is unique

- It is natural to ask for what matrices \mathbf{A} and vectors \bar{e} can we find a given \mathcal{A} in $\mathfrak{C}(\mathbf{A}, \bar{e})$, and at what vectors \bar{v} are its states?
- It can be shown that only one \mathbf{A} works
- Becker even found a way of computing \mathbf{A} given the automaton
- It can also be shown that for any \bar{e} , if \mathcal{A} is a subautomaton, its location in the structure is unique
- There are infinitely many choices of \bar{e} though, and the goal is to understand them.

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\bar{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\vec{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}
- Seemingly weird idea:

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\vec{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}
- Seemingly weird idea:
 - ▶ Can we take *polynomials* as our scalars instead?

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\vec{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}
- Seemingly weird idea:
 - ▶ Can we take *polynomials* as our scalars instead?
- Typically this is done by setting $x\vec{v} = \mathbf{A}\vec{v}$ for some linear transformation.

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\vec{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}
- Seemingly weird idea:
 - ▶ Can we take *polynomials* as our scalars instead?
- Typically this is done by setting $x\vec{v} = \mathbf{A}\vec{v}$ for some linear transformation.
- If only we had a obvious linear transformation floating around our structure that one might try...

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\vec{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}
- Seemingly weird idea:
 - ▶ Can we take *polynomials* as our scalars instead?
- Typically this is done by setting $x\vec{v} = \mathbf{A}\vec{v}$ for some linear transformation.
- If only we had a obvious linear transformation floating around our structure that one might try...
- For technical reasons, we'll use \mathbf{A}^{-1} instead of \mathbf{A} .

$\mathbb{Z}[x]$ -module

- We can multiply vectors $\bar{v} \in \mathbb{Z}^m$ by scalars in \mathbb{Z}
- Seemingly weird idea:
 - ▶ Can we take *polynomials* as our scalars instead?
- Typically this is done by setting $x\bar{v} = \mathbf{A}\bar{v}$ for some linear transformation.
- If only we had a obvious linear transformation floating around our structure that one might try...
- For technical reasons, we'll use \mathbf{A}^{-1} instead of \mathbf{A} .

Definition

For $p \in \mathbb{Z}[x]$ and $\bar{v} \in \mathfrak{C}(\mathbf{A}, \bar{e})$, put $p \cdot \bar{v} = (p(\mathbf{A}^{-1}))\bar{v}$

- Why should we care?

- Why should we care?

Theorem

for each $\bar{v} \in \mathbb{Z}^m$, there is $p_{\bar{v}} \in \mathbb{Z}[x]$ such that $p_{\bar{v}} \cdot e_1 = \bar{v}$

- Why should we care?

Theorem

for each $\bar{v} \in \mathbb{Z}^m$, there is $p_{\bar{v}} \in \mathbb{Z}[x]$ such that $p_{\bar{v}} \cdot e_1 = \bar{v}$

Theorem

If \bar{e} is an odd vector, then $\varphi_{\bar{e}} : \mathfrak{C}(\mathbf{A}, \bar{e}_1) \hookrightarrow \mathfrak{C}(\mathbf{A}, \bar{e})$ by $\varphi_{\bar{e}}(\bar{v}) = p_{\bar{e}} \cdot \bar{v}$ is an embedding, and preserves the group structure and the residuation structure.

- Why should we care?

Theorem

for each $\bar{v} \in \mathbb{Z}^m$, there is $p_{\bar{v}} \in \mathbb{Z}[x]$ such that $p_{\bar{v}} \cdot e_1 = \bar{v}$

Theorem

If \bar{e} is an odd vector, then $\varphi_{\bar{e}} : \mathfrak{C}(\mathbf{A}, \bar{e}_1) \hookrightarrow \mathfrak{C}(\mathbf{A}, \bar{e})$ by $\varphi_{\bar{e}}(\bar{v}) = p_{\bar{e}} \cdot \bar{v}$ is an embedding, and preserves the group structure and the residuation structure.

- This embedding is surjective if and only if $p_{\bar{e}}$ is a unit in $\mathbb{Z}[x]/\chi$, where χ is the characteristic polynomial of \mathbf{A}^{-1}

- Why should we care?

Theorem

for each $\bar{v} \in \mathbb{Z}^m$, there is $p_{\bar{v}} \in \mathbb{Z}[x]$ such that $p_{\bar{v}} \cdot e_1 = \bar{v}$

Theorem

If \bar{e} is an odd vector, then $\varphi_{\bar{e}} : \mathfrak{C}(\mathbf{A}, \bar{e}_1) \hookrightarrow \mathfrak{C}(\mathbf{A}, \bar{e})$ by $\varphi_{\bar{e}}(\bar{v}) = p_{\bar{e}} \cdot \bar{v}$ is an embedding, and preserves the group structure and the residuation structure.

- This embedding is surjective if and only if $p_{\bar{e}}$ is a unit in $\mathbb{Z}[x]/\chi$, where χ is the characteristic polynomial of \mathbf{A}^{-1}
- So different residuation vectors give groups which *extend* the group $\mathfrak{C}(\mathbf{A}, \bar{e}_1)$

- Why should we care?

Theorem

for each $\bar{v} \in \mathbb{Z}^m$, there is $p_{\bar{v}} \in \mathbb{Z}[x]$ such that $p_{\bar{v}} \cdot e_1 = \bar{v}$

Theorem

If \bar{e} is an odd vector, then $\varphi_{\bar{e}} : \mathfrak{C}(\mathbf{A}, \bar{e}_1) \hookrightarrow \mathfrak{C}(\mathbf{A}, \bar{e})$ by $\varphi_{\bar{e}}(\bar{v}) = p_{\bar{e}} \cdot \bar{v}$ is an embedding, and preserves the group structure and the residuation structure.

- This embedding is surjective if and only if $p_{\bar{e}}$ is a unit in $\mathbb{Z}[x]/\chi$, where χ is the characteristic polynomial of \mathbf{A}^{-1}
- So different residuation vectors give groups which *extend* the group $\mathfrak{C}(\mathbf{A}, \bar{e}_1)$
- Also, if $p_{\bar{e}}$ divides $p_{\bar{r}}$, then $\mathfrak{C}(\mathbf{A}, \bar{r})$ extends $\mathfrak{C}(\mathbf{A}, \bar{e})$.

Theorem

If \mathcal{A} is an automaton whose group is \mathbb{Z}^m , then for each odd state in \mathcal{A} , there is exactly one \bar{e} which locates that state at \bar{e}_1 in $\mathfrak{C}(\mathbf{A}, \bar{e})$. Further, if \bar{e} and \bar{r} are two such residuation vectors, they differ by a unit. This procedure is effective.

Theorem

If \mathcal{A} has a state located at $\bar{v} \in \mathfrak{C}(\mathbf{A}, \bar{e})$, then \bar{v} is located at $p \cdot \bar{v} \in \mathfrak{C}(\mathbf{A}, p \cdot \bar{e})$

- Incredibly, we now understand residuation vectors!

Theorem

If \mathcal{A} is an automaton whose group is \mathbb{Z}^m , then for each odd state in \mathcal{A} , there is exactly one \bar{e} which locates that state at \bar{e}_1 in $\mathfrak{C}(\mathbf{A}, \bar{e})$. Further, if \bar{e} and \bar{r} are two such residuation vectors, they differ by a unit. This procedure is effective.

Theorem

If \mathcal{A} has a state located at $\bar{v} \in \mathfrak{C}(\mathbf{A}, \bar{e})$, then \bar{v} is located at $p \cdot \bar{v} \in \mathfrak{C}(\mathbf{A}, p \cdot \bar{e})$

- Incredibly, we now understand residuation vectors!
- First find \bar{r} such that \mathcal{A} has a state at \bar{e}_1 .
- \mathcal{A} is a subautomaton of $\mathfrak{C}(\mathbf{A}, \bar{e})$ if and only if $p_{\bar{r}}$ divides $p_{\bar{e}}$
- Also, if \mathcal{A} is a subautomaton, then $qp_{\bar{r}} = p_{\bar{e}}$, and \mathcal{A} is located at $q \cdot \bar{e}_1$

- It turns out we can “scale by an infinite polynomial” to get a universal structure which contains *every* automaton (with the correct matrix **A**) at exactly one location.
- The construction is a bit involved, so we don’t have time to discuss it, but it is computable, and removes the need for the extra parameter \bar{e} .

Questions?