

Extensions of Abelian Automaton Groups

Chris Grossack
(Advisor: Klaus Sutner)

Background

For our purposes, a **Mealy Automaton** is a tuple $\mathcal{A} = (S, \tau)$ where S is the **State Set**, and $\tau : S \times \mathbf{2} \rightarrow S \times \mathbf{2}$ is the **transition function**. Given a state $s \in S$, we can treat it as a length preserving function $\underline{s} : \mathbf{2}^* \rightarrow \mathbf{2}^*$ as follows:

$$\begin{aligned}\underline{s}(\varepsilon) &= \varepsilon \\ \underline{s}(ax) &= a'\underline{s}'(x) \quad (\text{where } (s', a') = \tau(s, a))\end{aligned}$$

For us, each $\tau(s, -) : \mathbf{2} \rightarrow \mathbf{2}$ will be a permutation, as this will ensure inverses exist when we define $\mathcal{G}(\mathcal{A})$, the group of functions the automaton generates.

A state is called **Odd** if its permutation flips its input, and **Even** if it copies its input.


Groups

We define $\mathcal{G}(\mathcal{A})$ to be the group generated by $\{\underline{s} | s \in \mathcal{A}\}$. We will restrict ourselves to Abelian groups, and write our groups additively.

For $f \in \mathcal{G}(\mathcal{A})$, put $\partial_0 f$ as the unique function such that for all $w \in \mathbf{2}^*$, $f(0w) = (f(0))(\partial_0 f)(w)$. Define $\partial_1 f$ symmetrically.

In the abelian case, $\mathcal{G}(\mathcal{A})$ will always be \mathbb{Z}^m or $(\mathbb{Z}/2\mathbb{Z})^m$. We restrict our attention to the \mathbb{Z}^m case.

An Automaton: \mathcal{A}_2^3



$$f(0110) = 1f_0(110) = 11f(10) = 110f_1(0) = 1100$$

$$\partial_0 f = f_0 \text{ and } \partial_0 f_0 = \partial_1 f_0 = f$$

Main Theorems

Theorem. Every nontrivial abelian automaton \mathcal{A} can be located at \bar{e}_1 in some $\mathfrak{C}(\mathbf{A}, \bar{e})$

Theorem. If $rp = q$ in $\mathbb{Z}[x]$, then $\mathfrak{C}(\mathbf{A}, p \cdot \bar{e}_1) \hookrightarrow \mathfrak{C}(\mathbf{A}, q \cdot \bar{e}_1)$, with a canonical injection $\varphi_r : \bar{v} \mapsto r \cdot \bar{v}$. In particular, if r is a unit, then $p \cdot \mathcal{G} \cong q \cdot \mathcal{G}$. This map preserves both the group and residuation structure.

These allow us to completely understand the residuation vector \bar{e} .

- First find \bar{r} such that \mathcal{A} has a state at \bar{e}_1 .
- \mathcal{A} is a subautomaton of $\mathfrak{C}(\mathbf{A}, \bar{e})$ if and only if $p_{\bar{r}}$ divides $p_{\bar{e}}$
- Also, if \mathcal{A} is a subautomaton, then $qp_{\bar{r}} = p_{\bar{e}}$, and \mathcal{A} is located at $q \cdot \bar{e}_1$

Example

So $p \cdot \bar{v} \in \mathfrak{C}(\mathbf{A}, p \cdot \bar{e}_1)$, computes exactly the same function as $\bar{v} \in \mathfrak{C}(\mathbf{A}, \bar{e})$. However, most vectors cannot be written as $p \cdot \bar{v}$. What do they do as functions? We call such vectors (and their corresponding functions) **Fractional**, due to the following observation and theorem:


Let δ be the function computed by $\bar{e}_1 \in \mathfrak{C}(\mathbf{A}, \bar{e})$. Then $3\bar{e}_1 \in \mathfrak{C}(\mathbf{A}, 3 \cdot \bar{e}_1)$ should compute the same function. So we should expect the function $\bar{e}_1 \in \mathfrak{C}(\mathbf{A}, 3 \cdot \bar{e}_1)$ to behave like " $\frac{1}{3}\delta$ ", and in fact it does.

Initial Object: The Principal Automaton

So $\mathfrak{C}(\mathbf{A}, \bar{e}_1)$ is a subautomaton of every $\mathfrak{C}(\mathbf{A}, p \cdot \bar{e})$, and it is reasonable to ask if there is a corresponding finite automaton whose group embeds into the group generated by every other automaton with the same matrix.

Such a machine was found by Okano, and is called the **Principal Automaton** of a matrix. It is located at \bar{e}_1 in $\mathfrak{C}(\mathbf{A}, \bar{e}_1)$ for any matrix \mathbf{A} , and can be directly constructed by taking differences of the states in a given automaton \mathcal{A} .

Below is the result of this construction for \mathcal{A}_2^3 :



Complete Automaton

Theorem (Nekrashevych and Sidki). If \mathcal{G} is an automaton group and $\varphi : \mathcal{G} \rightarrow \mathbb{Z}^m$ is a group isomorphism, then there is a matrix \mathbf{A} of \mathbb{Q} -irreducible character and an odd vector \bar{e} such that if \mathbb{Z}^m is equipped with the following residuation structure, then φ preserves residuation:

If \bar{v} is even:

$$\partial_0 \bar{v} = \partial_1 \bar{v} = \mathbf{A} \bar{v}$$

If \bar{v} is odd:

$$\partial_0 \bar{v} = \mathbf{A}(\bar{v} - \bar{e})$$

$$\partial_1 \bar{v} = \mathbf{A}(\bar{v} + \bar{e})$$

Further, \mathbf{A} is unique up to conjugation, and can always be taken to be " $\frac{1}{2}$ -integral", meaning \mathbf{A} is of the form

$$\begin{pmatrix} \frac{a_{11}}{2} & a_{12} & \dots & a_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{a_{m1}}{2} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

where each $a_{ij} \in \mathbb{Z}$.

These matrices all have characteristic polynomial

$\chi = x^n + \frac{1}{2}g(x)$, where $g \in \mathbb{Z}[x]$ and has constant term ± 1 .

Definition. This construction is called $\mathfrak{C}(\mathbf{A}, \bar{e})$

Theorem. Every abelian automaton \mathcal{A} can be embedded in $\mathfrak{C}(\mathbf{A}, \bar{e})$ for some \mathbf{A} and \bar{e} .

It is natural to wonder what vectors \bar{e} admit a given automaton \mathcal{A} as a subautomaton of $\mathfrak{C}(\mathbf{A}, \bar{e})$. We answer this question here.

$\mathbb{Z}[x]$ -modules

$\mathcal{G}(\mathcal{A})$ naturally allows scalars in \mathbb{Z} by putting

$$n \cdot f = \underbrace{f + f + \dots + f}_{n \text{ times}}$$

We allow scalars in $\mathbb{Z}[x]$, polynomials with integer coefficients, by embedding $\mathcal{G}(\mathcal{A})$ in $\mathfrak{C}(\mathbf{A}, \bar{e})$, and putting $p \cdot \bar{v} = p(\mathbf{A}^{-1})\bar{v}$.

Theorem. Because \mathbf{A} has irreducible character, every vector \bar{v} is equal to $p_{\bar{v}} \cdot \bar{e}_1$ for some $p_{\bar{v}}$. Here \bar{e}_1 is the unit vector $(1, 0, \dots, 0)$.

Acknowledgements

This research would not exist without the advice of my advisor Dr Klaus Sutner. There aren't enough thanks for the hours of conversation I enjoyed.