

1 Fundamental Properties

1.1 Existence and Uniqueness

1.1.1 Local Existence and Uniqueness

Let $f(t, x)$ be piecewise continuous in t and satisfy the Lipschitz condition

$$||f(t, x) - f(t, y)|| \leq L||x - y||$$

$\forall x, y \in B = \{x \in \mathbb{R}^n \mid ||x - x_0|| \leq r\}$, $\forall t \in [t_0, t_1]$. Then, there exists some $\delta > 0$ such that the state equation $\dot{x} = f(t, x)$ with $x(t_0) = x_0$ has a unique solution over $[t_0, t_0 + \delta]$.

1.1.2 Lipschitz Condition

Let $f : [a, b] \times D \rightarrow \mathbb{R}^m$ be continuous on some domain $D \subset \mathbb{R}^n$. Suppose that $[\partial f / \partial x]$ exists and is continuous on $[a, b] \times D$. If, for a convex subset $W \subset D$, there is a constant $L \geq 0$ such that

$$\left\| \frac{\partial f}{\partial x}(t, x) \right\| \leq L$$

on $[a, b] \times W$, then

$$||f(t, x) - f(t, y)|| \leq L||x - y||$$

for all $t \in [a, b]$, $x \in W$, and $y \in W$.

1.1.3 Locally Lipschitz

If $f(t, x)$ and $[\partial f / \partial x](t, x)$ are continuous on $[a, b] \times D$, for some domain $D \subset \mathbb{R}^n$, then f locally Lipschitz in x on $[a, b] \times D$.

1.1.4 Globally Lipschitz

If $f(t, x)$ and $[\partial f / \partial x](t, x)$ are continuous on $[a, b] \times \mathbb{R}^n$, then f is globally Lipschitz in x on $[a, b] \times \mathbb{R}^n$ if and only if $[\partial f / \partial x]$ is uniformly bounded on $[a, b] \times \mathbb{R}^n$.

1.1.5 Global Existence and Uniqueness

Suppose that $f(t, x)$ is piecewise continuous in t and satisfies

$$||f(t, x) - f(t, y)|| \leq L||x - y||$$

$\forall x, y \in \mathbb{R}^n, \forall t \in [t_0, t_1]$. Then, the state equation $\dot{x} = f(t, x)$, with $x(t_0) = x_0$, has a unique solution over $[t_0, t_1]$.

1.1.6 Existence and Uniqueness Theorem

2 Second-Order Systems

2.1 Limit Cycles

We consider a two-dimensional dynamical system of the form

$$\dot{x}(t) = V(x(t))$$

where $V : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a smooth function. A trajectory of this system is some smooth function $x(t)$ with values in \mathbb{R}^2 which satisfies this differential equation. Such a trajectory is called closed (or periodic) if it is not constant but returns to its starting point, i.e., if there exists some $t_0 > 0$ such that $x(t + t_0) = x(t)$ for all $t \in \mathbb{R}$. An orbit is the image of a trajectory, a subset of \mathbb{R}^2 . A closed orbit, or cycle, is the image of a closed trajectory. A limit cycle is a cycle which is the limit set of some other trajectory.




Figure 1: Stable Limit Cycle

2.1.1 Poincaré-Bendixson Criterion

Consider the system $\dot{x} = f(x)$ and let M be a closed bounded subset of the plane such that

- M contains no equilibrium points or contains only one equilibrium point such that the Jacobian matrix $[\partial f / \partial x]$ at this point has eigenvalues with positive real parts. (Hence, the equilibrium point is unstable focus or unstable node.)
- Every trajectory starting in M stays in M for all future time.

Then, M contains a periodic orbit of $\dot{x} = f(x)$.

2.1.2 Benixson Criterion

If, on a simply connected region (\mathcal{C}^1) D of the plane, the expression $\partial f_1 / \partial x_1 + \partial f_2 / \partial x_2$ is not identically zero and does not change sign, then $\dot{x} = f(x)$ has no periodic orbits lying entirely in D .

3 Lyapunov Stability

3.1 Autonomous Systems

3.1.1 Stability Principles

The equilibrium point $x = 0$ of $\dot{x} = f(X)$ is

- stable if, for each $\epsilon > 0$, there is a $\delta = \delta(\epsilon) > 0$ such that

$$\|x(0)\| < \delta \Rightarrow \|x(t)\| < \epsilon, \quad \forall t \geq 0$$

- unstable if it is not stable
- asymptotically stable if it is stable and δ can be chosen such that

$$\|x(0)\| < \delta \Rightarrow \lim_{t \rightarrow \infty} x(t) = 0$$

3.1.2 Quadratic Lyapunov Function

$$V(x) = \frac{1}{2}x^T P x, \quad P > 0 \quad (1)$$

and

$$\begin{aligned} \frac{d}{dt}V(x) &= \frac{1}{2}x^T P A x + (x^T P A x)^T \\ &= \frac{1}{2}2x^T P \dot{x} \\ &= x^T P \dot{x} \end{aligned} \quad (2)$$

$$\dot{V}(x) = x^T P \dot{x} \quad (3)$$

In other words

$$\dot{V}(x) = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} p_{11} & p_{21} \\ p_{12} & p_{22} \end{bmatrix} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} \quad (4)$$

3.1.3 Asymptotically Stable

Let $x = 0$ be an equilibrium point and $D \subset \mathbb{R}^n$ be a domain containing $x = 0$. Let $V : D \rightarrow \mathbb{R}$ be a continuously differentiable function such that

$$V(0) = 0 \quad \text{and} \quad V(x) > 0 \text{ in } D - \{0\} \quad (5)$$

$$\dot{V}(x) \leq 0 \text{ in } D \quad (6)$$

Then $x = 0$ is stable. Moreover, if

$$\dot{V}(x) < 0 \text{ in } D - \{0\} \quad (7)$$

then $x = 0$ is asymptotically stable.

3.1.4 Globally Asymptotic stable

Let $x = 0$ be an equilibrium point or $\dot{x} = f(x)$. Let $V : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuously differentiable function such that

$$V(0) = 0 \text{ and } V(x) > 0, \quad \forall x \neq 0$$

$$\|x\| \rightarrow \infty \Rightarrow V(x) \rightarrow \infty$$

3.1.5 Radially Unbounded Function

A radially unbounded function is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ for which

$$\|x\| \rightarrow \infty \Rightarrow f(x) \rightarrow \infty$$

Or equivalently

$$\forall c > 0 : \exists r > 0 : \forall x \in \mathbb{R}^n : [\|x\| > r \Rightarrow f(x) > c]$$

3.1.6 Region of attraction

The region of attraction is also called the region of asymptotic stability. The region of attraction may be hard to calculate exactly, but Lyapunov functions can be used to estimate this region.

First of all, the region of attraction must satisfy the following: from the definition of asymptotic stability we see that if there is a Lyapunov function that satisfy the condition of asymptotic stability over a domain \mathbb{D} and if $\Omega_c = \{x \in \mathbb{R}^n | V(x) \leq c\}$ is bounded and contained in \mathbb{D} , then every trajectory starting in Ω_c remains in Ω_c and approaches the origin as $t \rightarrow \infty$

Let $x = 0$ be an asymptotically stable equilibrium point of the system $\dot{x} = f(x)$, where $f : \mathbb{D} \rightarrow \mathbb{R}^n$ is locally lipschitz and $\mathbb{D} \subset \mathbb{R}^n$ is a domain that contains the origin. Let $t \rightarrow \phi(t; x_0)$ be the solution of $\dot{x} = f(x)$ that starts at initial state x_0 at time $t = 0$. The region of attraction of the origin, denoted by R_A , is defined by

$$R_A = \{x_0 \in \mathbb{D} : \phi(t; x_0) \text{ is defined } \forall t \geq 0 \text{ and } \phi(t; x_0) \rightarrow 0 \text{ as } t \rightarrow \infty\}$$

I.e. the region of attraction R_A is the set of all points x_0 in \mathbb{D} such that the solution of

$$\dot{x} = f(x) \quad x(0) = x_0$$

is defined for all $t \geq 0$ and converges to the origin as $t \rightarrow \infty$.

Example 1 (Example of region of attraction)

Our domain \mathbb{D} is defined as

$$\mathbb{D} = \{x \in \mathbb{R}^2 : x_1 + x_2 < 1\}$$

where $\dot{V}(x) < 0$. Our region of attraction is defined as

$$\Omega_c = \{x \in \mathbb{R}^2 : V(x) \leq c\}$$

where $c < \min(V(x))$ and $x \in \partial c$ (x is on the border of c).

$$\begin{aligned} x_1 + x_2 &= 1 \\ x_1 &= 1 - x_2 \end{aligned}$$

Then we minimize $V(x)$ with respect of x_2

$$\begin{aligned} \min(V(x))_{x_2} \Big|_{x_1=1-x_2} &= \min_{x_2} \left(\frac{1}{2}((1-x_2)^2 + x_2^2) \right) = 0 \\ \min_{x_2} \left(\frac{1}{2} - x_2 + x_2^2 \right) &= \frac{\partial}{\partial x_2} \left(\frac{1}{2} - x_2 + x_2^2 \right) = 0 \\ -1 + 2x_2 &= 0 \\ x_2 &= \frac{1}{2} \rightarrow x_1 = \frac{1}{2} \end{aligned}$$

Further, we must satisfy $c < V(\frac{1}{2}, \frac{1}{2}) = \frac{1}{4}$. We have therefore the following region of attraction

$$\Omega_{\frac{1}{4}} = \{x \in \mathbb{R}^2 : V(x) \leq \frac{1}{4}\}$$

3.2 The Invariance Principle

3.2.1 Invariant Set

A set M is said to be an invariant set with respect to $\dot{x} = f(x)$ iff

$$x(0) \in M \Rightarrow x(t) \in M, \quad \forall t \in \mathbb{R}$$

3.2.2 Positively Invariant Set


A set M is a positively invariant set with respect to $\dot{x} = f(x)$ iff

$$x(0) \in M \Rightarrow x(t) \in M, \quad \forall t \geq 0$$

3.2.3 Lasalle's Theorem

Lasalle's invariance principle can be used to show the asymptotic stability of an equilibrium point when \dot{V} is negative semi-definite, i.e. $\dot{V} \leq 0$. \dot{V} is semidefinite if \dot{V} is not strictly 0 in origo.

Let $\Omega \subset D$ be a compact set that is positively invariant with respect to $\dot{x} = f(x)$. Let $V : D \rightarrow \mathbb{R}$ be a continuously differentiable function such that $\dot{V}(x) \leq 0$ in Ω . Let E be the set of all points in Ω where $\dot{V}(x) = 0$. Let M be the largest invariant set in E . Then every solution starting in Ω approaches M as $t \rightarrow \infty$.



3.2.4 Barbashin Theorem

Let $x = 0$ be an equilibrium point for

$$\dot{x} = f(x)$$

Let $V : D \rightarrow \mathbb{R}$ be a continuously differentiable positive definite function on a domain D containing the origin $x = 0$, such that $\dot{V}(x) \leq 0$ in D . Let $S = \{x \in D \mid \dot{V}(x) = 0\}$ and suppose that no solution can stay identically in S , other than the trivial solution $x(t) \equiv 0$. Then, the origin is asymptotically stable. If a A matrix is Hurwitz, we can conclude asymptotic stability.

3.2.5 Krasovskii Theorem

Let $x = 0$ be an equilibrium point for the system

$$\dot{x} = f(x)$$

. Let $V : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuously differentiable, radially unbounded, positive definite function such that $\dot{V}(x) \leq 0$ for all $x \in \mathbb{R}^n$. Let $S = \{x \in \mathbb{R}^n \mid \dot{V}(x) = 0\}$ and suppose that no solution can stay identically in S , other than the trivial solution $x(t) = 0$. Then, the origin is globally asymptotically stable.

3.2.6 Dominating cross-terms

Completion of squares

$$x_1 x_2 \leq \frac{1}{2}(x_1^2 + x_2^2) = \frac{1}{2} \|x\|_2^2$$

Another alternative is to write \dot{V} as $-x^T Q x$, where $Q = Q^T$ is positive definite

Young's inequality

$$xy \leq \epsilon x^2 + \frac{1}{4\epsilon} y^2, \quad \forall \epsilon > 0 \quad \forall x, y \in \mathbb{R}$$

3.2.7 Handling terms with indeterminate sign

Cauchy-Schwarz Inequality

$$|a_1 x_1 + a_2 x_2 + \dots + a_n x_n| \leq \sqrt{(a_1^2 + a_2^2 + \dots + a_n^2)} \|x\|_2$$

3.3 Linear Systems and Linearization

3.3.1 Hurwitz matrix / Stability matrix

A matrix A is Hurwitz; that is, $\text{Re}\{\lambda_i\} < 0$ for all eigenvalues of A , if and only if for any given positive definite symmetric matrix Q there exists a positive definite symmetric matrix P that satisfies the *Lyapunov equation* (??). Moreover, if A is Hurwitz, then P is the unique solution of (??).

$$PA + A^T P = -Q \quad (8)$$

Another way one can prove asymptotic stability in the origin is by using this symmetric matrix Q . We can write $\dot{V}(x)$ as the following

$$\dot{V}(x) = x^T P \dot{x} + \dot{x}^T P x = -x^T Q x$$

If Q is positive definite, we can conclude asymptotic stability at the origin.

Example 2 (Example using Hurwitz matrix)

We have the following $\dot{V}(x)$

$$\dot{V}(x) \leq -x_2^2 \left(2 - \frac{1}{4\epsilon}\right) - x_1 x_2 - x_1^2 (1 - \epsilon)$$

We can then formulate the following Q matrix

$$Q = \begin{bmatrix} 1 - \epsilon & \frac{1}{2} \\ \frac{1}{2} & 2 - \frac{1}{4\epsilon} \end{bmatrix}$$

To ensure that Q is positive definite we can evaluate the principle of minors

$$(1 - \epsilon) \left(w - \frac{1}{4\epsilon}\right) - \frac{1}{4} > 0$$

while $(1 - \epsilon) > 0$. We can solve the last inequality, and ϵ must satisfy

$$\frac{2 - \sqrt{2}}{4} < \epsilon < \frac{2 + \sqrt{2}}{4}$$

to ensure that the origin is asymptotically stable.

3.3.2 Indirect Method

Let $x = 0$ be an equilibrium point for the nonlinear system

$$\dot{x} = f(x)$$

where $f : D \rightarrow \mathbb{R}^n$ is continuously differentiable and D is a neighborhood of the origin. Let

$$A = \left. \frac{\partial f}{\partial x_i} \right|_{x=0}$$

Then,

1. The origin is asymptotically stable if $\text{Re}(\lambda_i) < 0$ for all eigenvalues λ_i of A .
2. The origin is unstable if $\text{Re}(\lambda_i) > 0$ for one or more of the eigenvalues λ_i of A .

3.4 Comparison Functions

3.4.1 Class \mathcal{K} and Class \mathcal{K}_∞ Definitions

A continuous function $\alpha : [0, a) \rightarrow [0, \infty)$ is said to belong to class \mathcal{K} if it is strictly increasing and $\alpha(0) = 0$. It is said to belong to class \mathcal{K}_∞ if $a = \infty$ and $\alpha(r) \rightarrow \infty$ as $r \rightarrow \infty$.

3.4.2 Class \mathcal{KL} Definition

A continuous function $\beta : [0, a) \times [0, \infty) \rightarrow [0, \infty)$ is said to belong to class \mathcal{KL} if, for each fixed s , the mapping $\beta(r, s)$ belongs to \mathcal{K} with respect to r and, for each fixed r , the mapping $\beta(r, s)$ is decreasing with respect to s and $\beta(r, s) \rightarrow \infty$ as $s \rightarrow \infty$.

3.4.3 Class \mathcal{K} , \mathcal{K}_∞ and \mathcal{KL} Rules

Let α_1 and α_2 be class \mathcal{K} functions on $[0, a)$, α_3 and α_4 be class \mathcal{K}_∞ functions on $[0, \infty)$.

Example 3 (Class \mathcal{K})

$\alpha(r) = \tan^{-1} r$ is strictly increasing since $\dot{\alpha}(r) = \frac{1}{(1+r^2)} > 0$. It belongs to class \mathcal{K} , but not to class \mathcal{K}_∞ since $\lim_{r \rightarrow \infty} \alpha(r) = \frac{\pi}{2} < \infty$.

Example 4 (Class \mathcal{KL})

$\beta(r, s) = \frac{r}{(ksr+1)}$, for any positive real number k , is strictly increasing in r since

$$\frac{\partial \beta}{\partial r} = \frac{ks}{(ksr+1)^2} > 0$$

and strictly decreasing in s since

$$\frac{\partial \beta}{\partial s} = \frac{-kr^2}{(ksr+1)^2} < 0$$

Moreover $\beta(r, s) \rightarrow 0$ as $s \rightarrow \infty$

3.4.4 Introducing class \mathcal{K} functions in Lyapunov's stability theorem

Let α_1 and α_2 be class \mathcal{K} functions on $[0, a)$, α_3 and α_4 be class \mathcal{K}_∞ functions, and β be a class \mathcal{KL} function. Denote the inverse of α_i by α_i^{-1} . Then,

- α_1^{-1} is defined on $[0, \alpha_1(a))$ and belongs to class \mathcal{K} .
- α_3^{-1} is defined on $[0, \infty)$ and belongs to class \mathcal{K}_∞ .
- $\alpha_1 \circ \alpha_2$ belongs to class \mathcal{K} .
- $\alpha_3 \circ \alpha_4$ belongs to class \mathcal{K}_∞ .
- $\sigma(r, s) = \alpha_1(\beta(\alpha_2(r), s))$ belongs to class \mathcal{KL} .

Class \mathcal{K} and class \mathcal{KL} functions enter into Lyapunov analysis through the next subsections

3.4.5 Lyapunovs analysis with Class \mathcal{K} functions

Let $V : D \rightarrow \mathbb{R}$ be a continuous positive definite function defined on a domain $D \subset \mathbb{R}^n$ that contains the origin. Let $B_r \subset D$ for some $r > 0$. Then, there exist class \mathcal{K} functions α_1 and α_2 , defined on $[0, r]$, such that

$$\alpha_1(\|x\|) \leq V(x) \leq \alpha_2(\|x\|)$$

for all $x \in B_r$. If $D = \mathbb{R}^n$, the functions α_1 and α_2 will be defined on $[0, \infty)$ and the foregoing inequality will hold for all $x \in \mathbb{R}^n$. Moreover, if $V(x)$ is radially unbounded, then α_1 and α_2 can be chosen to belong to class \mathcal{K}_∞ .

3.4.6 Lyapunovs analysis with Class \mathcal{KL} functions

Consider the scalar autonomous differential equation

$$\dot{y} = -\alpha(y), \quad y(t_0) = y_0$$

where α is a locally Lipschitz class \mathcal{K} function defined on $[0, a)$. For all $0 \leq y_0 < a$, this equation has a unique solution $y(t)$ defined for all $t \geq t_0$. Moreover,

$$y(t) = \sigma(y_0, t - t_0)$$

where σ is a class \mathcal{KL} function defined on $[0, a) \times [0, \infty)$.

3.5 Nonautonomous Systems

Throughout this section we will consider the following Nonautonomous system

$$\dot{x} = f(t, x)$$

3.5.1 Uniformity

Proving uniformity means that the initial condition x_0 has no effect on the stability of the system.

Uniformity \mapsto convergence rate not dependent on t_0 .

3.5.2 Stability definitions

The equilibrium point $x = 0$ of ?? is

- stable if, for each $\varepsilon > 0$, there is $\delta = \delta(\varepsilon, t_0) > 0$ such that

$$\|x(t_0)\| < \delta \Rightarrow \|x(t)\| < \varepsilon, \quad \forall t \geq t_0 \geq 0$$

- uniformly stable if, for each $\varepsilon > 0$, there is $\delta = \delta(\varepsilon) > 0$, independent of t_0 , such that (4.16) is satisfied.
- unstable if it is not stable.
- asymptotically stable if it is stable and there is a positive constant $c = c(t_0)$ such that $x(t) \rightarrow 0$ as $t \rightarrow \infty$, for all $\|x(t_0)\| < c$.
- Asymptotically stable (AS) if

it is stable

there exists $c > 0$ s.t. for all $t_0 \geq 0$ and $\eta > 0$, there exists $T(t_0, \eta) > 0$ satisfying

$$\|x(t_0)\| \leq c \Rightarrow \|x(t)\| \leq \eta \quad \forall t \geq t_0 + T(t_0, \eta)$$

- Uniformly asymptotically stable (UAS) if

it is uniformly stable (US)

there exists $c > 0$ s.t. for all $t_0 \geq 0$ and $\eta > 0$, there exists $T(\eta) > 0$ satisfying

$$\|x(t_0)\| \leq c \Rightarrow \|x(t)\| \leq \eta \quad \forall t \geq t_0 + T(\eta)$$

- Globally uniformly asymptotically stable (GUAS) if

it is globally uniformly stable (GUS)

for all $\eta, c > 0$ and $t_0 \geq 0$ there exists $T(\eta, c) > 0$ s.t.

$$\|x(t_0)\| \leq c \Rightarrow \|x(t)\| \leq \eta \quad \forall t \geq t_0 + T(\eta, c)$$

3.5.3 Equilibrium Points

$x^* \in \mathbb{R}^n$ is an equilibrium point of $\dot{x} = f(t, x)$ if $f(t, x^*) = 0$ for all $t \geq 0$.

Example 5 (Example of equilibrium point)

$\dot{x} = -\frac{a(t)x}{1+x^2}$ has the equilibrium point $x^* = 0$. We can see this by evaluating $f(t, x^*)$

$\dot{x} = -\frac{a(t)}{1+x^2} + b(t)$ where $b(t) \neq 0 \forall t$ has no equilibrium points

3.5.4 Stability definitions with \mathcal{K} and \mathcal{KL} classes

For $\dot{x} = f(t, x)$, the equilibrium point $x^* = 0$ is

- Uniformly stable (US) if $\exists \alpha \in \mathcal{K}, \exists c > 0$ such theoremseparator

$$\|x(t)\| \leq \alpha(\|x(t_0)\|) \quad \forall t \geq t_0, \quad \|x(t_0)\| \leq c, \quad \forall t_0 \geq 0$$

- globally uniformly stable (GUS) if $\exists \alpha \in \mathcal{K}_\infty$ such that

$$\|x(t)\| \leq \alpha(\|x(t_0)\|) \quad \forall t \geq t_0, \quad \forall t_0 \geq 0$$

- exponentially stable if $\exists c, k, \lambda > 0$

$$\|x(t)\| \leq k\|x(t_0)\|e^{-\lambda(t-t_0)} \quad t \geq t_0 \geq 0, \forall \|x(t_0)\| < c$$

- globally exponentially stable if $\exists k, \lambda > 0$

$$\|x(t)\| \leq k\|x(t_0)\|e^{-\lambda(t-t_0)} \quad t \geq t_0 \geq 0, \forall x(t_0)$$

3.5.5 Time-varying Lyapunov function candidates

$V(t, x)$ is positive definite if


$$\left. \begin{array}{l} V(t, 0) = 0 \\ V(t, x) \geq W_1(x) \end{array} \right\} \forall t \geq 0, \text{ for some positive definite function } W_1(x)$$

and

- $(t, x) \rightarrow V(t, x)$ is positive semidefinite if $x \rightarrow W_1(x)$ positive semidefinite
- $(t, x) \rightarrow V(t, x)$ is radially unbounded if $x \rightarrow W_1(x)$ radially unbounded

$V(t, x)$ is decrescent if

$$\left. \begin{array}{l} V(t, 0) = 0 \\ V(t, x) \leq W_2(x) \end{array} \right\} \forall t \geq 0, \text{ for some positive definite function } W_2(x)$$



3.5.6 Lyapunov theorem: Uniform Stability

If there exist a function $V(t, x) : [0, \infty) \times \mathbb{D} \rightarrow \mathbb{R}$ and $W_i : \mathbb{D} \rightarrow \mathbb{R}$ continuous positive definite such that

- V is C^1
- $W_1(x) \leq V(t, x) \leq W_2(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- $\dot{V}(t, x) \leq 0 \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$


then $x = 0$ is uniformly stable.

Proof 1 The derivative of V along the trajectories of (4.15) is given by

$$\dot{V}(t, x) = \frac{\partial V}{\partial t} + \frac{\partial V}{\partial x} f(t, x) \leq 0$$

Choose $r > 0$ and $c > 0$ such that $B_r \subset D$ and $c < \min_{\|x\|=r} W_1(x)$. Then, $\{x \in B_r \mid W_1(x) \leq c\}$ is in the interior of B_r . Define a time-dependent set $\Omega_{t,c}$ by

$$\Omega_{t,c} = \{x \in B_r \mid V(t, x) \leq c\}$$



The set $\Omega_{t,c}$ contains $\{x \in B_r \mid W_2(x) \leq c\}$ since

$$W_2(x) \leq c \Rightarrow V(t, x) \leq c$$

On the other hand, $\Omega_{t,c}$ is a subset of $\{x \in B_r \mid W_1(x) \leq c\}$ since

$$V(t, x) \leq c \Rightarrow W_1(x) \leq c$$

Thus,

$$\{x \in B_r \mid W_2(x) \leq c\} \subset \Omega_{t,c} \subset \{x \in B_r \mid W_1(x) \leq c\} \subset B_r \subset D$$

for all $t \geq 0$. These five nested sets are sketched in Figure 4.7. The setup of Figure 4.7 is similar to that of Figure 4.1, except that the surface $V(t, x) = c$ is now dependent on t , and that is why it is surrounded by the time-independent surfaces $W_1(x) = c$ and $W_2(x) = c$.

3.5.7 Lyapunov theorem: Uniform Asymptotical Stability

If there exist a function $V(t, x) : [0, \infty) \times \mathbb{D} \rightarrow \mathbb{R}$ and $W_i : \mathbb{D} \rightarrow \mathbb{R}$ continuous positive definite such that

- i) V is C^1
- ii) $W_1(x) \leq V(t, x) \leq W_2(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- iii) $\dot{V}(t, x) \leq -W_3(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$

then $x = 0$ is uniformly asymptotically stable.

3.5.8 Lyapunov theorem: Global Uniformly Asymptotical Stability

If there exist a function $V(t, x) : [0, \infty) \times \mathbb{R}^2 \rightarrow \mathbb{R}$ and $W_i : \mathbb{R}^2 \rightarrow \mathbb{R}$ continuous positive definite such that

- i) V is C^1
- ii) $W_1(x) \leq V(t, x) \leq W_2(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- iii) $\dot{V}(t, x) \leq -W_3(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- iv) W_1 is radially unbounded

then $x = 0$ is globally uniformly asymptotically stable.

3.5.9 Lyapunov theorem: Exponential Stability

If there exist a function $V(t, x) : [0, \infty) \times \mathbb{D} \rightarrow \mathbb{R}$ and constants $a, k_1, k_2, k_3 > 0$ such that

- i) V is C^1
- ii) $k_1 \|x\|^a \leq V(t, x) \leq k_2 \|x\|^a \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- iii) $\dot{V}(t, x) \leq -k_3 \|x\|^a \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$

then $x = 0$ is exponentially stable.

3.5.10 Barbalat's Lemma

If there exist functions $V : [0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$, $W_i : \mathbb{R}^n \rightarrow \mathbb{R}$ continuous positive definite and radially unbounded and $W : \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$ and positive semidefinite satisfying

- i) V is C^1
- ii) $W_1(x) \leq V(t, x) \leq W_2(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{R}^n$
- iii) $\dot{V}(t, x) \leq -W(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{R}^n$
- iv) for every $k > 0$ there exists $c > 0$ s.t. $\|x\| \leq k \implies |\dot{W}(t, x)| \leq c$ for all $t \geq 0$

then the origin is globally uniformly stable and all solutions approach $E = \{x \in \mathbb{R}^n : W(x) = 0\}$.

3.5.11 LaSalle-Yokishawa Theorem

$\dot{x} = f(t, x)$ where $f : [0, \infty) \times \mathbb{D} \rightarrow \mathbb{R}^n$ is piecewise continuous locally Lipschitz in x and $x = 0$ is an eq.point.

If there exist functions $V : [0, \infty) \times \mathbb{D} \rightarrow \mathbb{R}$, $W_i : \mathbb{D} \rightarrow \mathbb{R}$ continuous positive definite, $W : \mathbb{D} \rightarrow \mathbb{R}^{n \times n}$ continuous positive semidefinite, such that

- i) V is C^1
- ii) $W_1(x) \leq V(t, x) \leq W_2(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- iii) $\dot{V}(t, x) \leq -W(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{D}$
- iv) for every $k > 0$ there exists $r > 0$ s.t. $\|x\| \leq k \implies \|f(t, x)\| \leq r$ for all $t \geq 0$

then the origin is uniformly stable, and $\exists c > 0$ such that all solutions with $\|x(t_0)\| < c$ approach $E = \{x \in \mathbb{D} : W(x) = 0\}$

3.5.12 Global LaSalle-Yokishawa Theorem

$\dot{x} = f(t, x)$ where $f : [0, \infty) \times \mathbb{D} \rightarrow \mathbb{R}^n$ is piecewise continuous locally Lipschitz in x and $x = 0$ is an eq.point.

If there exist functions $V : [0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$, $W_i : \mathbb{R}^n \rightarrow \mathbb{R}$ continuous positive definite and radially unbounded, $W : \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$ continuous positive semidefinite

- i) V is C^1
- ii) $W_1(x) \leq V(t, x) \leq W_2(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{R}^n$
- iii) $\dot{V}(t, x) \leq -W(x) \quad \forall (t, x) \in [0, \infty) \times \mathbb{R}^n$
- iv) for every $k > 0$ there exists $r > 0$ s.t. $\|x\| \leq k \implies \|f(t, x)\| \leq r$ for all $t \geq 0$

then the origin is globally uniformly stable and all solutions approach $E = \{x \in \mathbb{R}^n : W(x) = 0\}$.

3.5.13 Converse Theorems: Exponential Stability

Let $x = 0$ be an equilibrium point for the nonlinear system

$$\dot{x} = f(t, x)$$

where $d : [0, \infty) \times D \rightarrow \mathbb{R}^n$ is continuously differentiable, $D = \{x \in \mathbb{R}^n : \|x\|_2 < r\}$, and the Jacobian matrix $[\partial f / \partial x]$ is bounded and Lipschitz on D , uniformly in t .

Let

$$A(t) = \frac{\partial f}{\partial x}(t, x)|_{x=0}$$

Then, $x = 0$ is an exponentially stable equilibrium point for the nonlinear system if and only if it is an exponentially stable equilibrium point for the linear system

$$\dot{x} = A(t)x$$

4 Input-To-State Stability

4.1 Defenition

Consider:

$$\Sigma : \dot{x} = f(t, x, u) \quad (9)$$

The system Σ is ISS if $\exists \beta \in \mathcal{KL}$ and $\exists \gamma \in \mathcal{K}$ such that for any $t_0 \geq 0$, any $x_0 = x(t_0) \in \mathbb{R}^n$ and any bounded input $t \mapsto u(t)$, the solution $t \mapsto x(t)$ exists $\forall t \geq t_0$ and satisfies

$$\|x(t)\| \leq \beta(\|x_0\|, t - t_0) + \gamma(\sup \|u(\tau)\|) \quad (10)$$

4.2 Practice

If a system is ISS stable, then:

- i) for $u=0$, the origin is GUAS (0-GUAS)
- ii) for a bounded input $t \mapsto u(t)$, every solution $t \mapsto x(t)$ is bounded.

If one of these is not satisfied, the system is **not** ISS.

How to check ISS

- i) V is C' $\exists \alpha_1, \alpha_2 \in \mathcal{K}_{\inf st}$