

Long COVID Research Analysis

Generated 2026-05-16 — 4666 papers analyzed, 470 with deep methodological extraction (v2 pipeline with calibration + provenance)

Long COVID research has generated an extensive but methodologically weak evidence base—dominated by Very Low to Low GRADE quality studies, pervasive vaccination confounding, definitional fragmentation across 39 distinct duration thresholds, and calibrated certainty reaching only 'possible' for all symptom prevalence estimates—leaving key clinical and policy questions about burden, mechanisms, and effective treatments substantially unresolved.

Executive Summary

This synthesis integrates 4,666 triaged papers and 470 deeply analyzed studies on Long COVID (post-acute sequelae of SARS-CoV-2, PASC). The evidence base is large but methodologically weak: of 470 deep papers, GRADE quality was Very Low in 186 (39.6%), Low in 265 (56.4%), and Moderate in only 19 (4.0%), with zero High-quality studies identified. Extraction confidence across deeply analyzed papers ranged from approximately 0.70 to 0.87, reflecting moderate-to-good abstract fidelity but substantial underlying study limitations. The calibrated certainty distribution —0 established, 0 probable (aggregate consensus), 147 possible,

and 2,799 speculative findings—signals that virtually no individual symptom prevalence estimate achieves high certainty at the cross-paper level. Fatigue is the most consistently reported symptom, appearing in 1,173 of 4,666 triaged papers (34.6%) and 48 deeply analyzed papers (mean extraction confidence 0.783), yet calibrated certainty remains only 'possible' due to definitional heterogeneity and absence of controlled baselines [1,2,3]. Vaccination before infection is associated with modestly reduced Long COVID risk across multiple large cohorts and meta-analyses, though heterogeneity renders pooled estimates imprecise [4,5,6]. The field is characterized by contested prevalence estimates (ranging from under 1% in administrative coding studies to over 70% in hospitalized cohorts), fragmented case definitions spanning at least 39 distinct duration thresholds, and pervasive unmeasured confounding by vaccination status, variant era, and baseline symptom burden.

Methods

This report synthesizes findings from **4666 scientific papers** about Long COVID retrieved from PubMed Central (and medRxiv preprints). Papers were queried for "long covid", "post-acute sequelae", "PASC", or "post-COVID condition", restricted to 2020–present.

A two-tier LLM extraction was performed:

1. **Triage** (Claude Haiku 4.5, all 4666 papers): abstract-level structured parsing with per-paper `extraction_confidence`.
2. **Deep extraction** (Claude Sonnet 4.6, top 470 papers): two-stage protocol — factual extraction (only what is explicitly stated) + methodological appraisal (NOS, GRADE, MCID, 8-

axis bias audit, phenotype mapping, calibrated certainty, literal-quote provenance).

Every claim in the deep extractions is backed by a **provenance record** (literal quote from the source paper, section, confidence). Findings are aggregated cross-paper into **calibrated certainty levels** (established / probable / possible / speculative / contradicted) per the GRADE framework adapted for this corpus.

This report is a **snapshot of the current data**. It is **not** a systematic review: no PRISMA flow, no peer review of conclusions, no formal risk-of-bias tool against established checklists beyond Newcastle-Ottawa scoring.

Findings

1. Calibrated Findings by Certainty

The propagated cross-paper consensus partitions findings into five evidence tiers:

Established

- The Long COVID scientific literature has grown rapidly and constitutes approximately 2.3% of all COVID-19 publications, with treatment and mechanism studies comprising 50.6% and 47.9% respectively—demonstrating that research investment has been substantial but unevenly distributed toward descriptive and mechanistic studies rather than interventional trials [\[7\]](#).

Probable

- Pre-infection COVID-19 vaccination is associated with a modest reduction in Long COVID risk, with a pooled odds ratio of approximately 0.54–0.77 across meta-analyses; however, extreme statistical heterogeneity ($I^2 \approx 96\text{--}98\%$) and unmeasured healthy vaccinee bias in all contributing observational studies prevent precise quantification, and causal inference requires substantial qualification [\[4,8,9,5\]](#).
- Long COVID risk and cumulative sequelae burden are substantially higher following hospitalized versus non-hospitalized acute COVID-19, with 3-year DALY burdens estimated at 642.8 versus 91.2 per 1,000 persons respectively in the largest VA cohort study ($n=5,341,996$); residual confounding by disease severity is acknowledged but the gradient is consistent across multiple independent analyses [\[10,11\]](#).
- PASC cumulative incidence declined across successive variant eras and with increasing vaccination: unvaccinated pre-Delta incidence was approximately 10.42 per 100 persons at one year, falling to 5.34 per 100 during Delta and 3.50 per 100 during Omicron among vaccinated individuals; these estimates are derived from a large register-based cohort ($n=441,583$) and are replicated directionally across multiple studies, though healthy vaccinee bias cannot be excluded [\[6,12,13\]](#).
- Female sex is consistently associated with higher Long COVID risk across study designs and populations, with adjusted risk ratios or odds ratios typically ranging from 1.3 to 2.0; this association is robust to adjustment for acute severity and appears partially independent of comorbidity burden, though residual confounding by health-seeking behaviour and symptom attribution remains plausible [\[14,15,16,17\]](#).

- Socioeconomic deprivation is independently associated with Long COVID risk, with the most deprived decile showing approximately 46% higher adjusted odds versus least deprived in a nationally representative UK study of 201,799 participants; this gradient is replicated across multiple high-income country datasets [\[18,19,20\]](#).
- Pulmonary rehabilitation is probably effective for improving physical capacity and lung function in Long COVID, with a meta-analysis of 17 RCTs (n=3,363) finding a mean 6-minute walk test improvement of 77.95 m (95% CI 50.91–104.99 m); the GRADE certainty is Moderate for this outcome [\[21\]](#).
- Immune-mediated inflammatory disease (IMID) patients have a 64% higher adjusted risk of PASC compared to matched non-IMID individuals (aHR 1.64, 95% CI 1.42–1.89) in a prospective nationwide Danish cohort of 77,667 individuals, suggesting that pre-existing immune dysregulation amplifies Long COVID susceptibility [\[22\]](#).
- Reinfection with SARS-CoV-2 is associated with increased Long COVID risk compared to primary infection; in a Danish nationwide register-based cohort of 4,418,544 individuals, absolute Long COVID risk was 1.16% versus lower after primary infection across landmark timepoints, with the relative risk directionally confirmed in multiple independent cohorts [\[23,24,25\]](#).
- Online cognitive-behavioral therapy probably reduces fatigue in Long COVID with moderate certainty evidence (MD –8.4, 95% CI –13.11 to –3.69), and combined physical-mental health rehabilitation probably improves physical function, based on a living systematic review of 3,695 participants [\[26\]](#).
- Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is a diagnosable sequela of SARS-CoV-2 infection, with an incidence rate of 2.66 per 100 person-years in infected participants versus 0.93 in propensity-matched uninfected

controls (HR 4.93, 95% CI 3.62–6.71) in the RECOVER-Adult prospective cohort (n=13,224) [27].

- Long COVID is associated with substantially increased healthcare utilization; individuals with coded Long COVID were over 8 times more likely to use any healthcare resource compared to matched comparators (OR 8.29), and annual incremental costs are estimated at USD 11,641 in adjusted analyses of the Medical Expenditure Panel Survey [28,29,30].
- Anosmia and dysgeusia are among the most COVID-specific persistent symptoms, with loss of smell showing an adjusted OR of 28.6 (95% CI not reported) versus the no-COVID group in a multi-cohort UK study of 42,450 participants; these associations are replicated across multiple controlled studies [31,13,32].

Possible

- Fatigue is the most commonly reported Long COVID symptom, appearing in approximately 34.6% of all indexed papers and with pooled prevalences ranging from 17–45% across meta-analyses, but calibrated certainty remains 'possible' due to absence of pre-infection baselines and inconsistent definitions in virtually all contributing studies [33,34,35].
- Paxlovid (nirmatrelvir-ritonavir) treatment during acute COVID-19 may reduce overall Long COVID incidence modestly (RR 0.986, 95% CI 0.953–1.020—non-significant overall, but RR 0.91 for cognitive symptoms, 95% CI 0.84–0.99) and reduce post-acute hospitalization (HR 0.70); channeling bias and computable phenotype limitations preclude causal inference [36,37,38].
- Pre-existing diabetes is associated with higher Long COVID risk (RR 1.60, 95% CI 1.12–2.29 in the UK Household Longitudinal Study; OR 1.37, non-significant after full

adjustment in SHARE); evidence is possible rather than probable due to inconsistent confounding control across studies [39,40].

- HIV infection may be associated with approximately twice the odds of Long COVID (OR 2.20, 95% CI 1.25–3.86) in a meta-analysis of five studies (n=39,405); this estimate is categorized as possible given Very Low GRADE certainty and absence of vaccination adjustment [41,42].
- Elevated BMI above approximately 22–23 kg/m² is associated with impaired recovery and higher PASC risk in a dose-dependent pattern, supported by multiple large cohort studies; however, causality is limited by absence of pre-infection BMI baselines in most studies and likely confounding by comorbidity [43,44,45].
- Long COVID is associated with increased cardiovascular risk including elevated hazard ratios for cardiomyopathy (0.15% absolute excess), deep vein thrombosis, and atrial arrhythmia across multiple large retrospective cohorts, though surveillance bias (more frequent monitoring in Long COVID patients) inflates these estimates [46,47,48].
- COVID-19 infection is associated with increased risk of incident gastrointestinal disorders post-acutely, including GERD (HR 1.36, 95% CI 1.18–1.56) and inflammatory liver disease (HR 1.73, 95% CI 1.31–2.26), in a multinational cohort of 22.7 million; replication and consistency support a possible causal role [49,50].
- Cognitive impairment following non-severe COVID-19 is possible, with meta-analyses finding medium effect sizes for memory impairment (g=−0.41) and moderate effects for language deficits; absence of pre-COVID baselines limits causal attribution [51,52].
- Dyslipidemia is a possible post-acute sequela, with COVID-19 survivors showing increased risk of incident total cholesterol

>200 mg/dL (HR 1.26, 95% CI 1.22–1.29) and triglycerides >150 mg/dL (HR 1.27, 95% CI 1.23–1.31) in a large VA cohort (n=51,919); the predominantly male, older veteran population limits generalizability [53].

- Pre-existing sleep disturbances are possibly associated with increased Long COVID risk (OR 1.20 for Long COVID from meta-analysis of eight studies) and poorer pre-infection sleep quality is associated with higher Long COVID risk in prospective cohorts; reverse causation and symptom overlap cannot be excluded [54,55,56].
- IL-6 receptor antagonists (tocilizumab/sarilumab) in rheumatoid arthritis patients are associated with approximately 60% lower 12-month Long COVID risk (aRR 0.42, 95% CI 0.27–0.64) in an emulated trial design; this is a possible but preliminary finding requiring replication with larger samples and active comparators [57].
- Long COVID is associated with meaningful functional limitations and disability, with 65% of self-identified Long COVID participants in the Johns Hopkins cohort classified with at least one disability; volunteer recruitment bias severely inflates this estimate [58,59].
- Long COVID is associated with increased odds of anxiety (OR ~2.25) and depression (OR ~2.26–2.35) in large cross-sectional and cohort studies, with prospective associations confirmed at approximately 3 years post-infection; the direction and magnitude of causality are uncertain due to baseline mental health confounding [60,61,62].

Speculative

- Viral reservoir persistence as a primary mechanism for Long COVID symptom generation is speculated based on associations between prolonged viral RNA shedding (RR 1.03 per unit, CITE:PMC10563605) and Long COVID risk, and

indirect evidence from antiviral treatment effects, but direct human evidence of persistent replication-competent virus causing symptoms remains absent from this corpus.

- Autonomic dysfunction (dysautonomia, POTS) as a dominant Long COVID phenotype is suggested by symptom clustering analyses identifying palpitations, orthostatic intolerance, and post-exertional malaise as a co-occurring cluster, but objective autonomic testing data with controlled comparators are absent from the majority of included studies.
- Microclot-mediated endotheliopathy as a mechanistic substrate for Long COVID is proposed based primarily on work from a single research group in South Africa and is not yet independently replicated with standardized methods across populations [\[2\]](#).
- Serotonin depletion as a mechanistic pathway linking viral persistence to neurocognitive Long COVID symptoms is a preliminary hypothesis referenced in the literature but not yet empirically evaluated in this corpus.
- Preliminary findings hint that SSRI use during acute COVID-19 may reduce Long COVID risk (aRR 0.704–0.922 in two retrospective studies), but confounding by indication, outcome misclassification, and absence of RCT evidence preclude any clinical recommendation [\[63,64\]](#).
- Ashwagandha root extract as a treatment for Long COVID fatigue is under investigation in an ongoing RCT (APRIL trial, n=2,500) but no efficacy data are available [\[65\]](#).
- Genetic susceptibility via HLA alleles and venous thromboembolism-related variants has been associated with Long COVID in preliminary GWAS and Mendelian randomization analyses, but genome-wide significance was achieved for only a single variant in a substantially underpowered study [\[66,67\]](#).

Contradicted

- The association between obstructive sleep apnea (OSA) and PASC is contradicted between two cross-sectional analyses using the same COPE dataset (n=24,803): PMC11144080 reports OSA associated with PASC (aOR range reported across PASC definitions), while PMC10793517 reports the same association using an overlapping analytical framework; the contradiction reflects definitional sensitivity rather than true contradiction, but the magnitude and direction vary by PASC definition applied, preventing a stable conclusion [68,69].
- The association between pre-existing diabetes and Long COVID risk is contradicted between two prospective cohort studies: the UK Household Longitudinal Study (n=11,669) finds pre-pandemic diabetes associated with 60% higher Long COVID risk (RR 1.60, 95% CI 1.12-2.29) [39], while the SHARE cohort study (n=4,004) finds no significant adjusted association [40]; likely causes include differences in cohort age distribution, variant era, and diabetes type ascertainment.
- Paxlovid effectiveness against Long COVID is contradicted between two large target trial emulations: Al-Aly et al. using PCORnet (n=445,738) finds no significant overall effect (RR 0.986, 95% CI 0.953-1.020) but a significant cognitive symptom effect [36], while Xie et al. using VA data (n=281,793) find a 26% overall Long COVID reduction (RR 0.74, 95% CI 0.72-0.77) [38]; likely causes include differences in PASC computable phenotype definitions, population characteristics (VA predominantly male veterans vs. mixed), and channeling bias magnitude.
- HIV infection's association with Long COVID is contradicted across studies: a meta-analysis of five small studies

(n=39,405) reports OR 2.20 (95% CI 1.25–3.86) [\[41\]](#), while a large multicenter study using PCORnet and N3C (n=4,682,251) finds a small but significant association using one phenotype definition but no association using ICD-10 coding [\[42\]](#); likely causes include phenotype definition sensitivity, low vaccination rates in PWH, and unmeasured ART regimen effects.

- The effect of pregnancy on Long COVID risk is contradicted between two large retrospective cohort studies: a PCORnet/N3C analysis (n=1,511,505) finds lower Long COVID risk in pregnant versus non-pregnant females (aHR 0.86, 95% CI 0.83–0.90) [\[70\]](#), while a separate PCORnet/N3C analysis (n=89,312) similarly finds lower PASC incidence at 180 days in pregnant females (30.8 vs 35.8 per 100 persons) [\[71\]](#); these are directionally consistent but a WHF global prospective cohort [\[72\]](#) finds no significant difference (adjusted prevalence 10.2% vs 10.6%), likely due to differences in PASC computable phenotype validity in pregnancy.

Quantitative breakdown (symptom-level, derived from deep extractions' calibration):

Certainty	# symptoms
Established	0
Probable	0
Possible	147
Speculative	2799
Contradicted	0

Symptom-level finding cards

Possible — sorted by (n_papers, mean_extraction_confidence) descending:

- **fatigue** — n=48 papers · extraction confidence: 78%
- **headache** — n=25 papers · extraction confidence: 81%
- **cough** — n=23 papers · extraction confidence: 78%
- **chest_pain** — n=18 papers · extraction confidence: 80%
- **shortness_of_breath** — n=16 papers · extraction confidence: 79%
- **anxiety** — n=15 papers · extraction confidence: 80%
- **dyspnea** — n=15 papers · extraction confidence: 78%
- **brain_fog** — n=13 papers · extraction confidence: 81%
- **dizziness** — n=13 papers · extraction confidence: 79%
- **fever** — n=13 papers · extraction confidence: 78%
- **joint_pain** — n=13 papers · extraction confidence: 77%
- **long_covid_overall** — n=12 papers · extraction confidence: 83%
- **depression** — n=12 papers · extraction confidence: 75%
- **palpitations** — n=11 papers · extraction confidence: 77%
- **muscle_pain** — n=10 papers · extraction confidence: 81%
- **myalgia** — n=10 papers · extraction confidence: 79%
- **abdominal_pain** — n=10 papers · extraction confidence: 79%
- **hair_loss** — n=10 papers · extraction confidence: 79%
- **insomnia** — n=7 papers · extraction confidence: 84%
- **long_covid_any** — n=7 papers · extraction confidence: 82%
- **sleep_disturbance** — n=7 papers · extraction confidence: 82%
- **cognitive_impairment** — n=7 papers · extraction confidence: 81%
- **arthralgia** — n=7 papers · extraction confidence: 80%
- **loss_of_taste** — n=6 papers · extraction confidence: 82%

- **sore_throat** — n=6 papers · extraction confidence: 82%
- **pain** — n=6 papers · extraction confidence: 79%
- **diarrhea** — n=6 papers · extraction confidence: 71%
- **loss_of_smell** — n=5 papers · extraction confidence: 82%
- **loss_of_appetite** — n=5 papers · extraction confidence: 82%
- **weight_loss** — n=5 papers · extraction confidence: 81%
- **difficulty_concentrating** — n=5 papers · extraction confidence: 78%
- **not_reported** — n=5 papers · extraction confidence: 76%
- **other_symptoms** — n=4 papers · extraction confidence: 86%
- **loss_of_taste_or_smell** — n=4 papers · extraction confidence: 85%
- **gastrointestinal_symptoms** — n=4 papers · extraction confidence: 83%
- **back_pain** — n=4 papers · extraction confidence: 83%
- **fatigue_overall** — n=4 papers · extraction confidence: 82%
- **sleep_disorders** — n=4 papers · extraction confidence: 82%
- **any_long_covid** — n=4 papers · extraction confidence: 82%
- **respiratory_symptoms** — n=4 papers · extraction confidence: 82%
- **rash** — n=4 papers · extraction confidence: 82%
- **nasal_congestion** — n=4 papers · extraction confidence: 82%
- **sleep_disturbances** — n=4 papers · extraction confidence: 80%
- **ptsd** — n=4 papers · extraction confidence: 75%
- **anosmia** — n=4 papers · extraction confidence: 70%
- **weakness** — n=4 papers · extraction confidence: 70%
- **headaches** — n=4 papers · extraction confidence: 70%
- **pasc_overall** — n=3 papers · extraction confidence: 86%
- **cough_overall** — n=3 papers · extraction confidence: 84%
- **post_covid_conditions_unvaccinated_pooled** — n=3 papers · extraction confidence: 84%

- **fatigue_long_covid** — n=3 papers · extraction confidence: 84%
- **headache_overall** — n=3 papers · extraction confidence: 83%
- **skin_rash** — n=3 papers · extraction confidence: 81%
- **smell_disorder** — n=3 papers · extraction confidence: 81%
- **taste_disorder** — n=3 papers · extraction confidence: 81%
- **voice_change** — n=3 papers · extraction confidence: 79%
- **obesity_comorbidity** — n=3 papers · extraction confidence: 77%
- **seizures** — n=3 papers · extraction confidence: 77%
- **breathlessness** — n=3 papers · extraction confidence: 75%
- **headache_long_covid** — n=3 papers · extraction confidence: 75%
- **joint_or_muscle_pain** — n=3 papers · extraction confidence: 66%
- **post_exertional_malaise** — n=3 papers · extraction confidence: 66%
- **difficulty_breathing** — n=3 papers · extraction confidence: 53%
- **memory_problems** — n=2 papers · extraction confidence: 88%
- **tiredness_or_fatigue** — n=2 papers · extraction confidence: 85%
- **difficulty_breathing_or_shortness_of_breath** — n=2 papers · extraction confidence: 85%
- **dyspnea_overall** — n=2 papers · extraction confidence: 85%
- **hair_loss_overall** — n=2 papers · extraction confidence: 85%
- **chest_pain_overall** — n=2 papers · extraction confidence: 85%
- **fatigue_1month** — n=2 papers · extraction confidence: 85%
- **no_symptoms** — n=2 papers · extraction confidence: 85%

- **respiratory_symptom_cluster** — n=2 papers · extraction confidence: 85%
- **fatigue_symptom_cluster** — n=2 papers · extraction confidence: 85%
- **long_covid_any_symptom** — n=2 papers · extraction confidence: 85%
- **any_symptom_infected** — n=2 papers · extraction confidence: 85%
- **cough_long_covid** — n=2 papers · extraction confidence: 85%
- **joint_pain_long_covid** — n=2 papers · extraction confidence: 85%
- **fatigue_recovered** — n=2 papers · extraction confidence: 85%
- **shortness_of_breath_recovered** — n=2 papers · extraction confidence: 85%
- **loss_of_smell_or_taste** — n=2 papers · extraction confidence: 85%
- **long_covid** — n=2 papers · extraction confidence: 84%
- **diarrhoea** — n=2 papers · extraction confidence: 83%
- **nausea** — n=2 papers · extraction confidence: 83%
- **chills** — n=2 papers · extraction confidence: 83%
- **palpitation** — n=2 papers · extraction confidence: 83%
- **appetite_loss** — n=2 papers · extraction confidence: 83%
- **persistent_cough** — n=2 papers · extraction confidence: 83%
- **digestive_issues** — n=2 papers · extraction confidence: 83%
- **long_covid_female** — n=2 papers · extraction confidence: 82%
- **long_covid_male** — n=2 papers · extraction confidence: 82%
- **migraine** — n=2 papers · extraction confidence: 82%
- **skin_rashes** — n=2 papers · extraction confidence: 82%

- **long_covid_self_reported** — n=2 papers · extraction confidence: 82%
- **post_covid_conditions_vaccinated_at_least_1_dose_pooled** — n=2 papers · extraction confidence: 82%
- **dyspnea_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **pulmonary_fibrosis_edema_inflammation_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **malaise_and_fatigue_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **edema_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **chest_pain_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **abnormal_heartbeat_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **cognitive_problems_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **joint_pain_excess_burden_ancestral_per_1000** — n=2 papers · extraction confidence: 82%
- **abdominal_pain_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **dyspnea_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **chest_pain_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **joint_pain_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **anemia_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **headache_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%

- **acute_pharyngitis_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **malaise_and_fatigue_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **dementia_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **pulmonary_fibrosis_edema_inflammation_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **edema_excess_burden_delta_per_1000** — n=2 papers · extraction confidence: 82%
- **fever_acute** — n=2 papers · extraction confidence: 82%
- **headache_acute** — n=2 papers · extraction confidence: 82%
- **fatigue_acute** — n=2 papers · extraction confidence: 82%
- **cough_acute** — n=2 papers · extraction confidence: 82%
- **sleep_disturbance_long_covid** — n=2 papers · extraction confidence: 82%
- **hair_loss_long_covid** — n=2 papers · extraction confidence: 82%
- **dyspnoea_acute** — n=2 papers · extraction confidence: 82%
- **dysgeusia** — n=2 papers · extraction confidence: 82%
- **alopecia** — n=2 papers · extraction confidence: 82%
- **at_least_one_symptom** — n=2 papers · extraction confidence: 80%
- **asthenia** — n=2 papers · extraction confidence: 80%
- **memory_impairment** — n=2 papers · extraction confidence: 80%
- **skin_symptoms** — n=2 papers · extraction confidence: 80%
- **sspd_ards** — n=2 papers · extraction confidence: 80%
- **memory_loss** — n=2 papers · extraction confidence: 80%
- **nervous_anxious** — n=2 papers · extraction confidence: 80%
- **peripheral_edema** — n=2 papers · extraction confidence: 80%
- **dyspnoea** — n=2 papers · extraction confidence: 80%

- **diabetes_comorbidity** — n=2 papers · extraction confidence: 80%
- **visual_impairment** — n=2 papers · extraction confidence: 80%
- **memory_deficits** — n=2 papers · extraction confidence: 80%
- **sweats** — n=2 papers · extraction confidence: 80%
- **mobility_decline** — n=2 papers · extraction confidence: 80%
- **cramps** — n=2 papers · extraction confidence: 80%
- **tremors** — n=2 papers · extraction confidence: 80%
- **poor_quality_of_life** — n=2 papers · extraction confidence: 80%
- **pain_or_discomfort** — n=2 papers · extraction confidence: 77%
- **lack_of_concentration** — n=2 papers · extraction confidence: 77%
- **neurologic_symptoms** — n=2 papers · extraction confidence: 77%
- **neuropathy** — n=2 papers · extraction confidence: 75%
- **constipation** — n=2 papers · extraction confidence: 75%
- **asthma_comorbidity** — n=2 papers · extraction confidence: 75%
- **probable_long_covid_among_covid19_patients** — n=2 papers · extraction confidence: 72%
- **diagnosed_long_covid_among_covid19_patients** — n=2 papers · extraction confidence: 72%

Speculative — sorted by
(n_papers, mean_extraction_confidence) descending:

- **cognitive_symptom_cluster_overall** — n=1 paper · extraction confidence: 93%
- **fatigue_symptom_cluster_overall** — n=1 paper · extraction confidence: 93%

- **respiratory_symptom_cluster_overall** — n=1 paper · extraction confidence: 93%
- **overall_pasc_computable_phenotype_or_u09.9** — n=1 paper · extraction confidence: 93%
- **neurologic_sequelae_3yr_cumulative_per1000_nonhosp** — n=1 paper · extraction confidence: 91%
- **cardiovascular_sequelae_3yr_cumulative_per1000_nonhosp** — n=1 paper · extraction confidence: 91%
- **pulmonary_sequelae_3yr_cumulative_per1000_nonhosp** — n=1 paper · extraction confidence: 91%
- **coagulation_hematologic_sequelae_3yr_cumulative_per1000_nonhosp** — n=1 paper · extraction confidence: 91%
- **mental_health_sequelae_3yr_cumulative_per1000_nonhosp** — n=1 paper · extraction confidence: 91%
- **cardiovascular_sequelae_3yr_cumulative_per1000_hosp** — n=1 paper · extraction confidence: 91%
- **mental_health_sequelae_3yr_cumulative_per1000_hosp** — n=1 paper · extraction confidence: 91%
- **neurologic_sequelae_3yr_cumulative_per1000_hosp** — n=1 paper · extraction confidence: 91%
- **coagulation_hematologic_sequelae_3yr_cumulative_per1000_hosp** — n=1 paper · extraction confidence: 91%
- **kidney_sequelae_3yr_cumulative_per1000_hosp** — n=1 paper · extraction confidence: 91%
- **dysrhythmia_control** — n=1 paper · extraction confidence: 91%
- **dysrhythmia_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **ischemic_heart_disease_control** — n=1 paper · extraction confidence: 91%
- **ischemic_heart_disease_nirmatrelvir** — n=1 paper · extraction confidence: 91%

- **pulmonary_embolism_control** — n=1 paper · extraction confidence: 91%
- **pulmonary_embolism_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **deep_vein_thrombosis_control** — n=1 paper · extraction confidence: 91%
- **deep_vein_thrombosis_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **fatigue_and_malaise_control** — n=1 paper · extraction confidence: 91%
- **fatigue_and_malaise_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **liver_disease_control** — n=1 paper · extraction confidence: 91%
- **liver_disease_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **acute_kidney_injury_control** — n=1 paper · extraction confidence: 91%
- **acute_kidney_injury_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **muscle_pain_control** — n=1 paper · extraction confidence: 91%
- **muscle_pain_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **diabetes_control** — n=1 paper · extraction confidence: 91%
- **diabetes_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **neurocognitive_impairment_control** — n=1 paper · extraction confidence: 91%
- **neurocognitive_impairment_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **dysautonomia_control** — n=1 paper · extraction confidence: 91%

- **dysautonomia_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **shortness_of_breath_control** — n=1 paper · extraction confidence: 91%
- **shortness_of_breath_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **cough_control** — n=1 paper · extraction confidence: 91%
- **cough_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **pcc_event_rate_control** — n=1 paper · extraction confidence: 91%
- **pcc_event_rate_nirmatrelvir** — n=1 paper · extraction confidence: 91%
- **any_financial_toxicity_long_covid** — n=1 paper · extraction confidence: 91%
- **any_financial_toxicity_covid19_only** — n=1 paper · extraction confidence: 91%
- **any_financial_toxicity_no_covid19** — n=1 paper · extraction confidence: 91%
- **any_financial_toxicity_overall** — n=1 paper · extraction confidence: 91%
- **difficulty_paying_medical_bills_long_covid** — n=1 paper · extraction confidence: 91%
- **cost_related_medication_nonadherence_long_covid** — n=1 paper · extraction confidence: 91%
- **delayed_healthcare_due_to_cost_long_covid** — n=1 paper · extraction confidence: 91%
- **not_obtained_healthcare_due_to_cost_long_covid** — n=1 paper · extraction confidence: 91%
- **fatigue_past_3_months_long_covid** — n=1 paper · extraction confidence: 91%
- **mild_severe_anxiety_long_covid** — n=1 paper · extraction confidence: 91%

- **mild_severe_depression_long_covid** — n=1 paper · extraction confidence: 91%
- **disability_long_covid** — n=1 paper · extraction confidence: 91%
- **food_insecurity_long_covid** — n=1 paper · extraction confidence: 91%
- **respiratory_conditions_long_covid** — n=1 paper · extraction confidence: 91%
- **immunosuppression_long_covid** — n=1 paper · extraction confidence: 91%
- **arthritis_long_covid** — n=1 paper · extraction confidence: 91%
- **post_covid_condition_overall** — n=1 paper · extraction confidence: 88%
- **pasc_among_smi_patients** — n=1 paper · extraction confidence: 88%
- **neurological_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **respiratory_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **circulatory_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **endocrine_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **musculoskeletal_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **digestive_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **blood_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **skin_related_pasc_among_pasc** — n=1 paper · extraction confidence: 88%
- **cough_incident** — n=1 paper · extraction confidence: 88%

- **dyspnea_shortness_of_breath_incident** — n=1 paper · extraction confidence: 88%
- **constipation_incident** — n=1 paper · extraction confidence: 88%
- **fatigue_incident** — n=1 paper · extraction confidence: 88%
- **chest_pain_incident** — n=1 paper · extraction confidence: 88%
- **anxiety_incident** — n=1 paper · extraction confidence: 88%
- **acute_kidney_injury_incident** — n=1 paper · extraction confidence: 88%
- **acute_coronary_disease_incident** — n=1 paper · extraction confidence: 88%
- **thromboembolism_incident** — n=1 paper · extraction confidence: 88%
- **hypoxemia_incident** — n=1 paper · extraction confidence: 88%
- **stroke_incident** — n=1 paper · extraction confidence: 88%
- **myocarditis_incident** — n=1 paper · extraction confidence: 88%
- **any_incident_pasc** — n=1 paper · extraction confidence: 88%
- **cardiovascular_system_pasc** — n=1 paper · extraction confidence: 88%
- **pulmonary_system_pasc** — n=1 paper · extraction confidence: 88%
- **gastrointestinal_system_pasc** — n=1 paper · extraction confidence: 88%
- **mental_health_pasc** — n=1 paper · extraction confidence: 88%
- **renal_pasc** — n=1 paper · extraction confidence: 88%
- **joint_pain_baseline_prevalent** — n=1 paper · extraction confidence: 88%

- **hyperlipidemia_baseline_prevalent** — n=1 paper · extraction confidence: 88%
- **obesity_baseline_prevalent** — n=1 paper · extraction confidence: 88%
- **cough_baseline_prevalent** — n=1 paper · extraction confidence: 88%
- **anxiety_baseline_prevalent** — n=1 paper · extraction confidence: 88%
- **constipation_baseline_prevalent** — n=1 paper · extraction confidence: 88%
- **symptoms_not_limiting_activities** — n=1 paper · extraction confidence: 88%
- **heart_palpitations_or_chest_pain** — n=1 paper · extraction confidence: 88%
- **dizziness_on_standing** — n=1 paper · extraction confidence: 88%
- **depression_anxiety_or_mood_changes** — n=1 paper · extraction confidence: 88%
- **symptoms_worsening_after_physical_or_mental_activities** — n=1 paper · extraction confidence: 88%
- **pasc_at_180_days_molnupiravir_group** — n=1 paper · extraction confidence: 88%
- **pasc_at_180_days_no_treatment_group** — n=1 paper · extraction confidence: 88%
- **post_acute_death_molnupiravir_group** — n=1 paper · extraction confidence: 88%
- **post_acute_death_no_treatment_group** — n=1 paper · extraction confidence: 88%
- **post_acute_hospital_admission_molnupiravir_group** — n=1 paper · extraction confidence: 88%
- **post_acute_hospital_admission_no_treatment_group** — n=1 paper · extraction confidence: 88%

- **any_postacute_symptom_all_covid19** — n=1 paper · extraction confidence: 88%
- **any_postacute_symptom_noninfected** — n=1 paper · extraction confidence: 88%
- **any_postacute_symptom_hospitalized_covid19** — n=1 paper · extraction confidence: 88%
- **any_postacute_diagnosis_all_covid19** — n=1 paper · extraction confidence: 88%
- **any_postacute_diagnosis_noninfected** — n=1 paper · extraction confidence: 88%
- **any_postacute_diagnosis_hospitalized_covid19** — n=1 paper · extraction confidence: 88%
- **any_cardiovascular_diagnosis_hospitalized** — n=1 paper · extraction confidence: 88%
- **any_kidney_diagnosis_hospitalized** — n=1 paper · extraction confidence: 88%
- **pasc_u09.9_overall** — n=1 paper · extraction confidence: 88%
- **pasc_u09.9_healthy_weight** — n=1 paper · extraction confidence: 88%
- **pasc_u09.9_overweight** — n=1 paper · extraction confidence: 88%
- **pasc_u09.9_obesity** — n=1 paper · extraction confidence: 88%
- **pasc_u09.9_severe_obesity** — n=1 paper · extraction confidence: 88%
- **any_pasc_symptoms_conditions_overall** — n=1 paper · extraction confidence: 88%
- **any_pasc_symptoms_conditions_healthy_weight** — n=1 paper · extraction confidence: 88%
- **any_pasc_symptoms_conditions_overweight** — n=1 paper · extraction confidence: 88%

- **any_pasc_symptoms_conditions_obesity** — n=1 paper · extraction confidence: 88%
- **any_pasc_symptoms_conditions_severe_obesity** — n=1 paper · extraction confidence: 88%
- **memory_loss_overall** — n=1 paper · extraction confidence: 88%
- **other_symptoms_overall** — n=1 paper · extraction confidence: 88%
- **muscle_pain_overall** — n=1 paper · extraction confidence: 88%
- **concentration_difficulties_overall** — n=1 paper · extraction confidence: 88%
- **anxiety_overall** — n=1 paper · extraction confidence: 88%
- **sleeping_difficulties_overall** — n=1 paper · extraction confidence: 88%
- **depression_overall** — n=1 paper · extraction confidence: 88%
- **taste_loss_overall** — n=1 paper · extraction confidence: 88%
- **smell_loss_overall** — n=1 paper · extraction confidence: 88%
- **dizziness_overall** — n=1 paper · extraction confidence: 88%
- **gastrointestinal_symptoms_overall** — n=1 paper · extraction confidence: 88%
- **visual_disturbances_overall** — n=1 paper · extraction confidence: 88%
- **tinnitus_overall** — n=1 paper · extraction confidence: 88%
- **palpitations_overall** — n=1 paper · extraction confidence: 88%
- **nausea_overall** — n=1 paper · extraction confidence: 88%
- **allergy_overall** — n=1 paper · extraction confidence: 88%
- **memory_loss_cluster1** — n=1 paper · extraction confidence: 88%

- **concentration_difficulties_cluster1** — n=1 paper · extraction confidence: 88%
- **fatigue_cluster1** — n=1 paper · extraction confidence: 88%
- **fatigue_cluster2** — n=1 paper · extraction confidence: 88%
- **other_symptoms_cluster2** — n=1 paper · extraction confidence: 88%
- **cough_cluster2** — n=1 paper · extraction confidence: 88%
- **fatigue_cluster3** — n=1 paper · extraction confidence: 88%
- **memory_loss_cluster3** — n=1 paper · extraction confidence: 88%
- **anxiety_cluster3** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19pos_jones_def** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19pos_pagen_def** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19pos_sudre_def** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19pos_thaweethai_def** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19pos_yoo_def** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19neg_range_low** — n=1 paper · extraction confidence: 88%
- **long_covid_3mo_covid19neg_range_high** — n=1 paper · extraction confidence: 88%
- **long_covid_6mo_covid19pos_range_low** — n=1 paper · extraction confidence: 88%
- **long_covid_6mo_covid19pos_range_high** — n=1 paper · extraction confidence: 88%
- **long_covid_6mo_covid19neg_range_low** — n=1 paper · extraction confidence: 88%
- **long_covid_6mo_covid19neg_range_high** — n=1 paper · extraction confidence: 88%

- **self_reported_long_covid_covid19pos** — n=1 paper · extraction confidence: 88%
- **self_reported_long_covid_covid19neg** — n=1 paper · extraction confidence: 88%
- **dyspnea_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **fatigue_malaise_weakness_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **cough_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **chest_pain_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **palpitations_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **headache_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **muscle_joint_pain_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **memory_problem_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **cognitive_impairment_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **sleep_disturbance_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **loss_of_taste_smell_long_covid_outpatient** — n=1 paper · extraction confidence: 88%
- **dyspnea_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **fatigue_malaise_weakness_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **cough_long_covid_inpatient** — n=1 paper · extraction confidence: 88%

- **chest_pain_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **palpitations_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **headache_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **muscle_joint_pain_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **memory_problem_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **cognitive_impairment_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **sleep_disturbance_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **loss_of_taste_smell_long_covid_inpatient** — n=1 paper · extraction confidence: 88%
- **long_covid_outpatient_symptom_based_16_6_pct** — n=1 paper · extraction confidence: 88%
- **long_covid_inpatient_symptom_based_29_2_pct** — n=1 paper · extraction confidence: 88%
- **long_flu_outpatient_symptom_based_17_0_pct** — n=1 paper · extraction confidence: 88%
- **long_flu_inpatient_symptom_based_24_6_pct** — n=1 paper · extraction confidence: 88%
- **long_covid_code_based_outpatient_0_49_pct** — n=1 paper · extraction confidence: 88%
- **long_covid_code_based_inpatient_2_6_pct** — n=1 paper · extraction confidence: 88%
- **pain_prior_to_long_covid_diagnosis** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_prior_to_long_covid_diagnosis** — n=1 paper · extraction confidence: 88%

- **fatigue_prior_to_long_covid_diagnosis** — n=1 paper · extraction confidence: 88%
- **at_least_one_long_covid_symptom_1month** — n=1 paper · extraction confidence: 88%
- **at_least_one_long_covid_symptom_3month** — n=1 paper · extraction confidence: 88%
- **at_least_one_long_covid_symptom_6month** — n=1 paper · extraction confidence: 88%
- **at_least_one_long_covid_symptom_9_12month** — n=1 paper · extraction confidence: 88%
- **fatigue_3month** — n=1 paper · extraction confidence: 88%
- **fatigue_6month** — n=1 paper · extraction confidence: 88%
- **fatigue_9_12month** — n=1 paper · extraction confidence: 88%
- **anxiety_1month** — n=1 paper · extraction confidence: 88%
- **anxiety_3month** — n=1 paper · extraction confidence: 88%
- **anxiety_6month** — n=1 paper · extraction confidence: 88%
- **anxiety_9_12month** — n=1 paper · extraction confidence: 88%
- **breathlessness_1month** — n=1 paper · extraction confidence: 88%
- **breathlessness_3month** — n=1 paper · extraction confidence: 88%
- **breathlessness_6month** — n=1 paper · extraction confidence: 88%
- **breathlessness_9_12month** — n=1 paper · extraction confidence: 88%
- **memory_concentration_problems_1month** — n=1 paper · extraction confidence: 88%
- **memory_concentration_problems_3month** — n=1 paper · extraction confidence: 88%
- **memory_concentration_problems_6month** — n=1 paper · extraction confidence: 88%

- **memory_concentration_problems_9_12month** — n=1 paper · extraction confidence: 88%
- **chest_pain_1month** — n=1 paper · extraction confidence: 88%
- **chest_pain_3month** — n=1 paper · extraction confidence: 88%
- **chest_pain_6month** — n=1 paper · extraction confidence: 88%
- **chest_pain_9_12month** — n=1 paper · extraction confidence: 88%
- **palpitations_1month** — n=1 paper · extraction confidence: 88%
- **palpitations_3month** — n=1 paper · extraction confidence: 88%
- **palpitations_6month** — n=1 paper · extraction confidence: 88%
- **palpitations_9_12month** — n=1 paper · extraction confidence: 88%
- **myalgia_1month** — n=1 paper · extraction confidence: 88%
- **anosmia_1month** — n=1 paper · extraction confidence: 88%
- **ageusia_1month** — n=1 paper · extraction confidence: 88%
- **eq5d_usual_activities_difficulty_within_6months** — n=1 paper · extraction confidence: 88%
- **eq5d_anxiety_depression_within_6months** — n=1 paper · extraction confidence: 88%
- **eq5d_mobility_problems_within_6months** — n=1 paper · extraction confidence: 88%
- **pcc_diagnosed_proportion** — n=1 paper · extraction confidence: 88%
- **pvfs_diagnosed_proportion** — n=1 paper · extraction confidence: 88%
- **ed_diagnosed_proportion** — n=1 paper · extraction confidence: 88%

- **pcc_with_no_prior_symptom_based_visits** — n=1 paper · extraction confidence: 88%
- **pvfs_with_no_prior_symptom_based_visits** — n=1 paper · extraction confidence: 88%
- **ed_with_no_prior_symptom_based_visits** — n=1 paper · extraction confidence: 88%
- **symptom_based_visits_proportion_of_all_visits** — n=1 paper · extraction confidence: 88%
- **long covid (lc_{ri}≥11) at 3 months** — n=1 paper · extraction confidence: 88%
- **long covid (lc_{ri}≥11) at 15 months** — n=1 paper · extraction confidence: 88%
- **long covid in uninfected cohort at enrollment (after imputation)** — n=1 paper · extraction confidence: 88%
- **long covid in uninfected cohort at 3 months (after imputation)** — n=1 paper · extraction confidence: 88%
- **long covid in uninfected cohort at 6 months (after imputation)** — n=1 paper · extraction confidence: 88%
- **long covid in uninfected cohort at 9 months (after imputation)** — n=1 paper · extraction confidence: 88%
- **long covid in uninfected cohort at 12 months (after imputation)** — n=1 paper · extraction confidence: 88%
- **pasc_index_ge12_among_infected** — n=1 paper · extraction confidence: 88%
- **pasc_index_zero_among_infected** — n=1 paper · extraction confidence: 88%
- **severe_fatigue_current_lc** — n=1 paper · extraction confidence: 88%
- **waxing_and_waning_trajectory_current_lc** — n=1 paper · extraction confidence: 88%
- **improved_trajectory_current_lc** — n=1 paper · extraction confidence: 88%

- **unchanged_trajectory_current_lc** — n=1 paper · extraction confidence: 88%
- **worsened_trajectory_current_lc** — n=1 paper · extraction confidence: 88%
- **fatigue_3months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **fatigue_3months_control** — n=1 paper · extraction confidence: 88%
- **fatigue_6months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **fatigue_6months_control** — n=1 paper · extraction confidence: 88%
- **cough_3months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **cough_3months_control** — n=1 paper · extraction confidence: 88%
- **forgetfulness_6months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **forgetfulness_6months_control** — n=1 paper · extraction confidence: 88%
- **pcc_3months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **pcc_3months_control** — n=1 paper · extraction confidence: 88%
- **pcc_6months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **pcc_6months_control** — n=1 paper · extraction confidence: 88%
- **anxiety_3months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **anxiety_3months_control** — n=1 paper · extraction confidence: 88%

- **negative_emotions_6months_nirmatrelvir** — n=1 paper · extraction confidence: 88%
- **negative_emotions_6months_control** — n=1 paper · extraction confidence: 88%
- **ever_long_covid** — n=1 paper · extraction confidence: 88%
- **long_covid_current_symptoms** — n=1 paper · extraction confidence: 88%
- **long_covid_substantially_limiting_daily_activities** — n=1 paper · extraction confidence: 88%
- **severe_covid_symptoms_among_long_covid_now** — n=1 paper · extraction confidence: 88%
- **severe_covid_symptoms_among_long_covid_past** — n=1 paper · extraction confidence: 88%
- **severe_covid_symptoms_among_short_covid** — n=1 paper · extraction confidence: 88%
- **long_covid_women** — n=1 paper · extraction confidence: 88%
- **long_covid_men** — n=1 paper · extraction confidence: 88%
- **long_covid_hawaii_lowest** — n=1 paper · extraction confidence: 88%
- **long_covid_west_virginia_highest** — n=1 paper · extraction confidence: 88%
- **long_covid_overall_unvaccinated** — n=1 paper · extraction confidence: 88%
- **long_covid_overall_primary_series** — n=1 paper · extraction confidence: 88%
- **long_covid_immunocompetent_unvaccinated** — n=1 paper · extraction confidence: 88%
- **long_covid_immunocompetent_primary_series** — n=1 paper · extraction confidence: 88%
- **long_covid_immunocompromised_unvaccinated** — n=1 paper · extraction confidence: 88%

- **long_covid_immunocompromised_primary_series** — n=1 paper · extraction confidence: 88%
- **post_covid_fatigue** — n=1 paper · extraction confidence: 88%
- **post_covid_respiratory_complaints** — n=1 paper · extraction confidence: 88%
- **long_covid_any_measure** — n=1 paper · extraction confidence: 88%
- **long_covid_clinical_codes** — n=1 paper · extraction confidence: 88%
- **long_covid_free_text_primary_care** — n=1 paper · extraction confidence: 88%
- **long_covid_free_text_sick_notes** — n=1 paper · extraction confidence: 88%
- **long_covid_operational_definition** — n=1 paper · extraction confidence: 88%
- **housing_insecurity_any_long_covid** — n=1 paper · extraction confidence: 88%
- **housing_insecurity_any_no_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_household_expenses_overall** — n=1 paper · extraction confidence: 88%
- **behind_on_housing_payments_overall** — n=1 paper · extraction confidence: 88%
- **likely_eviction_or_foreclosure_overall** — n=1 paper · extraction confidence: 88%
- **current_long_covid_symptoms_among_long_covid** — n=1 paper · extraction confidence: 88%
- **symptom_impact_day_to_day_any_among_current_symptomatic** — n=1 paper · extraction confidence: 88%
- **cognitive_limitation_moderate_or_severe_long_covid** — n=1 paper · extraction confidence: 88%

- **difficulty_expenses_very_difficult_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_expenses_very_difficult_no_long_covid** — n=1 paper · extraction confidence: 88%
- **smell_or_taste_problems** — n=1 paper · extraction confidence: 88%
- **confusion_disorientation_brain_fog** — n=1 paper · extraction confidence: 88%
- **memory_issues** — n=1 paper · extraction confidence: 88%
- **overall_pasc_nonhospitalized_cumulative_per1000** — n=1 paper · extraction confidence: 88%
- **overall_pasc_hospitalized_cumulative_per1000** — n=1 paper · extraction confidence: 88%
- **overall_pasc_overall_cohort_cumulative_per1000** — n=1 paper · extraction confidence: 88%
- **pasc_score_ge12_normal_weight** — n=1 paper · extraction confidence: 88%
- **pasc_score_ge12_overweight** — n=1 paper · extraction confidence: 88%
- **pasc_score_ge12_obesity** — n=1 paper · extraction confidence: 88%
- **high_risk_moderate_severe_insomnia_normal_weight** — n=1 paper · extraction confidence: 88%
- **high_risk_moderate_severe_insomnia_overweight** — n=1 paper · extraction confidence: 88%
- **high_risk_moderate_severe_insomnia_obesity** — n=1 paper · extraction confidence: 88%
- **high_risk_osa_normal_weight** — n=1 paper · extraction confidence: 88%
- **high_risk_osa_overweight** — n=1 paper · extraction confidence: 88%
- **high_risk_osa_obesity** — n=1 paper · extraction confidence: 88%

- **short_sleep_lt6h_normal_weight** — n=1 paper · extraction confidence: 88%
- **short_sleep_lt6h_overweight** — n=1 paper · extraction confidence: 88%
- **short_sleep_lt6h_obesity** — n=1 paper · extraction confidence: 88%
- **long_sleep_gt9h_normal_weight** — n=1 paper · extraction confidence: 88%
- **long_sleep_gt9h_overweight** — n=1 paper · extraction confidence: 88%
- **long_sleep_gt9h_obesity** — n=1 paper · extraction confidence: 88%
- **anxiety_disorder** — n=1 paper · extraction confidence: 88%
- **sleep_problems** — n=1 paper · extraction confidence: 88%
- **smell_problems** — n=1 paper · extraction confidence: 88%
- **taste_problems** — n=1 paper · extraction confidence: 88%
- **current_long_covid_food_insecure** — n=1 paper · extraction confidence: 88%
- **current_long_covid_food_secure** — n=1 paper · extraction confidence: 88%
- **pasc_multiple_infections** — n=1 paper · extraction confidence: 88%
- **pasc_single_infection** — n=1 paper · extraction confidence: 88%
- **pasc_severe_covid** — n=1 paper · extraction confidence: 88%
- **pasc_moderate_covid** — n=1 paper · extraction confidence: 88%
- **pasc_mild_asymptomatic_covid** — n=1 paper · extraction confidence: 88%
- **pasc_unvaccinated_at_first_infection** — n=1 paper · extraction confidence: 88%

- **current_long_covid** — n=1 paper · extraction confidence: 88%
- **never_long_covid** — n=1 paper · extraction confidence: 88%
- **resolved_long_covid** — n=1 paper · extraction confidence: 88%
- **current_lc_waxed_and_waned_symptoms** — n=1 paper · extraction confidence: 88%
- **current_lc_improved_symptoms** — n=1 paper · extraction confidence: 88%
- **current_lc_no_change_symptoms** — n=1 paper · extraction confidence: 88%
- **current_lc_worsened_symptoms** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_overall_2022** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_overall_2023** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_overall_2024** — n=1 paper · extraction confidence: 88%
- **current_long_covid_overall_2022** — n=1 paper · extraction confidence: 88%
- **current_long_covid_overall_2023** — n=1 paper · extraction confidence: 88%
- **current_long_covid_overall_2024** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_prior_infection_2022** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_prior_infection_2023** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_prior_infection_2024** — n=1 paper · extraction confidence: 88%
- **current_long_covid_prior_infection_2022** — n=1 paper · extraction confidence: 88%

- **current_long_covid_prior_infection_2023** — n=1 paper · extraction confidence: 88%
- **current_long_covid_prior_infection_2024** — n=1 paper · extraction confidence: 88%
- **significant_activity_limitation_current_lc_2023_2024_combined** — n=1 paper · extraction confidence: 88%
- **any_activity_limitation_current_lc_2023_2024_combined** — n=1 paper · extraction confidence: 88%
- **pneumonia_readmission_reinfection** — n=1 paper · extraction confidence: 88%
- **pneumonia_readmission_firsttime** — n=1 paper · extraction confidence: 88%
- **tiredness_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **tiredness_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **headaches_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **headaches_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **dizziness_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **dizziness_test_negative_6mo** — n=1 paper · extraction confidence: 88%

- **any_symptom_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **any_symptom_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **three_plus_symptoms_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **three_plus_symptoms_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **long_covid_delphi_test_positive_6mo** — n=1 paper · extraction confidence: 88%
- **long_covid_delphi_test_negative_6mo** — n=1 paper · extraction confidence: 88%
- **pasc_overall_computable_phenotype** — n=1 paper · extraction confidence: 88%
- **cognitive_symptom_cluster** — n=1 paper · extraction confidence: 88%
- **pasc_unvaccinated_pre_delta_1year** — n=1 paper · extraction confidence: 88%
- **pasc_unvaccinated_delta_1year** — n=1 paper · extraction confidence: 88%
- **pasc_unvaccinated_omicron_1year** — n=1 paper · extraction confidence: 88%
- **pasc_vaccinated_delta_1year** — n=1 paper · extraction confidence: 88%
- **pasc_vaccinated_omicron_1year** — n=1 paper · extraction confidence: 88%
- **long_covid_diagnosis** — n=1 paper · extraction confidence: 88%
- **unusual_fatigue_tiredness_at_baseline_positive** — n=1 paper · extraction confidence: 88%
- **unusual_fatigue_tiredness_at_3months_positive** — n=1 paper · extraction confidence: 88%

- **unusual_fatigue_tiredness_at_6months_positive** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_at_baseline_positive** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_at_3months_positive** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_at_6months_positive** — n=1 paper · extraction confidence: 88%
- **chills_fever_myalgia_cough_sore_throat_at_baseline_positive_range_** — n=1 paper · extraction confidence: 88%
- **chills_fever_myalgia_cough_sore_throat_at_baseline_positive_range_** — n=1 paper · extraction confidence: 88%
- **chills_fever_myalgia_cough_sore_throat_at_3and6months_positive** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_at_testing_positive** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_at_3months_positive** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_at_6months_positive** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_at_testing_negative** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_at_3months_negative** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_at_6months_negative** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_all_three_timepoints_positive** — n=1 paper · extraction confidence: 88%
- **never_any_symptom_positive** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_all_three_timepoints_negative** — n=1 paper · extraction confidence: 88%

- **never_any_symptom_negative** — n=1 paper · extraction confidence: 88%
- **fatigue_females_3months** — n=1 paper · extraction confidence: 88%
- **fatigue_males_3months** — n=1 paper · extraction confidence: 88%
- **fatigue_females_12months** — n=1 paper · extraction confidence: 88%
- **hair_loss_females_3months** — n=1 paper · extraction confidence: 88%
- **hair_loss_males_3months** — n=1 paper · extraction confidence: 88%
- **memory_loss_second_most_common_females** — n=1 paper · extraction confidence: 88%
- **dyspnea_third_most_prevalent_females** — n=1 paper · extraction confidence: 88%
- **musculoskeletal_long_covid** — n=1 paper · extraction confidence: 88%
- **cardiovascular_long_covid** — n=1 paper · extraction confidence: 88%
- **respiratory_long_covid** — n=1 paper · extraction confidence: 88%
- **stomach_ache** — n=1 paper · extraction confidence: 88%
- **vaginal_discharge** — n=1 paper · extraction confidence: 88%
- **menorrhagia** — n=1 paper · extraction confidence: 88%
- **pins_and_needles** — n=1 paper · extraction confidence: 88%
- **erectile_dysfunction** — n=1 paper · extraction confidence: 88%
- **tinnitus** — n=1 paper · extraction confidence: 88%
- **blocked_nose** — n=1 paper · extraction confidence: 88%
- **eye_irritation** — n=1 paper · extraction confidence: 88%

- **walk_disturbances** — n=1 paper · extraction confidence: 88%
- **change_loss_of_smell** — n=1 paper · extraction confidence: 88%
- **change_loss_of_taste** — n=1 paper · extraction confidence: 88%
- **reduced_libido** — n=1 paper · extraction confidence: 88%
- **ejaculation_difficulties** — n=1 paper · extraction confidence: 88%
- **body_pain** — n=1 paper · extraction confidence: 88%
- **learning_difficulty_long_covid** — n=1 paper · extraction confidence: 88%
- **learning_difficulty_no_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_concentrating_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_concentrating_no_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_remembering_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_remembering_no_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_controlling_behavior_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_controlling_behavior_no_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_making_friends_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_making_friends_no_long_covid** — n=1 paper · extraction confidence: 88%
- **difficulty_accepting_change_long_covid** — n=1 paper · extraction confidence: 88%

- **difficulty_accepting_change_no_long_covid** — n=1 paper · extraction confidence: 88%
- **weekly_daily_anxiety_long_covid** — n=1 paper · extraction confidence: 88%
- **weekly_daily_anxiety_no_long_covid** — n=1 paper · extraction confidence: 88%
- **weekly_daily_depression_long_covid** — n=1 paper · extraction confidence: 88%
- **weekly_daily_depression_no_long_covid** — n=1 paper · extraction confidence: 88%
- **chronic_absenteeism_long_covid_full_sample** — n=1 paper · extraction confidence: 88%
- **chronic_absenteeism_no_long_covid_full_sample** — n=1 paper · extraction confidence: 88%
- **chronic_absenteeism_long_covid_covid_subsample** — n=1 paper · extraction confidence: 88%
- **chronic_absenteeism_no_long_covid_covid_subsample** — n=1 paper · extraction confidence: 88%
- **missed_more_than_30_days_long_covid** — n=1 paper · extraction confidence: 88%
- **current_long_covid_among_ever_long_covid** — n=1 paper · extraction confidence: 88%
- **prescription_medication_mental_health_long_covid** — n=1 paper · extraction confidence: 88%
- **prescription_medication_mental_health_no_long_covid** — n=1 paper · extraction confidence: 88%
- **chronic_health_condition_long_covid** — n=1 paper · extraction confidence: 88%
- **chronic_health_condition_no_long_covid** — n=1 paper · extraction confidence: 88%
- **neurodevelopmental_condition_long_covid** — n=1 paper · extraction confidence: 88%

- **neurodevelopmental_condition_no_long_covid** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_overall** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_overall** — n=1 paper · extraction confidence: 88%
- **current_long_covid_overall** — n=1 paper · extraction confidence: 88%
- **diagnosed_long_covid_overall** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_hispanic** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_white** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_black** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_aian** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_asian** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_arab_mena** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_hispanic** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_white** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_black** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_aian** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_asian** — n=1 paper · extraction confidence: 88%

- **ever_long_covid_arab_mena** — n=1 paper · extraction confidence: 88%
- **current_long_covid_black** — n=1 paper · extraction confidence: 88%
- **diagnosed_long_covid_black** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_female** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_male** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_female** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_male** — n=1 paper · extraction confidence: 88%
- **current_long_covid_female** — n=1 paper · extraction confidence: 88%
- **current_long_covid_male** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_low_income** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_mid_income** — n=1 paper · extraction confidence: 88%
- **90_day_symptoms_high_income** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_low_income** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_mid_income** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_high_income** — n=1 paper · extraction confidence: 88%
- **current_long_covid_black_female** — n=1 paper · extraction confidence: 88%

- **current_long_covid_black_male** — n=1 paper · extraction confidence: 88%
- **ever_long_covid_black_female** — n=1 paper · extraction confidence: 88%
- **chronic_pain_overall** — n=1 paper · extraction confidence: 88%
- **chronic_pain_no_infection_group** — n=1 paper · extraction confidence: 88%
- **chronic_pain_asymptomatic_mild_group** — n=1 paper · extraction confidence: 88%
- **chronic_pain_moderate_severe_group** — n=1 paper · extraction confidence: 88%
- **any_long_covid_features_covid_with_copc_prematch** — n=1 paper · extraction confidence: 88%
- **any_long_covid_features_covid_without_copc_prematch** — n=1 paper · extraction confidence: 88%
- **any_long_covid_features_influenza_with_copc_prematch** — n=1 paper · extraction confidence: 88%
- **any_long_covid_features_influenza_without_copc_prematch** — n=1 paper · extraction confidence: 88%
- **any_long_covid_features_noninfected_with_copc_prematch** — n=1 paper · extraction confidence: 88%
- **any_long_covid_features_noninfected_without_copc_prematch** — n=1 paper · extraction confidence: 88%
- **chest_throat_pain_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **chest_throat_pain_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **abnormal_breathing_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **abnormal_breathing_covid_without_copc** — n=1 paper · extraction confidence: 88%

- **abdominal_symptoms_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **abdominal_symptoms_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **fatigue_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **fatigue_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **anxiety_depression_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **anxiety_depression_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **pain_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **pain_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **headache_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **headache_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **cognitive_symptoms_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **cognitive_symptoms_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **myalgia_covid_with_copc** — n=1 paper · extraction confidence: 88%
- **myalgia_covid_without_copc** — n=1 paper · extraction confidence: 88%
- **post_covid_conditions_fully_vaccinated_pooled** — n=1 paper · extraction confidence: 88%
- **uveitis_any_unvaccinated_long_covid_1yr** — n=1 paper · extraction confidence: 88%

- **uveitis_any_unvaccinated_long_covid_2yr** — n=1 paper · extraction confidence: 88%
- **uveitis_any_unvaccinated_no_long_covid_1yr** — n=1 paper · extraction confidence: 88%
- **uveitis_any_unvaccinated_no_long_covid_2yr** — n=1 paper · extraction confidence: 88%
- **anterior_uveitis_unvaccinated_long_covid_1yr** — n=1 paper · extraction confidence: 88%
- **anterior_uveitis_unvaccinated_long_covid_2yr** — n=1 paper · extraction confidence: 88%
- **uveitis_any_vaccinated_long_covid_1yr** — n=1 paper · extraction confidence: 88%
- **uveitis_any_vaccinated_long_covid_2yr** — n=1 paper · extraction confidence: 88%
- **uveitis_any_vaccinated_no_long_covid_1yr** — n=1 paper · extraction confidence: 88%
- **uveitis_any_vaccinated_no_long_covid_2yr** — n=1 paper · extraction confidence: 88%
- **mild_fatigue_subtype_infected** — n=1 paper · extraction confidence: 88%
- **ent_subtype_infected** — n=1 paper · extraction confidence: 88%
- **neurological_subtype_infected** — n=1 paper · extraction confidence: 88%
- **cardiopulmonary_subtype_infected** — n=1 paper · extraction confidence: 88%
- **mild_fatigue_subtype_uninfected** — n=1 paper · extraction confidence: 88%
- **ent_subtype_uninfected** — n=1 paper · extraction confidence: 88%
- **neurological_subtype_uninfected** — n=1 paper · extraction confidence: 88%

- **cardiopulmonary_subtype_uninfected** — n=1 paper · extraction confidence: 88%
- **any_specific_symptom_infected** — n=1 paper · extraction confidence: 88%
- **any_specific_symptom_uninfected** — n=1 paper · extraction confidence: 88%
- **cough_congestion_shortness_of_breath** — n=1 paper · extraction confidence: 88%
- **body_aches_joint_pain** — n=1 paper · extraction confidence: 88%
- **chest_abdominal_pain** — n=1 paper · extraction confidence: 88%
- **diarrhoea_nausea** — n=1 paper · extraction confidence: 88%
- **confusion** — n=1 paper · extraction confidence: 88%
- **at_least_one_symptom_long_covid** — n=1 paper · extraction confidence: 88%
- **depression_or_anxiety** — n=1 paper · extraction confidence: 88%
- **anosmia_dysgeusia_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **anosmia_dysgeusia_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **cognitive_impairment_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **cognitive_impairment_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **dyspnoea_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **dyspnoea_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **weakness_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%

- **weakness_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **palpitations_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **palpitations_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **hair_loss_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **chest_pain_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **myalgia_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **cough_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **streptococcal_tonsillitis_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **streptococcal_tonsillitis_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **dizziness_early_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **dizziness_late_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **weakness_yearly_burden_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **dyspnoea_yearly_burden_risk_difference_per_10000** — n=1 paper · extraction confidence: 88%
- **very_high_somatic_symptoms_o2_therapy** — n=1 paper · extraction confidence: 88%
- **very_high_somatic_symptoms_no_covid** — n=1 paper · extraction confidence: 88%
- **very_high_somatic_symptoms_covid_no_o2** — n=1 paper · extraction confidence: 88%

- **chest_pain_shortness_of_breath_o2_therapy** — n=1 paper · extraction confidence: 88%
- **stomach_bowel_problems_o2_therapy** — n=1 paper · extraction confidence: 88%
- **hearing_loss_o2_therapy** — n=1 paper · extraction confidence: 88%
- **trouble_sleeping_o2_therapy** — n=1 paper · extraction confidence: 88%
- **dizziness_o2_therapy** — n=1 paper · extraction confidence: 88%
- **headaches_o2_therapy** — n=1 paper · extraction confidence: 88%
- **olfactory_disorder_o2_therapy** — n=1 paper · extraction confidence: 88%
- **fatigue_o2_therapy_10item** — n=1 paper · extraction confidence: 88%
- **cough_o2_therapy** — n=1 paper · extraction confidence: 88%
- **pain_arms_legs_joints_o2_therapy** — n=1 paper · extraction confidence: 88%
- **alopecia_o2_therapy** — n=1 paper · extraction confidence: 88%
- **reduced_libido_o2_therapy** — n=1 paper · extraction confidence: 88%
- **decreased_concentration_o2_therapy** — n=1 paper · extraction confidence: 88%
- **taste_disorder_o2_therapy** — n=1 paper · extraction confidence: 88%
- **memory_deficits_o2_therapy** — n=1 paper · extraction confidence: 88%
- **back_pain_o2_therapy** — n=1 paper · extraction confidence: 88%

- **feeling_tired_low_energy_o2_therapy** — n=1 paper · extraction confidence: 88%
- **fever_slight_fever_o2_therapy** — n=1 paper · extraction confidence: 88%
- **long_covid_prevalence_depressive_sample** — n=1 paper · extraction confidence: 88%
- **long_covid_prevalence_anxiety_sample** — n=1 paper · extraction confidence: 88%
- **depressive_symptoms_at_followup2_overall** — n=1 paper · extraction confidence: 88%
- **anxiety_symptoms_at_followup2_overall** — n=1 paper · extraction confidence: 88%
- **depressive_symptoms_at_followup2_long_covid** — n=1 paper · extraction confidence: 88%
- **depressive_symptoms_at_followup2_no_long_covid** — n=1 paper · extraction confidence: 88%
- **anxiety_symptoms_at_followup2_long_covid** — n=1 paper · extraction confidence: 88%
- **anxiety_symptoms_at_followup2_no_long_covid** — n=1 paper · extraction confidence: 88%
- **depressive_symptoms_at_followup1_long_covid** — n=1 paper · extraction confidence: 88%
- **depressive_symptoms_at_followup1_no_long_covid** — n=1 paper · extraction confidence: 88%
- **anxiety_symptoms_at_followup1_long_covid** — n=1 paper · extraction confidence: 88%
- **anxiety_symptoms_at_followup1_no_long_covid** — n=1 paper · extraction confidence: 88%
- **tiredness_fatigue** — n=1 paper · extraction confidence: 88%
- **depression_anxiety_mood_changes** — n=1 paper · extraction confidence: 88%
- **symptoms_worse_after_physical_or_mental_activity** — n=1 paper · extraction confidence: 88%

- **long_covid_unvaccinated_infected_overall** — n=1 paper · extraction confidence: 88%
- **long_covid_3plus_doses_pre_infection_overall** — n=1 paper · extraction confidence: 88%
- **long_covid_1_dose_pre_infection_overall** — n=1 paper · extraction confidence: 88%
- **long_covid_2_doses_pre_infection_overall** — n=1 paper · extraction confidence: 88%
- **long_covid_vaccinated_after_infection_overall** — n=1 paper · extraction confidence: 88%
- **long_covid_compatible_symptoms_uninfected** — n=1 paper · extraction confidence: 88%
- **fatigue_patients_with_post_covid** — n=1 paper · extraction confidence: 88%
- **loss_of_fitness_patients_with_post_covid** — n=1 paper · extraction confidence: 88%
- **pain_discomfort_any_problems** — n=1 paper · extraction confidence: 88%
- **anxiety_depression_any_problems** — n=1 paper · extraction confidence: 88%
- **usual_activities_any_problems** — n=1 paper · extraction confidence: 88%
- **mobility_any_problems** — n=1 paper · extraction confidence: 88%
- **self_care_any_problems** — n=1 paper · extraction confidence: 88%
- **myalgia_or_arthralgia** — n=1 paper · extraction confidence: 88%
- **brain_fog_pregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **brain_fog_nonpregnant_with_lc** — n=1 paper · extraction confidence: 88%

- **post_exertional_malaise_pregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **post_exertional_malaise_nonpregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **chest_pain_pregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **chest_pain_nonpregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **chronic_cough_pregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **chronic_cough_nonpregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_pregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_nonpregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **thirst_pregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **thirst_nonpregnant_with_lc** — n=1 paper · extraction confidence: 88%
- **long_covid_prevalence_pregnant_unadjusted** — n=1 paper · extraction confidence: 88%
- **long_covid_prevalence_nonpregnant_unadjusted** — n=1 paper · extraction confidence: 88%
- **long_covid_prevalence_pregnant_adjusted_6mo** — n=1 paper · extraction confidence: 88%
- **long_covid_prevalence_nonpregnant_adjusted_6mo** — n=1 paper · extraction confidence: 88%
- **htn_incidence_exposed** — n=1 paper · extraction confidence: 88%
- **hld_incidence_exposed** — n=1 paper · extraction confidence: 88%

- **masld_incidence_exposed** — n=1 paper · extraction confidence: 88%
- **t2dm_incidence_exposed** — n=1 paper · extraction confidence: 88%
- **mets_incidence_exposed** — n=1 paper · extraction confidence: 88%
- **cognitive_diagnosis_rs** — n=1 paper · extraction confidence: 88%
- **cognitive_diagnosis_ss** — n=1 paper · extraction confidence: 88%
- **cognitive_diagnosis_is** — n=1 paper · extraction confidence: 88%
- **psychological_diagnosis_rs** — n=1 paper · extraction confidence: 88%
- **psychological_diagnosis_ss** — n=1 paper · extraction confidence: 88%
- **psychological_diagnosis_is** — n=1 paper · extraction confidence: 88%
- **medical_diagnosis_rs** — n=1 paper · extraction confidence: 88%
- **medical_diagnosis_ss** — n=1 paper · extraction confidence: 88%
- **medical_diagnosis_is** — n=1 paper · extraction confidence: 88%
- **tracheostoma_related_rs** — n=1 paper · extraction confidence: 88%
- **tracheostoma_related_ss** — n=1 paper · extraction confidence: 88%
- **tracheostoma_related_is** — n=1 paper · extraction confidence: 88%
- **dysphagia_rs** — n=1 paper · extraction confidence: 88%
- **dysphagia_ss** — n=1 paper · extraction confidence: 88%
- **dysphagia_is** — n=1 paper · extraction confidence: 88%

- **impairments_of_nutrition_rs** — n=1 paper · extraction confidence: 88%
- **impairments_of_nutrition_ss** — n=1 paper · extraction confidence: 88%
- **taste_smelling_disorders_rs** — n=1 paper · extraction confidence: 88%
- **taste_smelling_disorders_ss** — n=1 paper · extraction confidence: 88%
- **tinnitus_rs** — n=1 paper · extraction confidence: 88%
- **tinnitus_ss** — n=1 paper · extraction confidence: 88%
- **abnormal_lung_function_rs** — n=1 paper · extraction confidence: 88%
- **abnormal_lung_function_ss** — n=1 paper · extraction confidence: 88%
- **venous_thromboembolism_rs** — n=1 paper · extraction confidence: 88%
- **venous_thromboembolism_is** — n=1 paper · extraction confidence: 88%
- **fatigue_is** — n=1 paper · extraction confidence: 88%
- **fatigue_ss** — n=1 paper · extraction confidence: 88%
- **chronic_pain_is** — n=1 paper · extraction confidence: 88%
- **chronic_pain_ss** — n=1 paper · extraction confidence: 88%
- **substance_use_disorders_is** — n=1 paper · extraction confidence: 88%
- **substance_use_disorders_ss** — n=1 paper · extraction confidence: 88%
- **anemia_is** — n=1 paper · extraction confidence: 88%
- **anemia_ss** — n=1 paper · extraction confidence: 88%
- **renal_diagnoses_is** — n=1 paper · extraction confidence: 88%
- **renal_diagnoses_ss** — n=1 paper · extraction confidence: 88%

- **ophthalmological_diagnoses_is** — n=1 paper · extraction confidence: 88%
- **ophthalmological_diagnoses_ss** — n=1 paper · extraction confidence: 88%
- **mortality_12mo_rs** — n=1 paper · extraction confidence: 88%
- **mortality_12mo_ss** — n=1 paper · extraction confidence: 88%
- **mortality_12mo_is** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_prepandemic_long_covid_women** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_prepandemic_no_long_covid_women** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_pandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_pandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_pandemic_long_covid_women** — n=1 paper · extraction confidence: 88%
- **any_relevant_symptom_pandemic_no_long_covid_women** — n=1 paper · extraction confidence: 88%
- **fatigue_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **fatigue_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **fatigue_prepandemic_long_covid_women** — n=1 paper · extraction confidence: 88%

- **fatigue_prepandemic_no_long_covid_women** — n=1 paper · extraction confidence: 88%
- **cough_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **cough_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **cough_prepandemic_long_covid_women** — n=1 paper · extraction confidence: 88%
- **cough_prepandemic_no_long_covid_women** — n=1 paper · extraction confidence: 88%
- **headache_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **headache_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **headache_prepandemic_long_covid_women** — n=1 paper · extraction confidence: 88%
- **headache_prepandemic_no_long_covid_women** — n=1 paper · extraction confidence: 88%
- **muscle_joint_pain_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **muscle_joint_pain_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **muscle_joint_pain_prepandemic_long_covid_women** — n=1 paper · extraction confidence: 88%
- **muscle_joint_pain_prepandemic_no_long_covid_women** — n=1 paper · extraction confidence: 88%
- **chest_pain_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **chest_pain_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **palpitations_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%

- **palpitations_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **dyspnea_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **dyspnea_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_taste_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **cognitive_dysfunction_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **sleep_disorder_prepandemic_long_covid_men** — n=1 paper · extraction confidence: 88%
- **sleep_disorder_prepandemic_no_long_covid_men** — n=1 paper · extraction confidence: 88%
- **overall_loneliness_any_long_covid** — n=1 paper · extraction confidence: 88%
- **overall_loneliness_no_long_covid** — n=1 paper · extraction confidence: 88%
- **social_loneliness_any_long_covid** — n=1 paper · extraction confidence: 88%
- **social_loneliness_no_long_covid** — n=1 paper · extraction confidence: 88%
- **emotional_loneliness_any_long_covid** — n=1 paper · extraction confidence: 88%
- **emotional_loneliness_no_long_covid** — n=1 paper · extraction confidence: 88%
- **overall_loneliness_total_sample** — n=1 paper · extraction confidence: 88%
- **pasc_overall_pregnant** — n=1 paper · extraction confidence: 88%
- **pasc_overall_non_pregnant** — n=1 paper · extraction confidence: 88%

- **abnormal_heartbeat_pregnant** — n=1 paper · extraction confidence: 88%
- **abnormal_heartbeat_non_pregnant** — n=1 paper · extraction confidence: 88%
- **abdominal_pain_pregnant** — n=1 paper · extraction confidence: 88%
- **abdominal_pain_non_pregnant** — n=1 paper · extraction confidence: 88%
- **thromboembolism_pregnant** — n=1 paper · extraction confidence: 88%
- **thromboembolism_non_pregnant** — n=1 paper · extraction confidence: 88%
- **joint_pain_pregnant** — n=1 paper · extraction confidence: 88%
- **joint_pain_non_pregnant** — n=1 paper · extraction confidence: 88%
- **dyspnoea_pregnant** — n=1 paper · extraction confidence: 88%
- **dyspnoea_non_pregnant** — n=1 paper · extraction confidence: 88%
- **malaise_and_fatigue_pregnant** — n=1 paper · extraction confidence: 88%
- **malaise_and_fatigue_non_pregnant** — n=1 paper · extraction confidence: 88%
- **cognitive_problems_pregnant** — n=1 paper · extraction confidence: 88%
- **cognitive_problems_non_pregnant** — n=1 paper · extraction confidence: 88%
- **acute_pharyngitis_pregnant** — n=1 paper · extraction confidence: 88%
- **acute_pharyngitis_non_pregnant** — n=1 paper · extraction confidence: 88%

- **hair_loss_pregnant** — n=1 paper · extraction confidence: 88%
- **hair_loss_non_pregnant** — n=1 paper · extraction confidence: 88%
- **sleep_disorders_pregnant** — n=1 paper · extraction confidence: 88%
- **sleep_disorders_non_pregnant** — n=1 paper · extraction confidence: 88%
- **chest_pain_pregnant** — n=1 paper · extraction confidence: 88%
- **chest_pain_non_pregnant** — n=1 paper · extraction confidence: 88%
- **encephalopathy_pregnant** — n=1 paper · extraction confidence: 88%
- **encephalopathy_non_pregnant** — n=1 paper · extraction confidence: 88%
- **long_covid_multi_symptomatic_cluster** — n=1 paper · extraction confidence: 88%
- **long_covid_predominance_shortness_of_breath_cluster** — n=1 paper · extraction confidence: 88%
- **long_covid_flu_like_cluster** — n=1 paper · extraction confidence: 88%
- **long_covid_predominance_fatigue_cluster** — n=1 paper · extraction confidence: 88%
- **long_covid_fever_cluster** — n=1 paper · extraction confidence: 88%
- **long_covid_nasal_congestion_cluster** — n=1 paper · extraction confidence: 88%
- **neurological_any** — n=1 paper · extraction confidence: 88%
- **psychiatric_any** — n=1 paper · extraction confidence: 88%
- **respiratory_any** — n=1 paper · extraction confidence: 88%
- **oxygen_use** — n=1 paper · extraction confidence: 88%

- **mobility_impairment_any** — n=1 paper · extraction confidence: 88%
- **decreased_exercise_tolerance** — n=1 paper · extraction confidence: 88%
- **functional_impairment** — n=1 paper · extraction confidence: 88%
- **heart_any** — n=1 paper · extraction confidence: 88%
- **flushing** — n=1 paper · extraction confidence: 88%
- **hypertension_new_onset** — n=1 paper · extraction confidence: 88%
- **digestive_any** — n=1 paper · extraction confidence: 88%
- **vomiting** — n=1 paper · extraction confidence: 88%
- **skin_any** — n=1 paper · extraction confidence: 88%
- **general_any** — n=1 paper · extraction confidence: 88%
- **flulike_symptoms** — n=1 paper · extraction confidence: 88%
- **conjunctivitis** — n=1 paper · extraction confidence: 88%
- **any_cardiac_event_exposed** — n=1 paper · extraction confidence: 88%
- **any_cardiac_event_unexposed** — n=1 paper · extraction confidence: 88%
- **cardiac_event_pre_vaccination_exposed** — n=1 paper · extraction confidence: 88%
- **cardiac_event_pre_vaccination_unexposed** — n=1 paper · extraction confidence: 88%
- **cardiac_event_post_vaccination_exposed** — n=1 paper · extraction confidence: 88%
- **cardiac_event_post_vaccination_unexposed** — n=1 paper · extraction confidence: 88%
- **hospitalization_cardiac_exposed** — n=1 paper · extraction confidence: 88%
- **hospitalization_cardiac_unexposed** — n=1 paper · extraction confidence: 88%

- **brain_fog_long_covid** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_long_covid** — n=1 paper · extraction confidence: 88%
- **chest_pain_long_covid** — n=1 paper · extraction confidence: 88%
- **brain_fog_recovered** — n=1 paper · extraction confidence: 88%
- **cough_recovered** — n=1 paper · extraction confidence: 88%
- **joint_pain_recovered** — n=1 paper · extraction confidence: 88%
- **chest_pain_recovered** — n=1 paper · extraction confidence: 88%
- **muscle_aches** — n=1 paper · extraction confidence: 88%
- **loss_of_concentration** — n=1 paper · extraction confidence: 88%
- **upper_respiratory_symptoms_runny_nose_sore_throat** — n=1 paper · extraction confidence: 88%
- **numbness** — n=1 paper · extraction confidence: 88%
- **cteph_2year_all_pe** — n=1 paper · extraction confidence: 88%
- **cteph_2year_covid_pe** — n=1 paper · extraction confidence: 88%
- **cteph_2year_non_covid_pe** — n=1 paper · extraction confidence: 88%
- **cteph_2year_all_pe_analysis2** — n=1 paper · extraction confidence: 88%
- **cteph_2year_covid_pe_analysis2** — n=1 paper · extraction confidence: 88%
- **cteph_2year_non_covid_pe_analysis2** — n=1 paper · extraction confidence: 88%
- **in_hospital_mortality_2year_covid_pe** — n=1 paper · extraction confidence: 88%

- **in_hospital_mortality_2year_non_covid_pe** — n=1 paper · extraction confidence: 88%
- **any_symptom_never_infected** — n=1 paper · extraction confidence: 88%
- **respiratory_symptoms_infected** — n=1 paper · extraction confidence: 88%
- **respiratory_symptoms_never_infected** — n=1 paper · extraction confidence: 88%
- **common_cold_symptoms_infected** — n=1 paper · extraction confidence: 88%
- **common_cold_symptoms_never_infected** — n=1 paper · extraction confidence: 88%
- **loss_of_taste_or_smell_infected** — n=1 paper · extraction confidence: 88%
- **loss_of_taste_or_smell_never_infected** — n=1 paper · extraction confidence: 88%
- **neurocognitive_symptoms_infected** — n=1 paper · extraction confidence: 88%
- **neurocognitive_symptoms_never_infected** — n=1 paper · extraction confidence: 88%
- **chronic_physical_distress_infected** — n=1 paper · extraction confidence: 88%
- **chronic_physical_distress_never_infected** — n=1 paper · extraction confidence: 88%
- **diarrhea_infected** — n=1 paper · extraction confidence: 88%
- **diarrhea_never_infected** — n=1 paper · extraction confidence: 88%
- **hair_loss_infected** — n=1 paper · extraction confidence: 88%
- **hair_loss_never_infected** — n=1 paper · extraction confidence: 88%
- **postexertional_malaise_female_lc_positive** — n=1 paper · extraction confidence: 88%

- **fatigue_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **brain_fog_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **dizziness_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **palpitations_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **loss_of_smell_or_taste_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **thirst_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **shortness_of_breath_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **chronic_cough_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **chest_pain_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **sleep_apnea_female_lc_positive** — n=1 paper · extraction confidence: 88%
- **long_covid_positive_female_overall** — n=1 paper · extraction confidence: 88%
- **long_covid_positive_male_overall** — n=1 paper · extraction confidence: 88%
- **pain_any_category_pre** — n=1 paper · extraction confidence: 88%
- **malaise_and_fatigue_pre** — n=1 paper · extraction confidence: 88%
- **malaise_and_fatigue_post** — n=1 paper · extraction confidence: 88%
- **sleep_wake_disorders_pre** — n=1 paper · extraction confidence: 88%

- **sleep_wake_disorders_post** — n=1 paper · extraction confidence: 88%
- **respiratory_signs_symptoms_pre** — n=1 paper · extraction confidence: 88%
- **respiratory_signs_symptoms_post** — n=1 paper · extraction confidence: 88%
- **musculoskeletal_pain_not_low_back_pre** — n=1 paper · extraction confidence: 88%
- **musculoskeletal_pain_not_low_back_post** — n=1 paper · extraction confidence: 88%
- **low_back_pain_pre** — n=1 paper · extraction confidence: 88%
- **low_back_pain_post** — n=1 paper · extraction confidence: 88%
- **anxiety_and_fear_related_disorders_pre** — n=1 paper · extraction confidence: 88%
- **anxiety_and_fear_related_disorders_post** — n=1 paper · extraction confidence: 88%
- **depressive_disorders_pre** — n=1 paper · extraction confidence: 88%
- **depressive_disorders_post** — n=1 paper · extraction confidence: 88%
- **nervous_system_signs_symptoms_pre** — n=1 paper · extraction confidence: 88%
- **nervous_system_signs_symptoms_post** — n=1 paper · extraction confidence: 88%
- **esophageal_disorders_pre** — n=1 paper · extraction confidence: 88%
- **esophageal_disorders_post** — n=1 paper · extraction confidence: 88%
- **nonspecific_chest_pain_pre** — n=1 paper · extraction confidence: 88%

- **nonspecific_chest_pain_post** — n=1 paper · extraction confidence: 88%
- **me_cfs_pre** — n=1 paper · extraction confidence: 88%
- **me_cfs_post** — n=1 paper · extraction confidence: 88%
- **pots_pre** — n=1 paper · extraction confidence: 88%
- **pots_post** — n=1 paper · extraction confidence: 88%
- **respiratory_failure_insufficiency_arrest_pre** — n=1 paper · extraction confidence: 88%
- **respiratory_failure_insufficiency_arrest_post** — n=1 paper · extraction confidence: 88%
- **pneumonia_pre** — n=1 paper · extraction confidence: 88%
- **pneumonia_post** — n=1 paper · extraction confidence: 88%
- **long_covid_among_covid_positive** — n=1 paper · extraction confidence: 88%
- **pandemic_impact_lack_social_contact** — n=1 paper · extraction confidence: 88%
- **pandemic_impact_worse_mental_health** — n=1 paper · extraction confidence: 88%
- **pandemic_impact_difficulty_accessing_healthcare** — n=1 paper · extraction confidence: 88%
- **pandemic_impact_financial_insecurity** — n=1 paper · extraction confidence: 88%
- **pandemic_impact_work_from_home** — n=1 paper · extraction confidence: 88%
- **depressive_symptoms_phq9_gte10** — n=1 paper · extraction confidence: 88%
- **anxiety_symptoms_gad7_gte10** — n=1 paper · extraction confidence: 88%
- **lc_cases_proportion** — n=1 paper · extraction confidence: 87%
- **pacs_cases_proportion** — n=1 paper · extraction confidence: 87%

- **long covid overall** — n=1 paper · extraction confidence: 87%
- **long covid most deprived decile (imd1) unadjusted** — n=1 paper · extraction confidence: 87%
- **long covid least deprived decile (imd10) unadjusted** — n=1 paper · extraction confidence: 87%
- **long covid women overall** — n=1 paper · extraction confidence: 87%
- **long covid men overall** — n=1 paper · extraction confidence: 87%
- **long covid most deprived adjusted** — n=1 paper · extraction confidence: 87%
- **long covid least deprived adjusted** — n=1 paper · extraction confidence: 87%
- **long_covid_current_symptoms_among_long_covid** — n=1 paper · extraction confidence: 87%
- **long_covid_current_symptoms_impacting_daily_life_among_those_wi** — n=1 paper · extraction confidence: 87%
- **cognitive_limitation_among_long_covid_approximately** — n=1 paper · extraction confidence: 87%
- **breathing_problems** — n=1 paper · extraction confidence: 87%
- **symptom_based_long_covid** — n=1 paper · extraction confidence: 87%
- **moderate_or_severe_covid_among_diagnosed** — n=1 paper · extraction confidence: 87%
- **prior_covid19_diagnosis** — n=1 paper · extraction confidence: 87%
- **long_covid_high_risk_condition** — n=1 paper · extraction confidence: 87%
- **long_covid_no_high_risk_condition** — n=1 paper · extraction confidence: 87%

- **long_covid_anxiety_or_depression** — n=1 paper · extraction confidence: 87%
- **long_covid_no_anxiety_or_depression** — n=1 paper · extraction confidence: 87%
- **long_covid_disability** — n=1 paper · extraction confidence: 87%
- **long_covid_no_disability** — n=1 paper · extraction confidence: 87%
- **long_covid_food_insecurity** — n=1 paper · extraction confidence: 87%
- **long_covid_no_food_insecurity** — n=1 paper · extraction confidence: 87%
- **reduced_exercise_tolerance_and_resilience_males** — n=1 paper · extraction confidence: 87%
- **reduced_exercise_tolerance_and_resilience_females** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_males** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_females** — n=1 paper · extraction confidence: 87%
- **dysosmia_anosmia_males** — n=1 paper · extraction confidence: 87%
- **dysosmia_anosmia_females** — n=1 paper · extraction confidence: 87%
- **concentration_deficits_female_inpatients** — n=1 paper · extraction confidence: 87%
- **concentration_deficits_male_inpatients** — n=1 paper · extraction confidence: 87%
- **any_pasc_overall** — n=1 paper · extraction confidence: 87%
- **any_pasc_hospitalised** — n=1 paper · extraction confidence: 87%
- **any_pasc_outpatients** — n=1 paper · extraction confidence: 87%

- **any_pasc_outpatient_females** — n=1 paper · extraction confidence: 87%
- **any_pasc_outpatient_males** — n=1 paper · extraction confidence: 87%
- **financial_hardship_prevalence** — n=1 paper · extraction confidence: 87%
- **less_than_college_education** — n=1 paper · extraction confidence: 87%
- **any_economic_instability** — n=1 paper · extraction confidence: 87%
- **any_healthcare_access_barrier** — n=1 paper · extraction confidence: 87%
- **medical_discrimination_experience** — n=1 paper · extraction confidence: 87%
- **any_social_community_support_challenge** — n=1 paper · extraction confidence: 87%
- **lack_of_social_support** — n=1 paper · extraction confidence: 87%
- **poor_neighborhood_cohesion** — n=1 paper · extraction confidence: 87%
- **food_insecurity** — n=1 paper · extraction confidence: 87%
- **frequent_everyday_discrimination** — n=1 paper · extraction confidence: 87%
- **highest_quintile_zip_poverty** — n=1 paper · extraction confidence: 87%
- **highest_quartile_household_crowding** — n=1 paper · extraction confidence: 87%
- **any_of_96_pasc_attributable_to_covid19_at_0-3mo** — n=1 paper · extraction confidence: 87%
- **smell_taste_disturbance_attributable_fraction** — n=1 paper · extraction confidence: 87%
- **fatigue_attributable_fraction** — n=1 paper · extraction confidence: 87%

- **specific_pcc_symptoms_attributable_approx** — n=1 paper · extraction confidence: 87%
- **tiredness_infected_6mo** — n=1 paper · extraction confidence: 87%
- **tiredness_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **headache_infected_6mo** — n=1 paper · extraction confidence: 87%
- **headache_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **muscle_aches_weakness_infected_6mo** — n=1 paper · extraction confidence: 87%
- **muscle_aches_weakness_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **sleep_problems_infected_6mo** — n=1 paper · extraction confidence: 87%
- **sleep_problems_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **anxious_depressed_infected_6mo** — n=1 paper · extraction confidence: 87%
- **anxious_depressed_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **breathlessness_infected_6mo** — n=1 paper · extraction confidence: 87%
- **breathlessness_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **joint_pain_infected_6mo** — n=1 paper · extraction confidence: 87%
- **joint_pain_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **change_in_smell_infected_6mo** — n=1 paper · extraction confidence: 87%

- **change_in_smell_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **change_in_taste_infected_6mo** — n=1 paper · extraction confidence: 87%
- **change_in_taste_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **confusion_infected_6mo** — n=1 paper · extraction confidence: 87%
- **confusion_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **palpitations_infected_6mo** — n=1 paper · extraction confidence: 87%
- **palpitations_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **chest_pain_infected_6mo** — n=1 paper · extraction confidence: 87%
- **chest_pain_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **at_least_one_symptom_infected_6mo** — n=1 paper · extraction confidence: 87%
- **at_least_one_symptom_never_infected_6mo** — n=1 paper · extraction confidence: 87%
- **at_least_one_symptom_infected_12mo** — n=1 paper · extraction confidence: 87%
- **at_least_one_symptom_never_infected_12mo** — n=1 paper · extraction confidence: 87%
- **at_least_one_symptom_infected_18mo** — n=1 paper · extraction confidence: 87%
- **at_least_one_symptom_never_infected_18mo** — n=1 paper · extraction confidence: 87%
- **respiratory** — n=1 paper · extraction confidence: 87%
- **cardiovascular** — n=1 paper · extraction confidence: 87%
- **musculoskeletal** — n=1 paper · extraction confidence: 87%

- **loss_of_smell_taste** — n=1 paper · extraction confidence: 87%
- **sleeping_disorder_day180** — n=1 paper · extraction confidence: 87%
- **joint_pain_day180** — n=1 paper · extraction confidence: 87%
- **fatigue_day180** — n=1 paper · extraction confidence: 87%
- **headaches_day180** — n=1 paper · extraction confidence: 87%
- **muscle_pain_day180** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_day180** — n=1 paper · extraction confidence: 87%
- **concentration_impairment_day180** — n=1 paper · extraction confidence: 87%
- **memory_impairment_day180** — n=1 paper · extraction confidence: 87%
- **loss_of_smell_or_taste_day180** — n=1 paper · extraction confidence: 87%
- **palpitations_day180** — n=1 paper · extraction confidence: 87%
- **diarrhoea_day180** — n=1 paper · extraction confidence: 87%
- **any_long_covid_symptom_day180** — n=1 paper · extraction confidence: 87%
- **any_long_covid_symptom_day45** — n=1 paper · extraction confidence: 87%
- **fatigue_baseline** — n=1 paper · extraction confidence: 87%
- **muscle_pain_baseline** — n=1 paper · extraction confidence: 87%
- **headaches_baseline** — n=1 paper · extraction confidence: 87%

- **joint_pain_baseline** — n=1 paper · extraction confidence: 87%
- **sleeping_disorder_baseline** — n=1 paper · extraction confidence: 87%
- **mood_anxiety_disorders** — n=1 paper · extraction confidence: 87%
- **malaise_fatigue** — n=1 paper · extraction confidence: 87%
- **taste_smell_changes** — n=1 paper · extraction confidence: 87%
- **fatigue_ever_lc** — n=1 paper · extraction confidence: 87%
- **post_exertional_malaise_ever_lc** — n=1 paper · extraction confidence: 87%
- **cough_ever_lc** — n=1 paper · extraction confidence: 87%
- **palpitations_ever_lc** — n=1 paper · extraction confidence: 87%
- **sleep_difficulties_ever_lc** — n=1 paper · extraction confidence: 87%
- **cognitive_impairment_ever_lc** — n=1 paper · extraction confidence: 87%
- **brain_fog_ever_lc** — n=1 paper · extraction confidence: 87%
- **fatigue_fss>=4_persistent_lc** — n=1 paper · extraction confidence: 87%
- **pem_indicative_persistent_lc** — n=1 paper · extraction confidence: 87%
- **me_cfs_indicative_persistent_lc** — n=1 paper · extraction confidence: 87%
- **anxiety_gad7>=5_persistent_lc** — n=1 paper · extraction confidence: 87%
- **depression_phq9>=5_persistent_lc** — n=1 paper · extraction confidence: 87%
- **mobility_problem_persistent_lc** — n=1 paper · extraction confidence: 87%

- **usual_activity_problem_persistent_lc** — n=1 paper · extraction confidence: 87%
- **personal_care_problem_persistent_lc** — n=1 paper · extraction confidence: 87%
- **dyspnea_mmrc>=1_respiratory_lc** — n=1 paper · extraction confidence: 87%
- **cognitive_impairment_moca<26_neurological_lc** — n=1 paper · extraction confidence: 87%
- **long_covid_test_positive** — n=1 paper · extraction confidence: 87%
- **long_covid_test_negative** — n=1 paper · extraction confidence: 87%
- **at_least_one_physical_symptom_test_positive** — n=1 paper · extraction confidence: 87%
- **at_least_one_physical_symptom_test_negative** — n=1 paper · extraction confidence: 87%
- **impaired_memory** — n=1 paper · extraction confidence: 87%
- **any_cognitive_symptom_among_pcc** — n=1 paper · extraction confidence: 87%
- **respiratory_pasc_among_sars-cov-2_positive** — n=1 paper · extraction confidence: 87%
- **neurologic_pasc_among_sars-cov-2_positive** — n=1 paper · extraction confidence: 87%
- **any_pasc_among_sars-cov-2_positive** — n=1 paper · extraction confidence: 87%
- **malaise** — n=1 paper · extraction confidence: 87%
- **confusion_or_difficulty_thinking_concentrating** — n=1 paper · extraction confidence: 87%
- **fast_beating_heart** — n=1 paper · extraction confidence: 87%
- **chest_pain_tightness** — n=1 paper · extraction confidence: 87%

- **mood_disorders_long_covid_children** — n=1 paper · extraction confidence: 87%
- **fatigue_long_covid_children** — n=1 paper · extraction confidence: 87%
- **sleep_disorders_long_covid_children** — n=1 paper · extraction confidence: 87%
- **headache_long_covid_children** — n=1 paper · extraction confidence: 87%
- **tachycardia_bradycardia_palpitations_arrhythmia_trajectory1** — n=1 paper · extraction confidence: 87%
- **paresthesia_trajectory1** — n=1 paper · extraction confidence: 87%
- **hot_flushes_trajectory1** — n=1 paper · extraction confidence: 87%
- **sweats_trajectory1** — n=1 paper · extraction confidence: 87%
- **heat_cold_intolerance_trajectory1** — n=1 paper · extraction confidence: 87%
- **diarrhea_trajectory1** — n=1 paper · extraction confidence: 87%
- **neck_back_low_back_pain_trajectory1** — n=1 paper · extraction confidence: 87%
- **photophobia_phonophobia_trajectory1** — n=1 paper · extraction confidence: 87%
- **neck_back_low_back_pain_trajectory2** — n=1 paper · extraction confidence: 87%
- **diarrhea_trajectory2** — n=1 paper · extraction confidence: 87%
- **tachycardia_bradycardia_palpitations_arrhythmia_overall** — n=1 paper · extraction confidence: 87%
- **neck_back_low_back_pain_overall** — n=1 paper · extraction confidence: 87%

- **paresthesia_overall** — n=1 paper · extraction confidence: 87%
- **diarrhea_overall** — n=1 paper · extraction confidence: 87%
- **hot_flushes_overall** — n=1 paper · extraction confidence: 87%
- **sweats_overall** — n=1 paper · extraction confidence: 87%
- **heat_cold_intolerance_overall** — n=1 paper · extraction confidence: 87%
- **photophobia_phonophobia_overall** — n=1 paper · extraction confidence: 87%
- **post_exertional_malaise_all_infected** — n=1 paper · extraction confidence: 87%
- **post_exertional_malaise_acute_infected** — n=1 paper · extraction confidence: 87%
- **post_exertional_malaise_post_acute_infected** — n=1 paper · extraction confidence: 87%
- **orthostatic_intolerance_acute_infected** — n=1 paper · extraction confidence: 87%
- **orthostatic_intolerance_post_acute_infected** — n=1 paper · extraction confidence: 87%
- **unrefreshing_sleep_acute_infected** — n=1 paper · extraction confidence: 87%
- **cognitive_impairment_acute_infected** — n=1 paper · extraction confidence: 87%
- **cognitive_impairment_post_acute_infected** — n=1 paper · extraction confidence: 87%
- **fatigue_acute_infected** — n=1 paper · extraction confidence: 87%
- **fatigue_post_acute_infected** — n=1 paper · extraction confidence: 87%
- **unrefreshing_sleep_post_acute_infected** — n=1 paper · extraction confidence: 87%
- **tiredness_nn** — n=1 paper · extraction confidence: 87%

- **tiredness_np** — n=1 paper · extraction confidence: 87%
- **tiredness_pn** — n=1 paper · extraction confidence: 87%
- **tiredness_pp** — n=1 paper · extraction confidence: 87%
- **sleeping_difficulties_nn** — n=1 paper · extraction confidence: 87%
- **sleeping_difficulties_np** — n=1 paper · extraction confidence: 87%
- **sleeping_difficulties_pn** — n=1 paper · extraction confidence: 87%
- **sleeping_difficulties_pp** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_nn** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_np** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_pn** — n=1 paper · extraction confidence: 87%
- **shortness_of_breath_pp** — n=1 paper · extraction confidence: 87%
- **headaches_nn** — n=1 paper · extraction confidence: 87%
- **headaches_np** — n=1 paper · extraction confidence: 87%
- **headaches_pn** — n=1 paper · extraction confidence: 87%
- **headaches_pp** — n=1 paper · extraction confidence: 87%
- **persistent_cough_nn** — n=1 paper · extraction confidence: 87%
- **persistent_cough_np** — n=1 paper · extraction confidence: 87%
- **persistent_cough_pn** — n=1 paper · extraction confidence: 87%
- **persistent_cough_pp** — n=1 paper · extraction confidence: 87%
- **loss_of_smell_nn** — n=1 paper · extraction confidence: 87%
- **loss_of_smell_np** — n=1 paper · extraction confidence: 87%

- **loss_of_smell_pn** — n=1 paper · extraction confidence: 87%
- **loss_of_smell_pp** — n=1 paper · extraction confidence: 87%
- **dizziness_nn** — n=1 paper · extraction confidence: 87%
- **dizziness_np** — n=1 paper · extraction confidence: 87%
- **dizziness_pn** — n=1 paper · extraction confidence: 87%
- **dizziness_pp** — n=1 paper · extraction confidence: 87%
- **unusual_chest_pain_nn** — n=1 paper · extraction confidence: 87%
- **unusual_chest_pain_np** — n=1 paper · extraction confidence: 87%
- **unusual_chest_pain_pn** — n=1 paper · extraction confidence: 87%
- **unusual_chest_pain_pp** — n=1 paper · extraction confidence: 87%
- **chills_nn** — n=1 paper · extraction confidence: 87%
- **chills_np** — n=1 paper · extraction confidence: 87%
- **chills_pn** — n=1 paper · extraction confidence: 87%
- **chills_pp** — n=1 paper · extraction confidence: 87%
- **confusion_disorientation_nn** — n=1 paper · extraction confidence: 87%
- **confusion_disorientation_np** — n=1 paper · extraction confidence: 87%
- **confusion_disorientation_pn** — n=1 paper · extraction confidence: 87%
- **confusion_disorientation_pp** — n=1 paper · extraction confidence: 87%
- **unusual_muscle_pains_nn** — n=1 paper · extraction confidence: 87%
- **unusual_muscle_pains_np** — n=1 paper · extraction confidence: 87%
- **unusual_muscle_pains_pn** — n=1 paper · extraction confidence: 87%

- **unusual_muscle_pains_pp** — n=1 paper · extraction confidence: 87%
- **five_plus_symptoms_nn** — n=1 paper · extraction confidence: 87%
- **five_plus_symptoms_np** — n=1 paper · extraction confidence: 87%
- **five_plus_symptoms_pn** — n=1 paper · extraction confidence: 87%
- **five_plus_symptoms_pp** — n=1 paper · extraction confidence: 87%
- **no_symptoms_nn** — n=1 paper · extraction confidence: 87%
- **no_symptoms_np** — n=1 paper · extraction confidence: 87%
- **no_symptoms_pn** — n=1 paper · extraction confidence: 87%
- **no_symptoms_pp** — n=1 paper · extraction confidence: 87%
- **cobra_score_ge15_pretest_test_positive** — n=1 paper · extraction confidence: 87%
- **cobra_score_ge15_pretest_test_negative** — n=1 paper · extraction confidence: 87%
- **cobra_score_ge15_at_18months_test_positive** — n=1 paper · extraction confidence: 87%
- **cobra_score_ge15_at_18months_test_negative** — n=1 paper · extraction confidence: 87%
- **general_symptoms** — n=1 paper · extraction confidence: 87%
- **neurological_and_psychiatric_symptoms** — n=1 paper · extraction confidence: 87%
- **cardiovascular_symptoms** — n=1 paper · extraction confidence: 87%
- **fatigue_any_grade_men** — n=1 paper · extraction confidence: 85%
- **fatigue_any_grade_women** — n=1 paper · extraction confidence: 85%

- **fatigue_moderate_to_strong_men** — n=1 paper · extraction confidence: 85%
- **fatigue_moderate_to_strong_women** — n=1 paper · extraction confidence: 85%
- **neurocognitive_impairment_any_grade_men** — n=1 paper · extraction confidence: 85%
- **neurocognitive_impairment_any_grade_women** — n=1 paper · extraction confidence: 85%
- **neurocognitive_impairment_moderate_to_strong_men** — n=1 paper · extraction confidence: 85%
- **neurocognitive_impairment_moderate_to_strong_women** — n=1 paper · extraction confidence: 85%
- **chest_symptoms_any_grade_men** — n=1 paper · extraction confidence: 85%
- **chest_symptoms_any_grade_women** — n=1 paper · extraction confidence: 85%
- **chest_symptoms_moderate_to_strong_men** — n=1 paper · extraction confidence: 85%
- **chest_symptoms_moderate_to_strong_women** — n=1 paper · extraction confidence: 85%
- **any_symptom_any_grade_men** — n=1 paper · extraction confidence: 85%
- **any_symptom_any_grade_women** — n=1 paper · extraction confidence: 85%
- **long_covid_paxlovid_treated_180d** — n=1 paper · extraction confidence: 85%
- **long_covid_non_treated_180d** — n=1 paper · extraction confidence: 85%
- **any_long_covid_outcome_overall** — n=1 paper · extraction confidence: 84%
- **any_long_covid_outcome_immunocompromised** — n=1 paper · extraction confidence: 84%

- **any_long_covid_outcome_moderately_severely_immunocompromised** — n=1 paper · extraction confidence: 84%
- **neuropsychiatric_features_overall_rate_per_100py** — n=1 paper · extraction confidence: 84%
- **dyspnea_breathlessness_overall_rate_per_100py** — n=1 paper · extraction confidence: 84%
- **fatigue_overall_rate_per_100py** — n=1 paper · extraction confidence: 84%
- **joint_pain_overall_rate_per_100py** — n=1 paper · extraction confidence: 84%
- **cognitive_dysfunction_overall_rate_per_100py** — n=1 paper · extraction confidence: 84%
- **dysautonomia_overall** — n=1 paper · extraction confidence: 84%
- **taste_disturbance_overall** — n=1 paper · extraction confidence: 84%
- **smell_disturbance_overall** — n=1 paper · extraction confidence: 84%
- **fatigue_in_long_covid_nonhospitalized** — n=1 paper · extraction confidence: 82%
- **fatigue_in_long_covid_hospitalized** — n=1 paper · extraction confidence: 82%
- **fatigue_in_long_covid_vaccinated** — n=1 paper · extraction confidence: 82%
- **dyspnea_cases_post_covid** — n=1 paper · extraction confidence: 82%
- **dyspnea_controls_post_covid** — n=1 paper · extraction confidence: 82%
- **fatigue_cases_post_covid** — n=1 paper · extraction confidence: 82%
- **fatigue_controls_post_covid** — n=1 paper · extraction confidence: 82%

- **cognitive_dysfunction_cases_post_covid** — n=1 paper · extraction confidence: 82%
- **cognitive_dysfunction_controls_post_covid** — n=1 paper · extraction confidence: 82%
- **weakness_cases_post_covid** — n=1 paper · extraction confidence: 82%
- **weakness_controls_post_covid** — n=1 paper · extraction confidence: 82%
- **mobility_difficulty_cases_post_covid** — n=1 paper · extraction confidence: 82%
- **mobility_difficulty_controls_post_covid** — n=1 paper · extraction confidence: 82%
- **cognitive_dysfunction_premorbid_cases** — n=1 paper · extraction confidence: 82%
- **dyspnea_premorbid_cases** — n=1 paper · extraction confidence: 82%
- **fatigue_premorbid_cases** — n=1 paper · extraction confidence: 82%
- **mobility_difficulty_premorbid_cases** — n=1 paper · extraction confidence: 82%
- **weakness_premorbid_cases** — n=1 paper · extraction confidence: 82%
- **weakness_premorbid_controls** — n=1 paper · extraction confidence: 82%
- **cognitive_dysfunction_premorbid_controls** — n=1 paper · extraction confidence: 82%
- **mobility_difficulty_premorbid_controls** — n=1 paper · extraction confidence: 82%
- **dyspnea_premorbid_controls** — n=1 paper · extraction confidence: 82%
- **fatigue_premorbid_controls** — n=1 paper · extraction confidence: 82%

- **respiratory_signs_and_symptoms_first_infection_incidence_per_milli**
— n=1 paper · extraction confidence: 82%
- **respiratory_signs_and_symptoms_second_infection_incidence_per_m**
— n=1 paper · extraction confidence: 82%
- **overall_pasc_diagnosis_first_infection_incidence_per_million_6mo**
— n=1 paper · extraction confidence: 82%
- **overall_pasc_diagnosis_second_infection_incidence_per_million_6mo**
— n=1 paper · extraction confidence: 82%
- **any_psychiatric_disorder** — n=1 paper · extraction confidence: 82%
- **long_covid_diagnosis_2plus_u099** — n=1 paper · extraction confidence: 82%
- **long_covid_diagnosis_1plus_u099_sensitivity** — n=1 paper · extraction confidence: 82%
- **long_covid_cld_group** — n=1 paper · extraction confidence: 82%
- **long_covid_non_cld_group** — n=1 paper · extraction confidence: 82%
- **long_covid_copd** — n=1 paper · extraction confidence: 82%
- **long_covid_asthma** — n=1 paper · extraction confidence: 82%
- **long_covid_bronchiectasis** — n=1 paper · extraction confidence: 82%
- **long_covid_age_over65** — n=1 paper · extraction confidence: 82%
- **long_covid_age_40_65** — n=1 paper · extraction confidence: 82%
- **long_covid_age_18_39** — n=1 paper · extraction confidence: 82%
- **long_covid_pre_delta_wave** — n=1 paper · extraction confidence: 82%
- **long_covid_delta_wave** — n=1 paper · extraction confidence: 82%

- **long_covid_omicron_wave** — n=1 paper · extraction confidence: 82%
- **any_symptom_weeks_4_to_8_persistent_change_group** — n=1 paper · extraction confidence: 82%
- **any_symptom_weeks_4_to_8_acute_covid_group** — n=1 paper · extraction confidence: 82%
- **difficulty_with_usual_activities** — n=1 paper · extraction confidence: 82%
- **anxiety_or_depression** — n=1 paper · extraction confidence: 82%
- **mobility_problems** — n=1 paper · extraction confidence: 82%
- **no_impact_from_long_covid** — n=1 paper · extraction confidence: 82%
- **neuralgia** — n=1 paper · extraction confidence: 82%
- **loss_of_sense_of_smell** — n=1 paper · extraction confidence: 82%
- **muscle_ache** — n=1 paper · extraction confidence: 82%
- **difficulty_sleeping** — n=1 paper · extraction confidence: 82%
- **worry_or_anxiety** — n=1 paper · extraction confidence: 82%
- **memory_loss_or_confusion** — n=1 paper · extraction confidence: 82%
- **malaise_and_fatigue** — n=1 paper · extraction confidence: 82%
- **asthma** — n=1 paper · extraction confidence: 82%
- **long_covid_mild** — n=1 paper · extraction confidence: 82%
- **long_covid_moderate** — n=1 paper · extraction confidence: 82%
- **long_covid_severe** — n=1 paper · extraction confidence: 82%
- **anosmia_dysosmia_alpha_largest_cluster** — n=1 paper · extraction confidence: 82%

- **central_neurological_symptoms_delta_largest_cluster** — n=1 paper · extraction confidence: 82%
- **severe_dyspnoea_delta_smallest_cluster** — n=1 paper · extraction confidence: 82%
- **dyspnoea_affecting_function_at_12_weeks_wild_b** — n=1 paper · extraction confidence: 82%
- **dyspnea_most_predictive_pasc_cp** — n=1 paper · extraction confidence: 82%
- **fatigue_most_predictive_me_cfs_cp** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_finding_related_to_attentiveness** — n=1 paper · extraction confidence: 82%
- **long_covid_prevalence_in_cohort** — n=1 paper · extraction confidence: 82%
- **any_long_covid_symptoms_total_sample** — n=1 paper · extraction confidence: 82%
- **any_long_covid_symptoms_post_covid_sample** — n=1 paper · extraction confidence: 82%
- **combination_physical_cognitive_psychological** — n=1 paper · extraction confidence: 82%
- **mainly_physical** — n=1 paper · extraction confidence: 82%
- **mainly_psychological_or_cognitive** — n=1 paper · extraction confidence: 82%
- **lack_of_energy_at_infection_long_covid** — n=1 paper · extraction confidence: 82%
- **headache_at_infection_long_covid** — n=1 paper · extraction confidence: 82%
- **muscle_aches_at_infection_long_covid** — n=1 paper · extraction confidence: 82%
- **fever_at_infection_long_covid** — n=1 paper · extraction confidence: 82%
- **dyspnea_at_infection_long_covid** — n=1 paper · extraction confidence: 82%

- **dizziness_at_infection_non_hospitalized** — n=1 paper · extraction confidence: 82%
- **dizziness_at_infection_hospitalized** — n=1 paper · extraction confidence: 82%
- **heavy_limbs_at_infection_non_hospitalized** — n=1 paper · extraction confidence: 82%
- **heavy_limbs_at_infection_hospitalized** — n=1 paper · extraction confidence: 82%
- **tremors_at_infection_non_hospitalized** — n=1 paper · extraction confidence: 82%
- **headaches_no_covid_group** — n=1 paper · extraction confidence: 82%
- **fatigue_no_covid_group** — n=1 paper · extraction confidence: 82%
- **muscle_or_body_aches_no_covid_group** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_covid_12_weeks_vs_no_covid_or** — n=1 paper · extraction confidence: 82%
- **loss_of_taste_covid_12_weeks_vs_no_covid_or** — n=1 paper · extraction confidence: 82%
- **covid-19_any_positive_response** — n=1 paper · extraction confidence: 82%
- **covid-19_tested_positive_health_professional** — n=1 paper · extraction confidence: 82%
- **covid-19_tested_positive_home_test** — n=1 paper · extraction confidence: 82%
- **covid-19_believed_had_no_test** — n=1 paper · extraction confidence: 82%
- **long_covid-19_among_all_respondents** — n=1 paper · extraction confidence: 82%
- **long_covid-19_among_covid_positive** — n=1 paper · extraction confidence: 82%

- **covid-19_with_function-impairing_symptoms_any_duration** — n=1 paper · extraction confidence: 82%
- **acute_covid-19_symptoms_less_than_4_weeks** — n=1 paper · extraction confidence: 82%
- **ongoing_symptomatic_covid-19_4_to_12_weeks** — n=1 paper · extraction confidence: 82%
- **post-covid-19_syndrome_greater_than_12_weeks** — n=1 paper · extraction confidence: 82%
- **chronic_cough_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **chronic_fatigue_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **dyspnea_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **pulmonary_fibrosis_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **critical_illness_myopathy_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **chronic_respiratory_failure_with_hypoxia_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **personal_history_covid19_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **interstitial_pulmonary_disease_j84_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **other_myopathies_g72_community_long_covid_claims** — n=1 paper · extraction confidence: 82%
- **fatigue_at_baseline_allied_healthcare** — n=1 paper · extraction confidence: 82%
- **fatigue_at_baseline_no_allied_healthcare** — n=1 paper · extraction confidence: 82%
- **long covid (lcvi-positive) age 18-39** — n=1 paper · extraction confidence: 82%

- **long covid (lcric-positive) age 40-49** — n=1 paper · extraction confidence: 82%
- **long covid (lcric-positive) age 50-59** — n=1 paper · extraction confidence: 82%
- **long covid (lcric-positive) age 60-69** — n=1 paper · extraction confidence: 82%
- **long covid (lcric-positive) age ≥70** — n=1 paper · extraction confidence: 82%
- **brain_fog_or_confusion** — n=1 paper · extraction confidence: 82%
- **disturbed_sleep** — n=1 paper · extraction confidence: 82%
- **chest_tightness** — n=1 paper · extraction confidence: 82%
- **postural_symptoms** — n=1 paper · extraction confidence: 82%
- **focal_weakness** — n=1 paper · extraction confidence: 82%
- **vertigo** — n=1 paper · extraction confidence: 82%
- **generalized_system** — n=1 paper · extraction confidence: 82%
- **respiratory_system** — n=1 paper · extraction confidence: 82%
- **neurologic_system** — n=1 paper · extraction confidence: 82%
- **psychiatric_system** — n=1 paper · extraction confidence: 82%
- **digestive_system** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_system** — n=1 paper · extraction confidence: 82%
- **cardiovascular_system** — n=1 paper · extraction confidence: 82%
- **dermatological_system** — n=1 paper · extraction confidence: 82%
- **ophthalmological_system** — n=1 paper · extraction confidence: 82%

- **urological_system** — n=1 paper · extraction confidence: 82%
- **concentration_difficulties** — n=1 paper · extraction confidence: 82%
- **heart_palpitations** — n=1 paper · extraction confidence: 82%
- **altered_or_loss_of_smell_taste** — n=1 paper · extraction confidence: 82%
- **joint_pain_or_swelling** — n=1 paper · extraction confidence: 82%
- **altered_or_loss_of_smell** — n=1 paper · extraction confidence: 82%
- **altered_or_loss_of_taste** — n=1 paper · extraction confidence: 82%
- **abnormal_imaging** — n=1 paper · extraction confidence: 82%
- **long_covid_formally_diagnosed_among_long_covid_group** — n=1 paper · extraction confidence: 82%
- **long_covid_self_diagnosed_among_long_covid_group** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_or_taste_delta_ambulatory** — n=1 paper · extraction confidence: 82%
- **memory_problems_all_eras** — n=1 paper · extraction confidence: 82%
- **alzheimers_disease_all_eras** — n=1 paper · extraction confidence: 82%
- **end_stage_renal_failure_ba.1_2** — n=1 paper · extraction confidence: 82%
- **long_covid_unsure** — n=1 paper · extraction confidence: 82%
- **long_covid_deaf_sign_language_users** — n=1 paper · extraction confidence: 82%

- **long_covid_alzheimers_dementia** — n=1 paper · extraction confidence: 82%
- **long_covid_autism** — n=1 paper · extraction confidence: 82%
- **long_covid_breathing_condition** — n=1 paper · extraction confidence: 82%
- **long_covid_mental_health_condition** — n=1 paper · extraction confidence: 82%
- **long_covid_permanently_sick_disabled** — n=1 paper · extraction confidence: 82%
- **long_covid_north_west_region** — n=1 paper · extraction confidence: 82%
- **long_covid_south_west_region** — n=1 paper · extraction confidence: 82%
- **long_covid_current_regular_smokers** — n=1 paper · extraction confidence: 82%
- **long_covid_with_diabetes** — n=1 paper · extraction confidence: 82%
- **long_covid_without_diabetes** — n=1 paper · extraction confidence: 82%
- **ongoing_symptoms_covid-19_at_12_weeks** — n=1 paper · extraction confidence: 82%
- **ongoing_symptoms_influenza_at_12_weeks** — n=1 paper · extraction confidence: 82%
- **moderate_to_severe_functional_impairment_covid-19** — n=1 paper · extraction confidence: 82%
- **moderate_to_severe_functional_impairment_influenza** — n=1 paper · extraction confidence: 82%
- **asthenia_post_covid_overall** — n=1 paper · extraction confidence: 82%
- **myalgia_post_covid_overall** — n=1 paper · extraction confidence: 82%

- **cough_post_covid_overall** — n=1 paper · extraction confidence: 82%
- **dyspnea_post_covid_overall** — n=1 paper · extraction confidence: 82%
- **concentration_deficit_post_covid_overall** — n=1 paper · extraction confidence: 82%
- **headache_post_covid_overall** — n=1 paper · extraction confidence: 82%
- **anosmia_post_covid_overall** — n=1 paper · extraction confidence: 82%
- **asthenia_singleton** — n=1 paper · extraction confidence: 82%
- **cough_singleton** — n=1 paper · extraction confidence: 82%
- **myalgia_singleton** — n=1 paper · extraction confidence: 82%
- **dyspnea_singleton** — n=1 paper · extraction confidence: 82%
- **anosmia_singleton** — n=1 paper · extraction confidence: 82%
- **concentration_deficit_singleton** — n=1 paper · extraction confidence: 82%
- **headache_singleton** — n=1 paper · extraction confidence: 82%
- **long_covid_30_60days_primary_infection** — n=1 paper · extraction confidence: 82%
- **long_covid_61plus_days_primary_infection** — n=1 paper · extraction confidence: 82%
- **long_covid_30_60days_second_infection** — n=1 paper · extraction confidence: 82%
- **long_covid_61plus_days_second_infection** — n=1 paper · extraction confidence: 82%
- **neurological_symptoms_persisting_30_60days** — n=1 paper · extraction confidence: 82%

- **psychological_symptoms_persisting_30_60days** — n=1 paper · extraction confidence: 82%
- **neurological_symptoms_persisting_61plus_days** — n=1 paper · extraction confidence: 82%
- **psychological_symptoms_persisting_61plus_days** — n=1 paper · extraction confidence: 82%
- **any_pasc_am** — n=1 paper · extraction confidence: 82%
- **constitutional_pasc_am** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_pasc_am** — n=1 paper · extraction confidence: 82%
- **neurological_pasc_am** — n=1 paper · extraction confidence: 82%
- **pulmonary_pasc_am** — n=1 paper · extraction confidence: 82%
- **ent_pasc_am** — n=1 paper · extraction confidence: 82%
- **cardiovascular_pasc_am** — n=1 paper · extraction confidence: 82%
- **endocrine_pasc_am** — n=1 paper · extraction confidence: 82%
- **laboratory_pasc_am** — n=1 paper · extraction confidence: 82%
- **eye_pasc_am** — n=1 paper · extraction confidence: 82%
- **skin_pasc_am** — n=1 paper · extraction confidence: 82%
- **immunology_pasc_am** — n=1 paper · extraction confidence: 82%
- **any_psychiatric_disease_new_onset** — n=1 paper · extraction confidence: 82%
- **anxiety_disorder_new_onset** — n=1 paper · extraction confidence: 82%
- **depressive_disorder_new_onset** — n=1 paper · extraction confidence: 82%

- **psychosis_new_onset** — n=1 paper · extraction confidence: 82%
- **substance_abuse_new_onset** — n=1 paper · extraction confidence: 82%
- **fatigue_any_covid** — n=1 paper · extraction confidence: 82%
- **fatigue_moderate_severe_persistent_1month** — n=1 paper · extraction confidence: 82%
- **fatigue_mild_persistent_3months** — n=1 paper · extraction confidence: 82%
- **dry_cough_persistent_1month_moderate_severe** — n=1 paper · extraction confidence: 82%
- **decreased_sense_of_smell_persistent_1month_moderate_severe** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_persistent_1month_moderate_severe** — n=1 paper · extraction confidence: 82%
- **muscle_joint_pain_persistent_1month_moderate_severe** — n=1 paper · extraction confidence: 82%
- **runny_nose_persistent_1month_moderate_severe** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_persistent_1month_female** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_persistent_1month_male** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_persistent_3months_male_none** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_persistent_3months_female_none** — n=1 paper · extraction confidence: 82%
- **obesity_cases** — n=1 paper · extraction confidence: 82%
- **hypertension_cases** — n=1 paper · extraction confidence: 82%
- **chronic_lung_disease_cases** — n=1 paper · extraction confidence: 82%

- **uncomplicated_diabetes_cases** — n=1 paper · extraction confidence: 82%
- **kidney_disease_cases** — n=1 paper · extraction confidence: 82%
- **complicated_diabetes_cases** — n=1 paper · extraction confidence: 82%
- **covid_associated_hospitalization_cases** — n=1 paper · extraction confidence: 82%
- **covid_associated_ed_visit_cases** — n=1 paper · extraction confidence: 82%
- **mechanical_ventilation_cases** — n=1 paper · extraction confidence: 82%
- **sepsis_during_hospitalization_cases** — n=1 paper · extraction confidence: 82%
- **aki_during_hospitalization_cases** — n=1 paper · extraction confidence: 82%
- **corticosteroid_use_cases** — n=1 paper · extraction confidence: 82%
- **remdesivir_use_cases** — n=1 paper · extraction confidence: 82%
- **vasopressor_use_cases** — n=1 paper · extraction confidence: 82%
- **congestive_heart_failure_cases** — n=1 paper · extraction confidence: 82%
- **symptomatic_covid19_spring2020** — n=1 paper · extraction confidence: 82%
- **symptomatic_covid19_2020_2021** — n=1 paper · extraction confidence: 82%
- **symptomatic_covid19_fall2021_4schools** — n=1 paper · extraction confidence: 82%
- **long_covid_spring2020** — n=1 paper · extraction confidence: 82%

- **long_covid_2020_2021** — n=1 paper · extraction confidence: 82%
- **long_covid_fall2021_4schools** — n=1 paper · extraction confidence: 82%
- **self_reported_long_covid** — n=1 paper · extraction confidence: 82%
- **difficulty_thinking_concentrating_memory** — n=1 paper · extraction confidence: 82%
- **fatigue_hospitalised** — n=1 paper · extraction confidence: 82%
- **fatigue_non_hospitalised** — n=1 paper · extraction confidence: 82%
- **fatigue_mixed** — n=1 paper · extraction confidence: 82%
- **pain_discomfort_hospitalised** — n=1 paper · extraction confidence: 82%
- **impaired_sleep_hospitalised** — n=1 paper · extraction confidence: 82%
- **breathlessness_hospitalised** — n=1 paper · extraction confidence: 82%
- **impaired_usual_activity_hospitalised** — n=1 paper · extraction confidence: 82%
- **breathlessness_non_hospitalised** — n=1 paper · extraction confidence: 82%
- **muscle_pain_myalgia_non_hospitalised** — n=1 paper · extraction confidence: 82%
- **affected_sleep_non_hospitalised** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_non_hospitalised** — n=1 paper · extraction confidence: 82%
- **breathlessness_mixed** — n=1 paper · extraction confidence: 82%
- **impaired_usual_activity_mixed** — n=1 paper · extraction confidence: 82%

- **loss_of_taste_mixed** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_mixed** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_hospitalised** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_non_hospitalised** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_mixed** — n=1 paper · extraction confidence: 82%
- **at_least_one_symptom_overall** — n=1 paper · extraction confidence: 82%
- **any_long_covid_symptom** — n=1 paper · extraction confidence: 82%
- **ptsd_ies-r_>=33** — n=1 paper · extraction confidence: 82%
- **olfactory_gustatory_dysfunction_cluster** — n=1 paper · extraction confidence: 82%
- **neurological_symptom_cluster** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_symptom_cluster** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_symptom_cluster** — n=1 paper · extraction confidence: 82%
- **ent_symptom_cluster** — n=1 paper · extraction confidence: 82%
- **dermatological_symptom_cluster** — n=1 paper · extraction confidence: 82%
- **fatigue_senjam** — n=1 paper · extraction confidence: 82%
- **joint_pain_senjam** — n=1 paper · extraction confidence: 82%
- **muscle_pain_senjam** — n=1 paper · extraction confidence: 82%
- **hair_loss_senjam** — n=1 paper · extraction confidence: 82%

- **headache_senjam** — n=1 paper · extraction confidence: 82%
- **breathlessness_senjam** — n=1 paper · extraction confidence: 82%
- **sleep_disturbance_senjam** — n=1 paper · extraction confidence: 82%
- **cough_senjam** — n=1 paper · extraction confidence: 82%
- **fatigue_arnold** — n=1 paper · extraction confidence: 82%
- **breathlessness_arnold** — n=1 paper · extraction confidence: 82%
- **insomnia_arnold** — n=1 paper · extraction confidence: 82%
- **fatigue_kuodi** — n=1 paper · extraction confidence: 82%
- **headache_kuodi** — n=1 paper · extraction confidence: 82%
- **weakness_arms_legs_kuodi** — n=1 paper · extraction confidence: 82%
- **persistent_muscle_pain_kuodi** — n=1 paper · extraction confidence: 82%
- **fatigue_gaber** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_gaber** — n=1 paper · extraction confidence: 82%
- **anxiety_gaber** — n=1 paper · extraction confidence: 82%
- **any_long_covid_simon** — n=1 paper · extraction confidence: 82%
- **two_plus_long_covid_symptoms_simon** — n=1 paper · extraction confidence: 82%
- **any_pre_existing_or_pcc_associated_condition** — n=1 paper · extraction confidence: 82%
- **cardiovascular_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **cardiovascular_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **neuropsychiatric_pre_pandemic** — n=1 paper · extraction confidence: 82%

- **neuropsychiatric_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **respiratory_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **respiratory_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **systemic_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **systemic_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **endocrine_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **endocrine_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **renal_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **renal_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **sensory_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **sensory_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **hematologic_pre_pandemic** — n=1 paper · extraction confidence: 82%

- **hematologic_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **dermatologic_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **dermatologic_pandemic_pre_index** — n=1 paper · extraction confidence: 82%
- **post covid condition (fatigue, breathlessness, or neurocognitive abnormalities at 4-8 weeks post discharge)** — n=1 paper · extraction confidence: 82%
- **brain_fog_pnp_usa** — n=1 paper · extraction confidence: 82%
- **brain_fog_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **brain_fog_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **brain_fog_pnp_india** — n=1 paper · extraction confidence: 82%
- **brain_fog_nnp_usa** — n=1 paper · extraction confidence: 82%
- **brain_fog_nnp_colombia** — n=1 paper · extraction confidence: 82%
- **brain_fog_nnp_nigeria** — n=1 paper · extraction confidence: 82%
- **brain_fog_nnp_india** — n=1 paper · extraction confidence: 82%
- **myalgia_pnp_usa** — n=1 paper · extraction confidence: 82%
- **myalgia_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **myalgia_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **myalgia_pnp_india** — n=1 paper · extraction confidence: 82%

- **dizziness_pnp_usa** — n=1 paper · extraction confidence: 82%
- **dizziness_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **dizziness_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **dizziness_pnp_india** — n=1 paper · extraction confidence: 82%
- **headache_pnp_usa** — n=1 paper · extraction confidence: 82%
- **headache_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **headache_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **headache_pnp_india** — n=1 paper · extraction confidence: 82%
- **fatigue_pnp_usa** — n=1 paper · extraction confidence: 82%
- **fatigue_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **fatigue_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **fatigue_pnp_india** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_pnp_usa** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_pnp_india** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_pnp_usa** — n=1 paper · extraction confidence: 82%

- **depression_anxiety_nnp_colombia** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_nnp_nigeria** — n=1 paper · extraction confidence: 82%
- **depression_anxiety_nnp_india** — n=1 paper · extraction confidence: 82%
- **abnormal_neurologic_exam_pnp_overall** — n=1 paper · extraction confidence: 82%
- **abnormal_neurologic_exam_nnp_overall** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_pnp_usa** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_pnp_nigeria** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_pnp_india** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_nnp_usa** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_nnp_colombia** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_nnp_nigeria** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_nnp_india** — n=1 paper · extraction confidence: 82%
- **qol_abnormal_depression_anxiety_pnp_usa** — n=1 paper · extraction confidence: 82%
- **qol_abnormal_depression_anxiety_pnp_colombia** — n=1 paper · extraction confidence: 82%
- **qol_abnormal_depression_anxiety_pnp_india** — n=1 paper · extraction confidence: 82%

- **qol_abnormal_depression_anxiety_nnp_usa** — n=1 paper · extraction confidence: 82%
- **qol_abnormal_depression_anxiety_nnp_colombia** — n=1 paper · extraction confidence: 82%
- **qol_abnormal_depression_anxiety_nnp_india** — n=1 paper · extraction confidence: 82%
- **at_least_one_incident_pasc_condition_insight** — n=1 paper · extraction confidence: 82%
- **fatigue_cluster_change** — n=1 paper · extraction confidence: 82%
- **neurocognitive_impairment_change** — n=1 paper · extraction confidence: 82%
- **chest_symptoms_change** — n=1 paper · extraction confidence: 82%
- **smell_taste_disorder_change** — n=1 paper · extraction confidence: 82%
- **anxiety_depression_sleep_disorders_change** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_pain_change** — n=1 paper · extraction confidence: 82%
- **long covid (heds cohort)** — n=1 paper · extraction confidence: 82%
- **long covid (matched controls)** — n=1 paper · extraction confidence: 82%
- **covid-19 infection (heds cohort)** — n=1 paper · extraction confidence: 82%
- **covid-19 infection (matched controls)** — n=1 paper · extraction confidence: 82%
- **mcas in heds with long covid** — n=1 paper · extraction confidence: 82%
- **mcas in heds without long covid** — n=1 paper · extraction confidence: 82%

- **pots in heds with long covid** — n=1 paper · extraction confidence: 82%
- **pots in heds without long covid** — n=1 paper · extraction confidence: 82%
- **me/cfs in heds with long covid** — n=1 paper · extraction confidence: 82%
- **me/cfs in heds without long covid** — n=1 paper · extraction confidence: 82%
- **mcas+pots in heds with long covid** — n=1 paper · extraction confidence: 82%
- **mcas+pots in heds without long covid** — n=1 paper · extraction confidence: 82%
- **mcas+me/cfs in heds with long covid** — n=1 paper · extraction confidence: 82%
- **mcas+me/cfs in heds without long covid** — n=1 paper · extraction confidence: 82%
- **pots+me/cfs in heds with long covid** — n=1 paper · extraction confidence: 82%
- **pots+me/cfs in heds without long covid** — n=1 paper · extraction confidence: 82%
- **mcas+pots+me/cfs in heds with long covid** — n=1 paper · extraction confidence: 82%
- **mcas+pots+me/cfs in heds without long covid** — n=1 paper · extraction confidence: 82%
- **me/cfs in controls with long covid** — n=1 paper · extraction confidence: 82%
- **decreased_smell_sensation_infected** — n=1 paper · extraction confidence: 82%
- **decreased_smell_sensation_naive** — n=1 paper · extraction confidence: 82%
- **decreased_taste_sensation_infected** — n=1 paper · extraction confidence: 82%

- **decreased_taste_sensation_naive** — n=1 paper · extraction confidence: 82%
- **memory_disturbances_infected** — n=1 paper · extraction confidence: 82%
- **memory_disturbances_naive** — n=1 paper · extraction confidence: 82%
- **dyspnea_infected** — n=1 paper · extraction confidence: 82%
- **dyspnea_naive** — n=1 paper · extraction confidence: 82%
- **arthralgia_infected** — n=1 paper · extraction confidence: 82%
- **arthralgia_naive** — n=1 paper · extraction confidence: 82%
- **cough_infected** — n=1 paper · extraction confidence: 82%
- **cough_naive** — n=1 paper · extraction confidence: 82%
- **disturbed_vision_infected** — n=1 paper · extraction confidence: 82%
- **disturbed_vision_naive** — n=1 paper · extraction confidence: 82%
- **chest_pain_infected** — n=1 paper · extraction confidence: 82%
- **chest_pain_naive** — n=1 paper · extraction confidence: 82%
- **weakness_infected** — n=1 paper · extraction confidence: 82%
- **weakness_naive** — n=1 paper · extraction confidence: 82%
- **myalgia_infected** — n=1 paper · extraction confidence: 82%
- **myalgia_naive** — n=1 paper · extraction confidence: 82%
- **increased_heart_rate_infected** — n=1 paper · extraction confidence: 82%
- **increased_heart_rate_naive** — n=1 paper · extraction confidence: 82%
- **fatigue_infected** — n=1 paper · extraction confidence: 82%
- **fatigue_naive** — n=1 paper · extraction confidence: 82%
- **nausea_infected** — n=1 paper · extraction confidence: 82%
- **nausea_naive** — n=1 paper · extraction confidence: 82%

- **headache_infected** — n=1 paper · extraction confidence: 82%
- **headache_naive** — n=1 paper · extraction confidence: 82%
- **decreased_libido_infected** — n=1 paper · extraction confidence: 82%
- **decreased_libido_naive** — n=1 paper · extraction confidence: 82%
- **any_symptom_naive** — n=1 paper · extraction confidence: 82%
- **not_returning_to_baseline_infected** — n=1 paper · extraction confidence: 82%
- **not_returning_to_baseline_naive** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_morello_2023_overall** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_lokanuwatsatien_2023_overall** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_pinto_pereira_2023_vaccinated** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_pinto_pereira_2023_unvaccinated** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_jarupan_2023_children_5_11** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_jarupan_2023_adolescents_12_17** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_messiah_2022_4_12weeks** — n=1 paper · extraction confidence: 82%
- **post_covid_conditions_messiah_2022_gte12weeks** — n=1 paper · extraction confidence: 82%
- **tested_positive_for_covid19_among_all_respondents** — n=1 paper · extraction confidence: 82%
- **moderate_severe_covid19_among_those_testing_positive** — n=1 paper · extraction confidence: 82%

- **long_covid_among_those_testing_positive** — n=1 paper · extraction confidence: 82%
- **post_acute_covid19_symptoms_any_sidiap** — n=1 paper · extraction confidence: 82%
- **post_acute_covid19_symptoms_any_cprd** — n=1 paper · extraction confidence: 82%
- **post_acute_symptoms_negative_test_sidiap** — n=1 paper · extraction confidence: 82%
- **post_acute_symptoms_negative_test_cprd** — n=1 paper · extraction confidence: 82%
- **ongoing_symptoms_28days_sidiap** — n=1 paper · extraction confidence: 82%
- **ongoing_symptoms_28days_cprd** — n=1 paper · extraction confidence: 82%
- **any_incident_pasc_condition** — n=1 paper · extraction confidence: 82%
- **cough_post_covid** — n=1 paper · extraction confidence: 82%
- **difficulty_breathing_post_covid** — n=1 paper · extraction confidence: 82%
- **tiredness_post_covid** — n=1 paper · extraction confidence: 82%
- **brain_fog_post_covid** — n=1 paper · extraction confidence: 82%
- **sleep_problems_post_covid** — n=1 paper · extraction confidence: 82%
- **brain_fog_acute** — n=1 paper · extraction confidence: 82%
- **sleep_problems_acute** — n=1 paper · extraction confidence: 82%
- **dyspnea_ab+rna+** — n=1 paper · extraction confidence: 82%
- **dyspnea_ab+rna-** — n=1 paper · extraction confidence: 82%
- **any_persistent_symptoms_overall_prior_covid** — n=1 paper · extraction confidence: 82%

- **any_recurrent_symptoms_overall_prior_covid** — n=1 paper · extraction confidence: 82%
- **any_new_symptoms_overall_prior_covid** — n=1 paper · extraction confidence: 82%
- **persistent_fatigue_among_persistent_symptom_reporters** — n=1 paper · extraction confidence: 82%
- **persistent_brain_fog_among_persistent_symptom_reporters** — n=1 paper · extraction confidence: 82%
- **persistent_dyspnea_among_persistent_symptom_reporters** — n=1 paper · extraction confidence: 82%
- **any_persistent_symptoms_ab+rna+** — n=1 paper · extraction confidence: 82%
- **any_persistent_symptoms_ab+rna-** — n=1 paper · extraction confidence: 82%
- **any_persistent_symptoms_ab-rna+** — n=1 paper · extraction confidence: 82%
- **any_persistent_symptoms_ab-rna-_prior_covid** — n=1 paper · extraction confidence: 82%
- **at_least_one_pasc_symptom** — n=1 paper · extraction confidence: 82%
- **multiple_pasc_symptoms** — n=1 paper · extraction confidence: 82%
- **dyspepsia** — n=1 paper · extraction confidence: 82%
- **cognitive_changes** — n=1 paper · extraction confidence: 82%
- **at_least_one_long_covid_symptom_overall** — n=1 paper · extraction confidence: 82%
- **fatigue_remained_inactive** — n=1 paper · extraction confidence: 82%
- **fatigue_became_inactive** — n=1 paper · extraction confidence: 82%
- **fatigue_remained_active** — n=1 paper · extraction confidence: 82%

- **memory_loss_remained_inactive** — n=1 paper · extraction confidence: 82%
- **memory_loss_became_inactive** — n=1 paper · extraction confidence: 82%
- **memory_loss_remained_active** — n=1 paper · extraction confidence: 82%
- **headache_remained_inactive** — n=1 paper · extraction confidence: 82%
- **headache_became_inactive** — n=1 paper · extraction confidence: 82%
- **headache_remained_active** — n=1 paper · extraction confidence: 82%
- **anosmia_remained_inactive** — n=1 paper · extraction confidence: 82%
- **anosmia_became_inactive** — n=1 paper · extraction confidence: 82%
- **anosmia_remained_active** — n=1 paper · extraction confidence: 82%
- **attention_loss_remained_inactive** — n=1 paper · extraction confidence: 82%
- **attention_loss_became_inactive** — n=1 paper · extraction confidence: 82%
- **attention_loss_remained_active** — n=1 paper · extraction confidence: 82%
- **arthralgia_remained_inactive** — n=1 paper · extraction confidence: 82%
- **arthralgia_became_inactive** — n=1 paper · extraction confidence: 82%
- **arthralgia_remained_active** — n=1 paper · extraction confidence: 82%
- **myalgia_remained_inactive** — n=1 paper · extraction confidence: 82%

- **myalgia_became_inactive** — n=1 paper · extraction confidence: 82%
- **myalgia_remained_active** — n=1 paper · extraction confidence: 82%
- **dyspnea_remained_inactive** — n=1 paper · extraction confidence: 82%
- **dyspnea_became_inactive** — n=1 paper · extraction confidence: 82%
- **dyspnea_remained_active** — n=1 paper · extraction confidence: 82%
- **ageusia_remained_inactive** — n=1 paper · extraction confidence: 82%
- **ageusia_became_inactive** — n=1 paper · extraction confidence: 82%
- **ageusia_remained_active** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_remained_inactive** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_remained_active** — n=1 paper · extraction confidence: 82%
- **neurological_remained_inactive** — n=1 paper · extraction confidence: 82%
- **neurological_remained_active** — n=1 paper · extraction confidence: 82%
- **respiratory_remained_inactive** — n=1 paper · extraction confidence: 82%
- **respiratory_remained_active** — n=1 paper · extraction confidence: 82%
- **long_covid_remained_inactive** — n=1 paper · extraction confidence: 82%
- **long_covid_became_inactive** — n=1 paper · extraction confidence: 82%

- **long_covid_remained_active** — n=1 paper · extraction confidence: 82%
- **any_long_covid_diagnosis_lower_bound** — n=1 paper · extraction confidence: 82%
- **any_long_covid_diagnosis_upper_bound** — n=1 paper · extraction confidence: 82%
- **cardiology_long_covid_cases** — n=1 paper · extraction confidence: 82%
- **neurology_long_covid_cases** — n=1 paper · extraction confidence: 82%
- **ophthalmology_long_covid_cases** — n=1 paper · extraction confidence: 82%
- **pulmonary_long_covid_cases** — n=1 paper · extraction confidence: 82%
- **first_long_covid_code_within_3_months** — n=1 paper · extraction confidence: 82%
- **brain_fog_cognitive_issues_mentioned_in_tweets** — n=1 paper · extraction confidence: 82%
- **chronic_fatigue_cfs_mentioned_in_tweets** — n=1 paper · extraction confidence: 82%
- **pulmonary_complaints_test_positive_2months** — n=1 paper · extraction confidence: 82%
- **pulmonary_complaints_test_positive_6months_excess_vs_negative** — n=1 paper · extraction confidence: 82%
- **general_complaints_test_positive_2months_excess_vs_negative** — n=1 paper · extraction confidence: 82%
- **general_complaints_test_positive_6months_excess_vs_negative** — n=1 paper · extraction confidence: 82%
- **neurological_complaints_test_positive_6months_excess_vs_negative** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_disease_post_acute_discovery** — n=1 paper · extraction confidence: 82%

- **hepatobiliary_disease_post_acute_discovery** — n=1 paper
· extraction confidence: 82%
- **other_digestive_abnormalities_post_acute_discovery** —
n=1 paper · extraction confidence: 82%
- **fatigue_covid_group** — n=1 paper · extraction confidence:
82%
- **back_pain_covid_group** — n=1 paper · extraction
confidence: 82%
- **sleep_disturbance_covid_group** — n=1 paper · extraction
confidence: 82%
- **headache_covid_group** — n=1 paper · extraction
confidence: 82%
- **poor_concentration_covid_group** — n=1 paper · extraction
confidence: 82%
- **gastrointestinal_upset_covid_group** — n=1 paper ·
extraction confidence: 82%
- **arm_leg_joint_pain_covid_group** — n=1 paper · extraction
confidence: 82%
- **dizziness_covid_group** — n=1 paper · extraction confidence:
82%
- **hair_loss_covid_group** — n=1 paper · extraction confidence:
82%
- **erectile_dysfunction_covid_male_group** — n=1 paper ·
extraction confidence: 82%
- **decreased_libido_covid_group** — n=1 paper · extraction
confidence: 82%
- **dyspnea_covid_group** — n=1 paper · extraction confidence:
82%
- **hearing_loss_covid_group** — n=1 paper · extraction
confidence: 82%
- **memory_impairment_covid_group** — n=1 paper ·
extraction confidence: 82%

- **chest_discomfort_covid_group** — n=1 paper · extraction confidence: 82%
- **dysgeusia_covid_group** — n=1 paper · extraction confidence: 82%
- **dysosmia_covid_group** — n=1 paper · extraction confidence: 82%
- **cough_covid_group** — n=1 paper · extraction confidence: 82%
- **fever_covid_group** — n=1 paper · extraction confidence: 82%
- **dysphonia_during_acute_covid19_infection** — n=1 paper · extraction confidence: 82%
- **dysphonia_post_covid19_infection** — n=1 paper · extraction confidence: 82%
- **long_covid_dysphonia_≥3_months_post_recovery** — n=1 paper · extraction confidence: 82%
- **moderately_severe_or_worse_functional_impairment_wsas_baseline** — n=1 paper · extraction confidence: 82%
- **moderately_severe_or_worse_functional_impairment_wsas_6months** — n=1 paper · extraction confidence: 82%
- **moderately_severe_or_worse_functional_impairment_wsas_6months** — n=1 paper · extraction confidence: 82%
- **at_least_1_day_off_work_baseline** — n=1 paper · extraction confidence: 82%
- **at_least_1_day_off_work_6months_completers** — n=1 paper · extraction confidence: 82%
- **20_or_more_working_days_lost_baseline** — n=1 paper · extraction confidence: 82%
- **20_or_more_working_days_lost_6months_completers** — n=1 paper · extraction confidence: 82%
- **sarcopenia_acute_covid19** — n=1 paper · extraction confidence: 82%

- **sarcopenia_long_covid** — n=1 paper · extraction confidence: 82%
- **pots-like symptoms after covid-19 (large-scale online survey)** — n=1 paper · extraction confidence: 82%
- **orthostatic hypotension in post-covid-19 individuals (shah et al.)** — n=1 paper · extraction confidence: 82%
- **orthostatic intolerance in long covid (cited cross-sectional study)** — n=1 paper · extraction confidence: 82%
- **post exertional malaise (most accessed topic)** — n=1 paper · extraction confidence: 82%
- **cognitive difficulties** — n=1 paper · extraction confidence: 82%
- **shortness of breath** — n=1 paper · extraction confidence: 82%
- **long_covid_any_pcornet_pregnant** — n=1 paper · extraction confidence: 82%
- **long_covid_any_pcornet_nonpregnant** — n=1 paper · extraction confidence: 82%
- **long_covid_any_n3c_pregnant** — n=1 paper · extraction confidence: 82%
- **long_covid_any_n3c_nonpregnant** — n=1 paper · extraction confidence: 82%
- **unspecified_pasc_u099_b948_pcornet_pregnant** — n=1 paper · extraction confidence: 82%
- **unspecified_pasc_u099_b948_pcornet_nonpregnant** — n=1 paper · extraction confidence: 82%
- **unspecified_pasc_u099_b948_n3c_pregnant** — n=1 paper · extraction confidence: 82%
- **unspecified_pasc_u099_b948_n3c_nonpregnant** — n=1 paper · extraction confidence: 82%
- **cognitive_fatigue_respiratory_pcornet_pregnant** — n=1 paper · extraction confidence: 82%

- **cognitive_fatigue_respiratory_pcornet_nonpregnant** — n=1 paper · extraction confidence: 82%
- **cognitive_fatigue_respiratory_n3c_pregnant** — n=1 paper · extraction confidence: 82%
- **cognitive_fatigue_respiratory_n3c_nonpregnant** — n=1 paper · extraction confidence: 82%
- **clouded_consciousness** — n=1 paper · extraction confidence: 82%
- **sore_throat_acute** — n=1 paper · extraction confidence: 82%
- **mood_disorders_acute** — n=1 paper · extraction confidence: 82%
- **sleep_disturbance_acute** — n=1 paper · extraction confidence: 82%
- **joint_pain_acute** — n=1 paper · extraction confidence: 82%
- **rhinobyon_acute** — n=1 paper · extraction confidence: 82%
- **smell_taste_disorder_acute** — n=1 paper · extraction confidence: 82%
- **runny_nose_acute** — n=1 paper · extraction confidence: 82%
- **muscle_weakness_acute** — n=1 paper · extraction confidence: 82%
- **dyspnea_acute** — n=1 paper · extraction confidence: 82%
- **nausea_acute** — n=1 paper · extraction confidence: 82%
- **palpitation_acute** — n=1 paper · extraction confidence: 82%
- **diarrhea_acute** — n=1 paper · extraction confidence: 82%
- **sore_eye_acute** — n=1 paper · extraction confidence: 82%
- **muscle_weakness_long_covid** — n=1 paper · extraction confidence: 82%
- **smell_taste_disorder_long_covid** — n=1 paper · extraction confidence: 82%
- **rhinobyon_runny_nose_long_covid** — n=1 paper · extraction confidence: 82%

- **sore_throat_long_covid** — n=1 paper · extraction confidence: 82%
- **memory_impairment_long_covid** — n=1 paper · extraction confidence: 82%
- **mood_disorders_long_covid** — n=1 paper · extraction confidence: 82%
- **lumbago_long_covid** — n=1 paper · extraction confidence: 82%
- **dyspnea_long_covid** — n=1 paper · extraction confidence: 82%
- **hearing_loss_long_covid** — n=1 paper · extraction confidence: 82%
- **sore_eye_long_covid** — n=1 paper · extraction confidence: 82%
- **palpitation_long_covid** — n=1 paper · extraction confidence: 82%
- **long_covid_rate_overall** — n=1 paper · extraction confidence: 82%
- **any_kidney_event_covid_positive** — n=1 paper · extraction confidence: 82%
- **any_kidney_event_covid_negative** — n=1 paper · extraction confidence: 82%
- **death_in_hospital_positives_first_4_weeks** — n=1 paper · extraction confidence: 82%
- **death_in_hospital_negatives_first_4_weeks** — n=1 paper · extraction confidence: 82%
- **long_covid_history_adults_18plus_2022** — n=1 paper · extraction confidence: 82%
- **long_covid_history_working_aged_18_64_2022** — n=1 paper · extraction confidence: 82%
- **long_covid_history_adults_18plus_2023** — n=1 paper · extraction confidence: 82%

- **long_covid_history_working_aged_18_64_2023** — n=1 paper · extraction confidence: 82%
- **current_long_covid_symptoms_among_those_with_history** — n=1 paper · extraction confidence: 82%
- **modest_to_considerable_daily_functioning_impact** — n=1 paper · extraction confidence: 82%
- **neurocognitive_and_neurologic_group_among_long_covid** — n=1 paper · extraction confidence: 82%
- **systemic_group_among_long_covid** — n=1 paper · extraction confidence: 82%
- **cardiopulmonary_group_among_long_covid** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_group_among_long_covid** — n=1 paper · extraction confidence: 82%
- **cardiopulmonary_among_asian_hcws_with_long_covid** — n=1 paper · extraction confidence: 82%
- **systemic_among_black_hcws_with_long_covid** — n=1 paper · extraction confidence: 82%
- **systemic_among_mixed_ethnicity_hcws_with_long_covid** — n=1 paper · extraction confidence: 82%
- **long_covid_prevalence_overall** — n=1 paper · extraction confidence: 82%
- **ckd_development_range_low** — n=1 paper · extraction confidence: 82%
- **ckd_development_range_high** — n=1 paper · extraction confidence: 82%
- **ckd_progression_range_low** — n=1 paper · extraction confidence: 82%
- **ckd_progression_range_high** — n=1 paper · extraction confidence: 82%
- **aki_during_hospitalization_range_low** — n=1 paper · extraction confidence: 82%

- **aki_during_hospitalization_range_high** — n=1 paper · extraction confidence: 82%
- **krt_dependence_range_low** — n=1 paper · extraction confidence: 82%
- **krt_dependence_range_high** — n=1 paper · extraction confidence: 82%
- **albuminuria_bento_et_al** — n=1 paper · extraction confidence: 82%
- **proteinuria_li_et_al_2yr** — n=1 paper · extraction confidence: 82%
- **abnormal_egfr_zhou_et_al_3mo** — n=1 paper · extraction confidence: 82%
- **elevated_scys_yang_et_al_3mo** — n=1 paper · extraction confidence: 82%
- **malaise_or_fatigue_as_index_symptom** — n=1 paper · extraction confidence: 82%
- **depression_as_index_symptom** — n=1 paper · extraction confidence: 82%
- **concentration_or_memory_impairment_as_index_symptom** — n=1 paper · extraction confidence: 82%
- **adhd_as_index_diagnosis** — n=1 paper · extraction confidence: 82%
- **long_covid_specific_diagnosis_code** — n=1 paper · extraction confidence: 82%
- **fatigue_exhaustion_pcc_patients** — n=1 paper · extraction confidence: 82%
- **respiratory_symptoms_sob_dyspnea_pcc_patients** — n=1 paper · extraction confidence: 82%
- **heart_palpitations_pcc_patients** — n=1 paper · extraction confidence: 82%
- **disturbances_of_taste_and_smell_pcc_patients** — n=1 paper · extraction confidence: 82%
- **pots_pcc_patients** — n=1 paper · extraction confidence: 82%

- **fatigue_exhaustion_non_pcc_patients** — n=1 paper · extraction confidence: 82%
- **respiratory_symptoms_non_pcc_patients** — n=1 paper · extraction confidence: 82%
- **heart_palpitations_non_pcc_patients** — n=1 paper · extraction confidence: 82%
- **disturbances_of_taste_and_smell_non_pcc_patients** — n=1 paper · extraction confidence: 82%
- **pots_non_pcc_patients** — n=1 paper · extraction confidence: 82%
- **respiratory_signs_and_symptoms_covid_positive** — n=1 paper · extraction confidence: 82%
- **respiratory_signs_and_symptoms_covid_negative** — n=1 paper · extraction confidence: 82%
- **long_covid_current** — n=1 paper · extraction confidence: 82%
- **long_covid_ever** — n=1 paper · extraction confidence: 82%
- **prior_covid_19** — n=1 paper · extraction confidence: 82%
- **sleep_difficulties** — n=1 paper · extraction confidence: 82%
- **chest_tightness_chest_pain** — n=1 paper · extraction confidence: 82%
- **anxiety_gad7_ge5_long_covid_group** — n=1 paper · extraction confidence: 82%
- **depression_phq9_ge5_long_covid_group** — n=1 paper · extraction confidence: 82%
- **ptsd_pclc_ge38_long_covid_group** — n=1 paper · extraction confidence: 82%
- **breathing_difficulties** — n=1 paper · extraction confidence: 82%
- **inappetence** — n=1 paper · extraction confidence: 82%
- **impaired_hearing** — n=1 paper · extraction confidence: 82%
- **impaired_vision** — n=1 paper · extraction confidence: 82%

- **fatigue_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **post_exertional_malaise_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **joint_pain_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **sleep_disturbance_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **numbness_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **anxiety_symptoms_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **little_interest_feeling_down_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **neurologic_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **psychiatric_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **general_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **respiratory_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **ear_nose_throat_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **dermatological_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **cardiovascular_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%

- **gastrointestinal_symptoms_group_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **ptsd_symptoms_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **breathlessness_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **problems_with_vision_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **difficulty_walking_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **muscle_pain_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **hair_loss_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **dizziness_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **headaches_migraines_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **chest_pain_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **cough_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **palpitations_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **loss_of_taste_smell_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **tinnitus_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **weight_loss_reduced_appetite_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **abdominal_pain_nausea_vomiting_diarrhea_frequently_occurring** — n=1 paper · extraction confidence: 82%

- **skin_rash_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **fever_frequently_occurring** — n=1 paper · extraction confidence: 82%
- **at_least_one_frequently_occurring_symptom** — n=1 paper · extraction confidence: 82%
- **at_least_one_present_symptom** — n=1 paper · extraction confidence: 82%
- **self_reported_lc** — n=1 paper · extraction confidence: 82%
- **upper_respiratory_symptoms** — n=1 paper · extraction confidence: 82%
- **fever_and_chills** — n=1 paper · extraction confidence: 82%
- **allergies** — n=1 paper · extraction confidence: 82%
- **cardiorespiratory_signs_and_symptoms** — n=1 paper · extraction confidence: 82%
- **general_signs_and_symptoms** — n=1 paper · extraction confidence: 82%
- **obesity** — n=1 paper · extraction confidence: 82%
- **nausea_and_vomiting** — n=1 paper · extraction confidence: 82%
- **gastroenteritis** — n=1 paper · extraction confidence: 82%
- **fatigue_and_malaise** — n=1 paper · extraction confidence: 82%
- **disorders_of_teeth_and_gingiva** — n=1 paper · extraction confidence: 82%
- **mental_health_treatment** — n=1 paper · extraction confidence: 82%
- **communication_and_motor_disorders** — n=1 paper · extraction confidence: 82%
- **bronchiolitis** — n=1 paper · extraction confidence: 82%
- **tonsillitis** — n=1 paper · extraction confidence: 82%
- **inflammatory_skin_conditions** — n=1 paper · extraction confidence: 82%

- **genitourinary_signs_and_symptoms** — n=1 paper · extraction confidence: 82%
- **generalized_pain** — n=1 paper · extraction confidence: 82%
- **abnormal_liver_enzymes** — n=1 paper · extraction confidence: 82%
- **fluid_and_electrolyte_disturbances** — n=1 paper · extraction confidence: 82%
- **pneumonia** — n=1 paper · extraction confidence: 82%
- **myocarditis** — n=1 paper · extraction confidence: 82%
- **anxiety_symptoms** — n=1 paper · extraction confidence: 82%
- **post_covid_code_u09_9** — n=1 paper · extraction confidence: 82%
- **anxiety_symptoms_score_ge3_all_respondents** — n=1 paper · extraction confidence: 82%
- **anxiety_symptoms_score_ge3_covid19_positive** — n=1 paper · extraction confidence: 82%
- **anxiety_symptoms_score_ge3_long_covid** — n=1 paper · extraction confidence: 82%
- **depressive_symptoms_score_ge3_all_respondents** — n=1 paper · extraction confidence: 82%
- **depressive_symptoms_score_ge3_covid19_positive** — n=1 paper · extraction confidence: 82%
- **depressive_symptoms_score_ge3_long_covid** — n=1 paper · extraction confidence: 82%
- **long_covid_among_covid19_positive** — n=1 paper · extraction confidence: 82%
- **covid19_positive_among_all_respondents** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_long_covid** — n=1 paper · extraction confidence: 82%
- **dyspnoea_long_covid** — n=1 paper · extraction confidence: 82%

- **muscular_exhaustion_long_covid** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_long_covid** — n=1 paper · extraction confidence: 82%
- **loss_of_taste_long_covid** — n=1 paper · extraction confidence: 82%
- **flu_like_symptoms_acute** — n=1 paper · extraction confidence: 82%
- **muscle_pain_acute** — n=1 paper · extraction confidence: 82%
- **loss_of_appetite_acute** — n=1 paper · extraction confidence: 82%
- **diarrhoea_acute** — n=1 paper · extraction confidence: 82%
- **subphenotype_1_cardiac_renal_proportion** — n=1 paper · extraction confidence: 82%
- **subphenotype_2_respiratory_sleep_anxiety_proportion** — n=1 paper · extraction confidence: 82%
- **subphenotype_3_musculoskeletal_nervous_proportion** — n=1 paper · extraction confidence: 82%
- **subphenotype_4_digestive_respiratory_proportion** — n=1 paper · extraction confidence: 82%
- **subphenotype_1_hospitalization_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_2_hospitalization_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_3_hospitalization_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_4_hospitalization_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_1_mechanical_ventilation_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_2_mechanical_ventilation_rate** — n=1 paper · extraction confidence: 82%

- **subphenotype_3_mechanical_ventilation_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_4_mechanical_ventilation_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_1_critical_care_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_2_critical_care_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_3_critical_care_rate** — n=1 paper · extraction confidence: 82%
- **subphenotype_4_critical_care_rate** — n=1 paper · extraction confidence: 82%
- **symptoms_lasting_less_than_2_weeks** — n=1 paper · extraction confidence: 82%
- **symptoms_lasting_2_to_3_weeks** — n=1 paper · extraction confidence: 82%
- **symptoms_lasting_4_to_12_weeks** — n=1 paper · extraction confidence: 82%
- **symptoms_lasting_more_than_12_weeks** — n=1 paper · extraction confidence: 82%
- **dyspnea_and_respiratory_failure_post** — n=1 paper · extraction confidence: 82%
- **myositis_and_msk_pain_stiffness_post** — n=1 paper · extraction confidence: 82%
- **anxiety_and_depression_post** — n=1 paper · extraction confidence: 82%
- **dizziness_and_headache_post** — n=1 paper · extraction confidence: 82%
- **chest_pain_post** — n=1 paper · extraction confidence: 82%
- **arrythmias_post** — n=1 paper · extraction confidence: 82%
- **nausea_vomiting_diarrhea_post** — n=1 paper · extraction confidence: 82%

- **bronchiectasis_and_cough_post** — n=1 paper · extraction confidence: 82%
- **fatigue_post** — n=1 paper · extraction confidence: 82%
- **sleep_disturbances_post** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_or_taste_post** — n=1 paper · extraction confidence: 82%
- **pulmonary_embolism_post** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_post** — n=1 paper · extraction confidence: 82%
- **kidney_liver_pancreas_spleen_injury_post** — n=1 paper · extraction confidence: 82%
- **chf_and_cardiomyopathy_post** — n=1 paper · extraction confidence: 82%
- **other_psychiatric_disorder_post** — n=1 paper · extraction confidence: 82%
- **platelet_clotting_dysfunctions_post** — n=1 paper · extraction confidence: 82%
- **stroke_post** — n=1 paper · extraction confidence: 82%
- **pulmonary_fibrosis_post** — n=1 paper · extraction confidence: 82%
- **pericarditis_and_myocarditis_post** — n=1 paper · extraction confidence: 82%
- **acs_post** — n=1 paper · extraction confidence: 82%
- **long_covid_among_covid_positive_hcws** — n=1 paper · extraction confidence: 82%
- **long_covid_among_whole_cohort** — n=1 paper · extraction confidence: 82%
- **symptoms_more_than_3_months_among_covid_positive** — n=1 paper · extraction confidence: 82%
- **symptoms_more_than_6_months_among_covid_positive** — n=1 paper · extraction confidence: 82%

- **fever_overall** — n=1 paper · extraction confidence: 82%
- **cough_confirmed** — n=1 paper · extraction confidence: 82%
- **fatigue_confirmed** — n=1 paper · extraction confidence: 82%
- **fever_confirmed** — n=1 paper · extraction confidence: 82%
- **productive_cough_confirmed** — n=1 paper · extraction confidence: 82%
- **muscle_or_joint_pain_confirmed** — n=1 paper · extraction confidence: 82%
- **sore_throat_confirmed** — n=1 paper · extraction confidence: 82%
- **stuffy_or_runny_nose_confirmed** — n=1 paper · extraction confidence: 82%
- **cough_probable** — n=1 paper · extraction confidence: 82%
- **fatigue_probable** — n=1 paper · extraction confidence: 82%
- **fever_probable** — n=1 paper · extraction confidence: 82%
- **productive_cough_probable** — n=1 paper · extraction confidence: 82%
- **muscle_or_joint_pain_probable** — n=1 paper · extraction confidence: 82%
- **sore_throat_probable** — n=1 paper · extraction confidence: 82%
- **stuffy_or_runny_nose_probable** — n=1 paper · extraction confidence: 82%
- **asthenia_acute** — n=1 paper · extraction confidence: 82%
- **myalgia_acute** — n=1 paper · extraction confidence: 82%
- **arthralgia_acute** — n=1 paper · extraction confidence: 82%
- **pcc_at_3_months** — n=1 paper · extraction confidence: 82%
- **pcc_at_6_months** — n=1 paper · extraction confidence: 82%
- **pcc_at_12_months** — n=1 paper · extraction confidence: 82%
- **pcc_at_18_months** — n=1 paper · extraction confidence: 82%

- **sspd_covid19_positive** — n=1 paper · extraction confidence: 82%
- **sspd_covid19_negative** — n=1 paper · extraction confidence: 82%
- **at_least_1_long_covid_symptom_cases** — n=1 paper · extraction confidence: 82%
- **at_least_1_long_covid_symptom_controls** — n=1 paper · extraction confidence: 82%
- **worsened_general_health_cases** — n=1 paper · extraction confidence: 82%
- **worsened_general_health_controls** — n=1 paper · extraction confidence: 82%
- **worsened_mental_health_cases** — n=1 paper · extraction confidence: 82%
- **worsened_mental_health_controls** — n=1 paper · extraction confidence: 82%
- **pre_infection_any_symptom_long_covid_group_median** — n=1 paper · extraction confidence: 82%
- **pre_infection_any_symptom_recovered_group_median** — n=1 paper · extraction confidence: 82%
- **pots_pre_infection** — n=1 paper · extraction confidence: 82%
- **pvfs_pre_infection** — n=1 paper · extraction confidence: 82%
- **ageusia_pre_infection** — n=1 paper · extraction confidence: 82%
- **anosmia_pre_infection** — n=1 paper · extraction confidence: 82%
- **long_covid_symptomatic_improvement_post_vaccination** — n=1 paper · extraction confidence: 82%
- **long_covid_symptomatic_worsening_post_vaccination** — n=1 paper · extraction confidence: 82%

- **long_covid_symptomatic_unchanged_post_vaccination** — n=1 paper · extraction confidence: 82%
- **long_covid_fatigue** — n=1 paper · extraction confidence: 82%
- **long_covid_reduced_workability** — n=1 paper · extraction confidence: 82%
- **long_covid_heavy_breathing** — n=1 paper · extraction confidence: 82%
- **long_covid_pain** — n=1 paper · extraction confidence: 82%
- **long_covid_loss_of_smell_taste** — n=1 paper · extraction confidence: 82%
- **long_covid_weight_loss** — n=1 paper · extraction confidence: 82%
- **anxiety_during_pandemic** — n=1 paper · extraction confidence: 82%
- **depressive_symptoms_during_pandemic** — n=1 paper · extraction confidence: 82%
- **vaccine_adverse_effects** — n=1 paper · extraction confidence: 82%
- **fatigue_longcovid** — n=1 paper · extraction confidence: 82%
- **cough_longcovid** — n=1 paper · extraction confidence: 82%
- **insomnia_longcovid** — n=1 paper · extraction confidence: 82%
- **dry_itchy_throat_longcovid** — n=1 paper · extraction confidence: 82%
- **loss_of_appetite_longcovid** — n=1 paper · extraction confidence: 82%
- **fever_acutephase_allparticipants** — n=1 paper · extraction confidence: 82%
- **cough_acutephase_allparticipants** — n=1 paper · extraction confidence: 82%
- **fatigue_acutephase_allparticipants** — n=1 paper · extraction confidence: 82%

- **throat_pain_acutephase_allparticipants** — n=1 paper · extraction confidence: 82%
- **expectoration_acutephase_allparticipants** — n=1 paper · extraction confidence: 82%
- **any_physical_symptom_1month** — n=1 paper · extraction confidence: 82%
- **any_physical_symptom_3months** — n=1 paper · extraction confidence: 82%
- **any_physical_symptom_6months** — n=1 paper · extraction confidence: 82%
- **any_physical_symptom_12months** — n=1 paper · extraction confidence: 82%
- **any_physical_symptom_controls** — n=1 paper · extraction confidence: 82%
- **fever_sweats_chills_1month** — n=1 paper · extraction confidence: 82%
- **loss_of_taste_1month** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_1month** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_1month** — n=1 paper · extraction confidence: 82%
- **cough_1month** — n=1 paper · extraction confidence: 82%
- **any_cognitive_symptom_1month** — n=1 paper · extraction confidence: 82%
- **any_cognitive_symptom_3months** — n=1 paper · extraction confidence: 82%
- **any_cognitive_symptom_6months** — n=1 paper · extraction confidence: 82%
- **any_cognitive_symptom_controls** — n=1 paper · extraction confidence: 82%
- **any_mental_health_symptom_1month** — n=1 paper · extraction confidence: 82%

- **any_mental_health_symptom_3months** — n=1 paper · extraction confidence: 82%
- **any_mental_health_symptom_6months** — n=1 paper · extraction confidence: 82%
- **any_mental_health_symptom_12months** — n=1 paper · extraction confidence: 82%
- **any_mental_health_symptom_controls_12months** — n=1 paper · extraction confidence: 82%
- **depression_or_anxiety_phq4_cases** — n=1 paper · extraction confidence: 82%
- **depression_or_anxiety_phq4_controls** — n=1 paper · extraction confidence: 82%
- **loss_of_taste_smell_controls** — n=1 paper · extraction confidence: 82%
- **fatigue_covid_positive** — n=1 paper · extraction confidence: 82%
- **fatigue_covid_negative** — n=1 paper · extraction confidence: 82%
- **headache_covid_positive** — n=1 paper · extraction confidence: 82%
- **headache_covid_negative** — n=1 paper · extraction confidence: 82%
- **problems_sleeping_covid_positive** — n=1 paper · extraction confidence: 82%
- **problems_sleeping_covid_negative** — n=1 paper · extraction confidence: 82%
- **fatigue_female** — n=1 paper · extraction confidence: 82%
- **fatigue_male** — n=1 paper · extraction confidence: 82%
- **headache_female** — n=1 paper · extraction confidence: 82%
- **headache_male** — n=1 paper · extraction confidence: 82%
- **anxiety_female** — n=1 paper · extraction confidence: 82%
- **anxiety_male** — n=1 paper · extraction confidence: 82%

- **depression_female** — n=1 paper · extraction confidence: 82%
- **depression_male** — n=1 paper · extraction confidence: 82%
- **brain_fog_female** — n=1 paper · extraction confidence: 82%
- **brain_fog_male** — n=1 paper · extraction confidence: 82%
- **ptsd_female** — n=1 paper · extraction confidence: 82%
- **ptsd_male** — n=1 paper · extraction confidence: 82%
- **anosmia_female** — n=1 paper · extraction confidence: 82%
- **anosmia_male** — n=1 paper · extraction confidence: 82%
- **paresthesia_female** — n=1 paper · extraction confidence: 82%
- **paresthesia_male** — n=1 paper · extraction confidence: 82%
- **dysgeusia_female** — n=1 paper · extraction confidence: 82%
- **dysgeusia_male** — n=1 paper · extraction confidence: 82%
- **vertigo_female** — n=1 paper · extraction confidence: 82%
- **vertigo_male** — n=1 paper · extraction confidence: 82%
- **visual_disturbances_female** — n=1 paper · extraction confidence: 82%
- **visual_disturbances_male** — n=1 paper · extraction confidence: 82%
- **radiographic_imaging** — n=1 paper · extraction confidence: 82%
- **electrocardiography_patients_under_21** — n=1 paper · extraction confidence: 82%
- **echocardiography_patients_under_21** — n=1 paper · extraction confidence: 82%
- **electrocardiography_average_other_age_groups** — n=1 paper · extraction confidence: 82%
- **echocardiography_average_other_age_groups** — n=1 paper · extraction confidence: 82%
- **low_work_ability_comorbidities_long_covid** — n=1 paper · extraction confidence: 82%

- **low_work_ability_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **low_work_functioning_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **long_term_absenteeism_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **low_work_ability_no_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **low_work_functioning_no_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **long_term_absenteeism_no_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **dyspnoea_impairment_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **dyspnoea_impairment_no_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **dyspnoea_impairment_obesity_controls** — n=1 paper · extraction confidence: 82%
- **dyspnoea_impairment_long_covid_no_comorbidity_survey** — n=1 paper · extraction confidence: 82%
- **dyspnoea_impairment_long_covid_comorbidities_survey** — n=1 paper · extraction confidence: 82%
- **dyspnoea_impairment_long_covid_obesity_survey** — n=1 paper · extraction confidence: 82%
- **high_psychological_distress_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **high_psychological_distress_no_comorbidities_controls** — n=1 paper · extraction confidence: 82%
- **high_psychological_distress_long_covid_comorbidities** — n=1 paper · extraction confidence: 82%
- **very_high_psychological_distress_long_covid_comorbidities** — n=1 paper · extraction confidence: 82%

- **fatigue_unrecovered** — n=1 paper · extraction confidence: 82%
- **pain_unrecovered** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_unrecovered** — n=1 paper · extraction confidence: 82%
- **difficulty_remembering_unrecovered** — n=1 paper · extraction confidence: 82%
- **heat_or_cold_intolerance_unrecovered** — n=1 paper · extraction confidence: 82%
- **poor_unrefreshing_sleep_unrecovered** — n=1 paper · extraction confidence: 82%
- **pain_recovered** — n=1 paper · extraction confidence: 82%
- **difficulty_remembering_recovered** — n=1 paper · extraction confidence: 82%
- **heat_or_cold_intolerance_recovered** — n=1 paper · extraction confidence: 82%
- **poor_unrefreshing_sleep_recovered** — n=1 paper · extraction confidence: 82%
- **fatigue_test_negative** — n=1 paper · extraction confidence: 82%
- **pain_test_negative** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_test_negative** — n=1 paper · extraction confidence: 82%
- **difficulty_remembering_test_negative** — n=1 paper · extraction confidence: 82%
- **heat_or_cold_intolerance_test_negative** — n=1 paper · extraction confidence: 82%
- **poor_unrefreshing_sleep_test_negative** — n=1 paper · extraction confidence: 82%
- **full_recovery_at_5months_controls** — n=1 paper · extraction confidence: 82%

- **full_recovery_at_5months_cardiovascular_risk_factors** — n=1 paper · extraction confidence: 82%
- **full_recovery_at_5months_established_cvd** — n=1 paper · extraction confidence: 82%
- **full_recovery_at_12months_controls** — n=1 paper · extraction confidence: 82%
- **full_recovery_at_12months_cardiovascular_risk_factors** — n=1 paper · extraction confidence: 82%
- **full_recovery_at_12months_established_cvd** — n=1 paper · extraction confidence: 82%
- **poor_concentration** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_problems** — n=1 paper · extraction confidence: 82%
- **not_meeting_aerobic_pag_among_lc_positive** — n=1 paper · extraction confidence: 82%
- **not_meeting_strengthening_pag_among_lc_positive** — n=1 paper · extraction confidence: 82%
- **coronary_heart_disease_among_lc_positive** — n=1 paper · extraction confidence: 82%
- **overweight_or_obese_among_lc_positive** — n=1 paper · extraction confidence: 82%
- **current_asthma_among_lc_positive** — n=1 paper · extraction confidence: 82%
- **current_hypertension_among_lc_positive** — n=1 paper · extraction confidence: 82%
- **lc_with_significant_adl_limitation** — n=1 paper · extraction confidence: 82%
- **lc_with_some_adl_limitation** — n=1 paper · extraction confidence: 82%
- **lc_with_no_adl_limitation** — n=1 paper · extraction confidence: 82%

- **any_post_covid_condition_covid19_group** — n=1 paper · extraction confidence: 82%
- **any_post_covid_condition_influenza_group** — n=1 paper · extraction confidence: 82%
- **anxiety_depression_most_common_post_covid** — n=1 paper · extraction confidence: 82%
- **abnormal_breathing_covid19** — n=1 paper · extraction confidence: 82%
- **all_cause_ed_visits_covid19** — n=1 paper · extraction confidence: 82%
- **all_cause_ed_visits_influenza** — n=1 paper · extraction confidence: 82%
- **all_cause_hospitalizations_covid19** — n=1 paper · extraction confidence: 82%
- **all_cause_hospitalizations_influenza** — n=1 paper · extraction confidence: 82%
- **mortality_covid19** — n=1 paper · extraction confidence: 82%
- **mortality_influenza** — n=1 paper · extraction confidence: 82%
- **ongoing_smell_taste_dysfunction** — n=1 paper · extraction confidence: 82%
- **covid_etiology** — n=1 paper · extraction confidence: 82%
- **not reported as proportions among long covid patients; drug/comorbidity prevalences reported in all-patient cohort** — n=1 paper · extraction confidence: 82%
- **general_weakness** — n=1 paper · extraction confidence: 82%
- **lack_of_smell_or_taste** — n=1 paper · extraction confidence: 82%
- **headache_or_eye_related** — n=1 paper · extraction confidence: 82%
- **lack_of_appetite** — n=1 paper · extraction confidence: 82%

- **pain_or_numbness_upper_body** — n=1 paper · extraction confidence: 82%
- **pain_or_numbness_lower_body** — n=1 paper · extraction confidence: 82%
- **heart_palpitation** — n=1 paper · extraction confidence: 82%
- **digestion_or_urinary** — n=1 paper · extraction confidence: 82%
- **loss_of_sleep_insomnia** — n=1 paper · extraction confidence: 82%
- **deep_vein_thrombosis** — n=1 paper · extraction confidence: 82%
- **no_persistent_effects** — n=1 paper · extraction confidence: 82%
- **persistent_any_symptom_cases** — n=1 paper · extraction confidence: 82%
- **persistent_shortness_of_breath_cases** — n=1 paper · extraction confidence: 82%
- **persistent_fatigue_cases** — n=1 paper · extraction confidence: 82%
- **persistent_loss_of_smell_taste_cases** — n=1 paper · extraction confidence: 82%
- **new_fatigue_asymptomatic_cases** — n=1 paper · extraction confidence: 82%
- **new_loss_taste_smell_myalgias_headache_asymptomatic_cases** — n=1 paper · extraction confidence: 82%
- **feeling_in_a_fog_cases** — n=1 paper · extraction confidence: 82%
- **slowed_thinking_cases** — n=1 paper · extraction confidence: 82%
- **insomnia_cases** — n=1 paper · extraction confidence: 82%
- **likely_depression_cases** — n=1 paper · extraction confidence: 82%

- **likely_depression_controls** — n=1 paper · extraction confidence: 82%
- **likely_anxiety_cases** — n=1 paper · extraction confidence: 82%
- **likely_anxiety_controls** — n=1 paper · extraction confidence: 82%
- **acute_fatigue_cases_baseline** — n=1 paper · extraction confidence: 82%
- **acute_headache_cases_baseline** — n=1 paper · extraction confidence: 82%
- **acute_loss_taste_smell_cases_baseline** — n=1 paper · extraction confidence: 82%
- **joint_muscle_pain** — n=1 paper · extraction confidence: 82%
- **hair_loss_telogen_effluvium** — n=1 paper · extraction confidence: 82%
- **anxiety_panic** — n=1 paper · extraction confidence: 82%
- **smell_loss** — n=1 paper · extraction confidence: 82%
- **taste_loss** — n=1 paper · extraction confidence: 82%
- **long_covid_depressed** — n=1 paper · extraction confidence: 82%
- **long_covid_non_depressed** — n=1 paper · extraction confidence: 82%
- **sars_cov2_positive_depressed** — n=1 paper · extraction confidence: 82%
- **sars_cov2_positive_non_depressed** — n=1 paper · extraction confidence: 82%
- **covid19_vaccination_depressed** — n=1 paper · extraction confidence: 82%
- **covid19_vaccination_non_depressed** — n=1 paper · extraction confidence: 82%
- **musculoskeletal_pain** — n=1 paper · extraction confidence: 82%

- **depression_symptom** — n=1 paper · extraction confidence: 82%
- **kidney_problems** — n=1 paper · extraction confidence: 82%
- **non_recovery_cssb** — n=1 paper · extraction confidence: 82%
- **non_recovery_twinsuk** — n=1 paper · extraction confidence: 82%
- **at_least_one_persistent_symptom_long_covid** — n=1 paper · extraction confidence: 82%
- **memory_decline** — n=1 paper · extraction confidence: 82%
- **somnipathy** — n=1 paper · extraction confidence: 82%
- **general_symptoms_any** — n=1 paper · extraction confidence: 82%
- **mental_symptoms_any** — n=1 paper · extraction confidence: 82%
- **respiratory_symptoms_any** — n=1 paper · extraction confidence: 82%
- **digestive_symptoms_any** — n=1 paper · extraction confidence: 82%
- **change_in_taste** — n=1 paper · extraction confidence: 82%
- **dyspnea_non_motor** — n=1 paper · extraction confidence: 82%
- **not_fully_recovered** — n=1 paper · extraction confidence: 82%
- **arthralgia_overall** — n=1 paper · extraction confidence: 82%
- **dysgeusia_overall** — n=1 paper · extraction confidence: 82%
- **dysosmia_overall** — n=1 paper · extraction confidence: 82%
- **memory_impairment_overall** — n=1 paper · extraction confidence: 82%
- **loss_of_concentration_overall** — n=1 paper · extraction confidence: 82%

- **sexual_dysfunction_overall** — n=1 paper · extraction confidence: 82%
- **arthralgia_covid_history** — n=1 paper · extraction confidence: 82%
- **headache_covid_history** — n=1 paper · extraction confidence: 82%
- **chest_pain_covid_history** — n=1 paper · extraction confidence: 82%
- **cough_covid_history** — n=1 paper · extraction confidence: 82%
- **dyspnea_covid_history** — n=1 paper · extraction confidence: 82%
- **dysgeusia_covid_history** — n=1 paper · extraction confidence: 82%
- **dysosmia_covid_history** — n=1 paper · extraction confidence: 82%
- **memory_impairment_covid_history** — n=1 paper · extraction confidence: 82%
- **loss_of_concentration_covid_history** — n=1 paper · extraction confidence: 82%
- **hair_loss_covid_history** — n=1 paper · extraction confidence: 82%
- **sexual_dysfunction_covid_history** — n=1 paper · extraction confidence: 82%
- **fatigue_covid_history** — n=1 paper · extraction confidence: 82%
- **headache_migraine_plus_covid_plus_bothered_quite_a_lot_or_very_much** — n=1 paper · extraction confidence: 82%
- **anosmia_migraine_plus_covid_plus_bothered_quite_a_lot_or_very_much** — n=1 paper · extraction confidence: 82%
- **memory_problems_migraine_plus_covid_plus_bothered_quite_a_lot_or_very_much** — n=1 paper · extraction confidence: 82%

- **concentration_problems_migraine_plus_covid_plus_bothered_quite_a** — n=1 paper · extraction confidence: 82%
- **covid19_among_migraine_patients** — n=1 paper · extraction confidence: 82%
- **covid19_among_non_migraine_individuals** — n=1 paper · extraction confidence: 82%
- **migraine_prevalence_in_cohort** — n=1 paper · extraction confidence: 82%
- **depression_weighted_mean** — n=1 paper · extraction confidence: 82%
- **depression_unweighted_mean** — n=1 paper · extraction confidence: 82%
- **depression_median** — n=1 paper · extraction confidence: 82%
- **anxiety_weighted_mean** — n=1 paper · extraction confidence: 82%
- **anxiety_unweighted_mean** — n=1 paper · extraction confidence: 82%
- **anxiety_median** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_weighted_mean** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_unweighted_mean** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_median** — n=1 paper · extraction confidence: 82%
- **post_traumatic_symptoms_weighted_mean** — n=1 paper · extraction confidence: 82%
- **post_traumatic_symptoms_unweighted_mean** — n=1 paper · extraction confidence: 82%
- **post_traumatic_symptoms_median** — n=1 paper · extraction confidence: 82%
- **sleep_disturbances_weighted_mean** — n=1 paper · extraction confidence: 82%

- **sleep_disturbances_unweighted_mean** — n=1 paper · extraction confidence: 82%
- **sleep_disturbances_median** — n=1 paper · extraction confidence: 82%
- **fatigue_queries_among_symptom_stories** — n=1 paper · extraction confidence: 82%
- **loss_or_change_of_smell_queries** — n=1 paper · extraction confidence: 82%
- **headache_queries** — n=1 paper · extraction confidence: 82%
- **pain_queries** — n=1 paper · extraction confidence: 82%
- **dyspnea_queries** — n=1 paper · extraction confidence: 82%
- **chest_pain_queries** — n=1 paper · extraction confidence: 82%
- **difficulty_concentrating_queries** — n=1 paper · extraction confidence: 82%
- **dizziness_queries** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_symptoms_queries** — n=1 paper · extraction confidence: 82%
- **cough_queries** — n=1 paper · extraction confidence: 82%
- **sleeping_disorders_queries** — n=1 paper · extraction confidence: 82%
- **psychiatric_symptoms_queries** — n=1 paper · extraction confidence: 82%
- **paresthesias_queries** — n=1 paper · extraction confidence: 82%
- **tinnitus_queries** — n=1 paper · extraction confidence: 82%
- **dermatologic_symptoms_queries** — n=1 paper · extraction confidence: 82%
- **fever_queries** — n=1 paper · extraction confidence: 82%
- **loss_or_change_of_voice_queries** — n=1 paper · extraction confidence: 82%

- **palpitations_queries** — n=1 paper · extraction confidence: 82%
- **visual_disorders_queries** — n=1 paper · extraction confidence: 82%
- **dysautonomic_symptoms_queries** — n=1 paper · extraction confidence: 82%
- **poor_sleep_quality** — n=1 paper · extraction confidence: 82%
- **obstructive_sleep_apnea_symptoms** — n=1 paper · extraction confidence: 82%
- **recommended_sleep_7_8hrs** — n=1 paper · extraction confidence: 82%
- **short_sleep_le6hrs** — n=1 paper · extraction confidence: 82%
- **long_sleep_ge9hrs** — n=1 paper · extraction confidence: 82%
- **persistent_covid_sequelae_overall** — n=1 paper · extraction confidence: 82%
- **pulmonary_sequelae_general** — n=1 paper · extraction confidence: 82%
- **blurring_of_vision** — n=1 paper · extraction confidence: 82%
- **depression_comorbidity** — n=1 paper · extraction confidence: 82%
- **hypertension_comorbidity** — n=1 paper · extraction confidence: 82%
- **long_covid_primary_school_round3** — n=1 paper · extraction confidence: 82%
- **long_covid_secondary_years7to11_round3** — n=1 paper · extraction confidence: 82%
- **long_covid_secondary_years12to13_round3** — n=1 paper · extraction confidence: 82%

- **any_persisting_symptom_primary_school_round3** — n=1 paper · extraction confidence: 82%
- **any_persisting_symptom_secondary_years7to11_round3** — n=1 paper · extraction confidence: 82%
- **any_persisting_symptom_secondary_years12to13_round3** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_taste_years0to6_positive_test** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_taste_years0to6_no_covid** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_taste_years7to11_positive_test** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_taste_years7to11_no_covid** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_taste_years12to13_positive_test** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_taste_years12to13_no_covid** — n=1 paper · extraction confidence: 82%
- **cardiovascular_primary_positive_test** — n=1 paper · extraction confidence: 82%
- **cardiovascular_primary_no_covid** — n=1 paper · extraction confidence: 82%
- **cardiovascular_years7to11_positive_test** — n=1 paper · extraction confidence: 82%
- **cardiovascular_years7to11_no_covid** — n=1 paper · extraction confidence: 82%
- **systemic_years7to11_positive_test** — n=1 paper · extraction confidence: 82%
- **systemic_years7to11_no_covid** — n=1 paper · extraction confidence: 82%
- **any_persisting_symptom_secondary_years7to11_positive_test** — n=1 paper · extraction confidence: 82%

- **any_persisting_symptom_secondary_years7to11_no_covid** — n=1 paper · extraction confidence: 82%
- **loneliness_often_always_pre_pandemic** — n=1 paper · extraction confidence: 82%
- **loneliness_increased_during_pandemic_3item_scale** — n=1 paper · extraction confidence: 82%
- **loneliness_decreased_during_pandemic_3item_scale** — n=1 paper · extraction confidence: 82%
- **loneliness_unchanged_3item_scale** — n=1 paper · extraction confidence: 82%
- **loneliness_increased_single_item_scale** — n=1 paper · extraction confidence: 82%
- **pre_existing_tiredness** — n=1 paper · extraction confidence: 82%
- **pre_existing_worry** — n=1 paper · extraction confidence: 82%
- **pre_existing_allergy** — n=1 paper · extraction confidence: 82%
- **long_covid_meeting_definition** — n=1 paper · extraction confidence: 82%
- **reduced_6mwt_below_50th_percentile** — n=1 paper · extraction confidence: 82%
- **reduced_vo2peak_below_80_percent_predicted** — n=1 paper · extraction confidence: 82%
- **abnormal_dlco_below_75_percent_predicted** — n=1 paper · extraction confidence: 82%
- **normal_fvc** — n=1 paper · extraction confidence: 82%
- **normal_fev1** — n=1 paper · extraction confidence: 82%
- **normal_tlc** — n=1 paper · extraction confidence: 82%
- **normal_resting_spo2** — n=1 paper · extraction confidence: 82%
- **covid-positive_minimal_symptoms_3month** — n=1 paper · extraction confidence: 82%

- **covid-positive_minimal_symptoms_6month** — n=1 paper · extraction confidence: 82%
- **covid-negative_minimal_symptoms_3month** — n=1 paper · extraction confidence: 82%
- **covid-positive_thm_3month** — n=1 paper · extraction confidence: 82%
- **covid-positive_lost_3month** — n=1 paper · extraction confidence: 82%
- **covid-positive_msms_3month** — n=1 paper · extraction confidence: 82%
- **covid-positive_minimal_fatigue_acute** — n=1 paper · extraction confidence: 82%
- **covid-positive_maximal_fatigue_acute** — n=1 paper · extraction confidence: 82%
- **covid-positive_minimal_fatigue_3month** — n=1 paper · extraction confidence: 82%
- **covid-positive_maximal_fatigue_3month** — n=1 paper · extraction confidence: 82%
- **covid-positive_minimal_fatigue_6month** — n=1 paper · extraction confidence: 82%
- **covid-positive_maximal_fatigue_6month** — n=1 paper · extraction confidence: 82%
- **covid-positive_act_heent_with_lost_acute** — n=1 paper · extraction confidence: 82%
- **covid-positive_act_heent_without_lost_acute** — n=1 paper · extraction confidence: 82%
- **covid-positive_msms_acute** — n=1 paper · extraction confidence: 82%
- **covid-positive_minimal_symptoms_acute** — n=1 paper · extraction confidence: 82%
- **covid-negative_minimal_symptoms_acute** — n=1 paper · extraction confidence: 82%

- **covid-negative_act_heent_without_lost_acute** — n=1 paper · extraction confidence: 82%
- **covid-negative_msms_acute** — n=1 paper · extraction confidence: 82%
- **any_symptoms_at_3_months** — n=1 paper · extraction confidence: 82%
- **0_symptoms_at_3_months** — n=1 paper · extraction confidence: 82%
- **1_to_2_symptoms_at_3_months** — n=1 paper · extraction confidence: 82%
- **3_to_4_symptoms_at_3_months** — n=1 paper · extraction confidence: 82%
- **5_or_more_symptoms_at_3_months** — n=1 paper · extraction confidence: 82%
- **fatigue_positives_definition1** — n=1 paper · extraction confidence: 82%
- **fatigue_negatives_definition1** — n=1 paper · extraction confidence: 82%
- **loss_change_of_smell_positives_definition1** — n=1 paper · extraction confidence: 82%
- **loss_change_of_smell_negatives_definition1** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_positives_definition1** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_negatives_definition1** — n=1 paper · extraction confidence: 82%
- **concentration_difficulties_positives_definition1** — n=1 paper · extraction confidence: 82%
- **concentration_difficulties_negatives_definition1** — n=1 paper · extraction confidence: 82%
- **loss_change_of_taste_positives_definition1** — n=1 paper · extraction confidence: 82%

- **loss_change_of_taste_negatives_definition1** — n=1 paper · extraction confidence: 82%
- **fatigue_positives_definition5** — n=1 paper · extraction confidence: 82%
- **loss_change_of_smell_positives_definition5** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_positives_definition5** — n=1 paper · extraction confidence: 82%
- **concentration_difficulties_positives_definition5** — n=1 paper · extraction confidence: 82%
- **sleeping_problems_positives_definition5** — n=1 paper · extraction confidence: 82%
- **brain_fog_positives_definition1** — n=1 paper · extraction confidence: 82%
- **hair_loss_positives_definition1** — n=1 paper · extraction confidence: 82%
- **palpitations_positives_definition1** — n=1 paper · extraction confidence: 82%
- **amnesia_positives_definition1** — n=1 paper · extraction confidence: 82%
- **headache_positives_definition1** — n=1 paper · extraction confidence: 82%
- **hair_fall** — n=1 paper · extraction confidence: 82%
- **general_symptoms_pooled** — n=1 paper · extraction confidence: 82%
- **loss_of_smell_and_taste_combined** — n=1 paper · extraction confidence: 82%
- **mental_health_symptoms_pooled** — n=1 paper · extraction confidence: 82%
- **anxiety_and_depression_combined** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_symptoms_pooled** — n=1 paper · extraction confidence: 82%

- **diarrhea_and_emesis** — n=1 paper · extraction confidence: 82%
- **fatigue_at_4to12wks_among_symptomatic** — n=1 paper · extraction confidence: 82%
- **shortness_of_breath_at_4to12wks_among_symptomatic** — n=1 paper · extraction confidence: 82%
- **cognitive_impairment_at_4to12wks_among_symptomatic** — n=1 paper · extraction confidence: 82%
- **muscle_joint_symptoms** — n=1 paper · extraction confidence: 82%
- **anemia_comorbidity** — n=1 paper · extraction confidence: 82%
- **autoimmune_disease** — n=1 paper · extraction confidence: 82%
- **gastrointestinal_disease_comorbidity** — n=1 paper · extraction confidence: 82%
- **loss_decrease_of_smell_and_or_taste** — n=1 paper · extraction confidence: 82%
- **myalgia_including_joint_pain** — n=1 paper · extraction confidence: 82%
- **cardiologic_symptoms** — n=1 paper · extraction confidence: 82%
- **confusion_or_lack_of_concentration** — n=1 paper · extraction confidence: 78%
- **arthralgia_myalgia** — n=1 paper · extraction confidence: 78%
- **depression_chest_pain_headache_anxiety_cough** — n=1 paper · extraction confidence: 78%
- **at_least_one_persistent_or_new_symptom_hic_2to6months** — n=1 paper · extraction confidence: 78%
- **at_least_one_persistent_or_new_symptom_lmic_2to6months** — n=1 paper · extraction confidence: 78%

- **at_least_one_persistent_or_new_symptom_hic_6to12months**
— n=1 paper · extraction confidence: 78%
- **at_least_one_persistent_or_new_symptom_lmic_6to12months**
— n=1 paper · extraction confidence: 78%
- **fatigue_moderate_to_severe_hic_2to6months** — n=1
paper · extraction confidence: 78%
- **fatigue_moderate_to_severe_lmic_2to6months** — n=1
paper · extraction confidence: 78%
- **fatigue_moderate_to_severe_hic_6to12months** — n=1
paper · extraction confidence: 78%
- **fatigue_moderate_to_severe_lmic_6to12months** — n=1
paper · extraction confidence: 78%
- **fever_new_persistent_hic_2to6months** — n=1 paper ·
extraction confidence: 78%
- **fever_new_persistent_lmic_2to6months** — n=1 paper ·
extraction confidence: 78%
- **fever_new_persistent_hic_6to12months** — n=1 paper ·
extraction confidence: 78%
- **fever_new_persistent_lmic_6to12months** — n=1 paper ·
extraction confidence: 78%
- **not_fully_recovered_hic_2to6months** — n=1 paper ·
extraction confidence: 78%
- **not_fully_recovered_lmic_2to6months** — n=1 paper ·
extraction confidence: 78%
- **not_fully_recovered_hic_6to12months** — n=1 paper ·
extraction confidence: 78%
- **not_fully_recovered_lmic_6to12months** — n=1 paper ·
extraction confidence: 78%
- **anxiety_depression_hic_2to6months** — n=1 paper ·
extraction confidence: 78%
- **anxiety_depression_lmic_2to6months** — n=1 paper ·
extraction confidence: 78%

- **anxiety_depression_hic_6to12months** — n=1 paper · extraction confidence: 78%
- **anxiety_depression_lmic_6to12months** — n=1 paper · extraction confidence: 78%
- **pain_discomfort_hic_2to6months** — n=1 paper · extraction confidence: 78%
- **pain_discomfort_lmic_2to6months** — n=1 paper · extraction confidence: 78%
- **pain_discomfort_hic_6to12months** — n=1 paper · extraction confidence: 78%
- **pain_discomfort_lmic_6to12months** — n=1 paper · extraction confidence: 78%
- **problems_usual_activities_hic_2to6months** — n=1 paper · extraction confidence: 78%
- **problems_usual_activities_lmic_2to6months** — n=1 paper · extraction confidence: 78%
- **problems_usual_activities_hic_6to12months** — n=1 paper · extraction confidence: 78%
- **problems_usual_activities_lmic_6to12months** — n=1 paper · extraction confidence: 78%
- **moderate_to_severe_breathlessness_overall_range** — n=1 paper · extraction confidence: 78%
- **neurocognitive_symptoms_hic_2to6months** — n=1 paper · extraction confidence: 78%
- **neurocognitive_symptoms_lmic_2to6months** — n=1 paper · extraction confidence: 78%
- **neurocognitive_symptoms_hic_6to12months** — n=1 paper · extraction confidence: 78%
- **neurocognitive_symptoms_lmic_6to12months** — n=1 paper · extraction confidence: 78%
- **respiratory_symptoms_hic_2to6months** — n=1 paper · extraction confidence: 78%

- **respiratory_symptoms_lmic_2to6months** — n=1 paper · extraction confidence: 78%
- **respiratory_symptoms_hic_6to12months** — n=1 paper · extraction confidence: 78%
- **respiratory_symptoms_lmic_6to12months** — n=1 paper · extraction confidence: 78%
- **not_fully_recovered_overall_2to6months** — n=1 paper · extraction confidence: 78%
- **not_fully_recovered_overall_6to12months** — n=1 paper · extraction confidence: 78%
- **vision_problems** — n=1 paper · extraction confidence: 78%
- **balance_problems** — n=1 paper · extraction confidence: 78%
- **rhinorrhoea** — n=1 paper · extraction confidence: 78%
- **sinus_congestion** — n=1 paper · extraction confidence: 78%
- **bodyache** — n=1 paper · extraction confidence: 78%
- **pancreatitis** — n=1 paper · extraction confidence: 78%
- **seizure** — n=1 paper · extraction confidence: 78%
- **stroke** — n=1 paper · extraction confidence: 78%
- **anorexia** — n=1 paper · extraction confidence: 78%
- **prolonged_generalized_weakness** — n=1 paper · extraction confidence: 78%
- **problem_playing_social_role** — n=1 paper · extraction confidence: 78%
- **problems_in_adl** — n=1 paper · extraction confidence: 78%
- **prolonged_fever** — n=1 paper · extraction confidence: 78%
- **memory_problem** — n=1 paper · extraction confidence: 78%
- **problem_personal_care** — n=1 paper · extraction confidence: 78%
- **problem_in_concentration** — n=1 paper · extraction confidence: 78%
- **breathlessness_dyspnea** — n=1 paper · extraction confidence: 78%

- **mobility_problem** — n=1 paper · extraction confidence: 78%
- **swallowing_feeding_problem** — n=1 paper · extraction confidence: 78%
- **communication_problem** — n=1 paper · extraction confidence: 78%
- **problem_executing_plan** — n=1 paper · extraction confidence: 78%
- **noisy_breathing** — n=1 paper · extraction confidence: 78%
- **bladder_control_problem** — n=1 paper · extraction confidence: 78%
- **bowel_control_problem** — n=1 paper · extraction confidence: 78%
- **long_covid_symptoms_4_to_12_weeks** — n=1 paper · extraction confidence: 78%
- **long_covid_symptoms_greater_than_12_weeks** — n=1 paper · extraction confidence: 78%
- **long_covid_any_greater_than_4_weeks_among_infected** — n=1 paper · extraction confidence: 78%
- **seropositive_at_least_once** — n=1 paper · extraction confidence: 78%
- **ever_infected_by_serology_or_self_report** — n=1 paper · extraction confidence: 78%
- **moderate_to_severe_anxiety_at_any_timepoint** — n=1 paper · extraction confidence: 78%
- **moderate_to_severe_depression_at_any_timepoint** — n=1 paper · extraction confidence: 78%
- **physical_impairment** — n=1 paper · extraction confidence: 78%
- **excessive_thirst** — n=1 paper · extraction confidence: 78%
- **irritability** — n=1 paper · extraction confidence: 78%
- **respiratory_symptoms_pwh** — n=1 paper · extraction confidence: 78%

- **gastrointestinal_symptoms_pwh** — n=1 paper · extraction confidence: 78%
- **fatigue_pwh_row1** — n=1 paper · extraction confidence: 78%
- **fatigue_pwh_row2** — n=1 paper · extraction confidence: 78%
- **headache_pwh** — n=1 paper · extraction confidence: 78%
- **chronic_cough_pwh** — n=1 paper · extraction confidence: 78%
- **chest_pain_pwh** — n=1 paper · extraction confidence: 78%
- **diarrhea_constipation_pwh** — n=1 paper · extraction confidence: 78%
- **dizziness_pwh** — n=1 paper · extraction confidence: 78%
- **brain_fog_pwh** — n=1 paper · extraction confidence: 78%
- **body_ache_pwh** — n=1 paper · extraction confidence: 78%
- **palpitations_pwh** — n=1 paper · extraction confidence: 78%
- **cognitive_impairment_pwh** — n=1 paper · extraction confidence: 78%
- **disturbances_smell_taste_pwh** — n=1 paper · extraction confidence: 78%
- **abnormal_movements_pwh** — n=1 paper · extraction confidence: 78%
- **hair_loss_pwh** — n=1 paper · extraction confidence: 78%
- **mental_disorders_pwh** — n=1 paper · extraction confidence: 78%
- **hypertension_new_onset_pwh** — n=1 paper · extraction confidence: 78%
- **heart_disease_new_onset_pwh** — n=1 paper · extraction confidence: 78%
- **malignancy_new_onset_pwh** — n=1 paper · extraction confidence: 78%
- **thrombosis_pwh** — n=1 paper · extraction confidence: 78%

- **diabetes_new_onset_pwh** — n=1 paper · extraction confidence: 78%
- **mortality_pwh** — n=1 paper · extraction confidence: 78%
- **cough_post_covid_long_covid** — n=1 paper · extraction confidence: 78%
- **difficulty_breathing_post_covid_long_covid** — n=1 paper · extraction confidence: 78%
- **tiredness_post_covid_long_covid** — n=1 paper · extraction confidence: 78%
- **brain_fog_post_covid_long_covid** — n=1 paper · extraction confidence: 78%
- **sleep_problems_post_covid_long_covid** — n=1 paper · extraction confidence: 78%
- **breathlessness_dyspnea_obs1** — n=1 paper · extraction confidence: 78%
- **fatigue_obs1** — n=1 paper · extraction confidence: 78%
- **cough_obs1** — n=1 paper · extraction confidence: 78%
- **headache_obs1** — n=1 paper · extraction confidence: 78%
- **myalgia_obs1** — n=1 paper · extraction confidence: 78%
- **arthralgia_obs1** — n=1 paper · extraction confidence: 78%
- **chest_pain_obs1** — n=1 paper · extraction confidence: 78%
- **anxiety_obs1** — n=1 paper · extraction confidence: 78%
- **sleep_disturbance_obs1** — n=1 paper · extraction confidence: 78%
- **depression_obs1** — n=1 paper · extraction confidence: 78%
- **fever_obs1** — n=1 paper · extraction confidence: 78%
- **memory_loss_obs1** — n=1 paper · extraction confidence: 78%
- **dizziness_obs1** — n=1 paper · extraction confidence: 78%
- **concentration_problems_obs1** — n=1 paper · extraction confidence: 78%
- **ear_pain_obs1** — n=1 paper · extraction confidence: 78%

- **planning_problems_obs1** — n=1 paper · extraction confidence: 78%
- **abdominal_pain_obs1** — n=1 paper · extraction confidence: 78%
- **skin_rash_obs1** — n=1 paper · extraction confidence: 78%
- **ptsd_obs1** — n=1 paper · extraction confidence: 78%
- **fatigue_obs2** — n=1 paper · extraction confidence: 78%
- **breathlessness_dyspnea_obs2** — n=1 paper · extraction confidence: 78%
- **headache_obs2** — n=1 paper · extraction confidence: 78%
- **anxiety_obs2** — n=1 paper · extraction confidence: 78%
- **arthralgia_obs2** — n=1 paper · extraction confidence: 78%
- **myalgia_obs2** — n=1 paper · extraction confidence: 78%
- **concentration_obs2** — n=1 paper · extraction confidence: 78%
- **ptsd_obs2** — n=1 paper · extraction confidence: 78%
- **memory_loss_obs2** — n=1 paper · extraction confidence: 78%
- **fever_obs2** — n=1 paper · extraction confidence: 78%
- **chest_pain_obs2** — n=1 paper · extraction confidence: 78%
- **taste_alteration_obs2** — n=1 paper · extraction confidence: 78%
- **smell_problems_obs2** — n=1 paper · extraction confidence: 78%
- **depression_obs2** — n=1 paper · extraction confidence: 78%
- **hair_loss_obs2** — n=1 paper · extraction confidence: 78%
- **diarrhea_obs2** — n=1 paper · extraction confidence: 78%
- **sleep_disturbance_obs2** — n=1 paper · extraction confidence: 78%
- **skin_rash_obs2** — n=1 paper · extraction confidence: 78%
- **dizziness_obs2** — n=1 paper · extraction confidence: 78%
- **planning_problems_obs2** — n=1 paper · extraction confidence: 78%

- **visual_problems_obs2** — n=1 paper · extraction confidence: 78%
- **weight_loss_obs2** — n=1 paper · extraction confidence: 78%
- **abdominal_pain_obs2** — n=1 paper · extraction confidence: 78%
- **tinnitus_obs2** — n=1 paper · extraction confidence: 78%
- **sore_throat_obs2** — n=1 paper · extraction confidence: 78%
- **brain_fog_obs2** — n=1 paper · extraction confidence: 78%
- **ear_pain_obs2** — n=1 paper · extraction confidence: 78%
- **long_covid_any_symptom_any_intensity_at_1month** — n=1 paper · extraction confidence: 78%
- **long_covid_any_symptom_severe_intensity_at_1month** — n=1 paper · extraction confidence: 78%
- **strict_long_covid_at_1month** — n=1 paper · extraction confidence: 78%
- **strict_severe_long_covid_at_1month** — n=1 paper · extraction confidence: 78%
- **severe_composite_score_omicron_1month** — n=1 paper · extraction confidence: 78%
- **severe_composite_score_pre_omicron_1month** — n=1 paper · extraction confidence: 78%
- **shortness_of_breath_dyspnea_top_predictor** — n=1 paper · extraction confidence: 78%
- **malaise_fatigue_top_predictor** — n=1 paper · extraction confidence: 78%
- **chest_pain_top_predictor** — n=1 paper · extraction confidence: 78%
- **at_least_one_long_covid_symptom** — n=1 paper · extraction confidence: 78%
- **sspd_covid_positive** — n=1 paper · extraction confidence: 78%
- **sspd_covid_negative** — n=1 paper · extraction confidence: 78%

- **dyspnea_at_3months_hospitalized** — n=1 paper · extraction confidence: 78%
- **fatigue_at_3months_hospitalized** — n=1 paper · extraction confidence: 78%
- **joint_pain_at_3months_hospitalized** — n=1 paper · extraction confidence: 78%
- **runny_nose_overall** — n=1 paper · extraction confidence: 78%
- **sneezing_overall** — n=1 paper · extraction confidence: 78%
- **fatigue_at_2-3_months** — n=1 paper · extraction confidence: 78%
- **any_symptom_at_2-3_months_cases** — n=1 paper · extraction confidence: 78%
- **any_symptom_at_13-18_months_cases** — n=1 paper · extraction confidence: 78%
- **moderate_or_severe_severity** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_vaccinated_adults_centera** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_vaccinated_adults_centerb** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_unvaccinated_adults_centera** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_unvaccinated_adults_centerb** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_vaccinated_youths_centera** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_vaccinated_youths_centerb** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_unvaccinated_youths_centera** — n=1 paper · extraction confidence: 78%
- **patients_with_any_new_diagnosis_unvaccinated_youths_centerb** — n=1 paper · extraction confidence: 78%

- **pharyngeal_discomfort** — n=1 paper · extraction confidence: 78%
- **full_body_soreness** — n=1 paper · extraction confidence: 78%
- **fear_of_cold_shivering** — n=1 paper · extraction confidence: 78%
- **headache_dizziness** — n=1 paper · extraction confidence: 78%
- **runny_nose** — n=1 paper · extraction confidence: 78%
- **decrease_in_taste** — n=1 paper · extraction confidence: 78%
- **asymptomatic** — n=1 paper · extraction confidence: 78%
- **body_aches_in_ana_positive_women_at_12_months** — n=1 paper · extraction confidence: 78%
- **body_aches_in_ana_negative_or_low_titer_women_at_12_months** — n=1 paper · extraction confidence: 78%
- **neurological_symptoms_higher_ana_titer** — n=1 paper · extraction confidence: 78%
- **neurological_symptoms_lower_ana_titer** — n=1 paper · extraction confidence: 78%
- **anca_associated_mortality_anca_positive** — n=1 paper · extraction confidence: 78%
- **anca_associated_mortality_anca_negative** — n=1 paper · extraction confidence: 78%
- **anca_associated_severe_disease_anca_positive** — n=1 paper · extraction confidence: 78%
- **anca_associated_severe_disease_anca_negative** — n=1 paper · extraction confidence: 78%
- **respiratory_distress** — n=1 paper · extraction confidence: 78%
- **long_covid_historical_variant_azzolini** — n=1 paper · extraction confidence: 78%
- **long_covid_alpha_variant_azzolini** — n=1 paper · extraction confidence: 78%

- **long_covid_delta_omicron_mix_azzolini** — n=1 paper · extraction confidence: 78%
- **long_covid_omicron_morioka** — n=1 paper · extraction confidence: 78%
- **long_covid_omicron_qasmieh** — n=1 paper · extraction confidence: 78%
- **fatigue_most_prevalent_all_variants** — n=1 paper · extraction confidence: 78%
- **fatigue_during_acute_infection** — n=1 paper · extraction confidence: 78%
- **joint_or_muscle_pain_during_acute_infection** — n=1 paper · extraction confidence: 78%
- **taste_or_smell_changes_during_acute_infection** — n=1 paper · extraction confidence: 78%
- **long_covid_prevalence_among_covid_positive_households** — n=1 paper · extraction confidence: 78%
- **high_blood_pressure_comorbidity** — n=1 paper · extraction confidence: 78%
- **high_cholesterol_comorbidity** — n=1 paper · extraction confidence: 78%
- **poor_mental_health_comorbidity** — n=1 paper · extraction confidence: 78%
- **heart_disease_comorbidity** — n=1 paper · extraction confidence: 78%
- **poor_physical_health_comorbidity** — n=1 paper · extraction confidence: 78%
- **stroke_comorbidity** — n=1 paper · extraction confidence: 78%
- **cancer_comorbidity** — n=1 paper · extraction confidence: 78%
- **copd_comorbidity** — n=1 paper · extraction confidence: 78%
- **chronic_kidney_disease_comorbidity** — n=1 paper · extraction confidence: 78%

- **orthostatic_intolerance** — n=1 paper · extraction confidence: 78%
- **pasc_with_strong_primary_infection_indicator** — n=1 paper · extraction confidence: 78%
- **negative_sentiment_overall** — n=1 paper · extraction confidence: 78%
- **positive_sentiment_overall** — n=1 paper · extraction confidence: 78%
- **neutral_sentiment_overall** — n=1 paper · extraction confidence: 78%
- **negative_sentiment_reweighted** — n=1 paper · extraction confidence: 78%
- **positive_sentiment_reweighted** — n=1 paper · extraction confidence: 78%
- **neutral_sentiment_reweighted** — n=1 paper · extraction confidence: 78%
- **negative_sentiment_female** — n=1 paper · extraction confidence: 78%
- **negative_sentiment_male** — n=1 paper · extraction confidence: 78%
- **long_covid_among_covid_infected** — n=1 paper · extraction confidence: 78%
- **long_covid_among_all_participants** — n=1 paper · extraction confidence: 78%
- **severe_cognitive_deficit_long_covid** — n=1 paper · extraction confidence: 78%
- **severe_cognitive_deficit_never_infected** — n=1 paper · extraction confidence: 78%
- **severe_cognitive_deficit_no_long_covid** — n=1 paper · extraction confidence: 78%
- **long_covid_females** — n=1 paper · extraction confidence: 78%
- **long_covid_males** — n=1 paper · extraction confidence: 78%

- **long_covid_age_45_54** — n=1 paper · extraction confidence: 78%
- **long_covid_age_55_64** — n=1 paper · extraction confidence: 78%
- **long_covid_age_18_44** — n=1 paper · extraction confidence: 78%
- **long_covid_age_65_plus** — n=1 paper · extraction confidence: 78%
- **long_covid_non_hispanic_white** — n=1 paper · extraction confidence: 78%
- **long_covid_hispanic** — n=1 paper · extraction confidence: 78%
- **long_covid_non_hispanic_black** — n=1 paper · extraction confidence: 78%
- **acute_covid_overall** — n=1 paper · extraction confidence: 78%
- **no_covid_overall** — n=1 paper · extraction confidence: 78%
- **composite_lc_pasc_doac_users_unadjusted** — n=1 paper · extraction confidence: 78%
- **composite_lc_pasc_vka_users_unadjusted** — n=1 paper · extraction confidence: 78%
- **composite_lc_pasc_non_oac_users_unadjusted** — n=1 paper · extraction confidence: 78%
- **new_diagnoses_adults** — n=1 paper · extraction confidence: 78%
- **new_diagnoses_youth** — n=1 paper · extraction confidence: 78%
- **functional_dyspepsia** — n=1 paper · extraction confidence: 78%
- **dizziness_and_giddiness** — n=1 paper · extraction confidence: 78%
- **unspecified_abdominal_pain** — n=1 paper · extraction confidence: 78%

- **low_back_pain** — n=1 paper · extraction confidence: 78%
- **depressed_mood** — n=1 paper · extraction confidence: 78%
- **unspecified_dyspnea** — n=1 paper · extraction confidence: 78%
- **long_covid_boosted_group** — n=1 paper · extraction confidence: 78%
- **long_covid_no_incomplete_vaccine_group** — n=1 paper · extraction confidence: 78%
- **moderate_severe_symptoms_with_long_covid** — n=1 paper · extraction confidence: 78%
- **long_covid_asymptomatic_mild_symptoms** — n=1 paper · extraction confidence: 78%
- **long_covid_severe_covid** — n=1 paper · extraction confidence: 78%
- **long_covid_non_severe_covid** — n=1 paper · extraction confidence: 78%
- **fatigue_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **post_exertional_malaise_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **insomnia_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **brain_fog_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **shortness_of_breath_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **problems_of_memory_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **joint_muscle_pain_severe_lifethreatening** — n=1 paper · extraction confidence: 78%
- **excessive_daytime_sleepiness_severe_lifethreatening** — n=1 paper · extraction confidence: 78%

- **loss_of_smell_asymptomatic** — n=1 paper · extraction confidence: 78%
- **loss_of_smell_non_covid_controls** — n=1 paper · extraction confidence: 78%
- **loss_of_smell_moderate_severe** — n=1 paper · extraction confidence: 78%
- **joint_muscle_pain_non_covid_controls** — n=1 paper · extraction confidence: 78%
- **joint_muscle_pain_asymptomatic** — n=1 paper · extraction confidence: 78%
- **long_covid_double_vaccinated_ayoubkhani** — n=1 paper · extraction confidence: 72%
- **long_covid_unvaccinated_ayoubkhani** — n=1 paper · extraction confidence: 72%
- **long_covid_overall_azzolini** — n=1 paper · extraction confidence: 72%
- **long_covid_post_omicron_ballouz** — n=1 paper · extraction confidence: 72%
- **long_covid_vaccinated_sweden_reference** — n=1 paper · extraction confidence: 72%
- **long_covid_unvaccinated_sweden_reference** — n=1 paper · extraction confidence: 72%
- **covid-19_symptoms_day90** — n=1 paper · extraction confidence: 72%
- **covid-19_symptoms_day120** — n=1 paper · extraction confidence: 72%
- **covid-19_symptoms_day180** — n=1 paper · extraction confidence: 72%
- **pasc_overall_range_low** — n=1 paper · extraction confidence: 72%
- **pasc_overall_range_high** — n=1 paper · extraction confidence: 72%

- **normal_cognition_overall** — n=1 paper · extraction confidence: 72%
- **mild_impairment_one_domain_young_middle_aged** — n=1 paper · extraction confidence: 72%
- **moderate_impairment_two_domains_young_middle_aged** — n=1 paper · extraction confidence: 72%
- **severe_impairment_three_domains_young_middle_aged** — n=1 paper · extraction confidence: 72%
- **mild_impairment_one_domain_older_adults** — n=1 paper · extraction confidence: 72%
- **moderate_impairment_two_domains_older_adults** — n=1 paper · extraction confidence: 72%
- **severe_impairment_three_domains_older_adults** — n=1 paper · extraction confidence: 72%
- **persistent_symptoms_at_6_months** — n=1 paper · extraction confidence: 72%
- **long_covid_overall_prevalence_one_study** — n=1 paper · extraction confidence: 72%
- **persistent_symptoms_>20_weeks** — n=1 paper · extraction confidence: 72%
- **recovered_patients_with_symptoms_at_6_months** — n=1 paper · extraction confidence: 72%
- **persistent_symptoms_at_12_months_approximately** — n=1 paper · extraction confidence: 72%
- **fatigue_cfs_global_prevalence_prior_review** — n=1 paper · extraction confidence: 72%
- **fatigue_cfs_sub_saharan_africa_health_professionals** — n=1 paper · extraction confidence: 72%
- **post_covid_syndrome_hospitalized_patients_one_study** — n=1 paper · extraction confidence: 72%
- **nonhospitalized_patients_developing_post_covid** — n=1 paper · extraction confidence: 72%

- **hospitalized_patients_developing_post_covid** — n=1 paper · extraction confidence: 72%
- **vaccinated_individuals_developing_post_covid** — n=1 paper · extraction confidence: 72%
- **fatigue_3months** — n=1 paper · extraction confidence: 72%
- **fatigue_6months** — n=1 paper · extraction confidence: 72%
- **fatigue_9months** — n=1 paper · extraction confidence: 72%
- **fatigue_12months** — n=1 paper · extraction confidence: 72%
- **memory_decline_3months** — n=1 paper · extraction confidence: 72%
- **memory_decline_6months** — n=1 paper · extraction confidence: 72%
- **memory_decline_9months** — n=1 paper · extraction confidence: 72%
- **memory_decline_12months** — n=1 paper · extraction confidence: 72%
- **cough_expectoration_3months** — n=1 paper · extraction confidence: 72%
- **decreased_exercise_ability_3months** — n=1 paper · extraction confidence: 72%
- **decreased_exercise_ability_6months** — n=1 paper · extraction confidence: 72%
- **decreased_exercise_ability_9months** — n=1 paper · extraction confidence: 72%
- **decreased_exercise_ability_12months** — n=1 paper · extraction confidence: 72%
- **thirst_dry_mouth_3months** — n=1 paper · extraction confidence: 72%
- **sleep_disorders_3months** — n=1 paper · extraction confidence: 72%
- **sleep_disorders_6months** — n=1 paper · extraction confidence: 72%

- **sleep_disorders_9months** — n=1 paper · extraction confidence: 72%
- **sleep_disorders_12months** — n=1 paper · extraction confidence: 72%
- **brain_fog_3months** — n=1 paper · extraction confidence: 72%
- **brain_fog_6months** — n=1 paper · extraction confidence: 72%
- **brain_fog_9months** — n=1 paper · extraction confidence: 72%
- **brain_fog_12months** — n=1 paper · extraction confidence: 72%
- **fatigue_single_infection_3months** — n=1 paper · extraction confidence: 72%
- **memory_decline_single_infection_3months** — n=1 paper · extraction confidence: 72%
- **decreased_exercise_ability_single_infection_3months** — n=1 paper · extraction confidence: 72%
- **at_least_one_acute_covid19_symptom** — n=1 paper · extraction confidence: 72%
- **non_recovery_at_90_days_plus** — n=1 paper · extraction confidence: 72%
- **health_deterioration_vs_pre_covid** — n=1 paper · extraction confidence: 72%
- **anxiety_depression_increase** — n=1 paper · extraction confidence: 72%
- **mobility_issues_increase** — n=1 paper · extraction confidence: 72%
- **pain_discomfort_increase** — n=1 paper · extraction confidence: 72%
- **limitations_daily_activities_increase** — n=1 paper · extraction confidence: 72%

- **persistent_symptoms_at_one_year** — n=1 paper · extraction confidence: 72%
- **persistent_symptoms_post_discharge** — n=1 paper · extraction confidence: 72%
- **fatigue_and_exertional_dyspnea** — n=1 paper · extraction confidence: 72%
- **impaired_lung_diffusion** — n=1 paper · extraction confidence: 72%
- **elevated_d_dimer_twice_threshold** — n=1 paper · extraction confidence: 72%
- **new_onset_dyspnea_jutant** — n=1 paper · extraction confidence: 72%
- **new_onset_cough_jutant** — n=1 paper · extraction confidence: 72%
- **pe_in_dyspnea_patients_ekici** — n=1 paper · extraction confidence: 72%
- **cardiovascular_long_covid_murata** — n=1 paper · extraction confidence: 72%
- **current_long_covid_in_nhis_sample** — n=1 paper · extraction confidence: 72%
- **cvd_among_current_long_covid** — n=1 paper · extraction confidence: 72%
- **cvd_among_no_long_covid** — n=1 paper · extraction confidence: 72%
- **fatigue_lasting_over_8_weeks** — n=1 paper · extraction confidence: 72%
- **general_malaise_lasting_over_8_weeks** — n=1 paper · extraction confidence: 72%
- **memory_problems_lasting_over_8_weeks** — n=1 paper · extraction confidence: 72%
- **brain_fog_lasting_over_8_weeks** — n=1 paper · extraction confidence: 72%

- **memory_problems_lasting_over_6_months** — n=1 paper · extraction confidence: 72%
- **fatigue_lasting_over_6_months** — n=1 paper · extraction confidence: 72%
- **long_covid_in_endometriosis_patients** — n=1 paper · extraction confidence: 72%
- **long_covid_in_non_endometriosis_patients** — n=1 paper · extraction confidence: 72%
- **persistent_fatigue** — n=1 paper · extraction confidence: 72%
- **anosmia_dysgeusia** — n=1 paper · extraction confidence: 72%
- **pasc_cope_definition** — n=1 paper · extraction confidence: 72%
- **pasc_nice_definition** — n=1 paper · extraction confidence: 72%
- **pasc_recover_definition** — n=1 paper · extraction confidence: 72%
- **mobility_impairment_lc_u** — n=1 paper · extraction confidence: 72%
- **mobility_impairment_nc** — n=1 paper · extraction confidence: 72%
- **mobility_impairment_lc_r** — n=1 paper · extraction confidence: 72%
- **mobility_impairment_sc** — n=1 paper · extraction confidence: 72%
- **difficulty_seeing_lc_u** — n=1 paper · extraction confidence: 72%
- **difficulty_seeing_lc_r** — n=1 paper · extraction confidence: 72%
- **difficulty_seeing_nc** — n=1 paper · extraction confidence: 72%

- **difficulty_seeing_sc** — n=1 paper · extraction confidence: 72%
- **difficulty_hearing_lc_u** — n=1 paper · extraction confidence: 72%
- **difficulty_hearing_nc** — n=1 paper · extraction confidence: 72%
- **difficulty_hearing_lc_r** — n=1 paper · extraction confidence: 72%
- **difficulty_hearing_sc** — n=1 paper · extraction confidence: 72%
- **difficulty_concentrating_lc_u** — n=1 paper · extraction confidence: 72%
- **difficulty_concentrating_lc_r** — n=1 paper · extraction confidence: 72%
- **difficulty_concentrating_nc** — n=1 paper · extraction confidence: 72%
- **difficulty_concentrating_sc** — n=1 paper · extraction confidence: 72%
- **residual_symptoms_4_to_12_weeks** — n=1 paper · extraction confidence: 72%
- **mild_clinical_condition_acute_phase** — n=1 paper · extraction confidence: 72%
- **recurrent_infections** — n=1 paper · extraction confidence: 72%
- **high_vaccination_adherence** — n=1 paper · extraction confidence: 72%
- **chronic_disease_any** — n=1 paper · extraction confidence: 72%
- **systemic_arterial_hypertension** — n=1 paper · extraction confidence: 72%
- **diabetes_mellitus** — n=1 paper · extraction confidence: 72%
- **heart_disease** — n=1 paper · extraction confidence: 72%
- **cancer** — n=1 paper · extraction confidence: 72%

- **hiv** — n=1 paper · extraction confidence: 72%
- **dysphagia** — n=1 paper · extraction confidence: 72%
- **nausea_or_vomiting** — n=1 paper · extraction confidence: 72%
- **gastroesophageal_reflux** — n=1 paper · extraction confidence: 72%
- **dizziness_or_vertigo** — n=1 paper · extraction confidence: 72%
- **stress** — n=1 paper · extraction confidence: 72%
- **paresthesia** — n=1 paper · extraction confidence: 72%
- **smell_and_taste_problems** — n=1 paper · extraction confidence: 72%
- **weight_gain** — n=1 paper · extraction confidence: 72%
- **smi_overall** — n=1 paper · extraction confidence: 72%
- **smi_current_lc** — n=1 paper · extraction confidence: 72%
- **smi_previous_lc** — n=1 paper · extraction confidence: 72%
- **smi_infection_no_lc** — n=1 paper · extraction confidence: 72%
- **smi_non_infected** — n=1 paper · extraction confidence: 72%
- **cough_cases** — n=1 paper · extraction confidence: 72%
- **cough_controls** — n=1 paper · extraction confidence: 72%
- **pain_cases** — n=1 paper · extraction confidence: 72%
- **pain_controls** — n=1 paper · extraction confidence: 72%
- **gastrointestinal_symptoms_cases** — n=1 paper · extraction confidence: 72%
- **gastrointestinal_symptoms_controls** — n=1 paper · extraction confidence: 72%
- **anxiety_cases** — n=1 paper · extraction confidence: 72%
- **anxiety_controls** — n=1 paper · extraction confidence: 72%
- **persisting_chest_xray_changes** — n=1 paper · extraction confidence: 72%
- **arterial_hypertension_comorbidity** — n=1 paper · extraction confidence: 72%

- **hypothyroidism_comorbidity** — n=1 paper · extraction confidence: 72%
- **cardiac_arrhythmias_comorbidity** — n=1 paper · extraction confidence: 72%
- **mobility_impairment** — n=1 paper · extraction confidence: 72%
- **vomit** — n=1 paper · extraction confidence: 72%
- **encephalopathy** — n=1 paper · extraction confidence: 72%
- **stroke_associated** — n=1 paper · extraction confidence: 72%
- **haemorrhage** — n=1 paper · extraction confidence: 72%
- **ischaemic_infarcts** — n=1 paper · extraction confidence: 72%
- **ct_abnormalities_overall** — n=1 paper · extraction confidence: 72%
- **dlco_less_than_80_percent** — n=1 paper · extraction confidence: 72%
- **at_least_1_respiratory_symptom** — n=1 paper · extraction confidence: 72%
- **mmrc_dyspnea_scale_0** — n=1 paper · extraction confidence: 72%
- **fatigue_alpha_variant** — n=1 paper · extraction confidence: 72%
- **at_least_1_general_symptom_alpha_variant** — n=1 paper · extraction confidence: 72%
- **cough_alpha_variant** — n=1 paper · extraction confidence: 72%
- **dyspnea_alpha_variant** — n=1 paper · extraction confidence: 72%
- **at_least_1_symptom_omicron** — n=1 paper · extraction confidence: 72%
- **sleep_difficulty_omicron** — n=1 paper · extraction confidence: 72%

- **fatigue_omicron** — n=1 paper · extraction confidence: 72%
- **myalgia_omicron** — n=1 paper · extraction confidence: 72%
- **myalgia_wild_type** — n=1 paper · extraction confidence: 72%
- **sleep_difficulty_wild_type** — n=1 paper · extraction confidence: 72%
- **sleep_difficulty_delta** — n=1 paper · extraction confidence: 72%
- **anxiety_several_days_or_more_lc** — n=1 paper · extraction confidence: 72%
- **anxiety_several_days_or_more_sc** — n=1 paper · extraction confidence: 72%
- **anxiety_several_days_or_more_nc** — n=1 paper · extraction confidence: 72%
- **unable_to_stop_worrying_lc** — n=1 paper · extraction confidence: 72%
- **unable_to_stop_worrying_sc** — n=1 paper · extraction confidence: 72%
- **unable_to_stop_worrying_nc** — n=1 paper · extraction confidence: 72%
- **little_pleasure_or_interest_lc** — n=1 paper · extraction confidence: 72%
- **little_pleasure_or_interest_sc** — n=1 paper · extraction confidence: 72%
- **little_pleasure_or_interest_nc** — n=1 paper · extraction confidence: 72%
- **feeling_down_depressed_hopeless_lc** — n=1 paper · extraction confidence: 72%
- **feeling_down_depressed_hopeless_sc** — n=1 paper · extraction confidence: 72%
- **feeling_down_depressed_hopeless_nc** — n=1 paper · extraction confidence: 72%

- **cognitive_impairment_brain_fog** — n=1 paper · extraction confidence: 72%
- **worse_health_transition_frailty_group** — n=1 paper · extraction confidence: 72%
- **worse_health_transition_non_frailty_group** — n=1 paper · extraction confidence: 72%
- **lack_of_energy_longcovid** — n=1 paper · extraction confidence: 72%
- **headache_longcovid** — n=1 paper · extraction confidence: 72%
- **muscle_aches_longcovid** — n=1 paper · extraction confidence: 72%
- **fever_longcovid** — n=1 paper · extraction confidence: 72%
- **shortness_of_breath_dyspnea_longcovid** — n=1 paper · extraction confidence: 72%
- **dizziness_non_hospitalized** — n=1 paper · extraction confidence: 72%
- **dizziness_hospitalized** — n=1 paper · extraction confidence: 72%
- **heavy_limbs_non_hospitalized** — n=1 paper · extraction confidence: 72%
- **dyspnea_non_hospitalized** — n=1 paper · extraction confidence: 72%
- **tremors_non_hospitalized** — n=1 paper · extraction confidence: 72%
- **heavy_limbs_hospitalized** — n=1 paper · extraction confidence: 72%
- **at_least_1_long_covid_cluster_3_months** — n=1 paper · extraction confidence: 72%
- **persistent_fatigue_with_bodily_pain_or_mood_swings** — n=1 paper · extraction confidence: 72%
- **ongoing_respiratory_problems** — n=1 paper · extraction confidence: 72%

- **cognitive_problems** — n=1 paper · extraction confidence: 72%
- **women_aged_20_plus_any_cluster** — n=1 paper · extraction confidence: 72%
- **men_aged_20_plus_any_cluster** — n=1 paper · extraction confidence: 72%
- **both_sexes_under_20_any_cluster** — n=1 paper · extraction confidence: 72%
- **still_symptomatic_at_12_months_among_3month_cases** — n=1 paper · extraction confidence: 72%
- **long_covid_any_antihm_two_infections_vaccinated_preinfection** — n=1 paper · extraction confidence: 72%
- **long_covid_any_antihm_three_or_more_infections** — n=1 paper · extraction confidence: 72%
- **thrombosis_antihm_at_least_1_chronic_treatment** — n=1 paper · extraction confidence: 72%
- **thrombosis_noantihm_at_least_1_chronic_treatment** — n=1 paper · extraction confidence: 72%
- **thrombosis_population_over_60_most_cst_centers** — n=1 paper · extraction confidence: 72%
- **long_covid_symptom_clusters_overall** — n=1 paper · extraction confidence: 72%
- **pulmonary_symptoms** — n=1 paper · extraction confidence: 72%
- **cardiac_symptoms** — n=1 paper · extraction confidence: 72%
- **pasc_any_feature_test_positive** — n=1 paper · extraction confidence: 62%
- **pasc_any_feature_test_negative** — n=1 paper · extraction confidence: 62%
- **cardiac_subphenotype_covid_positive** — n=1 paper · extraction confidence: 62%

- **respiratory_subphenotype_covid_positive** — n=1 paper · extraction confidence: 62%
- **musculoskeletal_subphenotype_covid_positive** — n=1 paper · extraction confidence: 62%
- **cardiac_subphenotype_covid_negative** — n=1 paper · extraction confidence: 62%
- **respiratory_subphenotype_covid_negative** — n=1 paper · extraction confidence: 62%
- **musculoskeletal_subphenotype_covid_negative** — n=1 paper · extraction confidence: 62%
- **at_least_one_acute_covid_symptom** — n=1 paper · extraction confidence: 62%
- **long_covid_prevalence_by_who_definition** — n=1 paper · extraction confidence: 62%
- **cognitive_dysfunction** — n=1 paper · extraction confidence: 62%
- **muscle_pain_percentage_change** — n=1 paper · extraction confidence: 62%
- **hair_loss_percentage_change** — n=1 paper · extraction confidence: 62%
- **long_covid_diagnosis_ever** — n=1 paper · extraction confidence: 62%
- **long_covid_diagnosis_females** — n=1 paper · extraction confidence: 62%
- **long_covid_diagnosis_males** — n=1 paper · extraction confidence: 62%
- **long_covid_among_vaccinated** — n=1 paper · extraction confidence: 62%
- **cognitive_impairment_memory_loss** — n=1 paper · extraction confidence: 62%
- **liver_function_abnormalities** — n=1 paper · extraction confidence: 62%

- **depression_anxiety** — n=1 paper · extraction confidence: 62%
- **unrefreshing_sleep_or_sleep_disturbance_mecfs** — n=1 paper · extraction confidence: 62%
- **debilitating_fatigue_mecfs** — n=1 paper · extraction confidence: 62%
- **brain_fog_cognitive_difficulties_mecfs** — n=1 paper · extraction confidence: 62%
- **pain_muscles_joints_nerves_mecfs** — n=1 paper · extraction confidence: 62%
- **postexertional_malaise_mecfs** — n=1 paper · extraction confidence: 62%
- **light_noise_sensitivity_mecfs** — n=1 paper · extraction confidence: 62%
- **headache_mecfs** — n=1 paper · extraction confidence: 62%
- **ibs_mecfs** — n=1 paper · extraction confidence: 62%
- **temperature_sensitivity_mecfs** — n=1 paper · extraction confidence: 62%
- **dizziness_mecfs** — n=1 paper · extraction confidence: 62%
- **debilitating_fatigue_long_covid** — n=1 paper · extraction confidence: 62%
- **brain_fog_cognitive_difficulties_long_covid** — n=1 paper · extraction confidence: 62%
- **pain_muscles_joints_nerves_long_covid** — n=1 paper · extraction confidence: 62%
- **unrefreshing_sleep_long_covid** — n=1 paper · extraction confidence: 62%
- **breathlessness_long_covid** — n=1 paper · extraction confidence: 62%
- **postexertional_malaise_long_covid** — n=1 paper · extraction confidence: 62%
- **ibs_long_covid** — n=1 paper · extraction confidence: 62%

- **dizziness_long_covid** — n=1 paper · extraction confidence: 62%
- **ringing_in_ears_long_covid** — n=1 paper · extraction confidence: 62%
- **excessive_sweating_when_asleep_long_covid** — n=1 paper · extraction confidence: 62%
- **pasc_positive_at_6_months_omicron_era_subsample** — n=1 paper · extraction confidence: 62%
- **post_covid_symptoms_general** — n=1 paper · extraction confidence: 55%
- **one_or_more_persistent_symptoms_at_3_months** — n=1 paper · extraction confidence: 55%
- **at_least_one_persistent_long_covid_symptom** — n=1 paper · extraction confidence: 52%
- **postexertional_malaise** — n=1 paper · extraction confidence: 52%
- **change_in_smell_or_taste** — n=1 paper · extraction confidence: 52%
- **thirst** — n=1 paper · extraction confidence: 52%
- **chronic_cough** — n=1 paper · extraction confidence: 52%
- **sleep_apnea** — n=1 paper · extraction confidence: 52%
- **executive_deficits_above_clinical_threshold_sars-cov-2_positive** — n=1 paper · extraction confidence: 45%
- **executive_deficits_above_clinical_threshold_controls** — n=1 paper · extraction confidence: 45%
- **breathing_difficulties_covid19** — n=1 paper · extraction confidence: 42%
- **breathing_difficulties_ili_controls** — n=1 paper · extraction confidence: 42%
- **type2_diabetes_covid19** — n=1 paper · extraction confidence: 42%
- **type2_diabetes_ili_controls** — n=1 paper · extraction confidence: 42%

- **fatigue_covid19** — n=1 paper · extraction confidence: 42%
 - **fatigue_ili_controls** — n=1 paper · extraction confidence: 42%
 - **sleep_disturbances_covid19** — n=1 paper · extraction confidence: 42%
 - **sleep_disturbances_ili_controls** — n=1 paper · extraction confidence: 42%
 - **ageusia** — n=1 paper · extraction confidence: 42%
 - **impaired_focus** — n=1 paper · extraction confidence: 42%
 - **fatigue_highly_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **heart_conditions_highly_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **sleep_disturbances_highly_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **breathing_difficulties_highly_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **sleep_disturbances_secondary_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **fatigue_secondary_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **mental_health_secondary_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **cognition_secondary_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **type_2_diabetes_secondary_comorbid_class** — n=1 paper · extraction confidence: 31%
 - **stomach_pain** — n=1 paper · extraction confidence: 28%
-

2. Symptom Consensus (Triage-Level)

Across 4,666 triaged papers, fatigue is the most frequently indexed symptom (34.6% of papers), followed by anxiety (11.4%), depression (10.1%), dyspnea (9.4%), brain fog (7.5%), and cognitive impairment/dysfunction (combined ~13%). When cognitive symptom terms are deduplicated (brain fog, cognitive impairment, cognitive dysfunction, difficulty concentrating), neurocognitive complaints constitute a second major domain. Anosmia and ageusia, while appearing in fewer papers (2.7% for anosmia), are among the most COVID-specific symptoms identified in controlled studies, with odds ratios exceeding 4.0 in properly controlled analyses [31,32]. Post-exertional malaise appears in 3.4% of papers and is disproportionately represented in the deeply analyzed literature as a key phenotypic marker.

The findings_by_certainty partition yields zero established findings, zero probable findings at the aggregate consensus level, 147 possible findings, and 2,799 speculative findings. This distribution reflects that no symptom prevalence estimate survives cross-study synthesis with high certainty. The 'possible' tier (147 findings) is entirely symptom-prevalence based, with mean extraction confidence between 0.70 and 0.87. Speculative findings (2,799) encompass mechanistic attributions, subgroup-specific risk estimates, and phenotypic characterizations that rest on single studies or uncontrolled designs.

Prevalence estimates vary dramatically by ascertainment method: ICD-10-coded administrative studies report 0.2–3.9% [73,74,75], symptom-based EHR studies 15–45% [76,53], and self-report surveys 7–28% among adults with prior COVID-19 [17,77,78]. The Long-CISS Scotland prospective cohort of 198,096 participants estimated adjusted attributable prevalence at 6.6% (95% CI 6.3–6.9%) at six months [79], representing one of the more methodologically robust population estimates. These ranges are

not reconcilable without harmonized case definitions and controlled ascertainment.

Symptom	Studies reporting	Prevalence
fatigue	1173	34.6%
anxiety	386	11.4%
depression	341	10.1%
dyspnea	320	9.4%
brain fog	254	7.5%
cognitive impairment	245	7.2%
cognitive dysfunction	198	5.8%
headache	196	5.8%
cough	161	4.7%
persistent symptoms	141	4.2%
chest pain	133	3.9%
shortness of breath	132	3.9%
pain	122	3.6%
post-exertional malaise	114	3.4%
respiratory symptoms	108	3.2%
neurological symptoms	106	3.1%
insomnia	98	2.9%
anosmia	92	2.7%
sleep disturbance	81	2.4%
dizziness	80	2.4%
myalgia	79	2.3%

Symptom	Studies reporting	Prevalence
palpitations	72	2.1%
dyspnoea	70	2.1%
sleep disturbances	68	2.0%
breathlessness	63	1.9%
muscle pain	61	1.8%
gastrointestinal symptoms	59	1.7%
joint pain	59	1.7%
headaches	55	1.6%
exercise intolerance	53	1.6%

3. Definition Heterogeneity

Case definition fragmentation constitutes the most fundamental barrier to evidence synthesis in this field. Across the triaged corpus, at least 39 distinct symptom-duration thresholds were identified, with the most common being 12 weeks (n=429 papers), 4 weeks (n=246), and 26 weeks (n=85). This distribution reflects the coexistence of WHO 2021 (12-week threshold, symptoms arising within 3 months of infection, lasting at least 2 months), NICE (4-week threshold), and innumerable study-specific definitions. One study applying five published definitions to the same INSPIRE cohort found 3-month prevalence ranging from 30.84% to 91.02% [\[80\]](#), while a French national survey of 10,615 adults found prevalence ranging from 1.2% to 13.4% depending on definition applied [\[81\]](#).

Administrative coding studies using ICD-10 U09.9 or SNOMED codes capture only a fraction of clinical cases, with coded

prevalence consistently 5–50-fold lower than symptom-based estimates [76,82,73]. Computable phenotype studies using machine learning to expand case ascertainment identified 4-fold more PASC cases than ICD coding alone [83], but introduce their own circularity risks when trained on the same ICD codes [84]. Studies using study-specific definitions predominate the literature and are generally non-comparable. No study in this corpus used a fully validated, prospectively agreed consensus definition with objective clinical anchoring.

Threshold (weeks)	Number of studies
≥12 weeks	429
≥4 weeks	246
≥26 weeks	85
≥8 weeks	66
≥52 weeks	41
≥24 weeks	29
≥16 weeks	18
≥104 weeks	15
≥6 weeks	13
≥3 weeks	11
≥2 weeks	6
≥36 weeks	6
≥18 weeks	6
≥25 weeks	5
≥156 weeks	5
≥20 weeks	5

Threshold (weeks)	Number of studies
≥78 weeks	4
≥17 weeks	4
≥10 weeks	4
≥40 weeks	3
≥7 weeks	3
≥80 weeks	3
≥5 weeks	3
≥28 weeks	3
≥35 weeks	2
≥60 weeks	2
≥25.7 weeks	2
≥8.57 weeks	2
≥30 weeks	1
≥8.6 weeks	1
≥19 weeks	1
≥11 weeks	1
≥48 weeks	1
≥124 weeks	1
≥21 weeks	1
≥2.857 weeks	1
≥14 weeks	1
≥32 weeks	1
≥70 weeks	1

Study design distribution:

Design	Count
cohort	1206
review	935
cross_sectional	815
other	504
case_control	444
case_report	341
meta_analysis	202
RCT	166
qualitative	36
mixed_methods	5
systematic_review	4
scoping_review	3
systematic review	3
cost_of_illness	1
scoping review	1

4. Methodology Quality (NOS + GRADE)

Of 470 deeply analyzed papers, no study reached GRADE High quality, reflecting the absence of randomized controlled trial evidence for most prevalence and risk factor questions. The 19 Moderate-quality studies (4.0%) were predominantly large prospective cohorts or national register-based studies with active

comparator designs [85,10,79,22,11]. The mean Newcastle-Ottawa Scale score of 5.57 out of 9 indicates that most observational studies have meaningful methodological deficiencies in comparability or outcome ascertainment domains. The 265 Low-quality studies (56.4%) typically had adequate selection but inadequate outcome validation or unmeasured confounding. The 186 Very Low-quality studies (39.6%) included retrospective cross-sectional analyses, social media studies, small case series, and studies with fundamental design flaws such as absence of control groups, non-validated outcomes, or circular case definitions. GRADE downgrading was driven primarily by risk of bias (surveillance bias, selection bias, circular case definition), inconsistency (extreme heterogeneity, $I^2 > 90\%$ in most meta-analyses), and indirectness (non-representative populations, surrogate endpoints).

GRADE certainty across the 470 deeply-analysed papers:

GRADE level	Count
High	0
Moderate	19
Low	265
Very Low	186

Mean Newcastle-Ottawa score (observational subset): 5.57/9

5. Bias Audit Summary

Bias category	Frequency
Surveillance bias	411 of 470 papers (87.4%)

Bias category	Frequency
Baseline data absent	274 of 470 papers (58.3%)
No vaccination/variant adjustment	444 of 470 papers (94.5%)
Self-report bias (or ICD-code only)	284 of 470 papers (60.4%)
Circular case definition	149 of 470 papers (31.7%)

The bias audit reveals pervasive and overlapping methodological deficiencies. Surveillance bias affects 87.4% of deeply analyzed papers, reflecting the near-universal dependence on ICD coding, healthcare contact, or self-report for case ascertainment—all mechanisms that favor detection in individuals who seek care [86,87,88]. The worst-offender pattern is the administrative coding study that simultaneously defines exposure by COVID-19 diagnosis code and outcome by Long COVID code in the same EHR system, creating a structural positive correlation unrelated to true disease [74,89].

Vaccination adjustment is absent in 94.5% of deep papers—the single most consequential unmeasured confounder given vaccine effectiveness against Long COVID of approximately 23–37% in meta-analytic estimates [4,9]. Circular case definition affects 31.7% of studies, including instances where psychiatric disorders are treated as risk factors for PASC despite sharing symptomatic overlap with PASC diagnostic criteria [86,87]. Baseline absence of pre-infection symptom data in 58.3% of papers prevents distinction between new post-COVID symptoms and pre-existing conditions. Self-report bias in 60.4% of papers is compounded by pandemic-era hyperawareness of Long COVID as a diagnostic category, potentially inflating symptom attribution.

6. Mechanistic Phenotype Distribution

Mechanism	Papers
Unspecified	275
Mixed	109
Autonomic Metabolic	75
Autoimmunity	58
Vascular Endothelial	52
Viral Reservoir	25

The dominant phenotype classification across 470 deeply analyzed papers is 'unspecified' (n=275, 58.5%), reflecting that the majority of studies do not investigate or attribute biological mechanisms and rely solely on symptom-based or ICD-coded outcomes. Among studies with mechanism attribution, 'mixed' phenotypes (n=109) are most common, followed by autonomic/metabolic (n=75), autoimmunity (n=58), vascular/endothelial (n=52), and viral reservoir (n=25). This distribution reflects the field's conceptual fragmentation rather than mechanistic resolution: studies proposing autonomic/metabolic mechanisms cite dysautonomia, POTS, and post-exertional malaise clusters; autoimmunity studies reference persistent T-cell activation, autoantibody generation, and ME/CFS overlap; vascular/endothelial studies emphasize microclot formation, endotheliopathy, and coagulation abnormalities; and viral reservoir studies cite gastrointestinal and CNS persistence of SARS-CoV-2 RNA. None of these mechanistic frameworks has been validated through a sufficiently powered prospective study with objective biomarker endpoints in an unselected Long COVID population.

7. Recurring Methodological Errors

1. Absence of vaccination status adjustment (444 of 470 deep papers (94.5%))

Vaccination is a well-established modifier of Long COVID risk with estimated effectiveness of 23–37%, yet 94.5% of deeply analyzed papers failed to adjust for vaccination status in primary analyses. This introduces unmeasured confounding that can inflate or deflate risk estimates for any covariate correlated with vaccination uptake. This is particularly problematic in studies spanning 2021–2024.

Example papers: [\[2\]](#), [\[43\]](#), [\[90\]](#), [\[88\]](#), [\[4\]](#)

2. Circular case definition via ICD coding (149 of 470 deep papers (31.7%))

Studies that define Long COVID by ICD-10 U09.9 codes and simultaneously analyze risk factors coded in the same EHR create circular ascertainment: healthcare-engaged patients with more comorbidities receive more ICD codes, inflating both exposure and outcome measurements. Additionally, psychiatric symptom codes (fatigue, cognitive disturbance) overlap directly with Long COVID diagnostic criteria.

Example papers: [\[84\]](#), [\[87\]](#), [\[86\]](#), [\[91\]](#), [\[74\]](#)

3. Extreme statistical heterogeneity with uncritical pooling (Identified in all major meta-analyses across the corpus; affects approximately 30–40 of 202 meta-analyses in the triaged corpus)

Meta-analyses consistently report $I^2 \geq 90$ –99% yet proceed to present pooled estimates as clinically meaningful summary statistics. $I^2 = 99.5\%$ [\[2\]](#) and $I^2 \approx 98\%$ [\[4\]](#) render random-effects

pooled estimates mathematically valid but clinically uninterpretable as a central tendency. Fixed-effects models under such conditions are inappropriate; the correct response is subgroup analysis or narrative synthesis.

Example papers: [\[2\]](#), [\[4\]](#), [\[92\]](#), [\[93\]](#), [\[94\]](#)

4. Surveillance bias through healthcare utilization as outcome proxy (411 of 470 deep papers (87.4%))

When Long COVID is ascertained through healthcare contacts (clinic visits, prescriptions, ICD codes), patients with more frequent baseline healthcare engagement are systematically more likely to receive a Long COVID diagnosis independent of true disease burden. This creates a positive correlation between healthcare-associated exposures and Long COVID that does not reflect biological causation.

Example papers: [\[28\]](#), [\[86\]](#), [\[87\]](#), [\[73\]](#), [\[95\]](#)

5. Absence of uninfected or negative-tested control group (274 of 470 deep papers reporting baseline absence (58.3%); control group absent in approximately 100-120 deep papers)

Without a concurrent uninfected comparison group assessed with identical instruments at identical time points, it is impossible to distinguish COVID-19-attributable symptoms from background population symptom rates, pandemic-related psychosocial effects, or regression to the mean. Studies without controls produce absolute prevalence estimates that are uninterpretable as excess burden.

Example papers: [\[43\]](#), [\[2\]](#), [\[96\]](#), [\[97\]](#), [\[35\]](#)

6. Channeling bias and confounding by indication in treatment studies (Affects the majority of the 166 RCTs and treatment-focused retrospective cohorts; prominent in antiviral and immunomodulatory studies)

Antiviral prescriptions (Paxlovid, molnupiravir, remdesivir) and immunomodulatory therapies are channeled to patients with specific risk profiles, creating systematic differences between treated and untreated groups that propensity score matching cannot fully eliminate, particularly for unmeasured confounders such as acute disease trajectory and patient activation.

Example papers: [\[98\]](#), [\[99\]](#), [\[100\]](#), [\[101\]](#), [\[57\]](#)

7. Self-reported exposure and outcome with no clinical validation (284 of 470 deep papers (60.4%))

Studies relying entirely on self-reported COVID-19 diagnosis and self-reported Long COVID symptoms without PCR/serology confirmation or clinical adjudication cannot establish exposure status with confidence and are subject to differential misclassification if Long COVID awareness affects symptom attribution.

Example papers: [\[102\]](#), [\[103\]](#), [\[104\]](#), [\[105\]](#), [\[106\]](#)

8. Variant era confounding without stratification (Identified as a limitation in the majority of studies spanning multiple pandemic waves; explicitly flagged in bias audit for 444 papers for vaccination and variant confounding combined)

Studies spanning pre-Delta, Delta, and Omicron eras pool populations with markedly different Long COVID risk profiles (Omicron associated with lower risk in vaccinated individuals) without variant-stratified analyses, obscuring era-specific

estimates and producing pooled results that do not correspond to any real-world exposure scenario.

Example papers: [\[107\]](#), [\[108\]](#), [\[12\]](#), [\[4\]](#), [\[109\]](#)

8. Contradictions Between Papers

Overall Long COVID prevalence in African populations

[\[2\]](#): "Pooled Long COVID prevalence in African populations was 41% (95% CI 26–56%) with extreme heterogeneity ($I^2=99.5\%$)"

[\[93\]](#): "Cumulative incidence of Long COVID was 48.6% (95% CI 37.4–59.8%) across 25 studies with 29,213 African participants"

Likely cause: Different inclusion criteria and study sets; both estimates are statistically uninterpretable at $I^2 \geq 99\%$, but the divergence also reflects different country compositions and hospitalization rates in the included study pools

Paxlovid effectiveness against overall Long COVID incidence

[\[36\]](#): "Paxlovid treatment was not significantly associated with overall PASC incidence (RR 0.986, 95% CI 0.953–1.020) but reduced cognitive symptom cluster (RR 0.91, 95% CI 0.84–0.99) in a target trial emulation of 445,738 patients"

[\[38\]](#): "Nirmatrelvir use was associated with a 26% reduced relative risk of post-COVID condition (RR 0.74, 95% CI 0.72–0.77) in a retrospective cohort of 281,793 patients"

Likely cause: Differences in PASC computable phenotype definitions, study populations (PCORnet mixed payer vs. VA predominantly male veterans), channeling bias magnitude, and era (Omicron-dominant in both but different vaccination contexts)

HIV infection and Long COVID risk

[41]: "HIV infection was significantly associated with increased risk of Long COVID (OR 2.20, 95% CI 1.25–3.86) in a meta-analysis of 5 studies (n=39,405)"

[42]: "People with HIV had a small but statistically significant increase in odds of Long COVID using the computed phenotype definition but no significant association using ICD-10 code-based definition in a multicenter study (n=4,682,251)"

Likely cause: Phenotype definition sensitivity; the ICD-code outcome is subject to severe surveillance bias (PWH have more healthcare contacts) while the computed phenotype may partially adjust for this; the meta-analysis pools small heterogeneous studies with confounding by ART status and viral suppression

Effect of pre-existing diabetes on Long COVID risk

[39]: "Pre-pandemic diabetes was associated with a 60% higher risk of Long COVID in a prospective UK Household Longitudinal Study cohort of 11,669 participants (RR 1.60, 95% CI 1.12–2.29)"

[40]: "Pre-existing diabetes was not significantly associated with Long COVID after full adjustment in the SHARE

prospective cohort of 4,004 older adults aged 50+ (OR 1.37, non-significant after adjustment)"

Likely cause: Differences in cohort age distribution (SHARE restricted to ≥ 50), variant era (SHARE may have greater pre-Omicron overlap), diabetes type heterogeneity, and Long COVID definition (UKHLS used 4-week and 12-week thresholds; SHARE used symptom persistence)

Long COVID prevalence in cancer patients versus general population

[110]: "Among 76,807 patients with cancer during Omicron predominance, overall risk of postacute diagnoses was not significantly elevated (HR not significant) compared to test-negative cancer patients"

[111]: "Pooled prevalence of Long COVID in cancer patients was 23.52% (95% CI 12.14–40.64%) with extreme heterogeneity ($I^2=98.3\%$) across 6,653 participants"

Likely cause: The Omicron-era cohort study uses a test-negative active comparator design and overlap-weighted propensity scoring, substantially reducing apparent excess; the meta-analysis pools absolute prevalence across eras without control groups, making the estimates non-comparable

Long COVID risk in pregnant versus non-pregnant individuals

[70]: "Pregnancy at time of SARS-CoV-2 infection was associated with lower Long COVID risk (aHR 0.86, 95% CI 0.83–0.90) in a retrospective cohort of 1,511,505 females"

[71]: "SARS-CoV-2 infection during pregnancy was associated with lower PASC incidence at 180 days (30.8 per 100 vs 35.8 per 100 in non-pregnant females) in a retrospective cohort of 89,312"

Likely cause: While both find lower PASC in pregnant individuals, the RECOVER-Pregnancy prospective analysis **[72]** finds no statistically significant difference (10.2% vs 10.6%), suggesting the apparent protective effect in administrative data may reflect differential healthcare utilization surveillance bias rather than biological protection

9. Research Gaps

Randomized controlled trial evidence for Long COVID pharmacological interventions — Of 4,666 triaged papers, only 166 were classified as RCTs, and the living systematic review of interventions found moderate certainty evidence only for online CBT and combined rehabilitation; no pharmacological intervention has achieved high or moderate GRADE certainty for Long COVID outcomes. Metformin, antihistamines, anticoagulants, and antivirals are all being studied in observational frameworks with substantial channeling bias. See: **[26]**, **[112]**, **[113]**, **[114]**, **[115]**

Omicron-era Long COVID epidemiology with vaccination-stratified estimates — The majority of the literature predates or conflates Omicron and pre-Omicron eras. Studies specifically addressing Omicron-era Long COVID in vaccinated populations show substantially lower risk (3.50 per 100 person-years), but few studies provide vaccination-by-variant interaction analyses, preventing precise quantification of residual Long COVID burden under current epidemiological conditions. See: **[6]**, **[12]**, **[116]**, **[117]**, **[118]**

Validated, harmonized Long COVID case definition

accepted across studies — The 39 distinct duration thresholds and multiple case ascertainment strategies identified in this corpus render cross-study synthesis of prevalence estimates nearly impossible. A single validated case definition with objective clinical anchoring is absent from all 470 deeply analyzed studies. See: [\[80\]](#), [\[81\]](#), [\[119\]](#), [\[84\]](#), [\[120\]](#)

Biomarker-based prospective studies with pre-infection baselines — No study in this corpus assessed objective biomarkers (inflammatory, metabolic, autonomic, imaging) alongside pre-infection baselines and confirmed Long COVID endpoints in a general population. The absence of baseline data in 58.3% of deep papers is the primary barrier to causal inference about individual-level risk factors. See: [\[121\]](#), [\[122\]](#), [\[123\]](#), [\[52\]](#), [\[124\]](#)

Long COVID in low- and middle-income countries with adequate epidemiological design — Sub-Saharan African and South Asian Long COVID literature is dominated by single-centre hospitalized cohorts and small cross-sectional studies with extreme heterogeneity ($I^2=99\%$); LMIC-specific prevalence estimates ranging from 2% to 66.7% are uninterpretable. Community-based prospective cohorts with uninfected controls are essentially absent. See: [\[2\]](#), [\[93\]](#), [\[125\]](#), [\[126\]](#), [\[90\]](#)

Mechanistic studies linking specific biological pathways to clinically defined Long COVID phenotypes — Phenotype mapping assigned 'unspecified' to 58.5% of deeply analyzed studies and 'mixed' to 23.2%, reflecting that biological mechanisms are proposed narratively but not empirically tested with adequate experimental designs. No study in this corpus delivered prospective mechanistic biomarker data alongside validated Long COVID phenotype classification in a sufficiently powered general population sample. See: [\[127\]](#), [\[122\]](#), [\[27\]](#), [\[63\]](#), [\[128\]](#)

Pediatric Long COVID with pre-infection cognitive and functional baselines — Pediatric studies are particularly affected by the absence of pre-COVID baselines, with symptom prevalence in COVID-positive children often only marginally higher than in

COVID-negative children (e.g., 60.9% vs 43.2% with ≥ 1 symptom at 6 months, CITE:PMC10086284), and pandemic indirect effects on mental health confounding all comparisons. The RECOVER-Pediatrics infrastructure is designed to address this but has not yet yielded definitive results. See: [\[129\]](#), [\[130\]](#), [\[131\]](#), [\[132\]](#), [\[133\]](#)

Long-term (>2 years) natural history in community-based non-hospitalized populations — Most follow-up studies extend to 6–12 months; 2-year data are available primarily from hospitalized VA cohorts [\[11\]](#) and selected registries, with limited representation of non-hospitalized mild infection cases who constitute the vast majority of Long COVID burden. Trajectory analyses identifying persistent versus resolving versus worsening subgroups are limited to a few prospective cohorts. See: [\[11\]](#), [\[10\]](#), [\[134\]](#), [\[135\]](#), [\[136\]](#)

10. Research Recommendations

- Establish and mandate a single internationally harmonized Long COVID case definition with a minimum 12-week symptom duration threshold, objective functional anchoring (e.g., validated instrument such as the RECOVER LCRI), and confirmed SARS-CoV-2 infection, for use in all future prevalence and risk factor studies.
- Conduct adequately powered randomized controlled trials for candidate pharmacological interventions (antivirals, anticoagulants, immunomodulators, SSRIs) with pre-specified Long COVID endpoints defined by the harmonized case definition, standardized follow-up to at least 12 months, and pre-stratification by vaccination status and variant era.
- Fund and implement prospective cohort studies in LMIC settings with community-based recruitment, uninfected concurrent controls, pre-infection biospecimen collection, and

standardized Long COVID outcome ascertainment to address the critical evidence gap in non-high-income-country populations.

- Require mandatory pre-infection baseline symptom, cognitive, and functional assessments in all future prospective Long COVID studies to enable attribution of post-infection changes specifically to SARS-CoV-2 infection.
- Develop and validate objective biomarker panels (autonomic, inflammatory, metabolic, neuroimaging) capable of distinguishing Long COVID phenotypes in clinical settings, using longitudinal studies with matched uninfected controls and pre-infection baselines.
- Conduct variant-stratified and vaccination-stratified meta-analyses of Long COVID prevalence using individual participant data from existing cohorts to produce reliable era-specific and protection-specific prevalence estimates.
- Mandate vaccination status, number of doses, and variant era data as required covariates in all Long COVID observational studies, and provide data sharing through federated platforms (e.g., N3C, PCORnet) to enable pooled analyses.
- Invest in long-term follow-up (3–5 years) of established prospective cohorts (RECOVER-Adult, Long-CISS, EPILOC) with particular attention to recovery trajectories in community-based non-hospitalized individuals to characterize natural history.
- Develop and externally validate computable phenotypes for Long COVID that are tested against clinical adjudication as the reference standard, not against ICD codes, and that account for differential healthcare utilization across demographic groups.
- Prioritize research on pediatric Long COVID using pre-infection neuropsychological and functional baselines, pandemic-matched control groups, and longitudinal follow-up

through adolescence to disentangle SARS-CoV-2-specific effects from pandemic indirect effects.

Limitations of This Analysis

- **Sample size:** 4666 papers; the full Long COVID literature exceeds 8,900 publications.
- **Deep-extraction subset:** 470 papers received full methodological appraisal with provenance.
- **Full-text coverage:** only PMC Open Access papers have full text. The rest were analyzed at the abstract level — this is captured in the calibration layer (lower extraction_confidence and downgraded GRADE).
- **Date / language skew:** 2025-2026 bias from PubMed ordering; English-language only.
- **LLM extraction risk:** every structured field is LLM-generated. The provenance layer enables literal-quote verification; the calibration layer downgrades when the LLM is uncertain. Spot-check the provenance entries to validate.
- **Not a systematic review:** no PRISMA flow, no quality scoring against external checklists beyond integrated NOS/ GRADE per paper.

References

1. Unresolved reference: 10.1186/s12916-024-03476-6
2. Peter S. Nyasulu, Jacques L. Tamuzi, Rajiv T. Erasmus. Burden, causation, and particularities of Long COVID in African populations: A rapid systematic review Long COVID and chronic fatigue syndrome: a survey of elderly female

survivors in Egypt Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Attributes and predictors of long COVID Long COVID, a comprehensive systematic scoping review Deleterious outcomes in Long-Hauler COVID-19: the effects of SARS-CoV-2 on the CNS in chronic COVID syndrome Post-COVID-19 condition 3 months after hospitalisation with SARS-CoV-2 in South Africa: a prospective cohort study Postacute sequelae and adaptive immune responses in people with HIV recovering from SARS-COV-2 infection The PRISMA 2020 statement: an updated guideline for reporting systematic reviews Persistent clotting protein pathology in long COVID/post-acute sequelae of COVID-19 (PASC) is accompanied by increased levels of antiplasmin Proteomics of fibrin amyloid microclots in long COVID/post-acute sequelae of COVID-19 (PASC) shows many entrapped pro-inflammatory molecules that may also contribute to a failed fibrinolytic system Prevalence, characteristics and risk factors in a Moroccan cohort of Long-Covid-19 Detection and characterization of fibrin/amyloid microclots in patients with post-acute sequelae of Covid-19 Increased levels of inflammatory and endothelial biomarkers in blood of long COVID patients point to thrombotic endothelialitis Post-discharge symptoms among hospitalized COVID-19 patients in Nigeria: a single-center study 'Long COVID': persistent COVID-19 symptoms in survivors managed in Lagos State, Nigeria Prevalence of symptoms, comorbidities, fibrin amyloid microclots and platelet pathology in individuals with long COVID/post-acute sequelae of COVID-19 (PASC) Two-month follow-up of persons with SARS-CoV-2 infection-Zambia, September 2020: a cohort study Two months follow-up of patients with non-critical COVID-19 in Cape Town, South Africa Determinants of persistent post-COVID-19 symptoms: value of a novel

COVID-19 symptom score Determinants of COVID-19-related length of hospital stays and long COVID in Ghana: a cross-sectional analysis Factors associated with severe COVID-19 and post-acute COVID-19 syndrome in a cohort of people living with HIV on antiretroviral treatment and with undetectable HIV RNA Global prevalence of post-coronavirus disease 2019 (COVID-19) condition or long COVID: a meta-analysis and systematic review Undiagnosed comorbidities among individuals hospitalised with COVID-19 in South African public hospitals Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Long COVID after breakthrough SARS-CoV-2 infection Survival after severe COVID-19: long-term outcomes of patients admitted to an intensive care unit COVID-19 and cardiovascular disease: from basic mechanisms to clinical perspectives Hypercoagulation in COPD: the clot thickens Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: a systematic review IJID Regions. 2023. [PMC10477483](https://pubmed.ncbi.nlm.nih.gov/40147483/)

3. Zijun Xu, Wenyue Wang, Dexing Zhang, King Wa Tam, Yiqi Li, Dicken Cheong Chun Chan, Zuyao Yang, Samuel Yeung Shan Wong, et al.. Excess risks of long COVID symptoms compared with identical symptoms in the general population: A systematic review and meta-analysis of studies with control groups Understanding the burden of post-covid-19 condition. Disability-adjusted life years (DALYs) due to the direct health impact of COVID-19 in India, 2020. Persistence of long COVID symptoms in COVID-19 survivors worldwide and its potential pathogenesis-a systematic review and meta-analysis. Characteristics and predictors of acute and chronic post-COVID syndrome: A systematic review and meta-analysis. Prevalence of post-acute COVID-19 syndrome symptoms at

different follow-up periods: a systematic review and meta-analysis. Long-Term Sequelae of COVID-19: A Systematic Review and Meta-Analysis of One-Year Follow-Up Studies on Post-COVID Symptoms. Persistent symptoms following SARS-CoV-2 infection amongst children and young people: a meta-analysis of controlled and uncontrolled studies. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. Rapid reviews may produce different results to systematic reviews: a meta-epidemiological study. Influence of pilot and small trials in meta-analyses of behavioral interventions: a meta-epidemiological study. Development of a definition of postacute sequelae of SARS-CoV-2 infection. Single-port versus multiple-port robot-assisted radical prostatectomy: a systematic review and meta-analysis. Three risk of bias tools lead to opposite conclusions in observational research synthesis. Predicting the extent of heterogeneity in meta-analysis, using empirical data from the Cochrane Database of Systematic Reviews. Assessing heterogeneity in meta-analysis: Q statistic or I² index? Bias in meta-analysis detected by a simple, graphical test. A nationwide questionnaire study of post-acute symptoms and health problems after SARS-CoV-2 infection in Denmark. Post-acute symptoms four months after SARS-CoV-2 infection during the Omicron period: a nationwide Danish questionnaire study. One-Year Temporal Changes in Long COVID Prevalence and Characteristics: A Systematic Review and Meta-Analysis. Pooled prevalence of long COVID-19 symptoms at 12 months and above follow-up period: a systematic review and meta-analysis. The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis. Recovery and symptom trajectories up to two years after SARS-CoV-2 infection:

population based, longitudinal cohort study. Risk Factors Associated With Post–COVID-19 Condition: A Systematic Review and Meta-analysis. Demographic And Clinical Factors Associated With Long COVID. Factors Associated With Long COVID Symptoms in an Online Cohort Study. Symptoms and risk factors for long COVID in non-hospitalized adults. Understanding Kids and COVID. Long COVID: major findings, mechanisms and recommendations. Viral persistence, reactivation, and mechanisms of long COVID. Long COVID: mechanisms, risk factors and recovery. Pathophysiology and mechanism of long COVID: a comprehensive review. Awareness and care seeking for long COVID symptoms among Coronavirus disease survivors in Bahir Dar City, Northwest Ethiopia: phenomenological study. Awareness and perceptions of Long COVID among people in the REACT programme: Early insights from a pilot interview study. Representation of long COVID syndrome in the awareness of the population is revealed by Google Trends analysis. Long COVID: major findings, mechanisms and recommendations. COVID-19 Vaccination for the Prevention and Treatment of Long COVID: A Systematic Review and Meta-analysis. Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: A systematic review. Effect of covid-19 vaccination on long covid: systematic review. Publication and related bias in meta-analysis: Power of statistical tests and prevalence in the literature. Journal of Global Health. 2024. [PMC11317913](#)

4. Milena Adina Man, Daniela Rosca, Felix Bratosin, Ovidiu Fira-Mladinescu, Adrian Cosmin Ilie, Sonia-Roxana Burtic, Ariadna Petronela Fildan, Camelia Melania Fizedean, Adelina Maria Jianu, Rodica Anamaria Negrean, Monica Steluta Marc, et al.. Impact of Pre-Infection COVID-19 Vaccination on the Incidence and Severity of Post-COVID Syndrome: A

Systematic Review and Meta-Analysis The coronavirus global pandemic and its impacts on society Impact of the COVID-19 Pandemic on the Elective Surgery for Colorectal Cancer: Lessons to Be Learned The impact of COVID-19 on health and health systems The Epidemiology of Malignant Melanoma during the First Two Years of the COVID-19 Pandemic: A Systematic Review COVID-19 Clinical Features and Outcomes in Elderly Patients during Six Pandemic Waves The World Health Organization COVID-19 surveillance database A comprehensive review on COVID-19 vaccines: Development, effectiveness, adverse effects, distribution and challenges Safety & effectiveness of COVID-19 vaccines: A narrative review The effectiveness of COVID-19 vaccines in reducing the incidence, hospitalization, and mortality from COVID-19: A systematic review and meta-analysis The efficacy and effectiveness of the COVID-19 vaccines in reducing infection, severity, hospitalization, and mortality: A systematic review Real-world effectiveness of COVID-19 vaccines: A literature review and meta-analysis Effect of Vaccination against SARS-CoV-2 on Long COVID-19: A Narrative Review Long COVID: Where Are We in 2023? Post-COVID-19 Syndrome: Incidence, Risk Factor, and the Most Common Persisting Symptoms Covid-19 vaccine effectiveness against post-covid-19 condition among 589 722 individuals in Sweden: Population based cohort study Effect of covid-19 vaccination on long covid: Systematic review The PRISMA 2020 statement: An updated guideline for reporting systematic reviews PROSPERO: An International Register of Systematic Review Protocols Identification and diagnosis of long COVID-19: A scoping review Association of SARS-CoV-2 infection and persistence with long COVID Long COVID after breakthrough SARS-CoV-2 infection Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID

Symptom Study app: A prospective, community-based, nested, case-control study Risk of Long COVID in People Infected With Severe Acute Respiratory Syndrome Coronavirus 2 After 2 Doses of a Coronavirus Disease 2019 Vaccine: Community-Based, Matched Cohort Study Association Between BNT162b2 Vaccination and Long COVID After Infections Not Requiring Hospitalization in Health Care Workers Long COVID risk and pre-COVID vaccination in an EHR-based cohort study from the RECOVER program Post COVID-19 condition after Wildtype, Delta, and Omicron SARS-CoV-2 infection and prior vaccination: Pooled analysis of two population-based cohorts The presence of symptoms within 6 months after COVID-19: A single-center longitudinal study Rates and Factors Associated With Documentation of Diagnostic Codes for Long COVID in the National Veterans Affairs Health Care System Differences in Clinical Presentation With Long COVID After Community and Hospital Infection and Associations With All-Cause Mortality: English Sentinel Network Database Study Presence of symptoms 6 weeks after COVID-19 among vaccinated and unvaccinated US healthcare personnel: A prospective cohort study Six-month sequelae of post-vaccination SARS-CoV-2 infection: A retrospective cohort study of 10,024 breakthrough infections Prevalence and Severity of Symptoms 3 Months After Infection With SARS-CoV-2 Compared to Test-Negative and Population Controls in the Netherlands The Protective Effect of Coronavirus Disease 2019 (COVID-19) Vaccination on Postacute Sequelae of COVID-19: A Multicenter Study From a Large National Health Research Network Attributes and Predictors of Long COVID Risk Factors Associated with Long COVID Syndrome: A Retrospective Study Long-COVID Clinical Features and Risk Factors: A Retrospective Analysis of Patients from the STOP-COVID Registry of the PoLoCOV

Study Long-Term Health Symptoms and Sequelae Following SARS-CoV-2 Infection: An Evidence Map Long-term post-acute sequelae of COVID-19 infection: A retrospective, multi-database cohort study in Hong Kong and the UK Developing a Long COVID Phenotype for Postacute COVID-19 in a National Primary Care Sentinel Cohort: Observational Retrospective Database Analysis One-year follow-up CT findings in COVID-19 patients: A systematic review and meta-analysis Safety and efficacy of the BNT162b2 mRNA COVID-19 vaccine Efficacy, Immunogenicity and Safety of COVID-19 Vaccines: A Systematic Review and Meta-Analysis Long Covid —Mechanisms, Risk Factors, and Management Age, Sex and Previous Comorbidities as Risk Factors Not Associated with SARS-CoV-2 Infection for Long COVID-19: A Systematic Review and Meta-Analysis Efficacy of covid-19 vaccination on the symptoms of patients with long covid: A target trial emulation using data from the compare e-Cohort in France The effect of SARS-CoV-2 vaccination on post-acute sequelae of COVID-19 (PASC): A prospective cohort study Definition of Post-COVID-19 Condition Among Published Research Studies Health outcomes in people 2 years after hospitalisation with COVID-19: A longitudinal cohort study Global prevalence of post Coronavirus Disease 2019 (COVID-19) condition or long COVID: A meta-analysis and systematic review Long COVID: Complications, Underlying Mechanisms, and Treatment Strategies Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection A retrospective analysis of clinically confirmed long post-COVID vaccination syndrome Presence of viral spike protein and vaccinal spike protein in the blood serum of patients with long-COVID syndrome Vaccines. 2024. [PMC10893048](#)

5. M. Daniel Brannock, Robert F. Chew, Alexander J. Preiss, Emily C. Hadley, Signe Redfield, Julie A. McMurry, Peter J.

Leese, Andrew T. Girvin, Miles Crosskey, Andrea G. Zhou, Richard A. Moffitt, Michele Jonsson Funk, Emily R. Pfaff, Melissa A. Haendel, Christopher G. Chute, TilJohanna J.Farrukh M. StürmerLoombaKoraishy, Til Stürmer, Johanna J. Loomba, Farrukh M. Koraishy, JasminLorna E.LeoraStuart DiversThorpeHorwitzKatz, Jasmin Divers, Lorna E. Thorpe, Leora Horwitz, Stuart Katz, et al.. Long COVID risk and pre-COVID vaccination in an EHR-based cohort study from the RECOVER program Long COVID Risk and Pre-COVID Vaccination: An EHR-Based Cohort Study from the RECOVER Program Toward understanding COVID-19 recovery: National Institutes of Health Workshop on postacute COVID-19 Symptoms and functional impairment assessed 8 months after mild COVID-19 among health care workers 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study Case definition working group on post-COVID-19 condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Characterizing long COVID: deep phenotype of a complex condition Identifying who has long COVID in the USA: a machine learning approach using N3C data The National COVID Cohort Collaborative (N3C): rationale, design, infrastructure, and deployment Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine Effectiveness of severe acute respiratory syndrome coronavirus 2 messenger RNA vaccines for preventing coronavirus disease 2019 hospitalizations in the United States The protective effect of coronavirus disease 2019 (COVID-19) vaccination on postacute sequelae of COVID-19: a multicenter study from a large National Health Research Network Long COVID after breakthrough SARS-CoV-2 infection Risk of long COVID in people infected with severe acute respiratory syndrome

coronavirus 2 after 2 doses of a coronavirus disease 2019 vaccine: community-based, matched cohort study Six-month sequelae of post-vaccination SARS-CoV-2 infection: a retrospective cohort study of 10,024 breakthrough infections Characteristics and predictors of Long COVID among diagnosed cases of COVID-19 The effect of SARS-CoV-2 vaccination on post-acute sequelae of COVID-19 (PASC): a prospective cohort study Association of vaccination with the persistence of post-COVID symptoms Controlling for informed presence bias due to the number of health encounters in an electronic health record Clinical severity of, and effectiveness of mRNA vaccines against, covid-19 from omicron, delta, and alpha SARS-CoV-2 variants in the United States: prospective observational study Covid-19 vaccine effectiveness against the Omicron (B.1.1.529) variant COVID-19 vaccine waning and effectiveness and side-effects of boosters: a prospective community study from the ZOE COVID Study Overadjustment bias and unnecessary adjustment in epidemiologic studies Does prevention of risk behaviour in primary care require a gender-specific approach? A cross-sectional study Men and health help-seeking behaviour: literature review Age and gender differences in health service utilization Synergies between centralized and federated approaches to data quality: a report from the national COVID cohort collaborative Measurement error and misclassification in electronic medical records: methods to mitigate bias Advancing the science for active surveillance: rationale and design for the observational medical outcomes partnership PheKB: a catalog and workflow for creating electronic phenotype algorithms for transportability Constructing inverse probability weights for marginal structural models Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Long covid-mechanisms, risk factors, and management Marginal

structural models and causal inference in epidemiology
Variance estimation when using inverse probability of
treatment weighting (IPTW) with survival analysis
Influential cases in multilevel modeling: a methodological comment
Reassessing Schoenfeld residual tests of proportional hazards
in political science event history analyses
The strengthening the reporting of observational studies in epidemiology
(STROBE) statement: guidelines for reporting observational
studies Nature Communications. 2023. [PMC10201472](#)

6. Yan Xie, Taeyoung Choi, Ziyad Al-Aly. Postacute Sequelae of SARS-CoV-2 Infection in the Pre-Delta, Delta, and Omicron Eras Viral Variants, Vaccinations, and Long Covid - New Insights. High-dimensional characterization of post-acute sequelae of COVID-19. Long COVID: major findings, mechanisms and recommendations. Long COVID after breakthrough SARS-CoV-2 infection. Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status. Solving the puzzle of long Covid. Postacute sequelae of COVID-19 at 2 years. Long-term neurologic outcomes of COVID-19. Long-term cardiovascular outcomes of COVID-19. Risks and burdens of incident diabetes in long COVID: a cohort study. Risks of mental health outcomes in people with covid-19: cohort study. Comparative evaluation of clinical manifestations and risk of death in patients admitted to hospital with covid-19 and seasonal influenza: cohort study. Long-term outcomes following hospital admission for COVID-19 versus seasonal influenza: a cohort study. Association of treatment with nirmatrelvir and the risk of post-COVID-19 condition. Molnupiravir and risk of post-acute sequelae of Covid-19: cohort study. Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021. Acute and

postacute sequelae associated with SARS-CoV-2 reinfection. Kidney outcomes in long COVID. Multiple imputation of discrete and continuous data by fully conditional specification. Propensity score weighting for causal inference with multiple treatments Negative controls: a tool for detecting confounding and bias in observational studies. Unexplained post-acute infection syndromes. The immunology of long COVID. Long COVID and significant activity limitation among adults, by age — United States, June 1–13, 2022, to June 7–19, 2023. Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2. The New England Journal of Medicine. 2024. [PMC11687648](#)

7. Cristina Honorato-Cia, Elena Cacho-Asenjo, Antonio Martinez-Simon, Irene Aquerreta, Jorge M. Núñez-Córdoba. Long COVID Research, 2020–2024: A PubMed-Based Bibliometric Analysis WHO COVID-19 Dashboard Long COVID: Major findings, mechanisms and recommendations Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Characterizing Long COVID in Children and Adolescents Long Covid Defined Impact of long COVID on health-related quality-of-life: An OpenSAFELY population cohort study using patient-reported outcome measures (OpenPROMPT) Healthcare utilisation in people with long COVID: An OpenSAFELY cohort study Solving the puzzle of Long Covid Long COVID: A clinical update Bibliometric analysis of global scientific research on COVID-19 Bibliometric Analysis of Early COVID-19 Research: The Top 50 Cited Papers A bibliometric analysis using VOSviewer of publications on COVID-19 COVID-19 Pandemic Related Research in Africa: Bibliometric Analysis of Scholarly Output, Collaborations and Scientific Leadership A Systematic Review and Bibliometric Analysis of the Scientific Literature on the Early Phase of COVID-19 in Italy A bibliometric analysis of COVID-19 research in Africa

Indian Publications on SARS-CoV-2: A bibliometric study of WHO COVID-19 database Research trends in COVID-19 vaccine: A bibliometric analysis Bibliometric Analysis of International Scientific Production on Pharmacologic Treatments for SARS-CoV-2/COVID-19 During 2020 Global research on RNA vaccines for COVID-19 from 2019 to 2023: A bibliometric analysis Top 100 cited articles on diabetes mellitus and Covid-19: A bibliometric analysis A Bibliometric Analysis of Publications on COVID-19 and Older Adults Bibliometric analysis of global scientific literature on effects of COVID-19 pandemic on mental health Global research trends in COVID-19 with MRI and PET/CT: A scoping review with bibliometric and network analyses COVID-19 Publications in Family Medicine Journals in 2020: A PubMed-Based Bibliometric Analysis COVID-19 and urology: A bibliometric analysis of the literature A Global Overview of COVID-19 Research in the Pediatric Field: Bibliometric Review A bibliometric analysis of rheumatology and COVID-19 researches A bibliometric analysis of COVID-19 publications in neurology by using the visual mapping method Global Research Trends in Pediatric COVID-19: A Bibliometric Analysis COVID-19 publications in anaesthesiology journals: A bibliometric analysis COVID-19 Publications in Allergy Journals, 2020-2021: A Bibliometric Analysis Bibliometric evaluation of 2020-2022 publications on COVID-19-related cardiovascular disease Publications on COVID-19 in radiology journals in 2020 and 2021: Bibliometric citation and co-citation network analysis Global research trends in the COVID-19 and digestive disease: A review of visualization and bibliometric study Palliative care and COVID-19: A bibliometric analysis Global Trends and Research Hotspots in Long COVID: A Bibliometric Analysis The long COVID research literature A Bibliometric Analysis of Post-COVID-19

Syndrome Visual analysis of hotspots and trends in long COVID research based on bibliometric Scars of COVID-19: A bibliometric analysis of post-COVID-19 fibrosis Exploring the research landscape of COVID-19-induced olfactory dysfunction: A bibliometric study Complementary and Alternative Medicine for Long COVID: Scoping Review and Bibliometric Analysis PubMed Update: Clinical Queries Usability Study and Interface Updates Long COVID or Post-acute Sequelae of COVID-19 (PASC): An Overview of Biological Factors That May Contribute to Persistent Symptoms Long COVID Definitions and Models of Care: A Scoping Review Long COVID: Clinical Findings, Pathology, and Endothelial Molecular Mechanisms Pathogenic mechanisms of post-acute sequelae of SARS-CoV-2 infection (PASC) The immunology of long COVID SARS-CoV-2 reservoir in post-acute sequelae of COVID-19 (PASC) Mechanisms of long COVID and the path toward therapeutics Healthcare. 2025. [PMC11816931](#)

8. Felicia Ceban, Dana Kulzhabayeva, Nelson B. Rodrigues, Joshua D. Di Vincenzo, Hartej Gill, Mehala Subramaniapillai, Leanna M.W. Lui, Bing Cao, Rodrigo B. Mansur, Roger C. Ho, Matthew J. Burke, Taeho Greg Rhee, Joshua D. Rosenblat, Roger S. McIntyre, et al.. COVID-19 vaccination for the prevention and treatment of long COVID: A systematic review and meta-analysis Corrigendum to “COVID-19 vaccination for the prevention and treatment of long COVID: A systematic review and meta-analysis” [Brain Behav. Immun. 111 (2023) 211-229] Long Covid after Breakthrough COVID-19: the post-acute sequelae of breakthrough COVID-19 Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study Prevalence, characteristics, and predictors of Long COVID among

diagnosed cases of COVID-19 Symptoms After COVID-19 Vaccination in Patients With Persistent Symptoms After Acute Infection: A Case Series Systems vaccinology of the BNT162b2 mRNA vaccine in humans Changes in the trajectory of Long Covid symptoms following COVID-19 vaccination: community-based cohort study Covid-19 Breakthrough Infections in Vaccinated Health Care Workers Fatigue and cognitive impairment in Post-COVID-19 Syndrome: A systematic review and meta-analysis Registered clinical trials investigating treatment of long COVID: a scoping review and recommendations for research Virological and serological kinetics of SARS-CoV-2 Delta variant vaccine breakthrough infections: a multicentre cohort study Long covid—mechanisms, risk factors, and management Are mRNA covid 19 vaccines safe in long covid patients? A health care workers perspective COVID-19 vaccine breakthrough infections Trust in a COVID-19 vaccine in the U.S.: A social-ecological perspective Comparative systematic review and meta-analysis of reactogenicity, immunogenicity and efficacy of vaccines against SARS-CoV-2 Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement Onset and frequency of depression in post-COVID-19 syndrome: A systematic review SARS-CoV-2 vaccine breakthrough infections with the alpha variant are asymptomatic or mildly symptomatic among health care workers Effect of SARS-CoV-2 Vaccination on Symptoms from Post-Acute Sequelae of COVID-19: Results from the Nationwide VAXILONG Study WHO Clinical Case Definition Working Group on Post-COVID-19 Condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses The Impact of COVID Vaccination on Symptoms of

Long COVID. An International Survey of People with Lived Experience of Long COVID Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group Multiple early factors anticipate post-acute COVID-19 sequelae Association between immune dysfunction and COVID-19 breakthrough infection after SARS-CoV-2 vaccination in the US Long-Term Symptoms Among Adults Tested for SARS-CoV-2 — United States, January 2020–April 2021 Immune Response to Vaccination in Adults with Mental Disorders: A Systematic Review Brain, Behavior, and Immunity. 2023. [PMC10067136](#)

9. Alexandre R. Marra, Takaaki Kobayashi, Gustavo Yano Callado, Isabele Pardo, Maria Celidonio Gutfreund, Mariana Kim Hsieh, Vivian Lin, Mohammed Alsuhaibani, Shinya Hasegawa, Joseph Tholany, Eli N. Perencevich, Jorge L. Salinas, Michael B. Edmond, Luiz Vicente Rizzo, et al.. The effectiveness of COVID-19 vaccine in the prevention of post-COVID conditions: a systematic literature review and meta-analysis of the latest research Efficacy, immunogenicity and safety of COVID-19 vaccines: a systematic review and meta-analysis Real-world effectiveness of COVID-19 vaccines: a literature review and meta-analysis Association between mRNA vaccination and COVID-19 hospitalization and disease severity The efficacy and effectiveness of the COVID-19 vaccines in reducing infection, severity, hospitalization, and mortality: a systematic review Current evidence on efficacy of COVID-19 booster dose vaccination against the Omicron variant: a systematic review Efficacy of a fourth dose of Covid-19 mRNA vaccine against Omicron The effectiveness of coronavirus disease 2019 (COVID-19) vaccine in the prevention of post-COVID-19 conditions: a systematic literature review and meta-analysis Symptoms after

COVID-19 vaccination in patients with persistent symptoms after acute infection: a case series Do vaccines protect from long COVID? Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group The presence of symptoms within 6 months after COVID-19: a single-center longitudinal study COVID-19 long-term sequelae: Omicron versus Alpha and Delta variants Prevalence and severity of symptoms 3 months after infection with SARS-CoV-2 compared to test-negative and population controls in the Netherlands Post-acute sequelae after SARS-CoV-2 infection by viral variant and vaccination status: a multicenter cross-sectional study The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews Accuracy of administrative code data for the surveillance of healthcare-associated infections: a systematic review and meta-analysis Conducting meta-analyses in R with the metafor package Neuropsychiatric symptoms in post-COVID-19 long haulers Long COVID after breakthrough SARS-CoV-2 infection Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Characteristics and predictors of Long COVID among diagnosed cases of COVID-19 Risk of long COVID in people infected with severe acute respiratory syndrome Coronavirus 2 after 2 doses of a coronavirus disease 2019 vaccine: community-based, matched cohort

study Association between BNT162b2 vaccination and long COVID after infections not requiring hospitalization in health care workers Post COVID-19 condition after Wildtype, Delta, and Omicron SARS-CoV-2 infection and prior vaccination: pooled analysis of two population-based cohorts Post-acute symptoms 3-15 months after COVID-19 among unvaccinated and vaccinated individuals with a breakthrough infection Prevalence, characteristics and risk factors in a Moroccan cohort of Long-Covid-19 16 months follow up of patients' behavior and mild COVID-19 patterns in a large cohort of cancer patients during the pandemic Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study Rates and factors associated with documentation of diagnostic codes for long COVID in the National Veterans Affairs Health Care System A cohort study of post-COVID-19 condition across the Beta, Delta, and Omicron waves in South Africa: 6-month follow-up of hospitalized and nonhospitalized participants Association between BNT162b2 vaccination and reported incidence of post-COVID-19 symptoms: cross-sectional study 2020-21, Israel Risk factors for long coronavirus disease 2019 (long COVID) among healthcare personnel, Brazil, 2020-2022 Differences in clinical presentation with long COVID after community and hospital infection and associations with all-cause mortality: English sentinel network database study Presence of symptoms 6 weeks after COVID-19 among vaccinated and unvaccinated US healthcare personnel: a prospective cohort study One-year persistent symptoms and functional impairment in SARS-CoV-2 positive and negative individuals Prevalence and correlates of long COVID symptoms among US adults Vaccination against SARS-CoV-2 protects from morbidity, mortality and sequelae from COVID19 in patients with cancer Persistent COVID-19

symptoms at 6 months after onset and the role of vaccination before or after SARS-CoV-2 infection Burnout and long COVID among the UK nephrology workforce: results from a national survey investigating the impact of COVID-19 on working lives A comprehensive assessment of self-reported post COVID-19 symptoms among beneficiaries of hospital employee scheme at a tertiary healthcare institution in Northern India Real-world effectiveness of COVID-19 vaccines and anti-SARS-CoV-2 monoclonal antibodies against postacute sequelae of SARS-CoV-2: analysis of a COVID-19 observational registry for a diverse US metropolitan population Six-month sequelae of post-vaccination SARS-CoV-2 infection: a retrospective cohort study of 10,024 breakthrough infections Development of a definition of postacute sequelae of SARS-CoV-2 infection The protective effect of coronavirus disease 2019 (COVID-19) vaccination on postacute sequelae of COVID-19: a multicenter study from a large national health research network Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review Prevalence and correlates of long COVID symptoms among US adults Associated-onset symptoms and post-COVID-19 symptoms in hospitalized COVID-19 survivors infected with Wuhan, Alpha or Delta SARS-CoV-2 variant Definition of post-COVID-19 condition among published research studies Global prevalence of post Coronavirus Disease 2019 (COVID-19) condition or long COVID: a meta-analysis and systematic review Antimicrobial Stewardship & Healthcare Epidemiology : ASHE. 2023. [PMC10644173](#)

10. Miao Cai, Yan Xie, Eric J. Topol, Ziyad Al-Aly. Three-year outcomes of post-acute sequelae of COVID-19 High-dimensional characterization of post-acute sequelae of

COVID-19 Solving the puzzle of Long Covid Long COVID: major findings, mechanisms and recommendations Long-term cardiovascular outcomes of COVID-19 Risks of mental health outcomes in people with covid-19: cohort study Risks and burdens of incident diabetes in long COVID: a cohort study Long COVID after breakthrough SARS-CoV-2 infection Long-term neurologic outcomes of COVID-19 Kidney outcomes in long COVID Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Acute and postacute sequelae associated with SARS-CoV-2 reinfection Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Long-term gastrointestinal outcomes of COVID-19 Incidence, co-occurrence, and evolution of long-COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19 Neurological and psychiatric risk trajectories after SARS-CoV-2 infection: an analysis of 2-year retrospective cohort studies including 1 284 437 patients Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study Two-year health outcomes in hospitalized COVID-19 survivors in China Symptom burden and immune dynamics 6 to 18 months following mild severe acute respiratory syndrome coronavirus 2 infection (SARS-CoV-2): a case-control study Healthy outcomes of patients with COVID-19 two years after the infection: a prospective cohort study Postacute sequelae of COVID-19 at 2 years 3-year outcomes of discharged survivors of COVID-19 following the SARS-CoV-2 omicron (B.1.1.529) wave in 2022 in China: a longitudinal cohort study Unexplained post-acute infection syndromes Longitudinal analysis reveals high prevalence of Epstein-Barr virus associated with multiple sclerosis Exome-wide association study to identify rare variants influencing COVID-19 outcomes: results from the Host Genetics Initiative

Humans with inherited MyD88 and IRAK-4 deficiencies are predisposed to hypoxemic COVID-19 pneumonia Genetically determined thymic function affects strength and duration of immune response in COVID patients with pneumonia Genetic regulation of OAS1 nonsense-mediated decay underlies association with COVID-19 hospitalization in patients of European and African ancestries A second update on mapping the human genetic architecture of COVID-19 SARS-CoV-2 infection and persistence in the human body and brain at autopsy SARS-CoV-2 infection triggers pro-atherogenic inflammatory responses in human coronary vessels Ambient fine particulate matter air pollution and the risk of hospitalization among COVID-19 positive individuals: cohort study Temporal trends of COVID-19 mortality and hospitalisation rates: an observational cohort study from the US Department of Veterans Affairs Temporal trends in incidence rates of lower extremity amputation and associated risk factors among patients using veterans health administration services from 2008 to 2018 Acute kidney injury in a national cohort of hospitalized US veterans with COVID-19 Molnupiravir and risk of post-acute sequelae of covid-19: cohort study High-dimensional propensity score adjustment in studies of treatment effects using health care claims data A proportional hazards model for the subdistribution of a competing risk Inverse probability-of-censoring weights for the correction of time-varying noncompliance in the effect of randomized highly active antiretroviral therapy on incident AIDS or death Negative controls: a tool for detecting confounding and bias in observational studies Nature Medicine. 2024. [PMC11186764](#)

11. Benjamin Bowe, Yan Xie, Ziyad Al-Aly. Postacute sequelae of COVID-19 at 2 years Long COVID: major findings, mechanisms and recommendations High-dimensional

characterization of post-acute sequelae of COVID-19 Long-term cardiovascular outcomes of COVID-19 Risks of mental health outcomes in people with COVID-19 Risks and burdens of incident diabetes in long COVID: a cohort study Long COVID after breakthrough SARS-CoV-2 infection Long-term neurologic outcomes of COVID-19 Kidney outcomes in long COVID Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Acute and postacute sequelae associated with SARS-CoV-2 reinfection Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Long-term gastrointestinal outcomes of COVID-19 Neurological and psychiatric risk trajectories after SARS-CoV-2 infection: an analysis of 2-year retrospective cohort studies including 1 284 437 patients Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study Two-year health outcomes in hospitalized COVID-19 survivors in China Symptom burden and immune dynamics 6 to 18 months following mild severe acute respiratory syndrome coronavirus 2 infection (SARS-CoV-2): a case-control study Healthy outcomes of patients with COVID-19 two years after the infection: a prospective cohort study Post-hospital syndrome—an acquired, transient condition of generalized risk A national strategy for the ‘new normal’ of life with COVID Long COVID: long-term health outcomes and implications for policy and research Diabetes after SARS-CoV-2 infection Prevention of long COVID: progress and challenges Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019 Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Making neighborhood-

disadvantage metrics accessible - the Neighborhood Atlas
Acute and postacute sequelae associated with SARS-CoV-2
reinfection Comparative evaluation of clinical manifestations
and risk of death in patients admitted to hospital with
COVID-19 and seasonal influenza: cohort study Risk of
persistent and new clinical sequelae among adults aged 65
years and older during the post-acute phase of SARS-CoV-2
infection: retrospective cohort study Risk of clinical sequelae
after the acute phase of SARS-CoV-2 infection: retrospective
cohort study 6-month neurological and psychiatric outcomes
in 236 379 survivors of COVID-19: a retrospective cohort
study using electronic health records Acute kidney injury in a
national cohort of hospitalized US veterans with COVID-19
Temporal trends of COVID-19 mortality and hospitalisation
rates: an observational cohort study from the US Department
of Veterans Affairs Risk of death in patients hospitalized for
COVID-19 vs seasonal influenza in fall-winter 2022-2023
Incidence, co-occurrence, and evolution of long-COVID
features: a 6-month retrospective cohort study of 273,618
survivors of COVID-19 Nirmatrelvir and the risk of post-acute
sequelae of COVID-19 Molnupiravir and risk of hospital
admission or death in adults with Covid-19: emulation of a
randomized target trial using electronic health records
Nirmatrelvir and risk of hospital admission or death in adults
with Covid-19: emulation of a randomized target trial using
electronic health records Molnupiravir and risk of post-acute
sequelae of covid-19: cohort study New creatinine- and
cystatin C-based equations to estimate GFR without race
High-dimensional propensity score adjustment in studies of
treatment effects using health care claims data Short term
exposure to fine particulate matter and hospital admission
risks and costs in the Medicare population: time stratified,
case crossover study Best definitions of multimorbidity to

identify patients with high health care resource utilization
Comparison of critical drug-drug interaction listings: the
Department of Veterans Affairs medical system and standard
reference compendia Polypharmacy, drug-drug interactions,
and potentially inappropriate medications in older adults with
human immunodeficiency virus infection An introduction to
propensity score methods for reducing the effects of
confounding in observational studies Balance diagnostics for
comparing the distribution of baseline covariates between
treatment groups in propensity-score matched samples
Global, regional, and national disability-adjusted life-years
(DALYs) for 359 diseases and injuries and healthy life
expectancy (HALE) for 195 countries and territories,
1990–2017: a systematic analysis for the Global Burden of
Disease Study 2017 Estimated global proportions of
individuals with persistent fatigue, cognitive, and respiratory
symptom clusters following symptomatic COVID-19 in 2020
and 2021 Global burden of 369 diseases and injuries in 204
countries and territories, 1990–2019: a systematic analysis
for the Global Burden of Disease Study 2019 Consequences of
depletion of susceptibles for hazard ratio estimators based on
propensity scores Trajectory of long COVID symptoms after
COVID-19 vaccination: community based cohort study
Negative controls: a tool for detecting confounding and bias
in observational studies Nature Medicine. 2023.
[PMC10504070](#)

12. Peihong Guo, Jue Tao Lim, Liang En Wee, An Ting Tay, Yichen Zhai, Calvin J. Chiew, Benjamin Ong, David Chien Boon Lye, Kelvin Bryan Tan, et al.. Characterization of post-acute multi-organ sequelae following SARS-CoV-2 Infection in the Delta and Omicron Eras in a highly boosted population Autoimmune sequelae after Delta or Omicron variant SARS-CoV-2 infection in a highly vaccinated cohort Long-term Cardiovascular,

Cerebrovascular, and Other Thrombotic Complications in COVID-19 Survivors: A Retrospective Cohort Study Long-term cardiovascular, cerebrovascular, and thrombotic complications after SARS-CoV-2-Omicron infection: a retrospective cohort study Long-term multi-systemic complications after SARS-CoV-2 Omicron and Delta infection in children: a retrospective cohort study Vaccination and the risk of post-acute sequelae after COVID-19 in the omicron-predominant period Covid-19 vaccine effectiveness against post-covid-19 condition among 589 722 individuals in Sweden: population based cohort study Bivalent boosters and risk of postacute sequelae following vaccine-breakthrough SARS-CoV-2 Omicron infection: a cohort study Nirmatrelvir/ritonavir treatment and risk for postacute sequelae of COVID-19 in older Singaporeans Postacute sequelae of SARS-CoV-2 infection in the pre-Delta, Delta, and Omicron eras Long-term real-world protection afforded by third mRNA doses against symptomatic severe acute respiratory syndrome coronavirus 2 infections, coronavirus disease 19-related emergency attendances and hospitalizations amongst older Singaporeans during an Omicron XBB wave Risks of SARS-CoV-2 JN.1 infection and COVID-19-associated emergency department visits/hospitalizations following updated boosters and prior infection: a population-based cohort study Access to COVID-19 vaccination by socioeconomic status in older Singaporean adults: a population-based cohort study Vaccine effectiveness against Delta, Omicron BA.1, and BA.2 in a highly vaccinated Asian setting: a test-negative design study Singapore's health-care system: key features, challenges, and shifts Long-term neuropsychiatric sequelae of Delta versus Omicron SARS-CoV-2 infection Factors associated with delayed diagnosis of symptomatic adult COVID-19 cases presenting to primary

care: a population-wide study during transition from Delta to Omicron BA.1 in Singapore Oral antiviral utilisation among older adults with COVID-19 in primary care: a population-wide study during successive Omicron waves in Singapore Long covid defined Long-term cardiovascular outcomes of COVID-19 Acute and postacute sequelae associated with SARS-CoV-2 reinfection Charlson comorbidity index: ICD-9 update and ICD-10 translation A proportional hazards model for the subdistribution of a competing risk High-dimensional characterization of post-acute sequelae of COVID-19 Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID symptom study app: a prospective, community-based, nested, case-control study Post-acute sequelae of COVID-19 3 to 12 months after infection: delta vs Omicron Prevalence and risk factors for long COVID among adults in Scotland using electronic health records: a national, retrospective, observational cohort study BMC Global and Public Health. 2026. [PMC12874983](#)

13. Barak Mizrahi, Tamar Sudry, Natalie Flaks-Manov, Yoav Yehezkelli, Nir Kalkstein, Pinchas Akiva, Anat Ekka-Zohar, Shirley Shapiro Ben David, Uri Lerner, Maytal Bivas-Benita, Shira Greenfeld, et al.. Long covid outcomes at one year after mild SARS-CoV-2 infection: nationwide cohort study Characterising long COVID: a living systematic review 1-year outcomes in hospital survivors with COVID-19: a longitudinal cohort study 3-month, 6-month, 9-month, and 12-month respiratory outcomes in patients following COVID-19-related hospitalisation: a prospective study Long COVID in a prospective cohort of home-isolated patients Post-acute effects of SARS-CoV-2 infection in individuals not requiring hospital admission: a Danish population-based cohort study Long-COVID Syndrome? A Study on the Persistence of Neurological, Psychological and Physiological Symptoms Risk

of clinical sequelae after the acute phase of SARS-CoV-2 infection: retrospective cohort study 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records Assessment of the Frequency and Variety of Persistent Symptoms Among Patients With COVID-19: A Systematic Review Post-acute COVID-19 syndrome Post-covid syndrome in individuals admitted to hospital with covid-19: retrospective cohort study Six-month sequelae of post-vaccination SARS-CoV-2 infection: A retrospective cohort study of 10,024 breakthrough infections Post-COVID syndrome symptoms, functional disability, and clinical severity phenotypes in hospitalized and nonhospitalized individuals: A cross-sectional evaluation from a community COVID rehabilitation service Global Prevalence of Post-Coronavirus Disease 2019 (COVID-19) Condition or Long COVID: A Meta-Analysis and Systematic Review Long COVID: A growing problem in need of intervention Long covid-mechanisms, risk factors, and management Attributes and predictors of long COVID Sequelae in Adults at 6 Months After COVID-19 Infection Patient outcomes after hospitalisation with COVID-19 and implications for follow-up: results from a prospective UK cohort Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting Protection of BNT162b2 Vaccine Booster against Covid-19 in Israel Interim findings from first-dose mass COVID-19 vaccination roll-out and COVID-19 hospital

admissions in Scotland: a national prospective cohort study
Effectiveness of the Pfizer-BioNTech and Oxford-AstraZeneca
vaccines on covid-19 related symptoms, hospital admissions,
and mortality in older adults in England: test negative case-
control study Symptoms After COVID-19 Vaccination in
Patients With Persistent Symptoms After Acute Infection: A
Case Series 'Post-COVID-19 syndrome and humoral response
association after one year in vaccinated and unvaccinated
patients': authors' response COVID-19 community care in
Israel-a nationwide cohort study from a large health
maintenance organization Safety of the BNT162b2 mRNA
Covid-19 Vaccine in a Nationwide Setting Urban diversity and
epidemic resilience: The case of the COVID-19 Long COVID
after breakthrough SARS-CoV-2 infection Comparison of
Time-to-First Event and Recurrent-Event Methods in
Randomized Clinical Trials Post-COVID syndrome in non-
hospitalised patients with COVID-19: a longitudinal
prospective cohort study Persistent symptoms 1.5-6 months
after COVID-19 in non-hospitalised subjects: a population-
based cohort study Long COVID symptoms in SARS-CoV-2-
positive adolescents and matched controls
(LongCOVIDKidsDK): a national, cross-sectional study
Changes in the incidence of invasive disease due to
Streptococcus pneumoniae, *Haemophilus influenzae*, and
Neisseria meningitidis during the COVID-19 pandemic in 26
countries and territories in the Invasive Respiratory Infection
Surveillance Initiative: a prospective analysis of surveillance
data Incidence of common infectious diseases in Japan during
the COVID-19 pandemic Prevalence of primary non-alcoholic
fatty liver disease in a population-based study and its
association with biochemical and anthropometric measures
Prognostic non-invasive biomarkers for all-cause mortality in
non-alcoholic fatty liver disease: A systematic review and

meta-analysis Characteristics of SARS-CoV-2 and COVID-19 Severe Acute Respiratory Syndrome Coronavirus 2 Variants of Concern: A Perspective for Emerging More Transmissible and Vaccine-Resistant Strains Health system impacts of SARS-CoV - 2 variants of concern: a rapid review Defining the risk of SARS-CoV-2 variants on immune protection Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 The Impact of COVID Vaccination on Symptoms of Long COVID: An International Survey of People with Lived Experience of Long COVID The BMJ. 2023. [PMC9832503](#)

14. Mar Alvarez-Pedrerol, Sara Polo-Alonso, Rafel Ramos, Ruth Martí-Lluch, Mel·lina Pinsach-Abuin, Irene R Dégano, Roberto Elosua, Isaac Subirana, Álvaro Hernáez, Elisabet Selga, Eulàlia Puigdecenet, Josefina Pruneda-Paz, Clàudia Solà-Richarte, Marta Puigmulé, Alexandra Pérez, Xavier Nogués, Joan Ramon Masclans, Roberto Güerri-Fernández, Héctor Cubero-Gallego, Helena Tizon-Marcos, Beatriz Vaquerizo, Ramon Brugada, Anna Camps-Vilaró, Jaume Marrugat, et al.. Sex Differences in Long COVID Prevalence Over One year After the Acute Phase, and Related Risk Factors. The GINA-COVID Cohort Study Long COVID: major findings, mechanisms and recommendations Long COVID: lights and shadows on the clinical characterization of this emerging pathology WHO Coronavirus (COVID-19) Dashboard The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis Long COVID: a clinical update A clinical case definition of post-COVID-19 condition by a Delphi consensus National Academies of Sciences, Engineering, and Medicine Committee on Examining the Working Definition for Long Covid. Long Covid Defined Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Development of a Definition of Postacute Sequelae of

SARS-CoV-2 Infection The long-term health outcomes, pathophysiological mechanisms and multidisciplinary management of long COVID Long COVID still has no cure - so these patients are turning to research Impact of long COVID on health-related quality-of-life: an OpenSAFELY population cohort study using patient-reported outcome measures (OpenPROMPT) Long COVID basics Long COVID incidence in adults and children between 2020 and 2023: a real-world data study from the RECOVER Initiative Female gender is associated with long COVID syndrome: a prospective cohort study [Coronary risk estimation in Spain using a calibrated Framingham function] Predictors of adverse prognosis in COVID-19: a systematic review and meta-analysis More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Manifestacions persistents de la COVID-19 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study Post-acute COVID-19 syndrome. Incidence and risk factors: a Mediterranean cohort study Biases and limitations in observational studies of Long COVID prevalence and risk factors: a rapid systematic umbrella review Protective effect of COVID-19 vaccination against long COVID syndrome: a systematic review and meta-analysis Sex differences in sequelae from COVID-19 infection and in long COVID syndrome: a review Sex differences in symptomatology and immune profiles of Long COVID Determinants of the onset and prognosis of the post-COVID-19 condition: a 2-year prospective observational cohort study Determinants of post-acute COVID-19 syndrome among hospitalized severe COVID-19 patients: a 2-year follow-up study Prevalence and predictors of Post-Acute COVID-19 Syndrome (PACS) after hospital discharge: a cohort study with 4 months median follow-up Sex-based differences in severity and mortality in COVID-19 Sex differences in the

mortality rate for coronavirus disease 2019 compared to other causes of death Sexual dimorphism in COVID-19: potential clinical and public health implications The X chromosome in immune functions: when a chromosome makes the difference Sex differences in the response to viral infections: TLR8 and TLR9 ligand stimulation induce higher IL10 production in males Influence of X chromosome in sex-biased autoimmune diseases What causes long COVID? Case builds for rogue antibodies Autoantibodies against apolipoprotein A-1 after COVID-19 predict symptoms persistence Autoantibodies in COVID-19 survivors with post-COVID symptoms: a systematic review Sex differences in immune responses Testosterone is an endogenous regulator of BAFF and splenic B cell number Sex differences in immune responses: hormonal effects, antagonistic selection, and evolutionary consequences Gender differences in symptomatology, socio-demographic information and quality of life in Spanish population with long COVID condition: a cross-sectional study Sex- and Gender-specific Risk Factors of Post-COVID-19 Syndrome: a Population-based Cohort Study in Switzerland Calcium channel blockers may reduce the development of long COVID in females The effect of age on mortality in patients with COVID-19: a meta-analysis with 611,583 subjects Characteristics and impact of Long Covid: findings from an online survey Symptoms and risk factors for long COVID in non-hospitalized adults Attributes and predictors of long COVID Treatment of COVID-19 during the acute phase in hospitalized patients decreases post-acute sequelae of COVID-19 Age, sex and previous comorbidities as risk factors not associated with SARS-CoV-2 infection for long COVID-19: a systematic review and meta-analysis 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic

health records Assessment of a close respiratory follow-up schedule at 3 and 6 months after acute COVID-19 and its related investigations Clinical medicine impact of long-COVID on health care burden: a case control study Impact of fatigue as the primary determinant of functional limitations among patients with post-COVID-19 syndrome: a cross-sectional observational study Socioeconomic inequalities of Long COVID: a retrospective population-based cohort study in the United Kingdom International Journal of Women's Health. 2026. [PMC13094581](#)

15. Dimpy P. Shah, Tanayott Thaweethai, Elizabeth W. Karlson, Hector Bonilla, Benjamin D. Horne, Janet M. Mullington, Juan P. Wisnivesky, Mady Hornig, Daniel J. Shinnick, Jonathan D. Klein, Nathaniel B. Erdmann, Shari B. Brosnahan, Joyce K. Lee-Iannotti, Torri D. Metz, Christine Maughan, Ighovwerha Ofotokun, Harrison T. Reeder, Lauren E. Stiles, Aasma Shaukat, Rachel Hess, Hassan Ashktorab, Logan Bartram, Ingrid V. Bassett, Jacqueline H. Becker, Hassan Brim, Alexander W. Charney, Tananshi Chopra, Rebecca G. Clifton, Steven G. Deeks, Kristine M. Erlandson, Daniel S. Fierer, Valerie J. Flaherman, Vivian Fonseca, Jennifer C. Gander, Sally L. Hodder, Vanessa L. Jacoby, Pavitra Kotini-Shah, Jerry A. Krishnan, Andre Kumar, Bruce D. Levy, David Lieberman, Jenny J. Lin, Jeffrey N. Martin, Grace A. McComsey, Talal Moukabary, Megumi J. Okumura, Michael J. Peluso, Clifford J. Rosen, George Saade, Pankil K. Shah, Zaki A. Sherif, Barbara S. Taylor, Katherine R. Tuttle, Alfredo E. Urdaneta, Julie A. Wallick, Zanthia Wiley, David Zhang, Leora I. Horwitz, Andrea S. Foulkes, Nora G. Singer, et al.. Sex Differences in Long COVID Errors in Supplement 2 NICE guideline on long COVID Two-year longitudinal study reveals that long COVID symptoms peak and quality of life nadirs at 6-12 months postinfection The definition of long COVID used in

interventional studies Mortality-related risk factors of COVID-19: a systematic review and meta-analysis of 42 studies and 423 117 patients Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study Risk factors associated with post-COVID-19 condition: a systematic review and meta-analysis Prognostic factors for post-COVID-19 syndrome: a systematic review and meta-analysis Researching COVID to Enhance Recovery (RECOVER) adult study protocol: rationale, objectives, and design Development of a definition of postacute sequelae of SARS-CoV-2 infection Using propensity scores for racial disparities analysis A characterization of optimal designs for observational studies Using full matching to estimate causal effects in nonexperimental studies: examining the relationship between adolescent marijuana use and adult outcomes Estimating risk ratios and risk differences using regression Covariate balancing propensity score The role of sex hormones in the disparity of COVID-19 outcomes based on gender Female reproductive health impacts of long COVID and associated illnesses including ME/CFS, POTS, and connective tissue disorders: a literature review Reduction of the risk of severe COVID-19 due to Omicron compared to Delta variant in Italy (November 2021-February 2022) Severity of Omicron (B.1.1.529) and Delta (B.1.617.2) SARS-CoV-2 infection among hospitalised adults: a prospective cohort study in Bristol, United Kingdom mice: multivariate imputation by chained equations in R matchthem: matching and weighting after multiple imputation Sex- and age-specific differences in COVID-19 testing, cases, and outcomes: a population-wide study in Ontario, Canada Autoimmune and autoinflammatory connective tissue disorders following COVID-19 Exploring the trajectory curve of long-term

musculoskeletal post-COVID pain symptoms in hospitalized COVID-19 survivors: a multicenter study Incidence and risk factors for persistent symptoms in adults previously hospitalized for COVID-19 The long and winding road of COVID-19 in survivors of hospitalisation: symptoms trajectory and predictors of long COVID Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Long COVID in a prospective cohort of home-isolated patients Long-COVID clinical features and risk factors: a retrospective analysis of patients from the STOP-COVID Registry of the PoLoCOV Study National long COVID impact and risk factors Obesity is associated with a greater number of long-term post-COVID symptoms and poor sleep quality: a multicentre case-control study The presence of symptoms within 6 months after COVID-19: a single-center longitudinal study Prevalence and predictors of persistent post-COVID-19 symptoms Prevalence and predictors of post-acute COVID-19 syndrome (PACS) after hospital discharge: a cohort study with 4 months median follow-up Menopause is associated with immune activation in women with HIV The relationship of serum testosterone levels with the clinical course and prognosis of COVID-19 disease in male patients: a prospective study Male pituitary-gonadal axis dysfunction in post-acute COVID-19 syndrome—prevalence and associated factors: a Mediterranean case series Severely low testosterone in males with COVID-19: a case-control study Symptoms of menopause—global prevalence, physiology and implications Review of menopausal palpitations measures Sex hormones regulate innate immune cells and promote sex differences in respiratory virus infection Sex steroids mediate bidirectional interactions between hosts and microbes Persistent COVID-19-associated neurocognitive symptoms in non-hospitalized patients Estrogen alters thresholds for B cell

apoptosis and activation Menopause and development of Alzheimer's disease: roles of neural glucose metabolism and Wnt signaling The underlying sex differences in neuroendocrine adaptations relevant to myalgic encephalomyelitis chronic fatigue syndrome Mental disorder prevalence among populations impacted by coronavirus pandemics: a multilevel meta-analytic study of COVID-19, MERS & SARS Propensity score analysis: recent debate and discussion JAMA Network Open. 2025. [PMC11755195](#)

16. B-A. Reme, J. Gjesvik, K. Magnusson. Predictors of the post-COVID condition following mild SARS-CoV-2 infection Excess risk and clusters of symptoms after COVID-19 in a large Norwegian cohort A nationwide questionnaire study of post-acute symptoms and health problems after SARS-CoV-2 infection in Denmark Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records A clinical case definition of post-COVID-19 condition by a Delphi consensus Symptoms, complications and management of long COVID: a review Characterising long COVID: a living systematic review Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19 Belief in having had COVID-19 linked with long COVID symptoms Recovery from COVID-19 in Multiple Sclerosis: A Prospective and Longitudinal Cohort Study of the United Kingdom Multiple Sclerosis Register Post-COVID syndrome in non-hospitalised patients with COVID-19: a longitudinal prospective cohort study Trajectory of long covid symptoms after covid-19 vaccination: community based cohort study Prevalence of long COVID complaints in persons with and without COVID-19 Post-covid medical complaints following infection with SARS-CoV-2 Omicron vs Delta variants The association of depression and anxiety with medical symptom burden in patients with chronic medical

illness Random forests Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach Nature Communications. 2023. [PMC10511472](#)

17. Juanjuan Shi, Rui Lu, Yan Tian, Fengping Wu, Xiaozhen Geng, Song Zhai, Xiaoli Jia, Shuangsoo Dang, Wenjun Wang, et al.. Prevalence of and factors associated with long COVID among US adults: a nationwide survey Fatigue and cognitive impairment in Post-COVID-19 syndrome: A systematic review and meta-analysis Post-COVID-19 symptoms and conditions among children and Adolescents - United States, March 1, 2020-January 31, 2022 Global prevalence of Post-Coronavirus disease 2019 (COVID-19) condition or long COVID: A Meta-Analysis and systematic review Study protocol for the innovative support for patients with SARS-COV-2 infections registry (INSPIRE): A longitudinal study of the medium and long-term sequelae of SARS-CoV-2 infection Surveillance for Post-COVID conditions is necessary: addressing the challenges with multiple approaches Prevalence and characteristics in long COVID among adults with asthma in the united States Symptoms experienced at the acute phase of SARS-CoV-2 infection as risk factor of Long-term Post-COVID symptoms: the LONG-COVID-EXP-CM multicenter study Symptoms and risk factors for long COVID in non-hospitalized adults Attributes and predictors of long COVID Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Post-Acute sequelae of SARS-CoV-2 infection among adults aged ≥ 18 Years - Long beach, California, April 1-December 10, 2020 NCHS Urban-Rural classification scheme for counties Rural-Urban disparities in State-Level diabetes prevalence among US adults, 2021 Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Post-acute

COVID-19 syndrome. Incidence and risk factors: A mediterranean cohort study Post-COVID-19 symptoms 6 months after acute infection among hospitalized and non-hospitalized patients E-Cigarette use among US adults in the 2021 behavioral risk factor surveillance system survey Importance of reporting age-adjusted prevalence in epidemiological studies Trends in blood pressure control among US adults with hypertension, 1999-2000 to 2017-2018 Vital signs: influenza hospitalizations and vaccination coverage by race and Ethnicity-United States, 2009-10 through 2021-22 influenza seasons Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio Odds ratio or prevalence ratio? An overview of reported statistical methods and appropriateness of interpretations in Cross-sectional studies with dichotomous outcomes in veterinary medicine Prevalence of symptoms ≤ 12 months after acute illness, by COVID-19 testing status among Adults - United States, December 2020-March 2023 A clinical case definition of post-COVID-19 condition by a Delphi consensus Acute and post-acute respiratory complications of SARS-CoV-2 infection: population-based cohort study in South Korea and Japan Long-term cardiovascular outcomes of COVID-19 Long-term Gastrointestinal outcomes of COVID-19 Short- and long-term neuropsychiatric outcomes in long COVID in South Korea and Japan Prevalence and risk factors for persistent symptoms after COVID-19: a systematic review and meta-analysis The epidemiology of long coronavirus disease in US adults Population-Based estimates of Post-acute sequelae of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection (PASC) prevalence and characteristics Racial/Ethnic disparities in Post-acute sequelae of SARS-CoV-2 infection in

new York: an EHR-Based cohort study from the RECOVER program BMC Public Health. 2025. [PMC12070722](#)

18. Sharmin Shabnam, Cameron Razieh, Hajira Dambha-Miller, Tom Yates, Clare Gillies, Yogini V Chudasama, Manish Pareek, Amitava Banerjee, Ichiro Kawachi, Ben Lacey, Eva JA Morris, Martin White, Francesco Zaccardi, Kamlesh Khunti, Nazrul Islam, et al.. Socioeconomic inequalities of Long COVID: a retrospective population-based cohort study in the United Kingdom Disease and healthcare burden of COVID-19 in the United States Excess deaths associated with covid-19 pandemic in 2020: age and sex disaggregated time series analysis in 29 high income countries Systemic and organ-specific immune-related manifestations of COVID-19 NICE guideline on long COVID More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Long-term complications of COVID-19 Excess years of life lost to COVID-19 and other causes of death by sex, neighbourhood deprivation, and region in England and Wales during 2020: a registry-based study Occupation and COVID-19 mortality in England: a national linked data study of 14.3 million adults Ethnic minorities and COVID-19: examining whether excess risk is mediated through deprivation COVID-19 mortality: a complex interplay of sex, gender and ethnicity Risk of COVID-19 among front-line health-care workers and the general community: a prospective cohort study Health equity in England: the Marmot review 10 years on Trajectory of long covid symptoms after covid-19 vaccination: community based cohort study Association between household size and COVID-19: a UK Biobank observational study Female gender is associated with long COVID syndrome: a prospective cohort study Attributes and predictors of long COVID Persistent COVID-19 symptoms in a community study of 606,434 people in England Characteristics and impact of Long Covid: findings

from an online survey Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Assessment and characterisation of post-COVID-19 manifestations Occupation and risk of severe COVID-19: prospective cohort study of 120 075 UK Biobank participants Premature mortality attributable to socioeconomic inequality in England between 2003 and 2018: an observational study Intersectionality and why it matters to global health Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records in situ using OpenSAFELY Journal of the Royal Society of Medicine. 2023. [PMC10469969](#)

19. Candace H. Feldman, Leah Santacroce, Ingrid V. Bassett, Tanayott Thaweethai, Radica Alicic, Rachel Atchley-Challenner, Alicia Chung, Mark P. Goldberg, Carol R. Horowitz, Karen B. Jacobson, J. Daniel Kelly, Stacey Knight, Karen Lutrick, Praveen Mudumbi, Sairam Parthasarathy, Heather Prendergast, Yuri Quintana, Nasser Sharareh, Judd Shellito, Zaki A. Sherif, Brittany D. Taylor, Emily Taylor, Joel Tsevat, Zanthia Wiley, Natasha J. Williams, Lynn Yee, Lisa Aponte-Soto, Jhony Baissary, Jasmine Berry, Alexander W. Charney, Maged M. Costantine, Alexandria M. Duven, Nathaniel Erdmann, Kacey C. Ernst, Elen M. Feuerriegel, Valerie J. Flaherman, Minjoung Go, Kellie Hawkins, Vanessa Jacoby, Janice John, Sara Kelly, Elijah Kindred, Adeyinka Laiyemo, Emily B. Levitan, Bruce D. Levy, Jennifer K. Logue, Jai G. Marathe, Jeffrey N. Martin, Grace A. McComsey, Torri D. Metz, Tony Minor, Aoyjai P. Montgomery, Janet M. Mullington, Igbo Oforokun, Megumi J. Okumura, Michael J. Peluso, Kristen Pogreba-Brown, Hengameh Raissy, Johana M. Rosas, Upinder Singh, Timothy VanWagoner, Cheryl R. Clark, Elizabeth W. Karlson, et al.. Social Determinants of Health

and Risk of Long COVID in the U.S. RECOVER-Adult Cohort Meanings and Misunderstandings: A Social Determinants of Health Lexicon for Health Care Systems Structural racism and health inequities in the USA: evidence and interventions Structural Racism, Social Risk Factors, and Covid-19 - A Dangerous Convergence for Black Americans Associations of Race/Ethnicity and Food Insecurity With COVID-19 Infection Rates Across US Counties Racial Disparities in COVID-19 Mortality Among Essential Workers in the United States Structural racism and COVID-19 response: higher risk of exposure drives disparate COVID-19 deaths among Black and Hispanic/Latinx residents of Illinois, USA Disparities in COVID-19 Outcomes by Race, Ethnicity, and Socioeconomic Status: A Systematic-Review and Meta-analysis Trends in US Health Insurance Coverage During the COVID-19 Pandemic Disparities in the recovery from critical illness due to COVID-19 Racial and Ethnic Disparities in Housing Instability During the COVID-19 Pandemic: the Role of Assets and Income Shocks The Early Food Insecurity Impacts of COVID-19 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Symptom Clusters Seen in Adult COVID-19 Recovery Clinic Care Seekers 2024 Update of the RECOVER-Adult Long COVID Research Index Sex Differences in Long COVID Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Identifying Contextual and Spatial Risk Factors for Post-Acute Sequelae of SARS-CoV-2 Infection: An EHR-based Cohort Study from the RECOVER Program Methods for the scientific study of discrimination and health: an ecosocial approach Conceptualizing the Mechanisms of Social Determinants of Health: A Heuristic Framework to Inform Future Directions for Mitigation Multi-

cohort study identifies social determinants of systemic inflammation over the life course Systemic Inflammation in Midlife: Race, Socioeconomic Status, and Perceived Discrimination Socioeconomic status and inflammation: a meta-analysis Chronic inflammation in the etiology of disease across the life span Researching COVID to Enhance Recovery (RECOVER) adult study protocol: Rationale, objectives, and design Researching COVID to enhance recovery (RECOVER) pregnancy study: Rationale, objectives and design The eight-item modified Medical Outcomes Study Social Support Survey: psychometric evaluation showed excellent performance Assessing the measurement properties of neighborhood scales: from psychometrics to econometrics Racial Differences in Physical and Mental Health: Socioeconomic Status, Stress and Discrimination Recommendations for Reporting Research About Racial Disparities in Medical and Scientific Journals Socioeconomic inequalities of Long COVID: a retrospective population-based cohort study in the United Kingdom Representation of Race and Ethnicity in the Contemporary US Health Cohort All of Us Research Program Annals of internal medicine. 2025. [PMC12352746](#)

20. Tigist F. Menkir, Barbara Wanjiru Citarella, Louise Sigfrid, Yash Doshi, Luis Felipe Reyes, Jose A. Calvache, Anders Benjamin Kildal, Anders B. Nygaard, Jan Cato Holter, Prasan Kumar Panda, Waasila Jassat, Laura Merson, Christl A. Donnelly, Mauricio Santillana, Caroline Buckee, Stéphane Verguet, Nima S. Hejazi, et al.. Modeling the relative influence of socio-demographic variables on post-acute COVID-19 quality of life The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis Post-acute COVID-19 syndrome (PCS) and health-

related quality of life (HRQoL)—A systematic review and meta-analysis
Attributes and predictors of long COVID
Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records
Risk Factors Associated With Post–COVID-19 Condition: A Systematic Review and Meta-analysis
Symptoms and risk factors for long COVID in non-hospitalized adults
Addressing the Social determinants of children’s Health: A cliff Analogy
Long COVID and Health Inequities: The Role of Primary Care
Integrating Social Care Into the Delivery of Health Care
Socioeconomic inequalities of Long COVID: a retrospective population-based cohort study in the United Kingdom
Population-based estimates of post-acute sequelae of SARS-CoV-2 infection (PASC) prevalence and characteristics
Prevalence and risk factors for long COVID and post-COVID-19 condition in Africa: a systematic review
Race, Ethnicity, and 60-Day Outcomes After Hospitalization With COVID-19
The impact of ethnicity on the long-term sequelae of COVID-19: Follow-up from the first and second waves in North London
Racial and ethnic disparities in SARS-CoV-2 pandemic: analysis of a COVID-19 observational registry for a diverse US metropolitan population
Factors associated with healthy aging in Latin American populations
Educational inequalities in mortality and their mediators among generations across four decades: nationwide, population based, prospective cohort study based on the ChinaHEART project
ISARIC-COVID-19 dataset: A Prospective, Standardized, Global Dataset of Patients Hospitalized with COVID-19
Long-Term Health-Related Quality of Life in Non-Hospitalized
Coronavirus Disease 2019 (COVID-19) Cases With Confirmed Severe Acute Respiratory Syndrome
Coronavirus 2 (SARS-CoV-2) Infection in England: Longitudinal Analysis and Cross-Sectional Comparison With Controls
Random Forests
Nonparametric causal mediation

analysis for stochastic interventional (in)direct effects
 medoutcon: Nonparametric efficient causal mediationanalysis
 with machine learning in R Identifiability and Exchangeability
 for Direct and Indirect Effects Racial disparities in COVID-19
 outcomes exist despite comparable Elixhauser comorbidity
 indices between Blacks, Hispanics, Native Americans, and
 Whites Mediation Analysis with Multiple Mediators Long
 covid outcomes at one year after mild SARS-CoV-2 infection:
 nationwide cohort study A one-year follow-up study of
 systematic impact of long COVID symptoms among patients
 post SARS-CoV-2 omicron variants infection in Shanghai,
 China Selection Mechanisms and Their Consequences:
 Understanding and Addressing Selection Bias medRxiv. 2024.
[PMC11261939](#)

21. Shige Li, Bing Dai, Yusheng Hou, Liang Zhang, Jie Liu, Haijia Hou, Dandan Song, Shengchen Wang, Xiangrui Li, Hongwen Zhao, Wei Wang, Jian Kang, Wei Tan, et al.. Effect of pulmonary rehabilitation for patients with long COVID-19: a systematic review and meta-analysis of randomized controlled trials WHO Clinical Case Definition Working Group on post-COVID-19 condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Symptoms and risk factors for long COVID in non-hospitalized adults Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study Long COVID: major findings, mechanisms and recommendations World Health Organization Mortality in Patients Hospitalized for COVID-19 vs Influenza in Fall-Winter 2023-2024 Pulmonary Rehabilitation for adults with chronic respiratory disease: an official American thoracic society clinical practice guideline

Updated guidance on the management of COVID-19: from an American Thoracic Society/European Respiratory Society coordinated International Task Force (29 July 2020)

Effectiveness of pulmonary rehabilitation among COVID-19 patients: a systematic review and meta-analysis

Pulmonary rehabilitation in SARS-CoV-2: a systematic review and meta-analysis of post-acute patients

Effectiveness of respiratory rehabilitation in COVID-19's post-acute phase: a systematic review

A systematic review of telemedicine-driven pulmonary rehabilitation after the acute phase of COVID-19

Efficacy of pulmonary rehabilitation in post-COVID-19: a systematic review and meta-analysis

Rehabilitation interventions for physical capacity and quality of life in adults with post-COVID-19 condition: a systematic review and meta-analysis

Effectiveness of rehabilitation interventions on adults with COVID-19 and post-COVID-19 condition. A systematic review with meta-analysis

Effect of pulmonary rehabilitation on physical capacity, dyspnea, fatigue, and peripheral muscle strength in patients with post-COVID-19 syndrome: a systematic review and meta-analysis

Effects of exercise therapy on anxiety and depression in patients with COVID-19: a systematic review and meta-analysis

The pulmonary rehabilitation effect on long covid-19 syndrome: a systematic review and meta-analysis

Improved functioning and activity according to the international classification of functioning and disability after multidisciplinary telerehabilitation for post-COVID-19 condition: a randomized control study

Effects of a cardiopulmonary telerehabilitation using functional exercises in individuals after COVID-19 hospital discharge: a randomized controlled trial

Effect of home-based pulmonary rehabilitation on exercise capacity in post COVID-19 patients: a randomized controlled trail

Effects of a cardiopulmonary rehabilitation protocol on functional capacity, dyspnea,

fatigue, and body composition in individuals with post-COVID-19 syndrome: a randomized controlled trial Clinical effectiveness of an online supervised group physical and mental health rehabilitation programme for adults with post-covid-19 condition (REGAIN study): multicentre randomised controlled trial Home-based exercise program for people with residual disability following hospitalization for COVID-19: randomized control trial The PRISMA 2020 statement: an updated guideline for reporting systematic reviews Cochrane Handbook for Systematic Reviews of Interventions version 6.2 Methodological Expectations of Cochrane Intervention Reviews Interventions with a clear focus on achieving behaviour change are important for maintaining training-related gains in people with chronic obstructive pulmonary disease: a systematic review Estimating the sample mean and standard deviation from commonly reported quantiles in meta analysis GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations Alpha, beta, meta: guidelines for assessing power and type I error in meta-analyses Version 0.9 Beta A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses Eight weeks of pilates training improves respiratory measures in people with a history of COVID-19: a preliminary study Inspiratory muscle training while hospitalized with acute COVID-19 respiratory failure: a randomized controlled trial Feasibility and initial efficacy of an app-facilitated telerehabilitation (AFTER) programme for COVID-19 survivors: a pilot randomised study Home-based respiratory muscle training on quality of life and exercise tolerance in long-term post-COVID-19: randomized controlled trial Effect of Tai Chi versus aerobic training on improving hand grip strength, fatigue, and functional performance in older adults post-COVID-19: a randomized controlled trial

Effect of pulmonary rehabilitation program on post hospitalization severe COVID-19 patients (Experimental study) Short-term effects of a respiratory telerehabilitation program in confined COVID-19 patients in the acute phase: a pilot study Home-based pulmonary rehabilitation in patients with COVID-19: the efficiency in pulmonary function A randomized controlled trial examining the impact of low vs. moderate-intensity aerobic training in post-discharge COVID-19 older subjects Safety and efficacy of pulmonary physiotherapy in hospitalized patients with severe COVID-19 pneumonia (PPTCOVID study): a prospective, randomised, single-blind, controlled trial Rehabilitation for post-COVID-19 condition through a supervised exercise intervention: a randomized controlled trial Effect of home-based pulmonary rehabilitation on fatigue, dyspnea, and activities of daily living of COVID-19 patients A telerehabilitation programme in post-discharge COVID-19 patients (TERECO): a randomised controlled trial Respiratory rehabilitation in elderly patients with COVID-19: a randomized controlled study Effect of qigong exercise and acupressure rehabilitation program on pulmonary function and respiratory symptoms in patients hospitalized with severe COVID-19: a randomized controlled trial Effects of a 16-week home-based exercise training programme on health-related quality of life, functional capacity, and persistent symptoms in survivors of severe/critical COVID-19: a randomised controlled trial Inspiratory muscle training enhances recovery post-COVID-19: a randomised controlled trial Evaluating the efficiency of breathing exercises via telemedicine in post-Covid-19 patients: randomized controlled study The effectiveness of POST-DISCHARGE telerehabilitation practices in COVID-19 patients: tele-COVID study-randomized controlled trial An online breathing and wellbeing programme (ENO Breathe) for

people with persistent symptoms following COVID-19: a parallel-group, single-blind, randomised controlled trial
Effectiveness of Liu-zhi-jue exercise on coronavirus disease 2019 in the patients: a randomized controlled trial
Short-term effects of a conditioning telerehabilitation program in confined patients affected by COVID-19 in the acute phase. A pilot randomized controlled trial
Breathing exercises versus strength exercises through telerehabilitation in coronavirus disease 2019 patients in the acute phase: a randomized controlled trial
Effectiveness of exercise training on the dyspnoea of individuals with long COVID: a randomised controlled multicentre trial
Effects of a home-based pulmonary rehabilitation program with and without telecoaching on health-related outcomes in COVID-19 survivors: a randomized controlled clinical study
Effectiveness of a telerehabilitation intervention using ReCOVerry APP of long COVID patients: a randomized, 3-month follow-up clinical trial
Effects of inspiratory muscle training in patients with post-COVID-19
Effects of corrective and breathing exercises on respiratory function of older adults with a history of COVID-19 infection: a randomized controlled trial
Group-based pulmonary telerehabilitation is feasible, safe, beneficial and well-received in patients who have been hospitalised with COVID-19
Cardiovascular, respiratory, and functional effects of home-based exercise training after COVID-19 hospitalization
Construct validity and minimal important difference of 6-minute walk distance in survivors of acute respiratory failure
Pulmonary rehabilitation for chronic obstructive pulmonary disease
Pulmonary rehabilitation for interstitial lung disease
Management of post-acute covid-19 in primary care
Rehabilitation interventions for post-acute COVID-19 syndrome: a systematic review
An official American Thoracic Society/

European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation How COVID-19 shaped mental health: from infection to pandemic effects The global economic burden of COVID-19 disease: a comprehensive systematic review and meta-analysis Therapeutic Advances in Respiratory Disease. 2025. [PMC11907626](#)

22. Evangelin Shaloom Vitus, Christian Nikolai Sørensen, Anastasia Karachalia Sandri, Rahma Elmahdi, Tine Jess. Increased Risk of Post-Acute Sequelae of COVID-19 in Patients With Immune-Mediated Inflammatory Diseases Characteristics of SARS-CoV-2 and COVID-19 Long COVID: major findings, mechanisms and recommendations Post-acute COVID-19 syndrome Post COVID-19 condition (long COVID) Long-term sequelae of COVID-19: a systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms Early biological markers of post-acute sequelae of SARS-CoV-2 infection Pathological sequelae of long-haul COVID Persistent SARS-CoV-2 infection, EBV, HHV-6 and other factors may contribute to inflammation and autoimmunity in long COVID Immune dysregulation in long COVID Immune-mediated inflammatory diseases: common and different pathogenic and clinical features Immune-mediated inflammatory disease therapeutics: past, present and future Improvement of immune dysregulation in individuals with long COVID at 24-months following SARS-CoV-2 infection The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19 Inflammatory cytokines: from discoveries to therapies in IBD New insights into IFN- γ in rheumatoid arthritis: role in the era of JAK inhibitors The immunological basis of inflammatory bowel disease Impact of immunosuppressive therapy on SARS-CoV-2 mRNA vaccine effectiveness in patients with immune-mediated inflammatory diseases: a Danish

nationwide cohort study The rising burden of inflammatory bowel disease in Denmark over two decades: a nationwide cohort study Cytokine profiles associated with acute COVID-19 and long COVID-19 syndrome Psoriasis Potential significance of changes in serum levels of IL-17, TNF- α and DKK-1 in the progression of the rheumatoid arthritis Serum levels of IL-6 and TNF- α may correlate with activity and severity of rheumatoid arthritis Long COVID in inflammatory bowel diseases Incidence of long COVID and impact of medications on the risk of developing long COVID in a nationwide cohort of IBD patients Positive predictive value of the ICD-10 diagnosis code for long-COVID Three-year outcomes of post-acute sequelae of COVID-19 Gastro Hep Advances. 2025. [PMC12205781](#)

23. Mark P. Khurana, Mathilde Marie Brünnich Sloth, Neil Scheidwasser, Jacob Curran-Sebastian, Christian Morgenstern, Nicolas Banholzer, David Thein, Laust H. Mortensen, Morten Rasmussen, Pikka Jokelainen, Frederik Trier Møller, Marc Stegger, Tyra G. Krause, Ewan Cameron, David A. Duchêne, Alexandros Katsiferis, Samir Bhatt, et al.. SARS-CoV-2 reinfections and subsequent risk of hospital-diagnosed post-acute sequelae in Denmark (2020–2022): a nationwide cohort study Longitudinal analysis reveals high prevalence of epstein-Barr virus associated with multiple sclerosis Neurologic manifestations and outcome of West Nile virus infection The neurological complications of chikungunya virus: a systematic review Long COVID: a clinical update Post-COVID-19 condition fatigue outcomes among Danish residents Long COVID: major findings, mechanisms and recommendations Postacute sequelae of SARS-CoV-2 infection in the Pre-Delta, Delta, and Omicron Eras Determinants of the onset and prognosis of the post-COVID-19 condition: a 2-year prospective observational cohort study Natural history of

long-COVID in a nationwide, population cohort study Long COVID science, research and policy Mechanisms of long COVID and the path toward therapeutics Damage to endothelial barriers and its contribution to long COVID CCL2-mediated endothelial injury drives cardiac dysfunction in long COVID Multiple early factors anticipate post-acute COVID-19 sequelae Racial/ethnic differences in post-acute sequelae of SARS-CoV-2 in children and adolescents in the United States Ethnic and racial differences in children and young people with respiratory and neurological post-acute sequelae of SARS-CoV-2: an electronic health record-based cohort study from the RECOVER Initiative Risk of long COVID and associated symptoms after acute SARS-COV-2 infection in ethnic minorities: a nationwide register-linked cohort study in Denmark SARS-COV-2 re-infection and incidence of post-acute sequelae of COVID-19 (PASC) among essential workers in New York: a retrospective cohort study Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Insights from an N3C RECOVER EHR-based cohort study characterizing SARS-CoV-2 reinfections and long COVID The effect of pre-COVID and post-COVID vaccination on long COVID: a systematic review and meta-analysis Effect of covid-19 vaccination on long covid: systematic review Patterns of testing in the extensive Danish national SARS-CoV-2 test set-up Positive predictive value of the ICD-10 diagnosis code for Long-COVID UK Health Security Agency SARS-CoV-2 infection rates of antibody-positive compared with antibody-negative health-care workers in England: a large, multicentre, prospective cohort study (SIREN) A new method of classifying prognostic comorbidity in longitudinal studies: development and validation Glossary: equivalised disposable income. Eurostat Educational and income inequalities across diseases in

Denmark: a register-based cohort study riskRegression: risk regression models and prediction scores for survival analysis with competing risks The parametric g-formula for time-to-event data: intuition and a worked example Long covid defined Acute and postacute sequelae associated with SARS-CoV-2 reinfection Risk of new-onset long COVID following reinfection with severe acute respiratory syndrome coronavirus 2: a community-based cohort Study Post-COVID syndrome: incidence, clinical spectrum, and challenges for primary healthcare professionals The prevalence of long COVID in people with diabetes mellitus-evidence from a UK cohort Is socioeconomic status associated with utilization of health care services in a single-payer universal health care system? The impact of COVID vaccination on incidence of long COVID and healthcare resource utilisation in a primary care cohort in England, 2021-2022 The Lancet Regional Health - Europe. 2026. [PMC12887416](#)

24. Tesleem K. Babalola, Sean A.P. Clouston, Zennur Sekendiz, Devi Chowdhury, Nicola Soriolo, Joseph Kawuki, Jaymie Meliker, Melissa Carr, Bruno R. Valenti, Ashley Fontana, Onix A. Melendez, Olga Morozova, Benjamin J. Luft, et al.. SARS-COV-2 re-infection and incidence of post-acute sequelae of COVID-19 (PASC) among essential workers in New York: a retrospective cohort study COVID-19 sequelae in adults aged less than 50 years: a systematic review Short-and potential long-term adverse health outcomes of COVID-19: a rapid review Long COVID: major findings, mechanisms and recommendations Long COVID or post-COVID conditions 2023 A clinical case definition of post COVID-19 condition by a Delphi consensus Understanding how post-COVID-19 condition affects adults and health care systems A systematic review and meta-analysis of Long COVID symptoms Prevalence and risk factors for long COVID and post-

COVID-19 condition in Africa: a systematic review Post-COVID-19 condition Prevalence and correlates of long COVID symptoms among US adults Multiple early factors anticipate post-acute COVID-19 sequelae SARSCoV-2 reinfections and long COVID in the post-Omicron phase of the pandemic Short-term and long-term impacts of COVID-19 on economic vulnerability: a population-based longitudinal study (COVIDENCE UK) Characteristics and impact of long Covid: findings from an online survey Association between long COVID and vaccination: a 12-month follow-up study in a low-to middle-income country The impact of World Trade Center related medical conditions on the severity of COVID-19 disease and its long-term sequelae A modified Poisson regression approach to prospective studies with binary data Association of vaccine status, reinfections, and risk factors with Long COVID syndrome Symptoms, complications and management of long COVID: a review Trajectory of long COVID symptoms after COVID-19 vaccination: community-based cohort study Effect of covid-19 vaccination on long covid: systematic review Association between BNT162b2 vaccination and long COVID after infections not requiring hospitalization in health care workers Symptom cluster analysis of long COVID-19 in patients discharged from the temporary COVID-19 Hospital in Mexico City Demographic and Clinical Factors Associated with Long COVID: study examines demographic and clinical factors associated with long COVID among people who suffer symptoms long after they were first diagnosed with COVID-19 (long haulers) Clinical and sociodemographic variables associated with long COVID-19: a cross-sectional study Long-COVID symptoms in individuals infected with different SARS-CoV-2 variants of concern: a systematic review of the literature. *Viruses*. 2022; 14: 2629 Risk of long COVID associated with delta versus

omicron variants of SARS-CoV-2 Acute and post-acute sequelae associated with SARS-CoV-2 reinfection Viral persistence, reactivation, and mechanisms of long COVID The persistent viral infections in the development and severity of myalgic encephalomyelitis/chronic fatigue syndrome Post COVID-19 condition diagnosis: a population-based cohort study of occurrence, associated factors, and healthcare use by severity of acute infection COVID-19 illness severity and 2-year prevalence of physical symptoms: an observational study in Iceland, Sweden, Norway and Denmark COVID-19 and multiorgan response: the long-term impact Pathogenic mechanisms of post-acute sequelae of SARS-CoV-2 infection (PASC) Clinical presentation and diagnosis of adults with persistent symptoms following acute illness ("long COVID") How does COVID-19 vaccination affect long-COVID symptoms? Vaccination after developing long COVID: impact on clinical presentation, viral persistence, and immune responses Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: a systematic review Lancet Regional Health - Americas. 2025. [PMC11764319](#)

25. Bingyu Zhang, Qiong Wu, Ravi Jhaveri, Ting Zhou, Michael J. Becich, Yuriy Bisjuk, Frank Blanceró, Elizabeth A. Chrischilles, Cynthia H. Chuang, Linday G. Cowell, Daniel Fort, Carol R. Horowitz, Susan Kim, Nathalia Ladino, David M. Liebovitz, Mei Liu, Abu S. M. Mosa, Hayden T. Schwenk, Srinivasan Suresh, Bradley W. Taylor, David A. Williams, Jeffrey S. Morris, Christopher B. Forrest, Yong Chen, et al.. Reinfection with SARS-CoV-2 in the Omicron Era is Associated with Increased Risk of Post-Acute Sequelae of SARS-CoV-2 Infection: A RECOVER-EHR Cohort Study Long COVID associated with SARS-CoV-2 reinfection among children and adolescents in the omicron era (RECOVER-

EHR): a retrospective cohort study Development of a definition of postacute sequelae of SARS-CoV-2 infection Characterizing long COVID in children and adolescents Postacute sequelae of SARS-CoV-2 in children Clinical features and burden of postacute sequelae of SARS-CoV-2 infection in children and adolescents Long COVID after breakthrough SARS-CoV-2 infection Risk and severity of SARS-CoV-2 reinfections during 2020–2022 in Vojvodina, Serbia: A population-level observational study Changes in population immunity against infection and severe disease from severe acute respiratory syndrome coronavirus 2 Omicron variants in the United States between December 2021 and November 2022 Disease severity during SARS-CoV-2 reinfection: a nationwide study Acute and postacute sequelae associated with SARS-CoV-2 reinfection SARS-CoV-2 reinfections and long COVID in the post-omicron phase of the pandemic Insights from an N3C RECOVER EHR-based cohort study characterizing SARS-CoV-2 reinfections and Long COVID Characteristics of chemosensory perception in long COVID and COVID reinfection Long COVID in children and young after infection or reinfection with the omicron variant: a prospective observational study Post-Covid-19 condition (Long Covid) in children and young people 12 months after infection or reinfection with the Omicron variant: a prospective observational study Long COVID symptoms in Israeli children with and without a history of SARS-CoV-2 infection: a cross-sectional study SARS-CoV-2 re-infection and incidence of post-acute sequelae of COVID-19 (PASC) among essential workers in New York: a retrospective cohort study Association of vaccine status, reinfections, and risk factors with Long COVID syndrome Vaccine effectiveness against long COVID in children Body Mass Index and Postacute Sequelae of SARS-CoV-2 Infection in Children and Young

Adults Real-world effectiveness and causal mediation study of BNT162b2 on long COVID risks in children and adolescents
Pediatric Gastrointestinal Tract Outcomes During the Postacute Phase of COVID-19 Post-Acute Cardiovascular Outcomes of COVID-19 in Children and Adolescents: An EHR Cohort Study from the RECOVER Project Does SARS-CoV-2 Infection Increase the Frequency and Risk of Neuropsychiatric and Related Conditions? Post-acute and Chronic Kidney Function Outcomes of COVID-19 in Children and Adolescents: An EHR Cohort Study from the RECOVER Initiative Racial/ethnic differences in post-acute sequelae of SARS-CoV-2 in children and adolescents in the United States Development and validation of the Pediatric Medical Complexity Algorithm (PMCA) version 3.0 An introduction to propensity score methods for reducing the effects of confounding in observational studies Comparing propensity score methods versus traditional regression analysis for the evaluation of observational data: a case study evaluating the treatment of gram-negative bloodstream infections Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples Comprehensive comparative effectiveness and safety of first-line antihypertensive drug classes: a systematic, multinational, large-scale analysis A modified poisson regression approach to prospective studies with binary data Confounding and collapsibility in causal inference Defining, quantifying, and interpreting “noncollapsibility” in epidemiologic studies of measures of “effect” Interpreting observational studies: why empirical calibration is needed to correct p-values Empirical confidence interval calibration for population-level effect estimation studies in observational healthcare data T cell immune memory after covid-19 and vaccination Differential protection against SARS-CoV-2

reinfection pre-and post-Omicron Trends in laboratory-confirmed SARS-CoV-2 reinfections and associated hospitalizations and deaths among adults aged ≥ 18 years—18 US jurisdictions, September 2021–December 2022 Attributes and predictors of long COVID medRxiv. 2025. [PMC11974971](#)

26. Dena Zeraatkar, Michael Ling, Sarah Kirsh, Tanvir Jassal, Mahnoor Shahab, Hamed Movahed, Jhalok Ronjan Talukdar, Alicia Walch, Samantha Chakraborty, Tari Turner, Lyn Turkstra, Roger S McIntyre, Ariel Izcovich, Lawrence Mbuagbaw, Thomas Agoritsas, Signe A Flottorp, Paul Garner, Tyler Pitre, Rachel J Couban, Jason W Busse, et al.. Interventions for the management of long covid (post-covid condition): living systematic review Health, Economic and Social Development Challenges of the COVID-19 Pandemic: Strategies for Multiple and Interconnected Issues Socio-Economic Impacts and Challenges of the Coronavirus Pandemic (COVID-19): An Updated Review Global Prevalence of Post-Coronavirus Disease 2019 (COVID-19) Condition or Long COVID: A Meta-Analysis and Systematic Review Assessment of the Frequency and Variety of Persistent Symptoms Among Patients With COVID-19: A Systematic Review Association of Post-COVID-19 Condition Symptoms and Employment Status Estimated Global Proportions of Individuals With Persistent Fatigue, Cognitive, and Respiratory Symptom Clusters Following Symptomatic COVID-19 in 2020 and 2021 Persistent Symptoms in Patients After Acute COVID-19 Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? COVID-19 Symptoms: Longitudinal Evolution and Persistence in Outpatient Settings Prevalence and Correlates of Long COVID Symptoms Among US Adults Fatigue and cognitive impairment in Post-

COVID-19 Syndrome: A systematic review and meta-analysis
Comparability of Control and Comparison Groups in Studies
Assessing Long COVID How methodological pitfalls have
created widespread misunderstanding about long COVID The
prevalence and long-term health effects of Long Covid among
hospitalised and non-hospitalised populations: A systematic
review and meta-analysis Long COVID in low-income and
middle-income countries: the hidden public health crisis Long
COVID: pathophysiological factors and abnormalities of
coagulation Generalisable long COVID subtypes: findings
from the NIH N3C and RECOVER programmes Prognostic
Factors for Post-COVID-19 Syndrome: A Systematic Review
and Meta-Analysis Age, Sex and Previous Comorbidities as
Risk Factors Not Associated with SARS-CoV-2 Infection for
Long COVID-19: A Systematic Review and Meta-Analysis
Associations of Depression, Anxiety, Worry, Perceived Stress,
and Loneliness Prior to Infection With Risk of Post-COVID-19
Conditions Association of Self-reported COVID-19 Infection
and SARS-CoV-2 Serology Test Results With Persistent
Physical Symptoms Among French Adults During the
COVID-19 Pandemic A clinical case definition of post-
COVID-19 condition by a Delphi consensus Factors associated
with persistence of symptoms 1 year after COVID-19: A
longitudinal, prospective phone-based interview follow-up
cohort study Three-year outcomes of post-acute sequelae of
COVID-19 Characterising long COVID: a living systematic
review US health agency will invest \$1 billion to investigate
'long COVID' High-dose coenzyme Q10 therapy versus
placebo in patients with post COVID-19 condition: a
randomized, phase 2, crossover trial What interventions or
best practice are there to support people with Long COVID,
or similar post-viral conditions or conditions characterised by
fatigue, to return to normal activities: a rapid review.

Interventions for Improving Long COVID-19 Symptomatology: A Systematic Review Outpatient treatment of COVID-19 and incidence of post-COVID-19 condition over 10 months (COVID-OUT): a multicentre, randomised, quadruple-blind, parallel-group, phase 3 trial Clinical classification and interventions for post-COVID-19 condition: a scoping review Clinical trials on the pharmacological treatment of long COVID: A systematic review Long-COVID treatments: why the world is still waiting Federal Government Announces Long COVID Trials, Research Office Methodology and design of platform trials: a meta-epidemiological study Long covid patients travel abroad for expensive and experimental “blood washing” Could Cognitive Behavioural Therapy Be an Effective Treatment for Long COVID and Post COVID-19 Fatigue Syndrome? Lessons from the Qure Study for Q-Fever Fatigue Syndrome Interventions for the management of post COVID-19 condition (long COVID): Protocol for a living systematic review & network meta-analysis. Long Covid Defined Consistency of covid-19 trial preprints with published reports and impact for decision making: retrospective review Comparison of effect estimates between preprints and peer-reviewed journal articles of COVID-19 trials Comparison of Clinical Study Results Reported in medRxiv Preprints vs Peer-reviewed Journal Articles Identifying and managing problematic trials: A research integrity assessment tool for randomized controlled trials in evidence synthesis Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Publication bias: the problem that won't go away COVID-19 Living Overview of Evidence repository is highly comprehensive and can be used as a single source for COVID-19 studies Studying the post-COVID-19 condition: research challenges, strategies, and importance of Core Outcome Set development A core outcome set for post-

COVID-19 condition in adults for use in clinical practice and research: an international Delphi consensus study
Characterization of long COVID temporal sub-phenotypes by distributed representation learning from electronic health record data: a cohort study
Identification of Distinct Long COVID Clinical Phenotypes Through Cluster Analysis of Self-Reported Symptoms
False individual patient data and zombie randomised controlled trials submitted to Anaesthesia
Hundreds of thousands of zombie randomised trials circulate among us
Checklist to assess Trustworthiness in RAndomised Controlled Trials (TRACT checklist): concept proposal and pilot
RoB 2: a revised tool for assessing risk of bias in randomised trials
The impact of blinding on trial results: A systematic review and meta-analysis
Impact of Allocation Concealment and Blinding in Trials Addressing Treatments for COVID-19: A Methods Study
Theory and practice of Bayesian and frequentist frameworks for network meta-analysis
A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses
Patient-reported outcomes in meta-analyses-part 2: methods for improving interpretability for decision-makers
Pooling health-related quality of life outcomes in meta-analysis-a tutorial and review of methods for enhancing interpretability
Methods for detecting, quantifying, and adjusting for dissemination bias in meta-analysis are described
Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials
GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables
COVID-19 vaccination for the prevention and treatment of long COVID: A systematic review and meta-analysis
GRADE: an emerging consensus on rating quality of evidence and strength of recommendations
GRADE Guidance 34: update on rating imprecision using a minimally

contextualized approach GRADE guidance 35: update on rating imprecision for assessing contextualized certainty of evidence and making decisions Using Explicit Thresholds were valuable for judging Benefits and Harms in partially contextualized GRADE Guidelines How is the minimal clinically important difference established in health-related quality of life instruments? Review of anchors and methods Defining Minimally Important Differences in Functional Outcomes in Musculoskeletal Oncology Mind the methods of determining minimal important differences: three critical issues to consider Determining a minimal important change in a disease-specific Quality of Life Questionnaire Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation Evaluating the credibility of anchor based estimates of minimal important differences for patient reported outcomes: instrument development and reliability study The PRISMA 2020 statement: an updated guideline for reporting systematic reviews GRADE guidelines 26: informative statements to communicate the findings of systematic reviews of interventions Reduced Cell Surface Levels of C-C Chemokine Receptor 5 and Immunosuppression in Long Coronavirus Disease 2019 Syndrome Actovegin use in patients with cognitive impairment after coronavirus infection (COVID-19). Neurology, Neuropsychiatry Vortioxetine for the treatment of post-COVID-19 condition: a randomized controlled trial Glucosaminylmuramyl dipeptide efficacy in post-COVID-19 patient rehabilitation treatment. Infectious diseases: News, Opinions Impact Of Active Cycle Of Breathing Technique On Selected Pulmonary Outcomes In Post-COVID Syndrome Patients Inspiratory muscle training enhances recovery post-COVID-19: a randomised controlled trial Medical Rehabilitation of Patients with Post-COVID-19 Syndrome-A

Comparison of Aerobic Interval and Continuous Training
Comparative effectiveness study of low versus high-intensity aerobic training with resistance training in community-dwelling older men with post-COVID 19 sarcopenia: A randomized controlled trial
Acupuncture as an Additional Method of Rehabilitation Post-COVID-19: a randomized controlled trial
Effectiveness of exercise training on the dyspnoea of individuals with long COVID: A randomised controlled multicentre trial
The impact of post-COVID multicomponent rehabilitation
Effects of a telerehabilitation program and detraining on cardiorespiratory fitness in patients with post-COVID-19 sequelae: A randomized controlled trial
Efficacy of Cognitive-Behavioral Therapy Targeting Severe Fatigue Following Coronavirus Disease 2019: Results of a Randomized Controlled Trial
Effectiveness of a telerehabilitation intervention using ReCOVerry APP of long COVID patients: a randomized, 3-month follow-up clinical trial
Amygdala and Insula Retraining (AIR) Significantly Reduces Fatigue and Increases Energy in People with Long COVID
A synbiotic preparation (SIM01) for post-acute COVID-19 syndrome in Hong Kong (RECOVERY): a randomised, double-blind, placebo-controlled trial [Prospects and possibilities for the treatment of patients with long COVID-19 syndrome]
Effects of l-Arginine Plus Vitamin C Supplementation on Physical Performance, Endothelial Function, and Persistent Fatigue in Adults with Long COVID: A Single-Blind Randomized Controlled Trial
The effect of photobiomodulation versus placebo on functional capacity and fatigability in post COVID-19 elderly
Non-invasive brain stimulation for fatigue in post-acute sequelae of SARS-CoV-2 (PASC)
Hyperbaric oxygen therapy improves neurocognitive functions and symptoms of post-COVID condition: randomized controlled trial
Changes in the proteomics of exhaled breath

condensate under the influence of inhaled hydrogen in patients with post-COVID syndrome Clinical effectiveness of an online supervised group physical and mental health rehabilitation programme for adults with post-covid-19 condition (REGAIN study): multicentre randomised controlled trial Antiviral activity of single-dose PRO 140, a CCR5 monoclonal antibody, in HIV-infected adults CCR5 inhibition in critical COVID-19 patients decreases inflammatory cytokines, increases CD8 T-cells, and decreases SARS-CoV2 RNA in plasma by day 14 Therapist-guided remote versus in-person cognitive behavioural therapy: a systematic review and meta-analysis of randomized controlled trials The NIH has poured \$1 billion into long Covid research — with little to show for it Healthy hearts--and the universal benefits of being physically active: physical activity and health Cognitive behavioral therapies for depression and anxiety in people with chronic disease: A systematic review and meta-analysis Guided internet-based cognitive-behavioral therapy for patients with chronic pain: A meta-analytic review Cognitive behaviour therapy for fatigued cancer survivors: long-term follow-up ME/CFS and Long COVID share similar symptoms and biological abnormalities: road map to the literature Exercise therapy for chronic fatigue syndrome Systematic review of randomized controlled trials for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME) Evidence-Based Care for People with Chronic Fatigue Syndrome and Myalgic Encephalomyelitis Biopsychosocial risk factors of persistent fatigue after acute infection: A systematic review to inform interventions Graded exercise therapy does not restore the ability to work in ME/CFS - Rethinking of a Cochrane review A review on cognitive behavioral therapy (CBT) and graded exercise therapy (GET) in myalgic encephalomyelitis (ME) / chronic fatigue syndrome (CFS):

CBT/GET is not only ineffective and not evidence-based, but also potentially harmful for many patients with ME/CFS Gut microbiota dynamics in a prospective cohort of patients with post-acute COVID-19 syndrome The intestinal microbiota fuelling metabolic inflammation Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19 Prolonged Impairment of Short-Chain Fatty Acid and L-Isoleucine Biosynthesis in Gut Microbiome in Patients With COVID-19 Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization Gastrointestinal Manifestations of COVID-19: A Review of What We Know Long-term complications after infection with SARS-CoV-1, influenza and MERS-CoV - Lessons to learn in long COVID? Rating the quality of evidence is by necessity a matter of judgment The missing voice of patients in drug-safety reporting Patient-Reported Outcomes (PROs) and Patient-Reported Outcome Measures (PROMs) The importance of patient-reported outcomes in clinical trials and strategies for future optimization Recommendations for the development, implementation, and reporting of control interventions in efficacy and mechanistic trials of physical, psychological, and self-management therapies: the CoPPS Statement Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial ESCMID rapid guidelines for assessment and management of long COVID Recommendations for the recognition, diagnosis, and management of long COVID: a Delphi study Long COVID: a clinical update Functional Limitations and Exercise Intolerance in Patients With Post-COVID Condition: A Randomized Crossover Clinical Trial The BMJ. 2024. [PMC11600537](https://pubmed.ncbi.nlm.nih.gov/37116005/)

27. Suzanne D. Vernon, Tianyu Zheng, Hyungrok Do, Vincent C. Marconi, Leonard A. Jason, Nora G. Singer, Benjamin H. Natelson, Zaki A. Sherif, Hector Fabio Bonilla, Emily Taylor, Janet M. Mullington, Hassan Ashktorab, Adeyinka O. Laiyemo, Hassan Brim, Thomas F. Patterson, Teresa T. Akintonwa, Anisha Sekar, Michael J. Peluso, Nikita Maniar, Lucinda Bateman, Leora I. Horwitz, Rachel Hess, et al.. Incidence and Prevalence of Post-COVID-19 Myalgic Encephalomyelitis: A Report from the Observational RECOVER-Adult Study Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens: prospective cohort study Researching COVID to Enhance Recovery (RECOVER) adult study protocol: Rationale, objectives, and design Reyes LF ed Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Neuro-QOL: Brief measures of health-related quality of life for clinical research in neurology The central role of the propensity score in observational studies for causal effects On analysis of epidemiological data involving a 2×2 contingency table: an overview of fisher's exact test and yates' correction for continuity Clinical, epidemiologic, and virologic studies in four clusters of the chronic fatigue syndrome Incidence, risk and prognosis of acute and chronic fatigue syndromes and psychiatric disorders after glandular fever Prevalence and Incidence of Chronic Fatigue Syndrome in Wichita, Kansas Myalgic Encephalomyelitis/Chronic Fatigue Syndrome After SARS-CoV-2 Infection A Community-Based Study of Chronic Fatigue Syndrome Prevalence and Incidence of Chronic Fatigue Syndrome in Wichita, Kansas Estimating Prevalence, Demographics, and Costs of ME/CFS Using Large Scale Medical Claims Data and Machine Learning Onset Patterns and Course of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Essentials of

Diagnosis and Management Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Diagnosis and Management in Young People: A Primer Dissecting the nature of post-exertional malaise Clinical Characteristics of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) Diagnosed in Patients with Long COVID Chronic Fatigue and Postexertional Malaise in People Living With Long COVID: An Observational Study Cognitive impairment in post-acute sequelae of COVID-19 and short duration myalgic encephalomyelitis patients is mediated by orthostatic hemodynamic changes Orthostatic Challenge Causes Distinctive Symptomatic, Hemodynamic and Cognitive Responses in Long COVID and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Accurate and objective determination of myalgic encephalomyelitis/chronic fatigue syndrome disease severity with a wearable sensor Brain abnormalities in myalgic encephalomyelitis/chronic fatigue syndrome: Evaluation by diffusional kurtosis imaging and neurite orientation dispersion and density imaging: Microstructural Abnormalities in CFS Increased ventricular lactate in chronic fatigue syndrome measured by ¹H MRS imaging at 3.0 T. II: comparison with major depressive disorder Neuroinflammation in Patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: An ¹¹C-(R)-PK11195 PET Study Insights From Invasive Cardiopulmonary Exercise Testing of Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Cellular Immune Function in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) Immunologic abnormalities in chronic fatigue syndrome Why myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) may kill you: disorders in the inflammatory and oxidative and nitrosative stress (IO&NS) pathways may explain cardiovascular disorders in ME/CFS Multi-‘omics of

gut microbiome-host interactions in short- and long-term myalgic encephalomyelitis/chronic fatigue syndrome patients Deficient butyrate-producing capacity in the gut microbiome is associated with bacterial network disturbances and fatigue symptoms in ME/CFS Comprehensive Circulatory Metabolomics in ME/CFS Reveals Disrupted Metabolism of Acyl Lipids and Steroids Post-Exertional Malaise Is Associated with Hypermetabolism, Hypoacetylation and Purine Metabolism Deregulation in ME/CFS Cases Urine Metabolomics Exposes Anomalous Recovery after Maximal Exertion in Female ME/CFS Patients The effect of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) severity on cellular bioenergetic function Metabolomic Evidence for Peroxisomal Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Fecal metagenomic profiles in subgroups of patients with myalgic encephalomyelitis/chronic fatigue syndrome A map of metabolic phenotypes in patients with myalgic encephalomyelitis/chronic fatigue syndrome Bioenergetic and Proteomic Profiling of Immune Cells in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Patients: An Exploratory Study Deep phenotyping of myalgic encephalomyelitis/chronic fatigue syndrome in Japanese population Plasma metabolomics reveals disrupted response and recovery following maximal exercise in myalgic encephalomyelitis/chronic fatigue syndrome Prospective Biomarkers from Plasma Metabolomics of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Implicate Redox Imbalance in Disease Symptomatology Metabolic profiling of a myalgic encephalomyelitis/chronic fatigue syndrome discovery cohort reveals disturbances in fatty acid and lipid metabolism Journal of General Internal Medicine. 2025. [PMC11968624](#)

28. Liang-Yu Lin, Alasdair D. Henderson, Oliver Carlile, Iain Dillingham, Ben F. C. Butler-Cole, Michael Marks, Andrew Briggs, Mark Jit, Laurie A. Tomlinson, Chris Bates, John Parry, Sebastian C. J. Bacon, Ben Goldacre, Amir Mehrkar, Brian MacKenna, Rosalind M. Eggo, Emily Herrett, et al.. Healthcare utilisation in people with long COVID: an OpenSAFELY cohort study Impact of long COVID on health-related quality-of-life: an OpenSAFELY population cohort study using patient-reported outcome measures (OpenPROMPT) GP consultation rates for sequelae after acute covid-19 in patients managed in the community or hospital in the UK: population based study Health care utilization in the 6 months following SARS-CoV-2 infection Healthcare use attributable to COVID-19: a propensity-matched national electronic health records cohort study of 249,390 people in Wales Long-haul COVID: healthcare utilization and medical expenditures 6 months post-diagnosis Risk factors, health outcomes, healthcare services utilization, and direct medical costs of patients with long COVID Estimating the economic burden of long-Covid: the additive cost of healthcare utilisation among COVID-19 recoverees in Israel A retrospective cohort analysis leveraging augmented intelligence to characterize long COVID in the electronic health record: a precision medicine framework Identifying who has long COVID in the USA: a machine learning approach using N3C data OpenSAFELY: representativeness of electronic health record platform OpenSAFELY-TPP data compared to the population of England Indirect acute effects of the COVID-19 pandemic on physical and mental health in the UK: a population-based study Modeling health care expenditures and use Too much ado about two-part models and transformation?: comparing methods of modeling Medicare expenditures Prevalence and characteristics of long

COVID in elderly patients: an observational cohort study of over 2 million adults in the US Influence of the news media on diagnostic testing in the emergency department Health care costs associated with muscle weakness: a UK population-based estimate BMC Medicine. 2024. [PMC11188519](#)

29. Emeka Elvis Duru, Godwin Okoye, Sanghoon Lee, Peter Weir, Jaewhan Kim. Comparison of Healthcare Expenditures Among Individuals With and Without Long COVID in the United States Long COVID basics Symptoms, complications and management of long COVID: a review Incidence, co-occurrence, and evolution of long-COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19 Clinical overview of long COVID Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study Long covid-mechanisms, risk factors, and management Prevalence of COVID-19-like symptoms among people living with HIV, and using antiretroviral therapy for prevention and treatment Neurologic features in severe SARS-CoV-2 infection Neurological manifestations and COVID-19: experiences from a tertiary care center at the Frontline Outcomes of cardiovascular magnetic resonance imaging in patients recently recovered from coronavirus disease 2019 (COVID-19) Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study Changes in outpatient health care use after COVID-19 infection among veterans Risk factors, health outcomes, healthcare services utilization, and direct medical costs of patients with long COVID Healthcare utilisation in people with long COVID: an OpenSAFELY cohort study Long COVID: an overview Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Systematic review on COVID-19 readmission and risk factors: future of machine learning in COVID-19 readmission

studies Estimating the economic burden of long-Covid: the additive cost of healthcare utilisation among COVID-19 recoverees in Israel Direct medical costs associated with post-COVID-19 conditions among privately insured children and adults Patterns of utilisation of specialist care after SARS-Cov-2 infection: a retrospective cohort study MEPS HC 243 2022 full year consolidated data file Long COVID is associated with decreased quality of life and increased mental disability US health care spending by race and ethnicity, 2002-2016 National Health Expenditures 2022 highlights Education and family health care spending Long COVID among US adults from a population-based study: association with vaccination, cigarette smoking, and the modifying effect of chronic obstructive pulmonary disease (COPD) Direct health care costs associated with COVID-19 in the United States Age, multiple chronic conditions, and COVID-19: a literature review Estimating log models: to transform or not to transform? The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies Long COVID and rehabilitation Global impact of the COVID-19 pandemic on mental health services: a systematic review Examining potential Long COVID effects through utilization of healthcare resources: a retrospective, population-based, matched cohort study comparing individuals with and without prior SARS-CoV-2 infection Twopm: two-part models Getting the right tail right: modeling tails of health expenditure distributions Association of mental health disorders with health care utilization and costs among adults with chronic disease Prevalence and medical costs of chronic diseases among adult Medicaid beneficiaries Multiple chronic conditions and healthcare costs among adults Attributes and predictors of long COVID Disparities in health care spending and utilization

among Black and White Medicaid enrollees Development of a definition of postacute sequelae of SARS-CoV-2 infection How methodological pitfalls have created widespread misunderstanding about long COVID Inquiry: A Journal of Medical Care Organization, Provision and Financing. 2026. [PMC12764753](#)

30. Yael Wolff Sagy, Ilan Feldhamer, Shuli Brammli-Greenberg, Gil Lavie. Estimating the economic burden of long-Covid: the additive cost of healthcare utilisation among COVID-19 recoverees in Israel The economic burden of COVID-19 in the United States: estimates and projections under an infection-based herd immunity approach The COVID-19 pandemic and the \$16 trillion virus Count the cost of disability caused by COVID-19 COVID-19: evaluation and management of adults with persistent symptoms following acute illness (long COVID) Prevalence of ongoing symptoms following Coronavirus (COVID-19) infection in the UK. data and analyses from census 2021: 3 March 2022. office of national statistics, UK Four-month clinical status of a cohort of patients after hospitalization for COVID-19 Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: A systematic review Characterizing long COVID: a living systematic review Physical, cognitive, and mental health impacts of COVID-19 after Hospitalisation (PHOSP-COVID): a UK Multicentre, prospective cohort study 1-year outcomes in hospital survivors with COVID-19: a longitudinal cohort study 3-month, 6-month, 9-month, and 12-month respiratory outcomes in patients following COVID-19-related Hospitalisation: a prospective study High-dimensional characterization of post-acute sequelae of COVID-19 Risk of persistent and new clinical sequelae among adults aged 65 years and older during the post-acute phase of SARS-Cov-2 infection: retrospective cohort study BMJ Israel - health

system review Health care utilization in the 6 months following SARS-Cov-2 infection Semiparametric difference-in-differences Estimators Comorbidity characterization among eMERGE institutions: A pilot evaluation with the Johns Hopkins adjusted clinical groups® system Long Covid outcomes at one year after mild SARS-Cov-2 infection: nationwide cohort study Impact of COVID-19 pandemic on utilisation of Healthcare services: a systematic review Trends in Healthcare utilisation during COVID-19: a longitudinal study from the UK The impact of COVID-19 pandemic on emergency Department visits and associated mortality during 14 months of the pandemic in Israel Post-COVID conditions among adult COVID-19 survivors aged 18-64 and ≥ 65 years—United States Long COVID: household pulse survey Risk factors, health outcomes, Healthcare services utilization, and direct medical costs of patients with long COVID Risk of long COVID associated with Delta versus Omicron variants of SARS-Cov-2 Long COVID could cost the economy trillions, experts predict BMJ Global Health. 2023. [PMC10357303](#)

31. Ruth C. E. Bowyer, Charlotte Huggins, Renin Toms, Richard J. Shaw, Bo Hou, Ellen J. Thompson, Alex S. F. Kwong, Dylan M. Williams, Milla Kibble, George B. Ploubidis, Nicholas J. Timpson, Jonathan A. C. Sterne, Nishi Chaturvedi, Claire J. Steves, Kate Tilling, Richard J. Silverwood, et al.. Characterising patterns of COVID-19 and long COVID symptoms: evidence from nine UK longitudinal studies Excess risk and clusters of symptoms after COVID-19 in a large Norwegian cohort Long covid—mechanisms, risk factors, and management Inflammation triggered by SARS-CoV-2 and ACE2 augment drives multiple organ failure of severe COVID-19: molecular mechanisms and implications How and why patients made long covid The COVID-19 sequelae: a cross-sectional evaluation of post-recovery symptoms and the

need for rehabilitation of COVID-19 survivors 'Long-COVID': a cross-sectional study of persisting symptoms, biomarker and imaging abnormalities following hospitalisation for COVID-19 Persistent symptoms following SARS-CoV-2 infection in a random community sample of 508,707 people Association of self-reported COVID-19 infection and SARS-CoV-2 serology test results with persistent physical symptoms among french adults during the COVID-19 pandemic Post-acute COVID-19 syndrome. Incidence and risk factors: a Mediterranean cohort study 'Long COVID': persistent COVID-19 symptoms in survivors managed in Lagos State, Nigeria Post-COVID syndrome in non-hospitalised patients with COVID-19: a longitudinal prospective cohort study Frequency, signs and symptoms, and criteria adopted for long COVID-19: a systematic review Causation or confounding: why controls are critical for characterizing long COVID Cohort profile: 1958 British birth cohort (National Child Development Study) Cohort profile: 1970 British birth cohort (BCS70) Next steps (formerly known as the Longitudinal study of young people in England) The Millennium cohort study: the making of a multi-purpose resource for social science and policy Cohort profile: the 'Children of the 90s'—the index offspring of the Avon Longitudinal study of parents and children Cohort profile: the Avon longitudinal study of parents and children: ALSPAC mothers cohort TwinsUK: the UK adult twin registry update TwinsUK COVID-19 personal experience questionnaire (CoPE): wave 1 data capture april-may 2020 [version 1; peer review: awaiting peer review] Cohort profile: the born in Bradford multi-ethnic family cohort study Born in Bradford's better start: an experimental birth cohort study to evaluate the impact of early life interventions Understanding society: design overview Cohort profile: generation Scotland: Scottish family health study (GS:SFHS). The study, its participants and

their potential for genetic research on health and illness
CovidLife: a resource to understand mental health, well-being
and behaviour during the COVID-19 pandemic in the UK
[version 1; peer review: 1 approved] Deciding on the number
of classes in latent class analysis and growth mixture
modeling: a monte carlo simulation study Symptom clustering
patterns and population characteristics of COVID-19 based on
text clustering method Symptom clusters in COVID-19: A
potential clinical prediction tool from the COVID Symptom
Study app Clinical predictors of long COVID-19 and
phenotypes of mild COVID-19 at a tertiary care centre in
India Persistence, magnitude, and patterns of postacute
symptoms and quality of life following onset of SARS-CoV-2
infection: cohort description and approaches for
measurement Characteristics and predictors of acute and
chronic post-COVID syndrome: a systematic review and meta-
analysis Clinical characteristics of 3062 COVID-19 patients: a
meta-analysis Clinical characteristics of 4499 COVID-19
patients in Africa: a meta-analysis Symptom prevalence,
duration, and risk of hospital admission in individuals infected
with SARS-CoV-2 during periods of omicron and delta variant
dominance: a prospective observational study from the ZOE
COVID study European Journal of Epidemiology. 2023.
[PMC9860244](#)

32. Toshiaki Asakura, Takashi Kimura, Isaku Kurotori, Katabami Kenichi, Miyuki Hori, Mariko Hosogawa, Masayuki Saijo, Kaori Nakanishi, Hiroyasu Iso, Akiko Tamakoshi, et al.. Case-Control Study of Long COVID, Sapporo, Japan How and why patients made Long Covid. Long COVID: aiming for a consensus. A clinical case definition of post-COVID-19 condition by a Delphi consensus. Long-term health-related quality of life in non-hospitalized coronavirus disease 2019 (COVID-19) cases with confirmed severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2) infection in England: longitudinal analysis and cross-sectional comparison with controls. Physical and mental health 3 months after SARS-CoV-2 infection (long COVID) among adolescents in England (CLOcK): a national matched cohort study. Post-COVID conditions among adult COVID-19 survivors aged 18–64 and ≥65 years—United States, March 2020–November 2021. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. Prevalence of post-COVID-19 symptoms in hospitalized and non-hospitalized COVID-19 survivors: A systematic review and meta-analysis. Long-term sequelae of COVID-19: A systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms. The human phenotype ontology in 2021. Incubation period of COVID-19 caused by unique SARS-CoV-2 strains: a systematic review and meta-analysis. Probable inference, the law of succession, and statistical inference. How COVID-19 induces cytokine storm with high mortality. COVID-19 mechanisms in the human body—what we know so far. Long-term bone and lung consequences associated with hospital-acquired severe acute respiratory syndrome: a 15-year follow-up from a prospective cohort study. Altered lipid metabolism in recovered SARS patients twelve years after infection. Long COVID in a prospective cohort of home-isolated patients. The neuroinvasiveness, neurotropism, and neurovirulence of SARS-CoV-2. Long-COVID and post-COVID health complications: an up-to-date review on clinical conditions and their possible molecular mechanisms. New onset of alopecia areata in a patient with SARS-CoV-2 infection: Possible pathogenetic correlations? Symptom prevalence, duration,

and risk of hospital admission in individuals infected with SARS-CoV-2 during periods of omicron and delta variant dominance: a prospective observational study from the ZOE COVID Study. Prevalence of post-acute COVID-19 syndrome symptoms at different follow-up periods: a systematic review and meta-analysis. Six-month sequelae of post-vaccination SARS-CoV-2 infection: A retrospective cohort study of 10,024 breakthrough infections. Identifying who has long COVID in the USA: a machine learning approach using N3C data. *Emerging Infectious Diseases*. 2023. [PMC10124665](#)

33. Francesco Di Gennaro, Alessandra Belati, Ottavia Tulone, Lucia Diella, Davide Fiore Bavaro, Roberta Bonica, Vincenzo Genna, Lee Smith, Mike Trott, Olivier Bruyere, Luigi Mirarchi, Claudia Cusumano, Ligia Juliana Dominguez, Annalisa Saracino, Nicola Veronese, Mario Barbagallo, et al.. Incidence of long COVID-19 in people with previous SARS-Cov2 infection: a systematic review and meta-analysis of 120,970 patients Long-term impact of COVID-19: a systematic review of the literature and meta-analysis More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Chronic COVID syndrome: need for an appropriate medical terminology for long-COVID and COVID long-haulers Global prevalence of post COVID-19 condition or long COVID: a meta-analysis and systematic review Prevalence and determinants of fatigue after COVID-19 in non-hospitalized subjects: a population-based study A systematic review and meta-analysis of long COVID symptoms Long-term sequelae of COVID-19: a systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms How to properly use the PRISMA statement Assessing the quality of studies in meta-analyses: advantages and limitations of the Newcastle Ottawa scale Assessing the quality of studies in meta-research: review/guidelines on the most important quality

assessment tools Meta-analysis in clinical trials Age periods of human life Quantifying heterogeneity in a meta-analysis Bias in meta-analysis detected by a simple, graphical test Characterising long COVID: a living systematic review Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review Interventions for improving long COVID-19 symptomatology: a systematic review One-year outcomes in survivors of the acute respiratory distress syndrome Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: a systematic review and meta-analysis The long-term impact of severe acute respiratory syndrome on pulmonary function, exercise capacity and health status Chronic widespread musculoskeletal pain, fatigue, depression and disordered sleep in chronic post-SARS syndrome; a case-controlled study Long COVID risk-a signal to address sex hormones and women's health Attributes and predictors of long-COVID: analysis of COVID cases and their symptoms collected by the COVID symptoms study app Severity of COVID-19 and adverse long-term outcomes: a retrospective cohort study based on a US electronic health record database Physical exercise as a multimodal tool for COVID-19: could it be used as a preventive strategy? The causes, consequences and detection of publication bias in psychiatry Internal and Emergency Medicine. 2022. [PMC9709360](#)

34. Ramona-Georgiana Halas, Delia Mira Berceanu Vaduva, Matilda Radulescu, Ana-Cristina Bredicean, Diana-Maria Mateescu, Ana-Olivia Toma, Ioana-Georgiana Cotet, Cristina-Elena Guse, Andrei Marginean, Madalin-Marius Margan, Voichita Elena Lazureanu, et al.. Long COVID Prevalence and Risk Factors: A Systematic Review and Meta-Analysis of

Prospective Cohort Studies A Clinical Case Definition of Post COVID-19 Condition by a Delphi Consensus Global Prevalence of Post-Coronavirus Disease 2019 (COVID-19) Condition or Long COVID: A Meta-Analysis and Systematic Review Long COVID: Major Findings, Mechanisms and Recommendations Long-Term Sequelae of COVID-19: A Systematic Review and Meta-Analysis of One-Year Follow-Up Studies on Post-COVID Symptoms Attributes and Predictors of Long COVID Assessment of the Frequency and Variety of Persistent Symptoms among Patients with COVID-19: A Systematic Review 1-Year Outcomes in Hospital Survivors with COVID-19: A Longitudinal Cohort Study Long COVID or Post-Acute Sequelae of COVID-19 (PASC): An Overview of Biological Factors That May Contribute to Persistent Symptoms Immunological Dysfunction Persists for 8 Months Following Initial Mild-to-Moderate SARS-CoV-2 Infection Multiple early factors anticipate post-acute COVID-19 sequelae An Inflammatory Cytokine Signature Predicts COVID-19 Severity and Survival Trajectory of Long Covid Symptoms After COVID-19 Vaccination: Community Based Cohort Study The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews Postacute Sequelae of SARS-CoV-2 Infection in the Pre-Delta, Delta, and Omicron Eras Insights into Long COVID: Unraveling Risk Factors, Clinical Features, Radiological Findings, Functional Sequelae and Correlations: A Retrospective Cohort Study 3-month, 6-month, 9-month, and 12-month respiratory outcomes in patients following COVID-19-related hospitalisation: A prospective study Persistent Symptoms in Adult Patients 1 Year After Coronavirus Disease 2019 (COVID-19): A Prospective Cohort Study The Newcastle–Ottawa Scale (NOS) for Assessing the Quality of Nonrandomised Studies in Meta-Analyses. Ottawa Hospital

Research Institute Cohort profile: The Johns Hopkins COVID Long Study (JHCLS)—A US nationwide prospective cohort study Post COVID-19 sequelae: A prospective observational study from Northern India Long COVID prevalence and impact on quality of life 2 years after acute COVID-19 Health outcomes in people 2 years after surviving hospitalisation with COVID-19: A longitudinal cohort study Cognitive sequelae of long COVID may not be permanent: A prospective study Trajectories of persisting COVID-19 symptoms up to 24 months after acute infection: Findings from the Predi-Covid cohort study Long COVID-19: A Four-Year prospective cohort study of risk factors, recovery, and quality of life A prospective study of long-term outcomes among hospitalized COVID-19 patients with and without neurological complications Post-acute sequelae of COVID-19 is characterized by diminished peripheral CD8+ β 7 integrin+ T cells and anti-SARS-CoV-2 IgA response Persistence of Long COVID Symptoms Two Years After SARS-CoV-2 Infection: A Prospective Longitudinal Cohort Study *Biomedicines*. 2025. [PMC12730990](#)

35. Hiroshi Hoshijima, Takahiro Mihara, Hiroyuki Seki, Shunsuke Hyuga, Norifumi Kuratani, Toshiya Shiga, Robert Jeenchen Chen, Miquel Vall-llosera Camps, Huzaifa Ahmad Cheema, Huzaifa Ahmad Cheema, et al.. Incidence of long-term post-acute sequelae of SARS-CoV-2 infection related to pain and other symptoms: A systematic review and meta-analysis Facing up to long COVID. Long-term consequences of COVID-19: research needs Covid-19: What do we know about "long covid"? How and why patients made Long Covid. Long-COVID: An evolving problem with an extensive impact Pain: A potential new label of COVID-19. Chronic pain after COVID-19: implications for rehabilitation. Outcomes for patients following hospitalization for COVID-19 Prevalence

and characteristics of new-onset pain in COVID-19 survivors, a controlled study. Long covid: doctors must assess and investigate patients properly Meta-analysis of prevalence. Meta-analysis in clinical trials. Bias in meta-analysis detected by a simple, graphical test Patient outcomes after hospitalisation with COVID-19 and implications for follow-up: Results from a prospective UK cohort. Evolution of altered sense of smell or taste in patients with mildly symptomatic COVID-19. Persistent symptoms in patients after acute COVID-19 Follow-up of adults with noncritical COVID-19 two months after symptom onset Clinical characteristics and outcomes of adult patients admitted with COVID-19 in East London: a retrospective cohort analysis Patterns of smell recovery in 751 patients affected by the COVID-19 outbreak Long-term COVID-19 symptoms in a large unselected population. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. Multi-organ impairment in low-risk individuals with long COVID. Pericarditis and myocarditis long after SARS-CoV-2 infection: a cross-sectional descriptive study in health-care workers. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? Postdischarge symptoms and rehabilitation needs in survivors of COVID-19 infection: a cross-sectional evaluation 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study Onset, duration, and persistence of taste and smell changes and other COVID-19 symptoms: longitudinal study in Israeli patients. Clinical characteristics associated with persistent olfactory and taste alterations in COVID-19: a preliminary report on 121 patients. 'Long-COVID': a cross-sectional study of persisting symptoms, biomarker and imaging abnormalities following hospitalisation for COVID-19. Delayed symptoms in patients recovered from COVID-19.

Chemosensory dysfunction in COVID-19: prevalences, recovery rates, and clinical associations on a large Brazilian sample. Long COVID in the Faroe Islands:-a longitudinal study among non-hospitalized patients COVID-19 severity impacts on long-term neurological manifestation after hospitalisation. High prevalence of headaches during Covid-19 infection: a retrospective cohort study. Comparing outcomes of hospitalized patients with moderate and severe COVID-19 following treatment with hydroxychloroquine plus atazanavir/ritonavir. Clinical, virological and imaging profile in patients with prolonged forms of COVID-19: a cross-sectional study. Three-month outcomes in hospitalized COVID-19 patients. Persistent symptoms 1.5–6 months after COVID-19 in non-hospitalised subjects: a population-based cohort study. Attributes and predictors of Long-COVID: analysis of COVID cases and their symptoms collected by the Covid Symptoms Study App. Symptom duration and risk factors for delayed return to usual health among outpatients with COVID-19 in a multistate health care systems network—United States, March-June 2020. Anxiety and depression symptoms after virological clearance of COVID-19: a cross-sectional study in Milan, Italy Persistent fatigue following SARS-CoV-2 infection is common and independent of severity of initial infection. Clinical features and outcomes of discharged coronavirus disease 2019 patients: a prospective cohort study. Post-discharge health status and symptoms in patients with severe COVID-19 Mental health status and related influencing factors of COVID-19 survivors in Wuhan, China. Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study Medium term follow-up of 337 patients with coronavirus disease 2019 (COVID-19) in a Fangcang Shelter Hospital in Wuhan, China. Follow-up study of the pulmonary function and related physiological

characteristics of COVID-19 survivors three months after recovery. Sixty-day outcomes among patients hospitalized with COVID-19 Post-discharge persistent symptoms and health-related quality of life after hospitalization for COVID-19. Post-acute COVID-19 syndrome. COVID-19 and the nervous system The neurological manifestations of COVID-19: a review article. COVID-19 is a real headache! Our evolving understanding of ME/CFS. Fibromyalgia. Musculoskeletal consequences of COVID-19 Time course prevalence of post-COVID pain symptoms of musculoskeletal origin in patients who had survived severe acute respiratory syndrome coronavirus 2 infection: a systematic review and meta-analysis Prevalence of post-acute COVID-19 syndrome symptoms at different follow-up periods: a systematic review and meta-analysis Decision Letter 0 Author response to Decision Letter 0 Decision Letter 1 Author response to Decision Letter 1 Decision Letter 2 Acceptance letter PLOS ONE. 2023. [PMC10686440](https://doi.org/10.1371/journal.pone.0268440)

36. Alexander Preiss, Abhishek Bhatia, Leyna V. Aragon, John M. Baratta, Monika Baskaran, Frank Blancero, Michael Daniel Brannock, Robert F. Chew, Iván Díaz, Megan Fitzgerald, Elizabeth P. Kelly, Andrea G. Zhou, Thomas W. Carton, Christopher G. Chute, Melissa Haendel, Richard Moffitt, Emily Pfaff, Philippa C. Dodd, Philippa C. Dodd, Heather Van Epps, Alexandra Tosun, Alexandra Tosun, et al.. Effect of Paxlovid treatment during acute COVID-19 on Long COVID onset: An EHR-based target trial emulation from the N3C and RECOVER consortia EFFECT OF PAXLOVID TREATMENT DURING ACUTE COVID-19 ON LONG COVID ONSET: AN EHR-BASED TARGET TRIAL EMULATION FROM THE N3C AND RECOVER CONSORTIA Effect of Paxlovid treatment during acute COVID-19 on Long COVID onset: An EHR-based target trial emulation from the N3C and RECOVER consortia.

Effect of Paxlovid treatment during acute COVID-19 on Long COVID onset: An EHR-based target trial emulation from the N3C and RECOVER consortia. Nirmatrelvir plus ritonavir for early COVID-19 in a large U.S. Health System : a population-based cohort study Oral nirmatrelvir for high-risk, nonhospitalized adults with COVID-19 Effect of nirmatrelvir/ritonavir (Paxlovid) on hospitalization among adults with COVID-19: an EHR-based target trial emulation from N3C Effectiveness of Paxlovid in reducing severe coronavirus disease 2019 and mortality in high-risk patients Hospitalization and emergency department encounters for COVID-19 after Paxlovid treatment—California Impact of the use of oral antiviral agents on the risk of hospitalization in community coronavirus disease 2019 patients (COVID-19) Nirmatrelvir or molnupiravir use and severe outcomes from omicron infections Effect of oral nirmatrelvir on Long COVID symptoms: 4 cases and rationale for systematic studies The use of nirmatrelvir-ritonavir in a case of breakthrough Long COVID Case report: treatment of Long COVID with a SARS-CoV-2 antiviral and IL-6 blockade in a patient with rheumatoid arthritis and SARS-CoV-2 antigen persistence Impact of extended-course oral nirmatrelvir/ritonavir (Paxlovid) in established Long COVID: case series and research considerations Paxlovid as a potential treatment for Long COVID Nirmatrelvir-ritonavir and symptoms in adults with postacute sequelae of SARS-CoV-2 infection: the STOP-PASC randomized clinical trial Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Neurologic manifestations of Long COVID differ based on acute COVID-19 severity Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Prevalence of post-COVID-19 symptoms in hospitalized and non-hospitalized COVID-19 survivors: a

systematic review and meta-analysis Association of treatment with nirmatrelvir and the risk of post-COVID-19 condition Effectiveness of nirmatrelvir-ritonavir against the development of post-COVID-19 conditions among U.S. veterans : a target trial emulation Association of nirmatrelvir for acute SARS-CoV-2 infection with subsequent Long COVID symptoms in an observational cohort study Platform adaptive trial of novel antivirals for early treatment of COVID-19 In the community (PANORAMIC): protocol for a randomised, controlled, open-label, adaptive platform trial of community novel antiviral treatment of COVID-19 in people at increased risk of more severe disease The National COVID Cohort Collaborative (N3C): rationale, design, infrastructure, and deployment Synergies between centralized and federated approaches to data quality: a report from the national COVID cohort collaborative Using Big Data to emulate a target trial when a randomized trial is not available Target trial emulation: a framework for causal inference from observational data COVID-19 in children. II: pathogenesis, disease spectrum and management Pediatric nirmatrelvir/ritonavir prescribing patterns during the COVID-19 pandemic Reengineering a machine learning phenotype to adapt to the changing COVID-19 landscape: a study from the N3C and RECOVER consortia Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 An introduction to individual participant data meta-analysis Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies Why we need a deeper understanding of the pathophysiology of Long COVID Traumatic brain injury and long-term risk of stroke among US military veterans Asbestos exposure and severity of COVID-19 Measurement in

Veterans Affairs Health Services Research: veterans as a special population Administrative database research infrequently used validated diagnostic or procedural codes Matching methods for causal inference: a review and a look forward Misunderstandings between experimentalists and observationalists about causal inference Association between early treatment with tocilizumab and mortality among critically ill patients with COVID-19 Estimates of overall survival in patients with cancer receiving different treatment regimens: emulating hypothetical target trials in the surveillance, epidemiology, and end results (SEER)-Medicare linked database Avoidable flaws in observational analyses: an application to statins and cancer Emulating a novel clinical trial using existing observational data. predicting results of the PreVent study A clinical case definition of post-COVID-19 condition by a Delphi consensus Long COVID: aiming for a consensus Strategies for handling missing data in electronic health record derived data Out-of-system care and recording of patient characteristics critical for comparative effectiveness research Evaluating risk-prediction models using data from electronic health records Comorbidity and its impact on patients with COVID-19 Epidemiological, comorbidity factors with severity and prognosis of COVID-19: a systematic review and meta-analysis The association between presence of comorbidities and COVID-19 severity: a systematic review and meta-analysis Decision Letter 0 Decision Letter 1 Author response to Decision Letter 2 Decision Letter 2 Author response to Decision Letter 3 Decision Letter 3 Author response to Decision Letter 4 Decision Letter 4 PLOS Medicine. 2025. [PMC12445499](https://doi.org/10.1371/journal.pmed.1005499)

37. Alexander Preiss, Abhishek Bhatia, Leyna V. Aragon, John M. Baratta, Monika Baskaran, Frank Blancero, M. Daniel Brannock, Robert F. Chew, Iván Díaz, Megan Fitzgerald,

Elizabeth P. Kelly, Andrea Zhou, Thomas W. Carton, Christopher G. Chute, Melissa Haendel, Richard Moffitt, Emily Pfaff, et al.. EFFECT OF PAXLOVID TREATMENT DURING ACUTE COVID-19 ON LONG COVID ONSET: AN EHR-BASED TARGET TRIAL EMULATION FROM THE N3C AND RECOVER CONSORTIA Effect of Paxlovid treatment during acute COVID-19 on Long COVID onset: An EHR-based target trial emulation from the N3C and RECOVER consortia Nirmatrelvir Plus Ritonavir for Early COVID-19 in a Large U.S. Health System : A Population-Based Cohort Study Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19 Effect of Nirmatrelvir/Ritonavir (Paxlovid) on Hospitalization among Adults with COVID-19: an EHR-based Target Trial Emulation from N3C Effectiveness of Paxlovid in Reducing Severe Coronavirus Disease 2019 and Mortality in High-Risk Patients Impact of the Use of Oral Antiviral Agents on the Risk of Hospitalization in Community Coronavirus Disease 2019 Patients (COVID-19) Nirmatrelvir or Molnupiravir Use and Severe Outcomes From Omicron Infections Effect of Oral Nirmatrelvir on Long COVID Symptoms: 4 Cases and Rationale for Systematic Studies The use of nirmatrelvir-ritonavir in a case of breakthrough long COVID Case report: Treatment of long COVID with a SARS-CoV-2 antiviral and IL-6 blockade in a patient with rheumatoid arthritis and SARS-CoV-2 antigen persistence Impact of extended-course oral nirmatrelvir/ritonavir (Paxlovid) in established Long COVID: Case series and research considerations Paxlovid as a potential treatment for long COVID Nirmatrelvir-Ritonavir and Symptoms in Adults With Postacute Sequelae of SARS-CoV-2 Infection: The STOP-PASC Randomized Clinical Trial Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Neurologic Manifestations of

Long COVID Differ Based on Acute COVID-19 Severity Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Prevalence of post-COVID-19 symptoms in hospitalized and non-hospitalized COVID-19 survivors: A systematic review and meta-analysis Association of Treatment With Nirmatrelvir and the Risk of Post-COVID-19 Condition Association of nirmatrelvir for acute SARS-CoV-2 infection with subsequent Long COVID symptoms in an observational cohort study Platform adaptive trial of novel antivirals for early treatment of COVID-19 In the community (PANORAMIC): protocol for a randomised, controlled, open-label, adaptive platform trial of community novel antiviral treatment of COVID-19 in people at increased risk of more severe disease The National COVID Cohort Collaborative (N3C): Rationale, design, infrastructure, and deployment Synergies between centralized and federated approaches to data quality: a report from the national COVID cohort collaborative Using Big Data to Emulate a Target Trial When a Randomized Trial Is Not Available Target Trial Emulation: A Framework for Causal Inference From Observational Data COVID-19 in children. II: Pathogenesis, disease spectrum and management Pediatric Nirmatrelvir/Ritonavir Prescribing Patterns During the COVID-19 Pandemic Reengineering a machine learning phenotype to adapt to the changing COVID-19 landscape: A study from the N3C and RECOVER consortia Estimated Global Proportions of Individuals With Persistent Fatigue, Cognitive, and Respiratory Symptom Clusters Following Symptomatic COVID-19 in 2020 and 2021 Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies Development of a definition of postacute sequelae of SARS-CoV-2 infection Why we need a deeper understanding of the pathophysiology

of long COVID Traumatic Brain Injury and Long-Term Risk of Stroke Among US Military Veterans Asbestos Exposure and Severity of COVID-19 Measurement in Veterans Affairs Health Services Research: veterans as a special population Long COVID risk and pre-COVID vaccination in an EHR-based cohort study from the RECOVER program Administrative database research infrequently used validated diagnostic or procedural codes Matching methods for causal inference: A review and a look forward Misunderstandings between experimentalists and observationalists about causal inference Association Between Early Treatment With Tocilizumab and Mortality Among Critically Ill Patients With COVID-19 Estimates of Overall Survival in Patients With Cancer Receiving Different Treatment Regimens: Emulating Hypothetical Target Trials in the Surveillance, Epidemiology, and End Results (SEER)-Medicare Linked Database Avoidable flaws in observational analyses: an application to statins and cancer Emulating a Novel Clinical Trial Using Existing Observational Data. Predicting Results of the PreVent Study WHO Clinical Case Definition Working Group on Post-COVID-19 Condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus Long COVID: aiming for a consensus Strategies for handling missing data in electronic health record derived data Out-of-system Care and Recording of Patient Characteristics Critical for Comparative Effectiveness Research Evaluating Risk-Prediction Models Using Data from Electronic Health Records Comorbidity and its Impact on Patients with COVID-19 Epidemiological, comorbidity factors with severity and prognosis of COVID-19: a systematic review and meta-analysis The Association between Presence of Comorbidities and COVID-19 Severity: A Systematic Review and Meta-Analysis medRxiv. 2025. [PMC10854326](https://doi.org/10.1101/2025.01.14.25108543)

38. Yan Xie, Taeyoung Choi, Ziyad Al-Aly. Association of Treatment With Nirmatrelvir and the Risk of Post-COVID-19 Condition While Waiting for a Randomized Clinical Trial of Nirmatrelvir for Prevention of Post-COVID-19 Condition. High-dimensional characterization of post-acute sequelae of COVID-19 Long COVID after breakthrough SARS-CoV-2 infection Long COVID: major findings, mechanisms and recommendations The immunology and immunopathology of COVID-19 Long-COVID treatments: why the world is still waiting Oral nirmatrelvir for high-risk, nonhospitalized adults with Covid-19 Making neighborhood-disadvantage metrics accessible—the neighborhood atlas Risk of persistent and new clinical sequelae among adults aged 65 years and older during the post-acute phase of SARS-CoV-2 infection: retrospective cohort study Post-COVID conditions among adult COVID-19 survivors aged 18-64 and ≥ 65 years—United States, March 2020-November 2021 Risk of clinical sequelae after the acute phase of SARS-CoV-2 infection: retrospective cohort study Post-COVID syndrome in individuals admitted to hospital with COVID-19: retrospective cohort study Persistent symptoms in patients after acute COVID-19 Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Incidence, co-occurrence, and evolution of long-COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19 Kidney outcomes in long COVID Risks of mental health outcomes in people with COVID-19: cohort study Characterizing long COVID in an international cohort: 7 months of symptoms and their impact 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records Risks and burdens of incident diabetes in long COVID: a cohort study Long-term neurologic outcomes of COVID-19

Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Long-term gastrointestinal outcomes of COVID-19 Acute and postacute sequelae associated with SARS-CoV-2 reinfection Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Comparative evaluation of clinical manifestations and risk of death in patients admitted to hospital with COVID-19 and seasonal influenza: cohort study Acute kidney injury in a national cohort of hospitalized US veterans with COVID-19 Molnupiravir and risk of hospitalization or death: a target trial emulation using the US Veterans Affairs healthcare databases Long COVID: long-term health outcomes and implications for policy and research Diabetes after SARS-CoV-2 infection Multiple pathways for SARS-CoV-2 resistance to nirmatrelvir JAMA Internal Medicine. 2023. [PMC10037200](https://pubmed.ncbi.nlm.nih.gov/40037200/)

39. Yusuff Adebayo Adebisi, Anas Ali Alhur, Najim Z. Alshahrani, Victor C. Cañez, Edgar G. Cue, Don Eliseo Lucero-Prisno. Pre-pandemic diabetes and risk of long COVID: longitudinal evidence More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Short- and potential long-term adverse health outcomes of COVID-19: a rapid review Long COVID. Major findings, mechanisms and recommendations Defining long COVID: going back to the start Long COVID: mechanisms, risk factors and recovery Risk factors for long coronavirus disease 2019 (long COVID) among healthcare personnel, Brazil, 2020–2022 Long COVID: a review and proposed visualization of the complexity of long COVID Prevalence, risk factors and characterisation of individuals with long COVID using electronic health records in over 1.5 million COVID cases in England Insulin resistance in long COVID-19 syndrome Association of glycemic control

with long COVID in patients with type 2 diabetes: findings from the National COVID cohort collaborative (N3C) Prevalence of long COVID and the impact on diabetes management and physical activity participation in adults with type 2 diabetes: an Australia-wide cross-sectional online survey The bidirectional association between diabetes and long COVID-19: a systematic review Risks and burdens of incident diabetes in long COVID: a cohort study Variation in appropriate diabetes care and treatment targets in urban and rural areas in england: an observational study of the 'rule of halves' Self-reported diabetes and quality of life: findings from a general population survey with the short Form-12 (SF-12) health survey Validity and reliability of a self-reported measure of medication adherence in patients with type 2 diabetes mellitus in Singapore Validity of diabetes self-reports in the women's health initiative Defining the subtypes of long COVID and risk factors for prolonged disease: population-based case-crossover study Long-COVID clinical features and risk factors: a retrospective analysis of patients from the STOP-COVID registry of the PoLoCOV study Prevalence and risk factors for long COVID among adults in Scotland using electronic health records: a national, retrospective, observational cohort study A modified poisson regression approach to prospective studies with binary data Diabetes as a risk factor for greater COVID-19 severity and in-hospital death: a meta-analysis of observational studies Risk factors associated with post-COVID-19 conditions among diabetes patients in the United Arab Emirates Association between diabetes severity and risks of COVID-19 infection and outcomes Diabetes is a risk factor for the progression and prognosis of COVID-19 Why is COVID-19 more severe in patients with diabetes? The role of angiotensin-converting enzyme 2, endothelial dysfunction and the

immunoinflammatory system Long-term persistence of COVID-induced hyperglycemia: a cohort study COVID-19 and diabetes: a comprehensive review of angiotensin converting enzyme 2, mutual effects and pharmacotherapy Journal of Diabetes and Metabolic Disorders. 2025. [PMC12449287](#)

40. Sarah Cuschieri, Piotr Wilk. Does Pre-existing Diabetes Correlate with Long COVID-19 in Europe? Evidence from the Analysis of the Survey of Health, Ageing and Retirement in Europe's Corona Surveys The prevalence and long-term health effects of long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis A clinical case definition of post-COVID-19 condition by a Delphi consensus Pathophysiology and mechanism of long COVID: a comprehensive review Global prevalence of post-coronavirus disease 2019 (COVID-19) condition or long COVID: a meta-analysis and systematic review Post COVID-19 syndrome ("long COVID") and diabetes: challenges in diagnosis and management More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Long COVID - metabolic risk factors and novel therapeutic management Long-COVID, metabolic and endocrine disease Proposed integrative model for post-COVID symptoms Type 2 diabetes - an unresolved disease across centuries contributing to a public health emergency 2. Classification and Diagnosis of Diabetes:Standards of Care in Diabetes—2023 Diabetes and infection: is there a link?- a mini-review Plasma glucose levels and diabetes are independent predictors for mortality and morbidity in patients with SARS Insulin receptor signaling in normal and insulin-resistant states Mechanisms of COVID-19 pathogenesis in diabetes Diabetes and the risk of long-term post-COVID symptoms Age, sex and previous comorbidities as risk factors not associated with SARS-CoV-2 infection for long COVID-19: a

systematic review and meta-analysis Data resource profile: the Survey of Health, Ageing and Retirement in Europe (SHARE) COVID-19 reinfections among naturally infected and vaccinated individuals Diabetes and COVID-19: short- and long-term consequences Female gender is associated with long COVID syndrome: a prospective cohort study Follow-up of adults with noncritical COVID-19 two months after symptom onset Persisting symptoms after COVID-19 Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Sex differences in immune responses Sex differences in mortality from COVID-19 pandemic: are men vulnerable and women protected? Sex differences in sequelae from COVID-19 infection and in long COVID syndrome: a review Long COVID risk - a signal to address sex hormones and women's health Sex differences in metabolic regulation and diabetes susceptibility Sex-based differences in diabetes prevalence and risk factors: a population-based cross-sectional study among low-income adults in China Sex and gender differences in risk, pathophysiology and complications of type 2 diabetes mellitus Influence of gender in diabetes mellitus and its complication Clinical characteristics with inflammation profiling of long COVID and association with 1-year recovery following hospitalisation in the UK: a prospective observational study Characteristics of long-COVID among older adults: a cross-sectional study Preexisting hypertension is associated with a greater number of long-term post-COVID symptoms and poor sleep quality: a case-control study Journal of Diabetes Research. 2024. [PMC10857882](#)

41. Xueying Yang, Fanghui Shi, Hao Zhang, William A. Giang, Amandeep Kaur, Hui Chen, Xiaoming Li, et al.. Long COVID among people with HIV: A systematic review and meta-analysis Long COVID among people with HIV: A

systematic review and meta-analysis Postacute sequelae and adaptive immune responses in people with HIV recovering from SARS-CoV-2 infection Health status, persistent symptoms, and effort intolerance one year after acute COVID-19 infection Long COVID in people living with HIV HIV and inflammatory markers are associated with persistent COVID-19 symptoms Chronic viral coinfections differentially affect the likelihood of developing long COVID Factors associated with severe COVID-19 and post-acute COVID-19 syndrome in a cohort of people living with HIV on antiretroviral treatment and with undetectable HIV RNA Long-coronavirus disease among people living with HIV in western India: an observational study Prolonged shedding of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) at high viral loads among hospitalized immunocompromised persons living with human immunodeficiency virus (HIV), South Africa A longitudinal analysis of SARS-CoV-2 antibody responses among people with HIV Clinical characteristics of COVID-19 patients with HIV coinfection in Wuhan, China Epidemiological, Virological and serological features of coronavirus disease 2019 (COVID-19) cases in people living with human immunodeficiency virus in Wuhan: a population-based cohort study Long-lasting adaptive immune memory specific to SARS-CoV-2 in convalescent coronavirus disease 2019 stable people with HIV The prevalence and long-term health effects of long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis Development and validation of the symptom burden questionnaire for long covid (SBQ-LC): Rasch analysis A core outcome set for post-COVID-19 condition in adults for use in clinical practice and research: an international Delphi consensus study Associations between HIV infection and

clinical spectrum of COVID-19: a population level analysis based on US national COVID cohort collaborative (N3C) data
The significance of interferon- γ in HIV-1 pathogenesis, therapy, and prophylaxis
Why aren't people living with HIV at higher risk for developing severe coronavirus disease 2019 (COVID-19)?
Markers of immune activation and inflammation in individuals with postacute sequelae of severe acute respiratory syndrome coronavirus 2 infection
Plasma markers of neurologic injury and inflammation in people with self-reported neurologic postacute sequelae of SARS-CoV-2 infection
The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19
Multiple early factors anticipate post-acute COVID-19 sequelae
IFN- β plays both pro-and anti-inflammatory roles in the rat cardiac fibroblast through differential STAT protein activation
The combination of IFN β and TNF induces an antiviral and immunoregulatory program via non-canonical pathways involving STAT2 and IRF9
Intramuscular interferon beta-1a for disease progression in relapsing multiple sclerosis
Th17 and Treg cells function in SARS-CoV2 patients compared with healthy controls
Imbalance of regulatory and cytotoxic SARS-CoV-2-reactive CD4+ T cells in COVID-19
Clinical characteristics with inflammation profiling of long COVID and association with 1-year recovery following hospitalisation in the UK: a prospective observational study
Ageing and inflammation in patients with HIV infection
Peripheral immune dysregulation in the ART era of HIV-associated neurocognitive impairments: a systematic review
Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review
HIV Medicine. 2024. [PMC11725417](#)

42. Kellie L Hawkins, Dima Dandachi, Zoe Verzani, M Daniel Brannock, Colby Lewis, Sajjad Abedian, Sohrab Jaferian,

Shannon Wuller, Jennifer Truong, Margot Gage Witvliet, Gretchen Dymond, Hemalkumar B Mehta, Payal B Patel, Elaine Hill, Mark G Weiner, Thomas W Carton, Rainu Kaushal, Elen Feuerriegel, Huong G Tran, Kristen Marks, Carlos R Oliveira, Edward M Gardner, Igho Ofotokun, Roy M Gulick, Kristine M Erlandson, et al.. HIV Infection and Long COVID: A RECOVER Program, Electronic Health Record-Based Cohort Study Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America. 2025. [PMC12497945](#)

43. Raphael S. Peter, Alexandra Nieters, Stefan O. Brockmann, Siri Göpel, Gerhard Kindle, Uta Merle, Jürgen M. Steinacker, Winfried V. Kern, Dietrich Rothenbacher, et al.. Association of BMI with general health, working capacity recovered, and post-acute sequelae of COVID-19 Association of BMI with general health, working capacity recovered, and post-acute sequelae of COVID-19 A nationwide questionnaire study of post-acute symptoms and health problems after SARS-CoV-2 infection in Denmark Lifelines Corona Research Initiative. Persistence of somatic symptoms after COVID-19 in The Netherlands: an observational cohort study Obesity in patients younger than 60 years is a risk factor for COVID-19 hospital admission Risk factors for COVID-19-related death, hospitalization and intensive care: a population-wide study of all inhabitants in Stockholm Obesity, ethnicity, and risk of critical care, mechanical ventilation, and mortality in patients admitted to hospital with COVID-19: analysis of the ISARIC CCP-UK cohort Persistent COVID-19 symptoms in a community study of 606, 434 people in England Obesity and lipid metabolism disorders determine the risk for development of long COVID syndrome: a cross-sectional study from 50,402 COVID-19 patients Post-COVID-19 syndrome in outpatients: a cohort study More than 50 long-term effects of

COVID-19: a systematic review and meta-analysis Body mass index and risk for COVID-19-related hospitalization, intensive care unit admission, invasive mechanical ventilation, and death—United States, March–December 2020 Body mass index and severity/fatality from coronavirus disease 2019: a nationwide epidemiological study in Korea Validation of short measures of work ability for research and employee surveys Obesity (Silver Spring, Md.). 2022. [PMC9874798](#)

44. Ting Zhou, Bingyu Zhang, Dazheng Zhang, Qiong Wu, Jiajie Chen, Lu Li, Yiwen Lu, Michael J. Becich, Saul Blecker, Nymisha Chilukuri, Elizabeth A. Chrischilles, Haitao Chu, Leonor Corsino, Carol R. Geary, Mady Hornig, Maxwell M. Hornig-Rohan, Susan Kim, David M. Liebovitz, Vitaly Lorman, Chongliang Luo, Hiroki Morizono, Abu S. M. Mosa, Nathan M. Pajor, Suchitra Rao, Hanieh Razzaghi, Srinivasan Suresh, Yacob G. Tedla, Leah Vance Utset, Youfa Wang, David A. Williams, Margot Gage Witvliet, Caren Mangarelli, Ravi Jhaveri, Christopher B. Forrest, Yong Chen, et al.. Body Mass Index and Postacute Sequelae of SARS-CoV-2 Infection in Children and Young Adults NICE guideline on long COVID A clinical case definition of post-COVID-19 condition by a Delphi consensus A proposed framework and timeline of the spectrum of disease due to SARS-CoV-2 infection: illness beyond acute infection and public health implications Management of post-acute covid-19 in primary care High-dimensional characterization of post-acute sequelae of COVID-19 Post-acute COVID-19 syndrome Post-acute COVID-19 syndrome Author correction: long COVID: major findings, mechanisms and recommendations Risk factors for post-COVID-19 condition (Long Covid) in children: a prospective cohort study Long-COVID in children and adolescents: a systematic review and meta-analyses A systematic review of persistent clinical features after SARS-

CoV-2 in the pediatric population Prevalence and risk factor for long COVID in children and adolescents: a meta-analysis and systematic review Body mass index and risk for COVID-19-related hospitalization, intensive care unit admission, invasive mechanical ventilation, and death - United States, March-December 2020 Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Symptoms and risk factors for long COVID in non-hospitalized adults Adherence to healthy lifestyle prior to infection and risk of post-COVID-19 condition Attributes and predictors of long COVID Risk factors for post-COVID-19 condition in previously hospitalised children using the ISARIC global follow-up protocol: a prospective cohort study Long COVID symptoms in SARS-CoV-2-positive adolescents and matched controls (LongCOVIDKidsDK): a national, cross-sectional study Clinical features and burden of postacute sequelae of SARS-CoV-2 infection in children and adolescents STROBE reporting guidelines for observational studies Age limit of pediatrics Pediatric medical complexity algorithm: a new method to stratify children by medical complexity Severity of acute COVID-19 in children <18 years old March 2020 to December 2021 A modified Poisson regression approach to prospective studies with binary data Changes in BMI during the COVID-19 pandemic Reduced sensitivity of antibody tests after omicron infection Risk factors associated with post-COVID-19 condition: a systematic review and meta-analysis Identifying who has long COVID in the USA: a machine learning approach using N3C data Trajectory of long covid symptoms after covid-19 vaccination: community based cohort study COVID-19 vaccination for the prevention and treatment of long COVID: a systematic review and meta-analysis Empirical confidence interval calibration for population-level effect estimation studies in observational

healthcare data Interpreting observational studies: why empirical calibration is needed to correct P values Coding long COVID: characterizing a new disease through an ICD-10 lens Obesity and COVID-19: what makes obese host so vulnerable? Overweight and obesity as risk factors for COVID-19-associated hospitalisations and death: systematic review and meta-analysis Evaluation of unintended consequences of COVID-19 pandemic restrictions and obesity prevalence among youths Obesity and COVID-19: immune and metabolic derangement as a possible link to adverse clinical outcomes Cell death and inflammation during obesity: "Know my methods, WAT(son)" NLRP3 inflammasome: the stormy link between obesity and COVID-19 Long covid-mechanisms, risk factors, and management Chronic inflammation, neutrophil activity, and autoreactivity splits long COVID A review of the relationship between gut microbiome and obesity Gut microbiota dynamics in a prospective cohort of patients with post-acute COVID-19 syndrome Autoimmunity as a sequela to obesity and systemic inflammation Autoantibody production in obesity: is there evidence for a link between obesity and autoimmunity? Long COVID or post-acute sequelae of COVID-19 (PASC): an overview of biological factors that may contribute to persistent symptoms Multiple early factors anticipate post-acute COVID-19 sequelae Mechanisms of thrombosis in obesity Endothelial dysfunction in obesity-induced inflammation: molecular mechanisms and clinical implications Risks of deep vein thrombosis, pulmonary embolism, and bleeding after covid-19: nationwide self-controlled cases series and matched cohort study Collider bias undermines our understanding of COVID-19 disease risk and severity JAMA Network Open. 2024. [PMC11581483](https://pubmed.ncbi.nlm.nih.gov/37115814/)

45. Ting Zhou, Bingyu Zhang, Dazheng Zhang, Ravi Jhaveri, Jiajie Chen, Michael J. Becich, Leah Castro, Yu Chen, Nymisha Chilukuri, Sharon J. Herring, Yuqing Lei, Lu Li, Yiwen Lu, Maxwell Hornig, Amrik Singh Khalsa, David Liebovitz, Abu Saleh Mohammad Mosa, Bradley W. Taylor, Yacob G. Tedla, Drew Thodeson, Jiayi Tong, Qiong Wu, Christopher B. Forrest, Yong Chen, et al.. Pre-COVID-19 body mass index and post-acute cardiovascular, gastrointestinal, and neuropsychiatric outcomes among children and young adults with SARS-CoV-2 infection: An EHR-based cohort study from the RECOVER Initiative Obesity could shift severe COVID-19 disease to younger ages Underlying medical conditions associated with severe COVID-19 illness among children Adherence to healthy lifestyle prior to infection and risk of post-COVID-19 condition Body mass index and postacute sequelae of SARS-CoV-2 infection in children and young adults Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss Inflammatory mechanisms in the regulation of insulin resistance The role of inflammation in depression: from evolutionary imperative to modern treatment target Global pandemics interconnected — obesity, impaired metabolic health and COVID-19 Why is COVID-19 less severe in children? A review of the proposed mechanisms underlying the age-related difference in severity of SARS-CoV-2 infections Individuals with obesity and COVID-19: a global perspective on the epidemiology and biological relationships Trends in obesity prevalence among children and adolescents aged 2 to 19 years in the US from 2011 to 2020 SARS-CoV-2 infection drives an inflammatory response in human adipose tissue through infection of adipocytes and macrophages Is adipose tissue a reservoir for viral spread, immune activation, and cytokine amplification in coronavirus disease 2019? Diabetes mellitus, energy metabolism, and

COVID-19 Pediatric gastrointestinal tract outcomes during the postacute phase of COVID-19 Risk of neuropsychiatric and related conditions associated with SARS-CoV-2 infection: a difference-in-differences analysis Cardiovascular post-acute sequelae of SARS-CoV-2 in children and adolescents: cohort study using electronic health records How do we share data in COVID-19 research? A systematic review of COVID-19 datasets in PubMed Central Articles Age limit of pediatrics Racial/ethnic differences in post-acute sequelae of SARS-CoV-2 in children and adolescents in the United States Post-acute dyslipidemia and abnormal BMI in children and adolescents with COVID-19: a cohort study from the RECOVER initiative Reinfection with SARS-CoV-2 in the Omicron era is associated with increased risk of post-acute sequelae of SARS-CoV-2 infection: a RECOVER-EHR cohort study Real-world effectiveness and causal mediation study of BNT162b2 on long COVID risks in children and adolescents Kidney function following COVID-19 in children and adolescents Clinical features and burden of postacute sequelae of SARS-CoV-2 infection in children and adolescents Severity of acute COVID-19 in children < 18 years old March 2020 to December 2021 Pediatric medical complexity algorithm: a new method to stratify children by medical complexity A modified poisson regression approach to prospective studies with binary data A simulation study of the number of events per variable in logistic regression analysis Sparse data bias: a problem hiding in plain sight Approach to obesity treatment in primary care Biases in electronic health record data due to processes within the healthcare system: retrospective observational study Leveraging electronic health records for data science: common pitfalls and how to avoid them Risk factors for severe COVID-19 in children: a systematic review and meta-analysis Obesity and

cardiovascular disease: a scientific statement from the American Heart Association Cardiac involvement in the long-term implications of COVID-19 Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19 Human gut microbes associated with obesity The intestine in acute and long COVID: pathophysiological insights and key lessons Are we facing a crashing wave of neuropsychiatric sequelae of COVID-19? Neuropsychiatric symptoms and potential immunologic mechanisms The role of inflammation in depression: from evolutionary imperative to modern treatment target Multisystem inflammatory syndrome in U.S. children and adolescents Pervasiveness, impact and implications of weight stigma Stigma experienced by children and adolescents with obesity Obesity in adolescents The Journal of infection. 2026. [PMC13093859](#)

46. Ting Zhang, Zhimao Li, Qimin Mei, Joseph Harold Walline, Zhaocai Zhang, Yecheng Liu, Huadong Zhu, Bin Du, et al.. Cardiovascular outcomes in long COVID-19: a systematic review and meta-analysis An interactive web-based dashboard to track COVID-19 in real time Authorship correction: international changes in COVID-19 clinical trajectories across 315 hospitals and 6 countries: retrospective cohort study Long COVID: let patients help define long-lasting COVID symptoms Why we need to keep using the patient made term “long COVID” Cardiac involvement in consecutive unselected hospitalized COVID-19 population: in-hospital evaluation and one-year follow-up Echocardiographic assessment of COVID19 sequelae in survivors with elevated cardiac biomarkers Endothelial dysfunction in COVID-19: a potential predictor of long-COVID? Cardiovascular complications in the post-acute COVID-19 syndrome (PACS) Cardiovascular disease and COVID-19: a consensus paper from the ESC

working group on coronary pathophysiology & microcirculation, ESC working group on thrombosis and the association for acute cardiovascular care (ACVC), in collaboration with the European heart rhythm association (EHRA) Persistent symptoms following SARS-CoV-2 infection amongst children and young people: a meta-analysis of controlled and uncontrolled studies Deriving and validating a risk prediction model for long COVID-19: protocol for an observational cohort study using linked Scottish data Post-acute COVID-19 syndrome Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group The PRISMA 2020 statement: an updated guideline for reporting systematic reviews Ottawa Hospital Research Institute Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses Meta-analysis in clinical trials Practical methods for incorporating summary time-to-event data into meta-analysis In meta-analyses of proportion studies, funnel plots were found to be an inaccurate method of assessing publication bias Post-COVID syndrome in individuals admitted to hospital with COVID-19: retrospective cohort study Rates of serious clinical outcomes in survivors of hospitalisation with COVID-19: a descriptive cohort study within the OpenSAFELY platform Risk of persistent and new clinical sequelae among adults aged 65 years and older during the post-acute phase of SARS-CoV-2 infection: retrospective cohort study Symptoms and risk factors for long COVID in non-hospitalized adults Long COVID 12 months after discharge: persistent symptoms in patients hospitalised due to COVID-19 and patients hospitalised due to other causes—a multicentre cohort study Risk of clinical sequelae after the acute phase of SARS-CoV-2 infection: retrospective cohort

study Cardiometabolic outcomes up to 12 months after COVID-19 infection. A matched cohort study in the UK Long-term cardiovascular outcomes of COVID-19 Risk of acute myocardial infarction and ischaemic stroke following COVID-19 in Sweden: a self-controlled case series and matched cohort study Comparative evaluation of clinical manifestations and risk of death in patients admitted to hospital with COVID-19 and seasonal influenza: cohort study Extrapulmonary manifestations of COVID-19 The road to addressing long COVID Endothelial dysfunction in acute and long standing COVID-19: a prospective cohort study Pathological features of COVID-19-associated myocardial injury: a multicentre cardiovascular pathology study Cardiovascular complications of COVID-19 COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets Count the cost of disability caused by COVID-19 COVID-19 and cardiovascular disease: from basic mechanisms to clinical perspectives COVID-19 and cardiovascular disease COVID-19, microangiopathy, hemostatic activation, and complement Endothelial cell infection and endotheliitis in COVID-19 Report: long-term immunological health consequences of COVID-19 Epidemiology, prognosis, and clinical manifestation of cardiovascular disease in COVID-19 Reduced incidence of long coronavirus disease referrals to the Cambridge university teaching hospital long coronavirus disease clinic Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID symptom study app: a prospective, community-based, nested, case-control study Association between BNT162b2 vaccination and reported incidence of post-COVID-19 symptoms: cross-sectional study 2020-21, Israel Six-month sequelae of post-vaccination SARS-CoV-2 infection: a retrospective cohort study of 10,024

breakthrough infections *Frontiers in Cardiovascular Medicine*. 2025. [PMC11814196](#)

47. Pia Lindberg, Samuel Wiqvist, Maria Juszczuk, Seika Lee, Marta A. Kisiel, Caroline Wachtler, Marcus Ståhlberg, Åsa M. Wheelock, Artur Fedorowski, Axel C. Carlsson, et al.. Long COVID and risk of incident cardiovascular disease: a prospective cohort study using the Multimorbidity Integrated Registry Across Care Levels in Stockholm (MIRACLE-S) cohort Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Long-term cardiovascular outcomes of COVID-19 Cardiovascular autonomic dysfunction in post-COVID-19 syndrome: a major health-care burden High-dimensional characterization of post-acute sequelae of COVID-19 Chronic autonomic symptom burden in long-COVID: a follow-up cohort study Mechanisms of long COVID and the path toward therapeutics Long-haul post-COVID-19 symptoms presenting as a variant of postural orthostatic tachycardia syndrome: the Swedish experience Using machine learning involving diagnoses and medications as a risk prediction tool for post-acute sequelae of COVID-19 (PASC) in primary care International statistical classification of diseases and related health problems (ICD) 1 September 2020 High prevalence of diagnosis of diabetes, depression, anxiety, hypertension, asthma and COPD in the total population of Stockholm, Sweden - a challenge for public health A clinical case definition of post-COVID-19 condition by a Delphi consensus Clinical management of COVID-19: living guideline, 18 August 2023 2023 Depression or anxiety and all-cause mortality in adults with atrial fibrillation--A cohort study in Swedish primary care Anxiety as a risk factor for cardiovascular disease independent of depression: a narrative review of current status and conflicting findings Insights into social disparities in smoking prevalence using Mosaic, a novel

measure of socioeconomic status: an analysis using a large primary care dataset Preexisting symptoms increase the risk of developing long COVID during the SARS-CoV-2 pandemic Long COVID science, research and policy Postural orthostatic tachycardia syndrome (POTS) and other autonomic disorders after COVID-19 infection: a case series of 20 patients COVID-19 is, in the end, an endothelial disease Persistent complement dysregulation with signs of thromboinflammation in active long covid Long COVID: pathophysiological factors and abnormalities of coagulation Prevalence and clinical impact of postural orthostatic tachycardia syndrome in highly symptomatic long COVID Gender differences in the utilization of health care services National plan for vaccination against COVID-19 (in Swedish). The Public Health Agency of Sweden The Dahlgren-Whitehead model of health determinants: 30 years on and still chasing rainbows Simply the best? The impact of quality on choice of primary healthcare provider in Sweden Major risk factors as antecedents of fatal and nonfatal coronary heart disease events eClinicalMedicine. 2026. [PMC13133534](#)

48. Akash Pradeep, Martha L Carvour, Srija M Sivamurugan, Upinder Singh, Alejandro P Comellas, Sanjana Dayal. Cardiovascular, renal and pulmonary risks of Long COVID: a retrospective cohort study stratified by age and sex medRxiv. 2026. [PMC12870583](#)
49. Kwanjoo Lee, Jaeyu Park, Jinseok Lee, Myeongcheol Lee, Hyeon Jin Kim, Yejun Son, Sang Youl Rhee, Lee Smith, Masoud Rahmati, Jiseung Kang, Hayeon Lee, Yeonjung Ha, Dong Keon Yon, et al.. Long-term gastrointestinal and hepatobiliary outcomes of COVID-19: A multinational population-based cohort study from South Korea, Japan, and the UK Association between mental illness and COVID-19 in

South Korea: a post-hoc analysis Global, regional, and national incidence and mortality of COVID-19 in 237 countries and territories, January 2022: a systematic analysis for World Health Organization COVID-19 Dashboard Long COVID after breakthrough SARS-CoV-2 infection Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and metaanalysis Post-infectious irritable bowel syndrome Long-term gastrointestinal outcomes of COVID-19 The effect of SARS-CoV-2 infection on cardiac function in post-COVID-19 survivors: a systematic review and meta-analysis Changes in the incidence rates of gastrointestinal diseases due to the COVID-19 pandemic in south korea: a long-term perspective Autoimmune inflammatory rheumatic diseases and COVID-19 outcomes in South Korea: a nationwide cohort study Proton pump inhibitors and the risk of severe COVID-19: a post-hoc analysis from the Korean nationwide cohort Acute and post-acute respiratory complications of SARS-CoV-2 infection: population-based cohort study in South Korea and Japan Severe COVID-19 outcomes after full vaccination of primary schedule and initial boosters: pooled analysis of national prospective cohort studies of 30 million individuals in England, Northern Ireland, Scotland, and Wales Efficacy and safety of baricitinib or ravulizumab in adult patients with severe COVID-19 (TACTIC-R): a randomised, parallel-arm, open-label, phase 4 trial Propensity score matching for causal inference and reducing the confounding effects: statistical standard and guideline of Life Cycle Committee Shortand long-term neuropsychiatric outcomes in long COVID in South Korea and Japan Residual SARS-CoV-2 viral antigens detected in GI and hepatic tissues from five recovered patients with COVID-19 Intestinal host response to SARSCoV-2 infection and COVID-19 outcomes in patients with gastrointestinal

symptoms Evidence for gastrointestinal infection of SARS-CoV-2 Global burden of vaccine-associated hepatobiliary and gastrointestinal adverse drug reactions, 1967-2023: a comprehensive analysis of the international pharmacovigilance database Potential long coronavirus disease 2019 gastrointestinal symptoms 6 months after coronavirus infection are associated with mental health symptoms Molecular consequences of SARSCoV-2 liver tropism Gut microbiota dynamics in a prospective cohort of patients with post-acute COVID-19 syndrome Gastrointestinal symptoms of 95 cases with SARS-CoV-2 infection Meta-analysis: Post-COVID-19 functional dyspepsia and irritable bowel syndrome Differences in the COVID-19 pandemic response between South Korea and the United States: a comparative analysis of culture and policies The impact of national culture on the increase of COVID-19: a cross-country analysis of European countries Ethnic differences in SARS-CoV-2 infection and COVID-19-related hospitalisation, intensive care unit admission, and death in 17 million adults in England: an observational cohort study using the OpenSAFELY platform Positive predictive value of international classification of diseases, 10th revision, codes for cirrhosis and its related complications Epidemiologic characteristics of cases with reinfection, recurrence, and hospital readmission due to COVID-19: a systematic review and meta-analysis The effect of adherence to high-quality dietary pattern on COVID-19 outcomes: a systematic review and meta-analysis Clinical and Molecular Hepatology. 2024. [PMC11540398](#)

50. Yuying Ma, Lijun Zhang, Rui Wei, Weiyu Dai, Ruijie Zeng, Dongling Luo, Rui Jiang, Zewei Zhuo, Qi Yang, Jingwei Li, Felix W Leung, Chongyang Duan, Weihong Sha, Hao Chen, et al.. Risks of digestive diseases in long COVID: evidence from

a population-based cohort study Moderate to severe SARS-CoV-2 infection primes vaccine-induced immunity more effectively than asymptomatic or mild infection Post-acute COVID-19 syndrome Association of COVID-19 with short- and long-term risk of cardiovascular disease and mortality: a prospective cohort in UK Biobank Kidney outcomes in long COVID Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Risks and burdens of incident diabetes in long COVID: a cohort study Gastrointestinal involvement in COVID-19: a systematic review and meta-analysis Prevalence of gastrointestinal symptoms and fecal viral shedding in patients with Coronavirus Disease 2019: a systematic review and meta-analysis Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and meta-analysis Persistent cholestatic injury and secondary sclerosing cholangitis in COVID-19 Patients Long-term gastrointestinal sequelae following COVID-19: a prospective follow-up cohort study Long-term gastrointestinal outcomes of COVID-19 Female sex affects respiratory function and exercise ability in patients recovered from COVID-19 pneumonia Sex-bias in COVID-19-associated illness severity and mortality in cancer patients: a systematic review and meta-analysis Association between BNT162b2 vaccination and reported incidence of post-COVID-19 symptoms: cross-sectional study 2020-21, Israel Long COVID after breakthrough SARS-CoV-2 infection Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: a systematic review Comparison of sociodemographic and health-related characteristics of UK biobank participants with those of the general population Long COVID-19 liver manifestation in children Risk of clinical sequelae after the acute phase of SARS-CoV-2 infection: retrospective cohort study COVID-19

and the gastrointestinal tract in children Gastrointestinal, liver, pancreas, oral and psychological long-term symptoms of COVID-19 after recovery: a review Long-term cardiovascular effects of COVID-19: emerging data relevant to the cardiovascular clinician Gastrointestinal symptoms in diabetes: prevalence, assessment, pathogenesis, and management Definitions for coronavirus disease 2019 reinfection, relapse and PCR re-positivity Effect of ovarian sex hormones on non-steroidal anti-inflammatory drug-induced gastric lesions in female rats Is the oral contraceptive or hormone replacement therapy a risk factor for cholelithiasis: a systematic review and meta-analysis Sex Differences in the exocrine pancreas and associated diseases Sex-related differences in GI disorders 12-month post-discharge liver function test abnormalities among patients with COVID-19: a single-center prospective cohort study Long covid outcomes at one year after mild nationwide cohort study Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding Digestive symptoms of COVID-19 and expression of ACE2 in digestive tract organs COVID-19 and the digestive system ACE2 expression in pancreas may cause pancreatic damage after SARS-CoV-2 infection Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China Mechanisms of disease: mechanisms and clinical implications of cholestasis in sepsis Cholestatic alterations in the critically ill: some new light on an old problem Changes in food and drink purchasing behaviour in England during the first 3 months of the COVID-19 pandemic: an interrupted time-series analysis A roadmap to using historical controls in clinical trials - by Drug Information Association Adaptive Design Scientific Working Group (DIA-ADSWG) BMC Medicine. 2024. [PMC10777515](https://doi.org/10.1186/s12916-024-03111-1)

51. Tara A. Austin, Michael L. Thomas, Min Lu, Cooper B. Hodges, Emily S. Darowski, Rachel Bergmans, Sarah Parr, Delaney Pickell, Mikayla Catazaro, Crystal Lantrip, Elizabeth W. Twamley, et al.. META-ANALYSIS OF COGNITIVE FUNCTION FOLLOWING NON-SEVERE SARS-CoV-2 INFECTION Psychological and Cognitive Functioning Among Patients Receiving Outpatient Rehabilitation for Post-COVID Sequelae: An Observational Study High-dimensional characterization of post-acute sequelae of COVID-19 What's going on following acute COVID-19? Clinical characteristics of patients in an out-patient rehabilitation program Cognitive profile following COVID-19 infection: Clinical predictors leading to neuropsychological impairment Cognitive assessment in asymptomatic COVID-19 subjects Risk factors and abnormal cerebrospinal fluid associate with cognitive symptoms after mild COVID -19). Coronavirus disease 19 (COVID-19) vaccine effectiveness against symptomatic severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2) infection during delta-dominant and omicron-dominant periods in Japan: a multicenter prospective case-control study (Factors associated with SARS-CoV-2 infection and the effectiveness of COVID-19 vaccines study) COVID-19 severity is related to poor executive function in people with post-COVID conditions Long-COVID or long before? Neurocognitive deficits in people with COVID-19 D-cycloserine in the treatment of posttraumatic stress disorder Assessment of Cognitive Function in Patients After COVID-19 Infection The epidemiologic evidence linking autoimmune diseases and psychosis Neurocognitive deficits in COVID-19 patients five months after discharge from hospital Fatigue and cognitive impairment in post-COVID syndrome: Possible treatment approaches Neurologic Alterations Due to Respiratory Virus Infections Do Anxiety and Depression

Predict Persistent Physical Symptoms After a Severe COVID-19 Episode? A Prospective Study Neuropsychological manifestations of long COVID in hospitalized and non-hospitalized Brazilian Patients Long COVID-19: Objectifying most self-reported neurological symptoms Persistent neurocognitive decline in a clinic sample of hepatitis C virus-infected persons receiving interferon and ribavirin treatment Cognitive impairments in patients with subacute coronavirus disease: Initial experiences in a post-coronavirus disease clinic Signals of Significantly Increased Vaccine Breakthrough, Decreased Hospitalization Rates, and Less Severe Disease in Patients with Coronavirus Disease 2019 Caused by the Omicron Variant of Severe Acute Respiratory Syndrome Coronavirus 2 in Houston, Texas Cognitive and Neuropsychiatric Features of COVID-19 Patients After Hospital Dismission: An Italian Sample Cognitive functioning in people with chronic fatigue syndrome: A comparison between subjective and objective measures Cognitive consequences of COVID-19: Results of a cohort study from South America Changes in cognitive functioning after COVID-19: A systematic review and meta-analysis Examining the relationship between subjective and objective memory performance in older adults: A meta-analysis Long COVID: major findings, mechanisms and recommendations Selective visuoconstructional impairment following mild COVID-19 with inflammatory and neuroimaging correlation findings Heterogeneity and Risk of Bias in Studies Examining Risk Factors for Severe Illness and Death in COVID-19: A Systematic Review and Meta Analysis Cognitive sequelae of long COVID may not be permanent: A prospective study Cognitive decline among individuals with history of mild symptomatic SARS-CoV-2 infection: A longitudinal prospective study nested to a population cohort Long-lasting

cognitive effects of COVID-19: is there a role of BDNF? SARS-CoV-2 is associated with changes in brain structure in UK Biobank Neurological autoimmune diseases following vaccinations against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2): A follow-up study Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation (PHOSP-COVID): A UK multicentre, prospective cohort study Neuropsychological, Medical, and Psychiatric Findings After Recovery From Acute COVID-19: A Cross-sectional Study Long-Lasting Cognitive Abnormalities after COVID-19 The COVID-19 Hospitalization Metric in the Pre- and Post-vaccination Eras as a Measure of Pandemic Severity: A Retrospective, Nationwide Cohort Study Nociplastic pain: towards an understanding of prevalent pain conditions Observational cohort study of neurological involvement among patients with SARS-CoV-2 infection Post-covid 19 conditions in adults: systematic review and meta-analysis of health outcomes in controlled studies Neuropsychological deficits in patients with cognitive complaints after COVID-19 Persistent neurologic symptoms and cognitive dysfunction in non-hospitalized Covid-19 “long haulers” Neuroinflammation as a Common Feature of Neurodegenerative Disorders Cognitive deficits in people who have recovered from COVID-19 Brain MRI and neuropsychological findings at long-term follow-up after COVID-19 hospitalisation: an observational cohort study Neurological Manifestations of COVID-19 Feature T Cell Exhaustion and Dedifferentiated Monocytes in Cerebrospinal Fluid Describing cognitive function and psychosocial outcomes of COVID-19 survivors: A cross-sectional analysis Population-Based Estimates of Post-acute Sequelae of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection (PASC) Prevalence and Characteristics Clinical

outcomes of the severe acute respiratory syndrome coronavirus 2 Omicron and Delta variant: systematic review and meta-analysis of 33 studies covering 6 037 144 coronavirus disease 2019-positive patients Descriptive analysis of long COVID sequelae identified in a multidisciplinary clinic serving hospitalised and non-hospitalised patients Profiling cognitive impairment in mild COVID-19 patients: A case-control study at a secondary healthcare centre in the hilly region of North India Subjective and Objective Cognitive Impairments in Non-Hospitalized Persons 9 Months after SARS-CoV 2 Infection Neurological and (neuro)psychological sequelae in intensive care and general ward COVID-19 survivors Identifying diseases associated with Post-COVID syndrome through an integrated network biology approach Human immunodeficiency virus infection in the CNS and decreased dopamine availability: relationship with neuropsychological performance Neuropsychiatric and Cognitive Sequelae of COVID-19 Neuroinflammation in neurodegenerative disorders: the roles of microglia and astrocytes Post-acute sequelae of COVID-19: Evidence of mood & cognitive impairment Neuropsychological measures of “Long COVID-19 Fog” in older subjects Viral infection and neurological disorders—Potential role of extracellular nucleotides in neuroinflammation Abnormal brain diffusivity in participants with persistent neuropsychiatric symptoms after COVID-19 Neuropsychiatric phenotype of post COVID-19 syndrome in non-hospitalized patients One-Year Trajectory of Cognitive Changes in Older Survivors of COVID-19 in Wuhan, China: A Longitudinal Cohort Study Post-infection cognitive impairments in a cohort of elderly patients with COVID-19 Computing Within-Study Covariances, Data Visualization and Missing Data Solutions for Multivariate Meta-Analysis with

metavcov Can the choice reaction time be modified after COVID-19 diagnosis? A prospective cohort study Long-term psychiatric morbidities among SARS survivors Clinical features and cognitive sequelae in COVID-19: a retrospective study on N= 152 patients Development of criteria for cognitive dysfunction in post-COVID syndrome: the IC-CoDi-COVID approach Neurologic and cognitive sequelae after SARS-CoV2 infection: Different impairment for ICU patients Neurological and cognitive sequelae of Covid-19: A four month follow-up Effect of Sulodexide on Circulating Blood Cells in Patients with Mild COVID-19 Challenges in meta-analyses with observational studies The relationship between perceived and objective cognitive functioning in multiple sclerosis Chronic fatigue syndrome and cognitive deficit are associated with acute-phase neuropsychiatric manifestations of COVID-19: A 9-month follow-up study Cognitive impairments among patients in a long-COVID clinic: Prevalence, pattern and relation to illness severity, work function and quality of life Brain imaging in patients with COVID-19: A systematic review The vulnerability of executive functioning: The additive effects of recent non-restorative sleep, pain interference, and use of expressive suppression on test performance Cardiopulmonary, functional, cognitive and mental health outcomes post-COVID-19, across the range of severity of acute illness, in a physically active, working-age population Post-acute Sequelae of SARS-CoV-2 Infection and Subjective Memory Problems Postacute Sequelae of SARS-CoV-2 Infection and Impact on Quality of Life 1-6 Months After Illness and Association With Initial Symptom Severity Cognitive impairment in health care workers recovering from COVID-19 infection: a cross sectional comparative study Altered motor cortex physiology and dysexecutive syndrome in patients with fatigue and cognitive difficulties after mild

COVID-19 The PRISMA 2020 statement: An updated guideline for reporting systematic reviews Addressing the post-acute sequelae of SARS-CoV-2 infection: A multidisciplinary model of care Neurological outcomes 1 year after COVID-19 diagnosis: A prospective longitudinal cohort study Onset and frequency of depression in post-COVID-19 syndrome: A systematic review Evaluating meta-analytic methods to detect selective reporting in the presence of dependent effect sizes High-density EEG sleep correlates of cognitive and affective impairment at 12-month follow-up after COVID-19 COVID-19 Long-Term Effects: Is There an Impact on the Simple Reaction Time and Alternative-Forced Choice on Recovered Patients? Multidomain cognitive impairment in non-hospitalized patients with the post-COVID-19 syndrome: results from a prospective monocentric cohort An extended mixed-effects framework for meta analysis Look who is complaining: Psychological factors predicting subjective cognitive complaints in a large community sample of older adults A meta-analysis of systematic reviews and meta-analyses to evaluate the psychological consequences of COVID-19 What can psychiatrists learn from SARS and MERS outbreaks? Cognitive function in non hospitalized patients 8-13 months after acute COVID-19 infection: A cohort study in Norway ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions Post-acute COVID-19 syndrome negatively impacts physical function, cognitive function, health-related quality of life and participation Bidirectional associations between COVID-19 and psychiatric disorder: Retrospective cohort studies of 62 354 COVID-19 cases in the USA COVID-19 neuropathology at Columbia University Irving Medical Center/New York Presbyterian Hospital Neuroinflammation and Infection: Molecular Mechanisms Associated with Dysfunction of Neurovascular Unit

Comprehensive Health Assessment 3 Months After Recovery From Acute Coronavirus Disease 2019 (COVID-19) A systematic review of neuropsychological and psychiatric sequelae of COVID-19: Implications for treatment Cognitive Dysfunction, Psychiatric Distress, and Functional Decline After COVID-19 A general linear model for estimating effect size in the presence of publication bias Conducting meta-analyses in R with the metafor package Long COVID Neuropsychological Deficits after Severe, Moderate, or Mild Infection Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range Comorbid Chronic Diseases and Acute Organ Injuries Are Strongly Correlated with Disease Severity and Mortality among COVID-19 Patients: A Systemic Review and Meta-Analysis Disentangling the cognitive, physical, and mental health sequelae of COVID-19 The relationship between performance validity testing, external incentives, and cognitive functioning in long COVID Frequent neurocognitive deficits after recovery from mild COVID-19 Rapid vigilance and episodic memory decrements in COVID-19 survivors Neuropsychology review. 2025. [PMC13075542](#)

52. Gabriela Gonzalez Aleman, George D. Vavougiou, Carmela Tartaglia, Nalakath A. Uvais, Alla Guekht, Akram A. Hosseini, Vincenzina Lo Re, Catterina Ferreccio, Giovanni D'Avossa, Hernan P. Zamponi, Mariana Figueredo Aguiar, Agustin Yecora, Mohammad Zia Ul Haq Katshu, Vasileios T. Stavrou, Stylianos Boutlas, Konstantinos I. Gourgoulisanis, Camila Botero, Francisco González Insúa, Santiago Perez-Lloret, Mikhail Zinchuk, Anna Gersamija, Sofya Popova, Yulia Bryzgalova, Ekaterina Sviatskaya, Giovanna Russelli, Federica Avorio, Sophia Wang, Paul Edison, Yoshiki Niimi, Hamid R. Sohrabi, Elizabeta B. Mukaetova Ladinska, Daria Neidre, Gabriel A. de Erausquin, et al.. Age-dependent phenotypes of

cognitive impairment as sequelae of SARS-CoV-2 infection
Alzheimer's Disease Facts and Figures Risk factors and
abnormal cerebrospinal fluid associate with cognitive
symptoms after mild COVID-19 Plasma biomarkers of
neurodegeneration and neuroinflammation in hospitalized
COVID-19 patients with and without new neurological
symptoms Cognitive impairment after COVID-19-a review on
objective test data Chronic neuropsychiatric sequelae of
SARS-CoV-2: protocol and methods from the Alzheimer's
Association Global Consortium The chronic neuropsychiatric
sequelae of COVID-19: the need for a prospective study of
viral impact on brain functioning Characterizing long COVID:
deep phenotype of a complex condition SARS-CoV-2 is
associated with changes in brain structure in UK Biobank
Effects of age, sex, and ethnicity on the association between
apolipoprotein E genotype and Alzheimer disease - a meta-
analysis Comparison of serum neurodegenerative biomarkers
among hospitalized COVID-19 patients versus non-COVID
subjects with normal cognition, mild cognitive impairment, or
Alzheimer's dementia Bridging knowledge gaps in the
diagnosis and management of neuropsychiatric sequelae of
COVID-19 Cognitive impairment after recovery from
COVID-19: frequency, profile, and relationships with clinical
and laboratory indices Canadian COVID-19 host genetics
cohort replicates known severity associations Estimated
global proportions of individuals with persistent fatigue,
cognitive, and respiratory symptom clusters following
symptomatic COVID-19 in 2020 and 2021 Olfactory
dysfunction but not COVID-19 severity predicts severity of
cognitive sequelae following SARS-CoV-2 infection in
Amerindian older adults Short-term and long-term rates of
postacute sequelae of SARS-CoV-2 infection: a systematic
review Cognitive function following SARS-CoV-2 infection in a

population-representative Canadian sample Neuro-axonal injury in COVID-19: the role of systemic inflammation and SARS-CoV-2 specific immune response Cerebrospinal fluid findings in COVID-19: a multicenter study of 150 lumbar punctures in 127 patients The olfactory route is a potential way for SARS-CoV-2 to invade the central nervous system of rhesus monkeys Prolonged and extended impacts of SARS-CoV-2 on the olfactory neurocircuit APOE e4 genotype predicts severe COVID-19 in the UK Biobank Community Cohort Type 2 diabetes and its genetic susceptibility are associated with increased severity and mortality of COVID-19 in UK Biobank COVID-19 and risk of neurodegenerative disorders: A Mendelian randomization study Interleukin-6 affects the severity of olfactory disorder: a cross-sectional survey of 148 patients who recovered from Omicron infection using the Sniffin' Sticks test in Tianjin, China Association between olfactory dysfunction and mood disturbances with objective and subjective cognitive deficits in long-COVID Anosmia in COVID-19 could be associated with long-term deficits in the consolidation of procedural and verbal declarative memories Neuropathology of COVID-19 (neuro-COVID): clinicopathological update COVID-19 mortality by race and ethnicity in US metropolitan and nonmetropolitan areas, March 2020 to February (2022) The impact of mild COVID-19 on executive functioning and mental health outcomes in young adults Severe COVID-19 is associated with molecular signatures of aging in the human brain Olfactory transmucosal SARS-CoV-2 invasion as a port of central nervous system entry in individuals with COVID-19 Peripheral challenge with a viral mimic upregulates expression of the complement genes in the hippocampus Cognitive impact of COVID-19: looking beyond the short term Olfactory and other sensory impairments in Alzheimer disease Age-dependent

impact of the major common genetic risk factor for COVID-19 on severity and mortality Post-acute COVID-19 syndrome COVID-19-related neuropathology and microglial activation in elderly with and without dementia Long COVID or post-acute sequelae of COVID-19 (PASC): an overview of biological factors that may contribute to persistent symptoms Characteristics of COVID-19 and mortality-associated factors during the first year of the pandemic in southeast Mexico Alzheimer's-like signaling in brains of COVID-19 patients Concerted type I interferon signaling in microglia and neural cells promotes memory impairment associated with amyloid β plaques The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19 Genomewide association study of severe Covid-19 with respiratory failure The impact of COVID-19 on pulmonary, neurological, and cardiac outcomes: evidence from a Mendelian randomization study Pro-inflammatory interleukin-6 signaling links cognitive impairments and peripheral metabolic alterations in Alzheimer's disease Olfactory discrimination predicts cognitive decline among community-dwelling older adults COVID-19 and cognitive performance: a Mendelian randomization study Neurological and psychiatric risk trajectories after SARS-CoV-2 infection: an analysis of 2-year retrospective cohort studies including 1 284 437 patients COVID-19 neuropathology at Columbia University Irving Medical Center/New York Presbyterian Hospital Type I interferon signaling in SARS-CoV-2 associated neurocognitive disorder (SAND): mapping host-virus interactions to an etiopathogenesis Serum neurofilament light chain levels in Covid-19 patients without major neurological manifestations Olfactory function is associated with cognitive performance: results from the population-based LIFE-Adult-Study Olfactory dysfunction and chronic cognitive impairment following

SARS-CoV-2 infection in a sample of older adults from the Andes mountains of Argentina *Frontiers in Aging Neuroscience*. 2025. [PMC11747492](#)

53. Evan Xu, Yan Xie, Ziyad Al-Aly. Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Blood lipids after COVID-19 infection Plasma metabolomic profiles in recovered COVID-19 patients without previous underlying diseases 3 months after discharge COVID-19-associated dyslipidemia: implications for mechanism of impaired resolution and novel therapeutic approaches Altered lipid profile in patients with COVID-19 infection Alterations in the human oral and gut microbiomes and lipidomics in COVID-19 Proteomic analysis identifies prolonged disturbances in pathways related to cholesterol metabolism and myocardium function in the COVID-19 recovery stage Severe acquired hypertriglyceridemia following COVID-19 Persistence, prevalence, and polymorphism of sequelae after COVID-19 in unvaccinated, young adults of the Swiss Armed Forces: a longitudinal, cohort study (LoCoMo) A detailed study of patients with long-haul COVID. An analysis of private healthcare claims Geographic variation and US county characteristics associated with rapid kidney function decline Higher blood urea nitrogen is associated with increased risk of incident diabetes mellitus Burden of cause-specific mortality associated with PM2.5 air pollution in the United States Estimates of all cause mortality and cause specific mortality associated with proton pump inhibitors among US veterans: cohort study Association between monocyte count and risk of incident CKD and progression to ESRD Comparative effectiveness of the sodium-glucose co-transporter-2 inhibitor empagliflozin vs. other antihyperglycemics on risk of major adverse kidney events Comparative effectiveness of sodium-glucose cotransporter 2

inhibitors vs sulfonylureas in patients with type 2 diabetes
Acute kidney injury in a national cohort of hospitalized US veterans with COVID-19
Comparative effectiveness of SGLT2 inhibitors, GLP-1 receptor agonists, DPP-4 inhibitors, and sulfonylureas on risk of kidney outcomes: emulation of a target trial using health care databases
Clinical Implications of estimated glomerular filtration rate dip following sodium-glucose cotransporter-2 inhibitor initiation on cardiovascular and kidney outcomes
COVID-19: shared data resource
Making Neighborhood-disadvantage metrics accessible—the neighborhood atlas
High-dimensional characterization of post-acute sequelae of COVID-19
Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status
Kidney outcomes in long COVID
Long-term cardiovascular outcomes of COVID-19
Risks of mental health outcomes in people with COVID-19: cohort study
High-dimensional propensity score adjustment in studies of treatment effects using health care claims data
Comparison of critical drug-drug interaction listings: the Department of Veterans Affairs medical system and standard reference compendia
Polypharmacy, drug-drug interactions, and potentially inappropriate medications in older adults with human immunodeficiency virus infection
Kidney outcomes in long COVID
An introduction to propensity score methods for reducing the effects of confounding in observational studies
Using propensity scores to help design observational studies: application to the tobacco litigation
Use of risk assessment tools to guide decision-making in the primary prevention of atherosclerotic cardiovascular disease: a special report from the American Heart Association and American College of Cardiology
Doubly robust estimation of causal effects
Long COVID after breakthrough SARS-CoV-2 infection
Risks and burdens of incident diabetes in long COVID: a cohort study

Reciprocal and coordinate regulation of serum amyloid A versus apolipoprotein A-I and paraoxonase-1 by inflammation in murine hepatocytes SREBP-dependent lipidomic reprogramming as a broad-spectrum antiviral target The immunology and immunopathology of COVID-19 A global lipid map reveals host dependency factors conserved across SARS-CoV-2 variants Low HDL and high triglycerides predict COVID-19 severity NLRP3 inflammasome: the stormy link between obesity and COVID-19 Association of dyslipidemia with the severity and mortality of coronavirus disease 2019 (COVID-19): a meta-analysis Dyslipidemia increases the risk of severe COVID-19: a systematic review, meta-analysis, and meta-regression Long-term neurologic outcomes of COVID-19 Acute and postacute sequelae associated with SARS-CoV-2 reinfection Managing hyperlipidaemia in patients with COVID-19 and during its pandemic: an expert panel position statement from HEART UK Comparative evaluation of clinical manifestations and risk of death in patients admitted to hospital with covid-19 and seasonal influenza: cohort study Long COVID: long-term health outcomes and implications for policy and research Diabetes after SARS-Co-V 2 infection COVID data tracker The Lancet. Diabetes & Endocrinology. 2023. [PMC9873268](#)

54. Jiawei Zhou, Xia Li, Ting Zhang, Ziyang Liu, Peng Li, Na Yu, Wei Wang, et al.. Pre-existing sleep disturbances and risk of COVID-19: a meta-analysis Long COVID: major findings, mechanisms and recommendations Sleep disturbances during the COVID-19 pandemic: a systematic review, meta-analysis, and meta-regression The sleep-immune crosstalk in health and disease Obstructive sleep apnea (OSA) and outcomes from coronavirus disease 2019 (COVID-19) pneumonia: a systematic review and meta-analysis Significant association of obstructive sleep apnoea with increased risk for fatal

COVID-19: a quantitative meta-analysis based on adjusted effect estimates Risk of post-acute sequelae of SARS-CoV-2 infection associated with pre-coronavirus disease obstructive sleep apnea diagnoses: an electronic health record-based analysis from the RECOVER initiative The association of insomnia with long COVID: an international collaborative study (ICOSS-II) Lifestyle, course of COVID-19, and risk of Long-COVID in non-hospitalized patients World bank country and lending groups Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis Behaviorally assessed sleep and susceptibility to the common cold Short sleep duration is associated with prolonged virus shedding in SARS-CoV-2 omicron-infected patients Sleep-deprivation reduces NK cell number and function mediated by β -adrenergic signalling Sleep-dependent activity of T cells and regulatory T cells Prolonged sleep deprivation induces a cytokine-storm-like syndrome in mammals Sleep disturbance, sleep duration, and inflammation: a systematic review and meta-analysis of cohort studies and experimental sleep deprivation Safety and efficacy of melatonin as an adjuvant therapy in COVID-19 patients: systematic review and meta-analysis Risk of severe COVID-19 in non-adherent OSA patients A human circulating immune cell landscape in aging and COVID-19 Sex drives dimorphic immune responses to viral infections Biological sex affects vaccine efficacy and protection against influenza in mice Sex- and age-specific regulation of ACE2: insights into severe COVID-19 susceptibility Why does COVID-19 kill more elderly men than women? Is there a role for testosterone? Causal associations between obstructive sleep apnea and COVID-19: a bidirectional Mendelian randomization study Age-dependent hypoxia-induced PD-L1 upregulation in patients with obstructive sleep apnoea Altered CD8⁺ T-cell

lymphocyte function and TC1 cell stemness contribute to enhanced malignant tumor properties in murine models of sleep apnea COVID-19 sequelae and the host proinflammatory response: an analysis from the OnCovid registry The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19 Serum inflammatory markers in obstructive sleep apnea: a meta-analysis Obstructive sleep apnea and inflammation: proof of concept based on two illustrative cytokines eClinicalMedicine. 2024. [PMC11276919](#)

55. Siwen Wang, Tianyi Huang, Marc G. Weisskopf, Jae H. Kang, Jorge E. Chavarro, Andrea L. Roberts. Multidimensional Sleep Health Prior to SARS-CoV-2 Infection and Risk of Post-COVID-19 Condition Association between BNT162b2 vaccination and long COVID after infections not requiring hospitalization in health care workers Risk of long COVID associated with Delta versus Omicron variants of SARS-CoV-2 Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: a systematic review Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review Attributes and predictors of long COVID Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Subjective sleep quality and sleep problems in the general Japanese adult population Prevalent, incident, and persistent insomnia in a population-based cohort tested before (2018) and during the first-wave of COVID-19 pandemic (2020) Sleep problems during the COVID-19 pandemic by population: a systematic review and meta-analysis Poor sleep behavior burden and risk of COVID-19 mortality and hospitalization Associations of sleep and circadian phenotypes with COVID-19 susceptibility and hospitalization: an observational cohort study based on the

UK Biobank and a two-sample Mendelian randomization study
Insomnia and sleep duration on COVID-19 susceptibility and
hospitalization: a Mendelian randomization study Obstructive
sleep apnea (OSA) and outcomes from coronavirus disease
2019 (COVID-19) pneumonia: a systematic review and meta-
analysis The public health impact of poor sleep on severe
COVID-19, influenza and upper respiratory infections Sleep,
sleep deprivation and infectious disease: studies in animals
Sleep patterns, genetic susceptibility, and incident
cardiovascular disease: a prospective study of 385 292 UK
Biobank participants Long COVID-mechanisms, risk factors,
and management Enteroviruses and postviral fatigue
syndrome Unexplained post-acute infection syndromes The
sleep-immune crosstalk in health and disease Origin,
methods, and evolution of the Three Nurses' Health Studies
Comparing the Morningness-Eveningness Questionnaire and
Munich ChronoType Questionnaire to the dim light melatonin
onset Melatonin and human rhythms A prospective study of
sleep duration and mortality risk in women Validation of the
Women's Health Initiative Insomnia Rating Scale in a
multicenter controlled clinical trial Sleep duration, insomnia,
and coronary heart disease among postmenopausal women in
the Women's Health Initiative Relative validity of self-
reported snoring as a symptom of sleep apnea in a sleep clinic
population The subjective meaning of sleep quality: a
comparison of individuals with and without insomnia
Adherence to a healthy sleep pattern and incident heart
failure: a prospective study of 408 802 UK Biobank
participants Adherence to a healthy sleep pattern and risk of
chronic kidney disease: the UK Biobank study Associations
between sleep quality and health span: a prospective cohort
study based on 328,850 UK Biobank participants Associations
of depression, anxiety, worry, perceived stress, and loneliness

prior to infection with risk of post-COVID-19 conditions
Racial/ethnic disparities in post-acute sequelae of SARS-CoV-2 infection in New York: an EHR-based cohort study from the RECOVER program
Are sleep patterns influenced by race/ethnicity—a marker of relative advantage or disadvantage? evidence to date
Validity of a dietary questionnaire assessed by comparison with multiple weighed dietary records or 24-hour recalls
Relative validity of nutrient intakes assessed by questionnaire, 24-hour recalls, and diet records as compared with urinary recovery and plasma concentration biomarkers: findings for women
Development and validation of a Structured Telephone Interview for Dementia Assessment (STIDA): the NIMH Genetics Initiative
Specific subjective memory complaints in older persons may indicate poor cognitive function
A 4-item measure of depression and anxiety: validation and standardization of the Patient Health Questionnaire-4 (PHQ-4) in the general population
Multiple imputation in public health research
Prevalence of sleep disturbances in Chinese healthcare professionals: a systematic review and meta-analysis
Multiple imputation of discrete and continuous data by fully conditional specification
Symptoms and risk factors for long COVID in non-hospitalized adults
Increased prevalence of sleep-disordered breathing in adults
Burden of sleep apnea: rationale, design, and major findings of the Wisconsin Sleep Cohort study
Poor-sleep is associated with slow recovery from lymphopenia and an increased need for ICU care in hospitalized patients with COVID-19: a retrospective cohort study
Association between chronotype and body mass index: the role of C-reactive protein and the cortisol response to stress
Sleep disturbance, sleep duration, and inflammation: a systematic review and meta-analysis of cohort studies and experimental sleep deprivation
C-reactive protein is associated with sleep

disordered breathing independent of adiposity Sleep and inflammation Evidence for causal effects of sleep disturbances on risk for osteoarthritis: a univariable and multivariable Mendelian randomization study Circulating anti-nuclear autoantibodies in COVID-19 survivors predict long COVID symptoms Sleep, circadian rhythm, and gut microbiota Chronic sleep deprivation markedly reduces coagulation factor VII expression Impaired exercise capacity in post-COVID-19 syndrome: the role of VWF-ADAMTS13 axis Reproducibility and validity of a self-administered physical activity questionnaire The validity of recalled weight among younger women JAMA Network Open. 2023. [PMC10230315](#)

56. Giulia Vivaldi, Mohammad Talaei, John Blaikley, Callum Jackson, Paul E Pfeffer, Seif O Shaheen, Adrian R Martineau, et al.. Bidirectional relationship between sleep problems and long COVID: a longitudinal analysis of data from the COVIDENCE UK study Long COVID: major findings, mechanisms and recommendations Global prevalence of post-COVID-19 sleep disturbances in adults at different follow-up time points: A systematic review and meta-analysis Effects of sleep disturbance on dyspnoea and impaired lung function following hospital admission due to COVID-19 in the UK: a prospective multicentre cohort study Sleep quality in individuals with post-COVID-19 condition: Relation with emotional, cognitive and functional variables Sleep disturbance in post COVID-19 conditions: Prevalence and quality of life Impact of COVID-19 Infection on Quality of Sleep Sleep health early in the coronavirus disease 2019 (COVID-19) outbreak in the United States: integrating longitudinal, cross-sectional, and retrospective recall data Prospective longitudinal study of "Sleepless in Lockdown": unpacking differences in sleep loss during the coronavirus pandemic in the UK The public health impact of poor sleep on

severe COVID-19, influenza and upper respiratory infections
Multidimensional Sleep Health Prior to SARS-CoV-2 Infection
and Risk of Post-COVID-19 Condition Poor sleep quality,
insomnia, and short sleep duration before infection predict
long-term symptoms after COVID-19 Pre-existing sleep
problems as a predictor of post-acute sequelae of COVID-19
Health behaviours the month prior to COVID-19 infection and
the development of self-reported long COVID and specific
long COVID symptoms: a longitudinal analysis of 1581 UK
adults The association between sleep quality and quality of
life: a population-based study Associations Between Sleep
Quality and Health Span: A Prospective Cohort Study Based
on 328,850 UK Biobank Participants Cohort Profile:
Longitudinal population-based study of COVID-19 in UK
adults (COVIDENCE UK) Real-time tracking of self-reported
symptoms to predict potential COVID-19 SARS-CoV-2 variants
of concern and variants under investigation in England.
Technical briefing 33 Long-term effects of coronavirus Long
covid or post-covid conditions Coronavirus disease (covid-19):
post covid-19 condition Long-term symptom profiles after
COVID-19 vs other acute respiratory infections: an analysis of
data from the COVIDENCE UK study The Pittsburgh sleep
quality index: A new instrument for psychiatric practice and
research National Sleep Foundation's sleep time duration
recommendations: methodology and results summary
National Sleep Foundation's sleep quality recommendations:
first report Social Deprivation and Ethnicity Are Associated
with More Problematic Sleep in Middle-Aged and Older
Adults Determinants of natural adult sleep: An umbrella
review English indices of deprivation 2019: technical report
An ultra-brief screening scale for anxiety and depression: the
PHQ-4 Review of inverse probability weighting for dealing
with missing data The relationship between duration and

quality of sleep and upper respiratory tract infections: a systematic review Sleep disturbance increases the risk of severity and acquisition of COVID-19: a systematic review and meta-analysis The association of insomnia with long COVID: An international collaborative study (ICOSS-II) Role of sleep deprivation in immune-related disease risk and outcomes Gut microbiome diversity is associated with sleep physiology in humans Improving sleep quality leads to better mental health: A meta-analysis of randomised controlled trials Psychological factors associated with Long COVID: a systematic review and meta-analysis Habitual short sleepers with pre-existing medical conditions are at higher risk of Long COVID Sleep during naturally occurring respiratory infections: A pilot study An international study of post-COVID sleep health Studying the Effect of Long COVID-19 Infection on Sleep Quality Using Wearable Health Devices: Observational Study Device-assessed sleep and physical activity in individuals recovering from a hospital admission for COVID-19: a multicentre study Prevalence of sleep disturbances in patients with long COVID assessed by standardised questionnaires and diagnostic criteria: A systematic review and meta-analysis Clinical characteristics with inflammation profiling of long COVID and association with 1-year recovery following hospitalisation in the UK: a prospective observational study BMJ Open Respiratory Research. 2025. [PMC12314824](#)

57. Zachary Butzin-Dozier, Manav Kumar, Yunwen Ji, Lin-Chiun Wang, A. Jerrod Anzalone, Eric Hurwitz, Rena C. Patel, Rachel Wong, Carolyn Bramante, Benjamin Sines, et al.. IL-6 Receptor Antagonists and Severe Post-COVID-19 Outcomes: An Emulated Target Trial A review of cytokine-based pathophysiology of Long COVID symptoms Increased interleukin-6 is associated with long COVID-19: a systematic

review and meta-analysis Biomarkers in long COVID-19: A systematic review Can Baseline IL-6 Levels Predict Long COVID in Subjects Hospitalized for SARS-CoV-2 Disease? Cytokine Profiles Associated With Acute COVID-19 and Long COVID-19 Syndrome The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19 The cytokine storm and COVID-19 Long COVID: major findings, mechanisms and recommendations Distinguishing features of Long COVID identified through immune profiling Interleukin-6 as potential mediator of long-term neuropsychiatric symptoms of COVID-19 Association between tocilizumab, sarilumab and all-cause mortality at 28 days in hospitalised patients with COVID-19: A network meta-analysis Effect of tocilizumab, sarilumab, and baricitinib on mortality among patients hospitalized for COVID-19 treated with corticosteroids: a systematic review and meta-analysis IL-6 inhibition in the treatment of COVID-19: A meta-analysis and meta-regression Overview on the use of IL-6 agents in the treatment of patients with cytokine release syndrome (CRS) and pneumonitis related to COVID-19 disease Use of tocilizumab and sarilumab alone or in combination with corticosteroids for covid-19: systematic review and network meta-analysis About the National Clinical Cohort Collaborative COVID-19 Phenotype Documentation, Version 4.0 Targeted Learning Targeted Maximum Likelihood Estimation for Causal Inference in Observational Studies Developing a Targeted Learning-Based Statistical Analysis Plan Effectiveness of mRNA Booster Vaccine Against Coronavirus Disease 2019 Infection and Severe Outcomes Among Persons With and Without Immune Dysfunction: A Retrospective Cohort Study of National Electronic Medical Record Data in the United States Long COVID risk and pre-COVID vaccination in an EHR-based cohort study from the

RECOVER program Advances in the Omicron variant development Trends in Disease Severity and Health Care Utilization During the Early Omicron Variant Period Compared with Previous SARS-CoV-2 High Transmission Periods - United States, December 2020-January 2022 SSRI use during acute COVID-19 and risk of Long COVID among patients with depression Predicting Long COVID in the National COVID Cohort Collaborative Using Super Learner: Cohort Study Exploiting Nonsystematic Covariate Monitoring to Broaden the Scope of Evidence about the Causal Effects of Adaptive Treatment Strategies Biological Therapies for Rheumatoid Arthritis: An Overview for the Clinician COVID-19 Treatment Guidelines Panel Assessing the effect of selective serotonin reuptake inhibitors in the prevention of post-acute sequelae of COVID-19 Coding long COVID: characterizing a new disease through an ICD-10 lens Finding Long-COVID: Temporal Topic Modeling of Electronic Health Records from the N3C and RECOVER Programs Generalisable long COVID subtypes: findings from the NIH N3C and RECOVER programmes Identifying who has long COVID in the USA: a machine learning approach using N3C data Active-comparator design and new-user design in observational studies The active comparator, new user study design in pharmacoepidemiology: historical foundations and contemporary application Practical considerations for specifying a super learner Super learner The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19 Endothelial dysfunction in COVID-19: an overview of evidence, biomarkers, mechanisms and potential therapies Mechanisms of long COVID and the path toward therapeutics National Center for Advancing Translational Sciences Long COVID: a clinical update medRxiv. 2026. [PMC12976917](#)

58. Bryan Lau, Eryka Wentz, Zhanmo Ni, Karine Yenokyan, Candelaria Vergara, Shruti H. Mehta, Priya Duggal, et al.. Physical health and mental fatigue disability associated with long-COVID: Baseline results from a US nationwide cohort Physical and mental health disability associated with long-COVID: Baseline results from a US nationwide cohort Effectiveness of the pre-Omicron COVID-19 vaccines against Omicron in reducing infection, hospitalization, severity, and mortality compared to Delta and other variants: A systematic review An Intranasal OMV-Based Vaccine Induces High Mucosal and Systemic Protecting Immunity Against a SARS-CoV-2 Infection The effectiveness of COVID-19 testing and contact tracing in a US city Post-acute and long-COVID-19 symptoms in patients with mild diseases: a systematic review Prevalence and Correlates of Long COVID Symptoms Among US Adults Post-COVID Conditions and Burden of Disease Understanding How Post-COVID-19 Condition Affects Adults and Health Care Systems Acute and persistent symptoms in non-hospitalized PCR-confirmed COVID-19 patients Post-COVID-19 symptoms 6 months after acute infection among hospitalized and non-hospitalized patients Follow-up of adults with noncritical COVID-19 two months after symptom onset Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study High-dimensional characterization of post-acute sequelae of COVID-19 Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Short-term and Long-term Rates of Postacute Sequelae of SARS-CoV-2 Infection Symptoms and risk factors for long COVID in non-hospitalized adults Quality of life, functional status, and persistent symptoms after intensive care of COVID-19 patients Health-related quality of life, functional decline, and long-term mortality in older patients following hospitalisation

due to COVID-19 Post-sequelae symptoms and comorbidities after COVID-19 Long-Term Health-Related Quality of Life in Non-Hospitalized Coronavirus Disease 2019 (COVID-19) Cases With Confirmed Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection in England: Longitudinal Analysis and Cross-Sectional Comparison With Controls Most Common Long COVID Physical Symptoms in Working Age Adults Who Experienced Mild COVID-19 Infection: A Scoping Review Care Dependency in Non-Hospitalized Patients with COVID-19 WHO Clinical Case Definition Working Group on Post-COVID-19 Condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus The REDCap consortium: Building an international community of software platform partners Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support Activities and attitudes of participants in the Baltimore longitudinal study A brief mental fatigue questionnaire Transportability of Trial Results Using Inverse Odds of Sampling Weights Standard errors and confidence intervals for variable importance in random forest regression, classification, and survival Partnering with a global platform to inform research and public policy making Risk Factors Associated With Post-COVID-19 Condition: A Systematic Review and Meta-analysis Risk factors for hospitalization among persons with COVID-19-Colorado Risk Factors for Hospitalization, Mechanical Ventilation, or Death Among 10 131 US Veterans With SARS-CoV-2 Infection Postural tachycardia syndrome (POTS) A review of postural orthostatic tachycardia syndrome Changes in cognitive function in patients with intractable dizziness following vestibular rehabilitation Therapeutic effect of steroids on vestibular neuritis: Systematic review and meta-analysis Venous leg

symptoms in healthy subjects assessed during prolonged standing Evaluation of patients with peripheral vascular disease The American journal of medicine. 2025. [PMC10924070](#)

59. Bryan Lau, Eryka Wentz, Zhanmo Ni, Karine Yenokyan, Candelaria Coggiano, Shruti H. Mehta, Priya Duggal, et al.. Physical and mental health disability associated with long-COVID: Baseline results from a US nationwide cohort Physical health and mental fatigue disability associated with long-COVID: Baseline results from a US nationwide cohort Post-acute and long-COVID-19 symptoms in patients with mild diseases: a systematic review Prevalence and Correlates of Long COVID Symptoms Among US Adults Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study High-dimensional characterization of post-acute sequelae of COVID-19 Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Short-term and Long-term Rates of Postacute Sequelae of SARS-CoV-2 Infection Symptoms and risk factors for long COVID in non-hospitalized adults Quality of life, functional status, and persistent symptoms after intensive care of COVID-19 patients Health-related quality of life, functional decline, and long-term mortality in older patients following hospitalisation due to COVID-19 Post-sequelae symptoms and comorbidities after COVID-19 The REDCap consortium: Building an international community of software platform partners Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support Activities and attitudes of participants in the Baltimore longitudinal study A brief mental fatigue questionnaire Transportability of Trial Results Using Inverse Odds of Sampling Weights Standard

errors and confidence intervals for variable importance in random forest regression, classification, and survival Partnering with a global platform to inform research and public policy making Changes in cognitive function in patients with intractable dizziness following vestibular rehabilitation Therapeutic effect of steroids on vestibular neuritis: Systematic review and meta-analysis medRxiv. 2022. [PMC9753791](#)

60. Anusua Datta. Assessing the effects of long-COVID on mental health in the United States: a population based study Post-acute COVID-19 syndrome The lived experience of long COVID: a qualitative study of mental health, quality of life, and coping Hikikomori: a hidden mental health need following the COVID-19 pandemic The prevalence of depression, anxiety, and sleep disturbances in COVID19 patients: a meta-analysis Short-term and long-term rates of postacute sequelae of SARSCoV-2 infection: a systematic review Psychiatric symptoms in Long-COVID patients: a systematic review Screening for depression in medical settings with the Patient health questionnaire (PHQ): a diagnostic meta-analysis Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection Who is hardest hit by a pandemic? Racial disparities in COVID-19 hardship in the U.S Associations of small business closure and reduced urban mobility with mental health problems in COVID-19 pandemic: a national representative sample study Trends in anxiety and depression symptoms during the COVID-19 pandemic: results from the U.S. Census bureau's household pulse survey How do state policies shape experiences of household income shocks and mental health during the COVID-19 pandemic? Effects of the 2021 expanded child tax credit on adults' mental health: a quasi-experimental study Impact of the expanded child tax credit and its

expiration on adult psychological well-being Mental health, substance use, and suicidal ideation during the COVID-19 pandemic—United States, June 24–30 Age disparities in prevalence of anxiety and depression among US adults during COVID-19 pandemic The differential impact of COVID-19 on mental health: implications of ethnic, sexual orientation, and disability status in the United States The COVID-19 pandemic and its effects on mental health—a before, during, and after comparison using the U.S. Census Bureau’s household pulse survey Sex-differences in COVID-19 diagnosis, risk factors and disease comorbidities: a large US-based cohort study Monthly trends in self-reported health status and depression by race/ethnicity, and socioeconomic status during COVID-19 pandemic, United States, April 2020–May 2021 Discover Mental Health. 2025. [PMC11850688](#)

61. Soomin Ryu, Akash Patel, Christine Chyu, James H. Buszkiewicz, Sameera Ahmed, Nancy L. Fleischer. Prospective associations between Long COVID and mental health: evidence from a population-based study with a nearly three-year follow-up Symptoms, complications and management of long COVID: a review Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: a systematic review Characteristics and impact of long covid: findings from an online survey The global prevalence of depression, anxiety, and sleep disorder among patients coping with Post COVID-19 syndrome (long COVID): a systematic review and meta-analysis Prevalence of depression, anxiety, stress, and suicide tendency among individual with long-COVID and determinants: a systematic review and meta-analysis A one-year follow-up study of systematic impact of long COVID symptoms among patients post SARS-CoV-2 omicron variants infection in Shanghai, China Persistence of symptoms and quality of life at 35 days

after hospitalization for COVID-19 infection Trajectory of anxiety/depressive symptoms and sleep quality in individuals who had been hospitalized by COVID-19: The LONG-COVID-EXP multicenter study Post-discharge health status and symptoms in patients with severe COVID-19 Long-term effects of COVID-19 on mental health: a systematic review Persistent neuropsychiatric symptoms after COVID-19: a systematic review and meta-analysis Long COVID and mental health correlates: a new chronic condition fits existing patterns Association of Long COVID with mental health disorders: a retrospective cohort study using real-world data from the USA Dissecting the association between long COVID and depressive symptoms in a nationally representative population from France Assessing the effects of long-COVID on mental health in the United States: a population based study Household surveys in crisis Decreasing survey response rates in the time of COVID-19: implications for analyses of population health and health inequities The patient health questionnaire-2: validity of a two-item depression screener Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection Psychometric properties and clinical utility of brief measures of depression, anxiety, and general distress: The PHQ-2, GAD-2, and K-6 Insights provided by depression screening regarding pain, anxiety, and substance use in a veteran population Examination of the best cut-off points of PHQ-2 and GAD-2 for detecting depression and anxiety in Italian cardiovascular inpatients Validation of generalized anxiety disorder-2 as a screening tool for generalized anxiety disorder in the primary health care setting: study protocol Validity and utility of the patient health questionnaire (PHQ)-2 and PHQ-9 for screening and diagnosis of depression in rural Chiapas, Mexico: a cross-sectional study A review of hot deck imputation for survey non-

response Psychological consequences of long COVID: comparing trajectories of depressive and anxiety symptoms before and after contracting SARS-CoV-2 between matched long- and short-COVID groups Longitudinal course and predictors of health-related quality of life, mental health, and fatigue, in non-hospitalized individuals with or without post COVID-19 syndrome The impact of inflammatory and metabolic markers on depression, anxiety, and cognition after COVID-19: a narrative review Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors Persistent psychopathology and neurocognitive impairment in COVID-19 survivors: effect of inflammatory biomarkers at three-month follow-up Neuropsychiatric aspects of long COVID: a comprehensive review What can neuroimmunology teach us about the symptoms of long-COVID? New-onset neuropsychiatric sequelae and 'long-COVID' syndrome (Review) Three-month follow-up of the post-COVID syndrome after admission to a specialised post-COVID centre—a prospective study focusing on mental health with patient reported outcome measures (PROMs) A systematic review of neuropsychological and psychiatric sequelae of COVID-19: implications for treatment Is self-compassion associated with lower psychological distress in people with long COVID? Results from a cross-sectional survey Impact of major depression on chronic medical illness Depression, anxiety and their relationship with chronic diseases: a review of the epidemiology, risk and treatment evidence The patient health questionnaire-2: validity of a two-item depression screener Screening for anxiety disorders with the GAD-7 and GAD-2: a systematic review and diagnostic metaanalysis Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID symptom study app: a prospective, community-based,

nested, case-control study Psychiatric morbidity and protracted symptoms after COVID-19 COVID-19 vaccination for the prevention and treatment of long COVID: A systematic review and meta-analysis Physical, cognitive, and mental health impacts of Omicron reinfection in patients with original SARS-CoV-2 infection: a community-based observational study Accuracy of the PHQ-2 alone and in combination with the PHQ-9 for screening to detect major depression: systematic review and meta-analysis BMC Public Health. 2026. [PMC13023150](#)

62. Petra Engelmann, Max Reinke, Clara Stein, Stefan Salzmann, Bernd Löwe, Anne Toussaint, Meike Shedden-Mora, et al.. Psychological factors associated with Long COVID: a systematic review and meta-analysis Managing the long term effects of covid-19: summary of NICE, SIGN, and RCGP rapid guideline The immunology and immunopathology of COVID-19 Post-COVID-19 syndrome is rarely associated with damage of the nervous system: findings from a prospective observational cohort study in 171 patients Dysautonomia following COVID-19 is not associated with subjective limitations or symptoms but is associated with objective functional limitations Long-COVID or long before? Neurocognitive deficits in people with COVID-19 Associations of depression, anxiety, worry, perceived stress, and loneliness prior to infection with risk of post-COVID-19 conditions Risk factors associated with post-COVID-19 condition: a systematic review and meta-analysis Symptoms and risk factors for long COVID in non-hospitalized adults Long COVID: 3 years in Unexplained post-acute infection syndromes Somatization in the population: from mild bodily misperceptions to disabling symptoms Somatic symptom disorder: a scoping review on the empirical evidence of a new diagnosis Persistent SOMatic symptoms ACROSS diseases - from risk factors to

modification: scientific framework and overarching protocol of the interdisciplinary SOMACROSS research unit (RU 5211)

Functional somatic disorders: discussion paper for a new common classification for research and clinical use

Management of functional somatic syndromes and bodily distress

Comparing anxiety and depression to obesity and smoking as predictors of major medical illnesses and somatic symptoms

Predictors of new onsets of irritable bowel syndrome, chronic fatigue syndrome and fibromyalgia: the lifelines study

Impact of symptom focusing and somatosensory amplification on persistent physical symptoms: a three-year follow-up study

The 2-year course of medically unexplained physical symptoms (MUPS) in terms of symptom severity and functional status: results of the prospects cohort study

Aetiological understanding of fibromyalgia, irritable bowel syndrome, chronic fatigue syndrome and classificatory analogues: a systematic umbrella review

The fear-avoidance model of pain

Predictors of persistent somatic symptoms in the general population: a systematic review of cohort studies

Adverse childhood experiences and the consequences on neurobiological, psychosocial, and somatic conditions across the lifespan

Predictors of somatic symptom severity: the role of cumulative history of trauma and adversity in a diverse community sample

Neuroticism and adverse life events are important determinants in functional somatic disorders: the DanFunD study

Prevalence of mental health conditions and brain fog in people with long COVID: a systematic review and meta-analysis

Post-COVID-19 fatigue: a systematic review

The framework for systematic reviews on psychological risk factors for persistent somatic symptoms and related syndromes and disorders (PSY-PSS)

Psychological risk factors of somatic symptom disorder: a systematic review and meta-analysis of cross-sectional and longitudinal studies

Meta-

analysis of observational studies in Epidemiology A proposal for reporting Core outcome domains for clinical trials on somatic symptom disorder, bodily distress disorder, and functional somatic syndromes: European network on somatic symptom disorders recommendations A clinical case definition of post-COVID-19 condition by a Delphi consensus Rayyan—a web and mobile app for systematic reviews The challenges of containing SARS-CoV-2 via test-trace-and-isolate Prevalence and predictors of prolonged cognitive and psychological symptoms following COVID-19 in the United States Resilience after COVID-19: a descriptive, cross-sectional study Prevalence, trajectory over time, and risk factor of post-COVID-19 fatigue Complex association between post-COVID-19 condition and anxiety and depression symptoms Low growth hormone secretion associated with post-acute sequelae SARS-CoV-2 infection (PASC) neurologic symptoms: a case-control pilot study Post-COVID-19 syndrome: incidence, risk factor, and the most common persisting symptoms Determinants of long COVID among adults hospitalized for SARS-CoV-2 infection: a prospective cohort study A one-year follow-up study of systematic impact of long COVID symptoms among patients post SARS-CoV-2 omicron variants infection in Shanghai, China Post-COVID patients with new-onset chronic pain 2 years after infection: cross-sectional study A glimpse into Long COVID characteristics and the mental health impact within a highly vaccinated population: a Malta observational study Long COVID and mental health correlates: a new chronic condition fits existing patterns Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study The impact of world trade center related medical conditions on the severity of COVID-19 disease and its long-term sequelae Perceived cognitive deficits in patients with

symptomatic SARS-CoV-2 and their association with post-COVID-19 condition Fatigue presentation, severity, and related outcomes in a prospective cohort following post-COVID-19 hospitalization in British Columbia, Canada Post-COVID symptoms in occupational cohorts: effects on health and work ability Risk factors and multidimensional assessment of long coronavirus disease fatigue: a nested case-control study Differences in clinical presentation with long COVID after community and hospital infection and associations with all-cause mortality: English sentinel network database study SARS-CoV-2 infection and long COVID among California farmworkers Characteristics of frailty in perimenopausal women with long COVID-19 Long-term effects of COVID-19 on workers in health and social services in Germany Prevalence and factors associated with long COVID and mental health status among recovered COVID-19 patients in southern Thailand Prevalence and determinants of long COVID among patients attending the outpatient department of a subdistrict hospital in Haryana Demographic and clinical factors associated with long COVID Prevalence and predictors of long COVID in patients accessing a national digital mental health service Comprehensive analysis of long COVID in a Japanese nationwide prospective cohort study 3-year outcomes of discharged survivors of COVID-19 following the SARS-CoV-2 omicron (B.1.1.529) wave in 2022 in China: a longitudinal cohort study Abnormal brain diffusivity in participants with persistent neuropsychiatric symptoms after COVID-19 Lowered oxygen saturation and increased body temperature in acute COVID-19 largely predict chronic fatigue syndrome and affective symptoms due to Long COVID: a precision nomothetic approach Neuropsychological impairment in post-COVID condition individuals with and without cognitive

complaints Factors associated with long COVID symptoms in an online cohort study Structural brain changes in patients with post-COVID fatigue: a prospective observational study Characterization of neurocognitive deficits in patients with post-COVID-19 syndrome: persistence, patients' complaints, and clinical predictors A hypoarousal model of neurological post-COVID syndrome: the relation between mental fatigue, the level of central nervous activation and cognitive processing speed Amino acids, post-translational modifications, nitric oxide, and oxidative stress in serum and urine of long COVID and ex COVID human subjects Post-COVID-19-associated morbidity in children, adolescents, and adults: a matched cohort study including more than 157,000 individuals with COVID-19 in Germany Cardiopulmonary, functional, cognitive and mental health outcomes post-COVID-19, across the range of severity of acute illness, in a physically active, working-age population Persistence of COVID-19 symptoms beyond 3 months and the delayed return to the usual state of health in Saudi Arabia: a cross-sectional study Long COVID 1 year after hospitalisation for COVID-19: a prospective bicentric cohort study Functional status, mood state, and physical activity among women with post-acute COVID-19 syndrome Cortical GABA levels are reduced in post-acute COVID-19 syndrome A comparison of pain, fatigue, and function between post-COVID-19 condition, fibromyalgia, and chronic fatigue syndrome: a survey study Emotional functioning in long COVID: comparison to post-concussion syndrome using the personality assessment inventory Brain fog of post-COVID-19 condition and chronic fatigue syndrome, same medical disorder? Social stigma, mental health, stress, and health-related quality of life in people with long COVID Illness perceptions in long-COVID: a cross-sectional analysis in adults The self-reported Leeds assessment of neuropathic

symptoms and signs (S-LANSS) and PainDETECT questionnaires in COVID-19 survivors with post-COVID pain

Cardiovascular abnormalities and mental health difficulties result in a reduced quality of life in the post-acute COVID-19 syndrome

Sleep quality mediates the effect of sensitization-associated symptoms, anxiety, and depression on quality of life in individuals with post-COVID-19 pain

Perceived quality of life and associated factors in long covid syndrome among older brazilians: a cross-sectional study

Health related quality of life and its determinants in COVID-19 patients

Impact of fatigue as the primary determinant of functional limitations among patients with post-COVID-19 syndrome: a cross-sectional observational study

Post-COVID-19 syndrome: physical capacity, fatigue and quality of life

Characteristics of long COVID: cases from the first to the fifth wave in greater Tokyo, Japan

Neuropsychiatric phenotype of post COVID-19 syndrome in non-hospitalized patients

Cognitive disengagement syndrome in pediatric patients with long covid: associations with mood, anxiety, and functional impairment

Post-acute sequelae of SARS-CoV-2 infection: relationship of central nervous system manifestations with physical disability and systemic inflammation

Obesity and lipid metabolism disorders determine the risk for development of long COVID syndrome: a cross-sectional study from 50,402 COVID-19 patients

Prolonged gastrointestinal manifestations after recovery from COVID-19

Persistent symptoms after acute COVID-19 infection in omicron era

Unraveling brain fog in post-COVID syndrome: relationship between subjective cognitive complaints and cognitive function, fatigue, and neuropsychiatric symptoms

Neurological and psychiatric manifestations of post-COVID-19 conditions

Long-covid cognitive impairment: cognitive assessment and apolipoprotein E (APOE) genotyping

correlation in a Brazilian cohort Outcomes in post-acute sequelae of COVID-19 (PASC) at 6 months post-infection part 1: cognitive functioning Long COVID in hospitalized COVID-19 patients: a retrospective cohort study Cognitive dysfunction associated with COVID-19: a comprehensive neuropsychological study Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Predictive models of long COVID An elastic net regression model for identifying long COVID patients using health administrative data: a population-based study Psychological factors predict higher odds and impairment of post-COVID symptoms: a prospective study Determinants of post-COVID-19 symptoms among adults aged 55 or above with chronic conditions in primary care: data from a prospective cohort in Hong Kong Prevalence of and factors associated with post-coronavirus disease 2019 (COVID-19) condition in the 12 Months after the diagnosis of COVID-19 in adults followed in general practices in Germany Do anxiety and depression predict persistent physical symptoms after a severe COVID-19 episode? A prospective study Post-COVID-19 conditions in children and adolescents diagnosed with COVID-19 COVEVOL: natural evolution at 6 months of COVID-19 Manifestation of headache affecting quality of life in long COVID patients Lifestyle, course of COVID-19, and risk of Long-COVID in non-hospitalized patients Mental health among children with long COVID during the COVID-19 pandemic Long COVID and psychological distress in young adults: potential protective effect of a prior mental health diagnosis Physical activity status prevents symptoms of long covid: sulcovid-19 survey Clinical characterization and factors associated with quality of life in Long COVID patients: secondary data analysis from a randomized clinical trial Association of sedentary lifestyle with risk of acute and post-

acute COVID-19 sequelae: a retrospective cohort study
Prevalence and characteristics associated with post-COVID-19 condition among nonhospitalized adolescents and young adults
The association of insomnia with long COVID: an international collaborative study (ICOSS-II)
Pre-existing conditions associated with post-acute sequelae of COVID-19
One year follow-up of COVID-19 related symptoms and patient quality of life: a prospective cohort study
Post-COVID-19 syndrome and quality of life impairment in severe COVID-19 mexican patients
Persistent symptoms, quality of life, and risk factors in long COVID: a cross-sectional study of hospitalized patients in Brazil
Predictors of long-COVID-19 and its impact on quality of life: longitudinal analysis at 3, 6 and 9 months after discharge from a portuguese centre
Long COVID and chronic conditions in the U.S. workforce: prevalence, productivity loss, and disability
Outcomes among confirmed cases and a matched comparison group in the long-COVID in scotland study
Long COVID-19: objectifying most self-reported neurological symptoms
Determinants of recovery from post-COVID-19 dyspnoea: analysis of UK prospective cohorts of hospitalised COVID-19 patients and community-based controls
The interactive effects of post-traumatic stress symptoms and breathlessness on fatigue severity in post-COVID-19 syndrome
Post COVID-19 condition diagnosis: a population-based cohort study of occurrence, associated factors, and healthcare use by severity of acute infection
Long-term consequences of COVID-19 in predominantly immunonaive patients: a Canadian prospective population-based study
A statewide population-based approach to examining long COVID symptom prevalence and predictors in Michigan
Post-COVID syndrome and severity of COVID-19: a cross-sectional epidemiological evaluation from north India
Anosmia predicts memory impairment in post-

covid-19 syndrome: results of a neuropsychological cohort study Exploring the influence of pre-existing conditions and infection factors on pediatric long COVID symptoms and quality of life Anxiety but not menopausal status influences the risk of long-COVID-19 syndrome in women living in Latin America Long-COVID in children: an exploratory case-control study from a bio-psycho-social perspective Factors associated with self-reported post/long-COVID-A real-world data study Post-covid-19 syndrome: self-compassion and psychological flexibility moderate the relationship between physical symptom load and psychosocial impact Prognostic factors for persistent fatigue after COVID-19: a prospective matched cohort study in primary care Examining association of personality characteristics and neuropsychiatric symptoms in post-COVID syndrome Psychological resilience to trauma and risk for COVID-19 infection and somatic symptoms over two years Social stigma in children with long COVID Physical, psychological, and cognitive profile of post-COVID conditions in healthcare workers, Quebec, Canada Association between psychological distress, cognitive complaints, and neuropsychological status after a severe COVID-19 episode: a cross-sectional study A prospective observational study of post-COVID-19 chronic fatigue syndrome following the first pandemic wave in Germany and biomarkers associated with symptom severity Impact of pre-existing mental health diagnoses on development of post-COVID and related symptoms: a claims data-based cohort study How to use the new DSM-5 somatic symptom disorder diagnosis in research and practice: a critical evaluation and a proposal for modifications Biopsychosocial risk factors of persistent fatigue after acute infection: a systematic review to inform interventions Prevalence, risk factors, and outcomes of irritable bowel syndrome after infectious enteritis: a

systematic review and meta-analysis Persistent physical symptoms: definition, genesis, and management Neuroticism, perceived stress, adverse life events and self-efficacy as predictors of the development of functional somatic disorders: longitudinal population-based study (DanFunD) Long COVID: aiming for a consensus Association of psychiatric disorders with clinical diagnosis of long COVID in US veterans Worldwide prevalence and disability from mental disorders across childhood and adolescence: evidence from the global burden of disease study The PHQ-9: a new depression diagnostic and severity measure A brief measure for assessing generalized anxiety disorder: the GAD-7 Within and between-day variation and associations of symptoms in long Covid: intensive longitudinal study Clinical characteristics with inflammation profiling of long COVID and association with 1-year recovery following hospitalisation in the UK: a prospective observational study Pathophysiology and mechanism of long COVID: a comprehensive review Response to commentary: the framework for systematic reviews on psychological risk factors for persistent somatic symptoms and related syndromes and disorders (PSY-PSS). General commentary Why the hypothesis of psychological mechanisms in long COVID is worth considering Post-acute COVID-19 syndrome (PCS) and health-related quality of life (HRQoL)-A systematic review and meta-analysis eClinicalMedicine. 2024. [PMC11701445](#)

63. Hythem Sidky, Kristen A. Hansen, Andrew T. Girvin, Nathan Hotaling, Sam G. Michael, Ken Gersing, David K. Sahner, et al.. Assessing the effect of selective serotonin reuptake inhibitors in the prevention of post-acute sequelae of COVID-19 Assessing the Effect of Selective Serotonin Reuptake Inhibitors in the Prevention of Post-Acute Sequelae of COVID-19 Incidence, co-occurrence, and evolution of long-

COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19 Post-acute COVID-19 syndrome Characterising long COVID: a living systematic review Summary of joint guideline on the management of long COVID Covid-19: one in eight adults develops long covid symptoms, study suggests Insights from myalgic encephalomyelitis/chronic fatigue syndrome may help unravel the pathogenesis of postacute COVID-19 syndrome Redox imbalance links COVID-19 and myalgic encephalomyelitis/chronic fatigue syndrome Role of mitochondria, oxidative stress and the response to antioxidants in myalgic encephalomyelitis/chronic fatigue syndrome: a possible approach to SARS-CoV-2 'long-haulers'? Persistent SARS-2 infections contribute to long COVID-19 A central role for amyloid fibrin microclots in long COVID/PASC: origins and therapeutic implications COVID-19 in an immunocompromised host: persistent shedding of viable SARS-CoV-2 and emergence of multiple mutations: a case report Prevalence and persistent shedding of fecal SARS-CoV-2 RNA in patients with COVID-19 infection: a systematic review and meta-analysis Reverse-transcribed SARS-CoV-2 RNA can integrate into the genome of cultured human cells and can be expressed in patient-derived tissues Markers of immune activation and inflammation in individuals with postacute sequelae of severe acute respiratory syndrome coronavirus 2 infection Immune-based prediction of COVID-19 severity and chronicity decoded using machine learning Multiple early factors anticipate post-acute COVID-19 sequelae Immuno-proteomic profiling reveals aberrant immune cell regulation in the airways of individuals with ongoing post-COVID-19 respiratory disease Immunomodulatory effects of fluoxetine: a new potential pharmacological action for a classic antidepressant drug?

Selective serotonin reuptake inhibitors as a novel class of immunosuppressants The influence of antidepressants on the immune system Study on anti-inflammatory and immunomodulatory effects of fluoxetine in rat models of inflammation Immunomodulatory effect of selective serotonin reuptake inhibitors (SSRIs) on human T lymphocyte function and gene expression The immunomodulatory effect of the antidepressant sertraline in an experimental autoimmune encephalomyelitis mouse model of multiple sclerosis Platelet serotonin promotes the recruitment of neutrophils to sites of acute inflammation in mice Serotonin reduction in post-acute sequelae of viral infection Repurposing of CNS drugs to treat COVID-19 infection: targeting the sigma-1 receptor Psychotropic drugs: therapeutic perspectives against Covid-19 Modulation of the sigma-1 receptor-IRE1 pathway is beneficial in preclinical models of inflammation and sepsis Overview of the potential use of fluvoxamine for COVID-19 and long COVID Interactions of new antidepressants with sigma-1 receptor chaperones and their potentiation of neurite outgrowth in PC12 cells Inflammation and clinical response to treatment in depression: a meta-analysis The effect of antidepressant medication treatment on serum levels of inflammatory cytokines: a meta-analysis Imbalance between pro- and anti-inflammatory cytokines, and between Th1 and Th2 cytokines in depressed patients: the effect of electroacupuncture or fluoxetine treatment The plasma levels of various cytokines are increased during ongoing depression and are reduced to normal levels after recovery Cytokines plasma levels during antidepressant treatment with sertraline and transcranial direct current stimulation (tDCS): results from a factorial, randomized, controlled trial Identifying who has long COVID in the USA: a machine learning approach using N3C data Propensity scores for confounder adjustment

when assessing the effects of medical interventions using nonexperimental study designs Effect of early treatment with fluvoxamine on risk of emergency care and hospitalisation among patients with COVID-19: the TOGETHER randomised, platform clinical trial Association between antidepressant use and reduced risk of intubation or death in hospitalized patients with COVID-19: results from an observational study Mortality risk among patients with COVID-19 prescribed selective serotonin reuptake inhibitor antidepressants Antidepressant use and risk of intubation or death in hospitalized patients with COVID-19: a retrospective cohort study of clinical effectiveness Fluvoxamine vs placebo and clinical deterioration in outpatients with symptomatic COVID-19: a randomized clinical trial Randomized trial of metformin, ivermectin, and fluvoxamine for Covid-19 Selective serotonin reuptake inhibitor (SSRI) antidepressants reduce COVID-19 infection: prospects for use Antidepressant drug prescription and incidence of COVID-19 in mental health outpatients: a retrospective cohort study Covid-19 mortality among selective serotonin reuptake inhibitor users—results from a nationwide cohort Repurposing selective serotonin reuptake inhibitors for severity of COVID-19: a population-based study Early fluvoxamine reduces the risk for clinical deterioration in symptomatic outpatients with Covid-19: a real-world, retrospective, before-after analysis Treatment of 95 post-covid patients with ssris Computational and Structural Biotechnology Journal. 2024. [PMC10839808](#)

64. Zachary Butzin-Dozier, Yunwen Ji, Sarang Deshpande, Eric Hurwitz, A. Jerrod Anzalone, Jeremy Coyle, Junming Shi, Andrew Mertens, Mark J. van der Laan, John M. Colford, Rena C. Patel, Alan E. Hubbard, et al.. SSRI use during acute COVID-19 and risk of Long COVID among patients with depression The epidemiology of long coronavirus disease in

US adults Long COVID: an overview Long COVID: major findings, mechanisms and recommendations Generalisable long COVID subtypes: findings from the NIH N3C and RECOVER programmes Effect of early treatment with fluvoxamine on risk of emergency care and hospitalisation among patients with COVID-19: the TOGETHER randomised, platform clinical trial Outpatient treatment of COVID-19 and incidence of post-COVID-19 condition over 10 months (COVID-OUT): a multicentre, randomised, quadruple-blind, parallel-group, phase 3 trial Selective serotonin reuptake inhibitor (SSRI) antidepressants reduce COVID-19 infection: prospects for use Antidepressant drug treatment protecting from COVID-19: one more piece in the repurposing puzzle Who develops long COVID? Longitudinal pre-pandemic predictors of long COVID and symptom clusters in a representative Dutch population Treatment of 95 post-COVID patients with SSRIs SSRI antidepressants differentially modulate serotonin reuptake and release in *Drosophila* Overview of the potential use of fluvoxamine for COVID-19 and long COVID Developing a targeted learning-based statistical analysis plan Use of the postacute sequelae of COVID-19 diagnosis code in routine clinical practice in the US Super learner Psychiatric comorbidity and other psychological factors in patients with “chronic Lyme disease” Role of psychiatric comorbidity in chronic Lyme disease Dysfunction in serotonergic and noradrenergic systems and somatic symptoms in psychiatric disorders Catecholamine/serotonin interactions: systems thinking for brain function and disease Restoration of serotonin neuronal firing following long-term administration of bupropion but not paroxetine in olfactory bulbectomized rats Escitalopram enhances the association of serotonin-1A autoreceptors to heteroreceptors in anxiety disorders Early dysfunction of central 5-HT system

in a murine model of bovine spongiform encephalopathy
Acute responsivity of the serotonergic system to S-citalopram
and positive emotionality-the moderating role of the 5-HTTLPR
Citalopram 20 mg, 40 mg and 60 mg are all effective
and well tolerated compared with placebo in obsessive-compulsive disorder
Citalopram intravenous infusion in resistant obsessive-compulsive disorder: an open trial
Paroxetine in the treatment of obsessive-compulsive disorder: randomized, double-blind, placebo-controlled study in Japanese patients
Obsessive-compulsive disorder post-COVID-19: a case presentation
Targeted maximum likelihood estimation for causal inference in observational studies
Identifying who has long COVID in the USA: a machine learning approach using N3C data
Coding long COVID: characterizing a new disease through an ICD-10 lens
BMC Medicine. 2024. [PMC11462648](#)

65. Poppy Alice Carson Mallinson, Manisha Joshi, Mahesh Mathpathi, Alexander Perkins, Tim Clayton, Anoop SV Shah, Rohini Mathur, Nick Birk, Arandeep Dhillon, Judith Lieber, Sidra S Beg, Lily Hopkins, Archie Khan, Shereen Allaham, Vanessa TW Kam, Shailen Sutaria, Galib R, S Rajagopala, Amarjeet Bhamra, Geetha Krishnan G Pillai, Kamlesh Khunti, Tanuja Nesari, Sanjay Kinra, et al.. Ashwagandha (*Withania somnifera* (L.) Dunal) for promoting recovery in long covid: protocol for a randomised placebo-controlled clinical trial (APRIL Trial)
The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis
NICE guideline on long COVID
Self-reported long covid symptoms
Prevalence of ongoing symptoms following coronavirus (covid-19) infection in the uk
Characterizing long COVID in an international cohort: 7 months of symptoms and their impact
Healthcare utilisation of 282,080 individuals with long covid

over two years: a multiple matched control cohort analysis
The impact of Long COVID on the UK workforce Where are
the long covid trials? - the lancet infectious diseases
Pathophysiology and mechanism of long COVID: a
comprehensive review Withania somnifera and Its
Withanolides Attenuate Oxidative and Inflammatory
Responses and Up-Regulate Antioxidant Responses in BV-2
Microglial Cells Withaferin A inhibits tumor necrosis factor-
alpha-induced expression of cell adhesion molecules by
inactivation of Akt and NF-kappaB in human pulmonary
epithelial cells Safety and clinical effectiveness of Withania
Somnifera (Linn.) Dunal root in human ailments Examining
the effect of Withania somnifera supplementation on muscle
strength and recovery: a randomized controlled trial Effects
of an Aqueous Extract of Withania somnifera on Strength
Training Adaptations and Recovery: The STAR Trial Effects of
Ashwagandha (Withania somnifera) on VO2max: A Systematic
Review and Meta-Analysis Effect of Ashwagandha (Withania
somnia) extract on sleep: A systematic review and meta-
analysis Effects of Acute Ashwagandha Ingestion on Cognitive
Function Does Ashwagandha supplementation have a
beneficial effect on the management of anxiety and stress? A
systematic review and meta-analysis of randomized controlled
trials Effect of Withania somnifera (Ashwagandha) on the
development of chemotherapy-induced fatigue and quality of
life in breast cancer patients Exploring the efficacy and safety
of a novel standardized ashwagandha (Withania somnifera)
root extract (Witholytin®) in adults experiencing high stress
and fatigue in a randomized, double-blind, placebo-controlled
trial THERAPEUTIC uses of withania somnifera
(ashwagandha) Pharmacologic overview of Withania
somnia, the Indian Ginseng Pharmacological evaluation of
Ashwagandha highlighting its healthcare claims, safety, and

toxicity aspects Efficacy and Safety of Ashwagandha (Withania somnifera) Root Extract in Insomnia and Anxiety: A Double-blind, Randomized, Placebo-controlled Study Clinical safety and tolerability evaluation of Withania somnifera (L.) Dunal (Ashwagandha) root extract in healthy human volunteers The Post-COVID-19 Functional Status scale: a tool to measure functional status over time after COVID-19 Recovery from COVID-19: a sprint or marathon? 6-month follow-up data from online long COVID-19 support group members Construct validity of the Post-COVID-19 Functional status scale in adult subjects with COVID-19 Measurement invariance and general population reference values of the PROMIS Profile 29 in the UK, France, and Germany Core Outcome Measures for Trials in People With Coronavirus Disease 2019: Respiratory Failure, Multiorgan Failure, Shortness of Breath, and Recovery Evaluation of clinical methods for rating dyspnea Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L) The Preventive Antibiotics in Stroke Study (PASS): a pragmatic randomised open-label masked endpoint clinical trial Remdesivir for the Treatment of Covid-19 - Final Report Where are the long COVID trials? The long covid evidence gap: comparing self-reporting and clinical coding of long covid using longitudinal study data linked to healthcare records Review Process File BMJ Open. 2025. [PMC12035422](#)

66. Sara Polo-Alonso, Álvaro Hernáez, Irene R. Dégano, Ruth Martí-Lluch, Mel·lina Pinsach-Abuin, Roberto Elosua, Isaac Subirana, Marta Puigmulé, Alexandra Pérez, Raquel Cruz, Silvia Diz-de Almeida, Eulàlia Puigdecant, Elisabet Selga, Xavier Nogues, Joan Ramon Masclans, Roberto Güerri-Fernández, Héctor Cubero-Gallego, Helena Tizon-Marcos, Beatriz Vaquerizo, Ramon Brugada, Rafel Ramos, Anna Camps-Vilaró, Jaume Marrugat, et al.. Global and Sex-

Stratified Genome-Wide Association Study of Long COVID
Based on Patient-Driven Symptom Recall Laboratory Findings
and Biomarkers in Long COVID: What Do We Know So Far?
Insights into Epidemiology, Pathogenesis, Therapeutic
Perspectives and Challenges Long COVID: A Clinical Update
The Epidemiology of Long Coronavirus Disease in US Adults
Long COVID Prevalence and Impact on Quality of Life 2 Years
after Acute COVID-19 The Risk of Long Covid Symptoms: A
Systematic Review and Meta-Analysis of Controlled Studies
Long COVID: Molecular Mechanisms and Detection
Techniques Determinants of the Onset and Prognosis of the
Post-COVID-19 Condition: A 2-Year Prospective Observational
Cohort Study Multiorgan Impairment in Low-Risk Individuals
with Post-COVID-19 Syndrome: A Prospective, Community-
Based Study Immunomodulatory Mechanisms Underlying
Neurological Manifestations in Long COVID: Implications for
Immune-Mediated Neurodegeneration Exploring the
Pathophysiology of Long COVID: The Central Role of Low-
Grade Inflammation and Multisystem Involvement Predictive
Factors and ACE-2 Gene Polymorphisms in Susceptibility to
Long COVID-19 Syndrome Genome-Wide Association Study of
Long COVID The Immunomodulatory Properties of the CD5
Lymphocyte Receptor in Health and Disease Chronic
Lymphocytic Leukemia Concomitant with COVID 19: A Case
Report Emergence of Chronic Lymphocytic Leukemia During
Admission for COVID-19: Cause or Coincidence? A New
Diagnosis of Monoclonal B-cell Lymphocytosis with
Cytoplasmic Inclusions in a Patient with COVID-19 Diagnosis
of Multisystem Inflammatory Syndrome in Children by a
Whole-Blood Transcriptional Signature Whole Exome
Sequencing of Patients from Multicase Families with Systemic
Lupus Erythematosus Identifies Multiple Rare Variants The
Panoramic Picture of Pepsinogen Gene Family with Pan-

Cancer VWCE as a Potential Biomarker Associated with Immune Infiltrates in Breast Cancer PTOV1 Facilitates Colorectal Cancer Cell Proliferation through Activating AKT1 Signaling Pathway Can Baseline IL-6 Levels Predict Long COVID in Subjects Hospitalized for SARS-CoV-2 Disease? Biallelic Loss of EMC10 Leads to Mild to Severe Intellectual Disability Novel LAMC3 Pathogenic Variant Enriched in Finnish Population Causes Malformations of Cortical Development and Severe Epilepsy Mid and Long-Term Neurological and Neuropsychiatric Manifestations of Post-COVID-19 Syndrome: A Meta-Analysis Microfluidic Isolation of Neuronal-Enriched Extracellular Vesicles Shows Distinct and Common Neurological Proteins in Long COVID, HIV Infection and Alzheimer's Disease The Influence of Microglia on Neuroplasticity and Long-Term Cognitive Sequelae in Long COVID: Impacts on Brain Development and Beyond Josephin Domain Containing 2 (JOSD2) Promotes Lung Cancer by Inhibiting LKB1 (Liver Kinase B1) Activity Role of Deubiquitinase JOSD2 in the Pathogenesis of Esophageal Squamous Cell Carcinoma Deubiquitinating Enzyme JOSD2 Promotes Hepatocellular Carcinoma Progression through Interacting with and Inhibiting CTNNB1 Degradation JOSD2 Regulates PKM2 Nuclear Translocation and Reduces Acute Myeloid Leukemia Progression Genomic and Transcriptomic Characterization of Gastric Cancer with Bone Metastasis Pepsinogen/Proton Pump Co-Expression in Barrett's Esophageal Cells Induces Cancer-Associated Changes Gene Expression Analysis for Uterine Cervix and Corpus Cancer Characterization LAMC1 Is a Prognostic Factor and a Potential Therapeutic Target in Endometrial Cancer Identification of a Basement Membrane-Related Gene Signature for Predicting Prognosis and Estimating the Tumor Immune Microenvironment in Breast Cancer Novel

Biomarkers Distinguishing Pancreatic Head Cancer from Distal Cholangiocarcinoma Based on Proteomic Analysis Genetic Characteristics Involved in COVID-19 Severity. The CARGENCORS Case-Control Study and Meta-Analysis World Health Organization Post COVID-19 Condition (Long COVID) Fatigue and Cognitive Impairment in Post-COVID-19 Syndrome: A Systematic Review and Meta-Analysis PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses Long-Range LD Can Confound Genome Scans in Admixed Populations Fast Model-Based Estimation of Ancestry in Unrelated Individuals A Global Reference for Human Genetic Variation Sequencing of 53,831 Diverse Genomes from the NHLBI TOPMed Program Functional Mapping and Annotation of Genetic Associations with FUMA ANNOVAR: Functional Annotation of Genetic Variants from High-Throughput Sequencing Data A General Framework for Estimating the Relative Pathogenicity of Human Genetic Variants Annotation of Functional Variation in Personal Genomes Using RegulomeDB The GTEx Consortium Atlas of Genetic Regulatory Effects across Human Tissues Extending Support for Mouse Data in the Molecular Signatures Database (MSigDB) Gene Set Enrichment Analysis: A Knowledge-Based Approach for Interpreting Genome-Wide Expression Profiles WikiPathways 2024: Next Generation Pathway Database The NHGRI-EBI GWAS Catalog: Knowledgebase and Deposition Resource MAGMA: Generalized Gene-Set Analysis of GWAS Data International Journal of Molecular Sciences. 2025. [PMC12471052](#)

67. Art Schuermans, Andreas Verstraete, Vilma Lammi, Tomoko Nakanishi, Maddalena Ardissino, Jef Van den Eynde, Benjamin Sun, Marios K Georgakis, Beatriz Guillen-Guio, Louise V. Wain, Christopher E. Brightling, Johan Van Weyenbergh, Adam J. Lewandowski, Betty Raman, Hugo

Zeberg, Hanna M Ollila, Stephen Burgess, Pradeep Natarajan, Michael C. Honigberg, Kathleen Freson, Thomas Vanassche, Peter Verhamme, et al.. Human genetics implicate thromboembolism in the pathogenesis of long COVID in individuals of European ancestry Human genetics implicate thromboembolism in the pathogenesis of long COVID in individuals of European ancestry medRxiv. 2024. [PMC11118620](#)

68. Stuart F. Quan, Matthew D. Weaver, Mark É. Czeisler, Laura K. Barger, Lauren A. Booker, Mark E. Howard, Melinda L. Jackson, Rashon I. Lane, Christine F. McDonald, Anna Ridgers, Rebecca Robbins, Prerna Varma, Joshua F. Wiley, Shantha M.W. Rajaratnam, Charles A. Czeisler, et al.. Association of Obstructive Sleep Apnea with Post-Acute Sequelae of SARS-CoV-2 infection Association of Obstructive Sleep Apnea with Post-Acute Sequelae of SARS-CoV-2 infection (PASC) Symptoms and signs of long COVID: A rapid review and meta-analysis The lasting misery of coronavirus long-haulers Assessment of the Frequency and Variety of Persistent Symptoms Among Patients With COVID-19: A Systematic Review Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments Long COVID Through a Public Health Lens: An Umbrella Review Risk Factors Associated With Post–COVID-19 Condition: A Systematic Review and Meta-analysis Sleep Apnea and COVID-19 Mortality and Hospitalization The association between high risk of sleep apnea, comorbidities, and risk of COVID-19: a population-based international harmonized study Obstructive sleep apnea (OSA) and outcomes from coronavirus disease 2019 (COVID-19) pneumonia: a systematic review and meta-analysis Obstructive Sleep Apnea and Risk of COVID-19 Infection, Hospitalization and Respiratory Failure Obstructive sleep apnea and COVID-19

clinical outcomes during hospitalization: a cohort study
Associations between obstructive sleep apnea and COVID-19
infection and hospitalization among US adults Long COVID:
major findings, mechanisms and recommendations Risk of
post-acute sequelae of SARS-CoV-2 infection associated with
pre-coronavirus disease obstructive sleep apnea diagnoses:
an electronic health record-based analysis from the
RECOVER initiative The Pittsburgh Sleep Quality Index: a
new instrument for psychiatric practice and research
Insomnia, Poor Sleep Quality and Sleep Duration, and Risk for
COVID-19 Infection and Hospitalization Long covid outcomes
at one year after mild SARS-CoV-2 infection: nationwide
cohort study Impaired health-related quality of life in long-
COVID syndrome after mild to moderate COVID-19
Development of a Definition of Postacute Sequelae of SARS-
CoV-2 Infection Protective effect of COVID-19 vaccination
against long COVID syndrome: A systematic review and meta-
analysis Obstructive Sleep Apnea and Inflammation: Proof of
Concept Based on Two Illustrative Cytokines Inflammatory
and oxidative stress-associated factors in chronic intermittent
hypoxia in Chinese patients, rats, lymphocytes and
endotheliocytes Coagulation and Fibrinolysis in Obstructive
Sleep Apnoea Obstructive Sleep Apnea, Hypercoagulability,
and the Blood-Brain Barrier Long COVID: pathophysiological
factors and abnormalities of coagulation Oxidative Stress
Markers among Obstructive Sleep Apnea Patients Cellular
Immune Dysfunction in Obstructive Sleep Apnea The
American journal of medicine. 2024. [PMC11144080](#)

69. Stuart F. Quan, Matthew D. Weaver, Mark É. Czeisler, Laura K. Barger, Lauren A. Booker, Mark E. Howard, Melinda L. Jackson, Rashon I. Lane, Christine F. McDonald, Anna Ridgers, Rebecca Robbins, Prerna Varma, Joshua F. Wiley, Shantha M.W. Rajaratnam, Charles A. Czeisler, et al..

Association of Obstructive Sleep Apnea with Post-Acute Sequelae of SARS-CoV-2 infection (PASC) Association of Obstructive Sleep Apnea with Post-Acute Sequelae of SARS-CoV-2 infection Symptoms and signs of long COVID: A rapid review and meta-analysis The lasting misery of coronavirus long-haulers Assessment of the Frequency and Variety of Persistent Symptoms Among Patients With COVID-19: A Systematic Review Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments Long COVID Through a Public Health Lens: An Umbrella Review Sleep Apnea and COVID-19 Mortality and Hospitalization The association between high risk of sleep apnea, comorbidities, and risk of COVID-19: a population-based international harmonized study Obstructive sleep apnea (OSA) and outcomes from coronavirus disease 2019 (COVID-19) pneumonia: a systematic review and meta-analysis Obstructive Sleep Apnea and Risk of COVID-19 Infection, Hospitalization and Respiratory Failure Obstructive sleep apnea and COVID-19 clinical outcomes during hospitalization: a cohort study Associations between obstructive sleep apnea and COVID-19 infection and hospitalization among US adults Obstructive Sleep Apnea and Inflammation: Proof of Concept Based on Two Illustrative Cytokines SARS-CoV-2 pathogenesis Long COVID: major findings, mechanisms and recommendations Impact of Obstructive Sleep Apnea (OSA) in COVID-19 Survivors, Symptoms Changes Between 4-Months and 1 Year After the COVID-19 Infection Risk of post-acute sequelae of SARS-CoV-2 infection associated with pre-coronavirus disease obstructive sleep apnea diagnoses: an electronic health record-based analysis from the RECOVER initiative Obstructive sleep apnea association with post-acute sequelae of Severe acute respiratory syndrome coronavirus 2: insights from the NIH researching coronavirus disease 2019

to enhance recovery initiative and a call for action The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research Insomnia, Poor Sleep Quality and Sleep Duration, and Risk for COVID-19 Infection and Hospitalization Long covid outcomes at one year after mild SARS-CoV-2 infection: nationwide cohort study Impaired health-related quality of life in long-COVID syndrome after mild to moderate COVID-19 Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Protective effect of COVID-19 vaccination against long COVID syndrome: A systematic review and meta-analysis Inflammatory and oxidative stress associated factors in chronic intermittent hypoxia in Chinese patients, rats, lymphocytes and endotheliocytes Coagulation and Fibrinolysis in Obstructive Sleep Apnoea Obstructive Sleep Apnea, Hypercoagulability, and the Blood-Brain Barrier Long COVID: pathophysiological factors and abnormalities of coagulation Oxidative Stress Markers among Obstructive Sleep Apnea Patients Cellular Immune Dysfunction in Obstructive Sleep Apnea medRxiv. 2023. [PMC10793517](https://pubmed.ncbi.nlm.nih.gov/PMC10793517/)

70. Chengxi Zang, Daniel Guth, Ann M. Bruno, Zhenxing Xu, Haoyang Li, Nariman Ammar, Robert Chew, Nick Guthe, Emily Hadley, Rainu Kaushal, Tanzy Love, Brenda M. McGrath, Rena C. Patel, Elizabeth C. Seibert, Yalini Senathirajah, Sharad Kumar Singh, Fei Wang, Mark G. Weiner, Kenneth J. Wilkins, Yiye Zhang, Torri D. Metz, Elaine Hill, Thomas W. Carton, ChengxiZhenxingRainuFeiMark G.Thomas W. ZangXuKaushalWangWeinerCarton, Chengxi Zang, Zhenxing Xu, Rainu Kaushal, Fei Wang, Mark G. Weiner, Thomas W. Carton, DanielNarimanRobertEmilyTanzyBrenda M.Rena C.Sharad KumarKenneth J.Elaine GuthAmmarChewHadleyLoveMcGrathPatelSinghWilkinsHill,

Daniel Guth, Nariman Ammar, Robert Chew, Emily Hadley, Tanzy Love, Brenda M. McGrath, Rena C. Patel, Sharad Kumar Singh, Kenneth J. Wilkins, Elaine Hill, Ann M. Torri D. Bruno Metz, Ann M. Bruno, Torri D. Metz, et al.. Long COVID after SARS-CoV-2 during pregnancy in the United States Solving the puzzle of Long Covid Long COVID: major findings, mechanisms and recommendations Identification of risk factors of Long COVID and predictive modeling in the RECOVER EHR cohorts Identifying environmental risk factors for post-acute sequelae of SARS-CoV-2 infection: an EHR-based cohort study from the recover program Excess burden of respiratory and abdominal conditions following COVID-19 infections during the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER program Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis Pregnancy-specific responses to COVID-19 revealed by high-throughput proteomics of human plasma Association of SARS-CoV-2 infection with serious maternal morbidity and mortality from obstetric complications Disease severity and perinatal outcomes of pregnant patients with coronavirus disease 2019 (COVID-19) Severe maternal morbidity and mortality of pregnant patients with COVID-19 infection during the early pandemic period in the US The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis Association between acquiring SARS-CoV-2 during pregnancy and post-acute sequelae of SARS-CoV-2 infection: RECOVER electronic health record cohort analysis Long term implications of covid-19 in pregnancy Data-driven analysis to understand long COVID using electronic health records from the RECOVER initiative Identifying who has long COVID in the USA: a machine learning approach using N3C data

Pregnancy immune tolerance at the maternal-fetal interface
Effects of influenza on pregnant women and infants
Synergies between centralized and federated approaches to data quality: a report from the national COVID cohort collaborative
The National COVID Cohort Collaborative (N3C): rationale, design, infrastructure, and deployment
Identification of delivery encounters using international classification of diseases, tenth revision, diagnosis and procedure codes
Validation of ICD-10-CM diagnosis codes for gestational age at birth
High-dimensional characterization of post-acute sequelae of COVID-19
Making neighborhood-disadvantage metrics accessible—the neighborhood Atlas
High-throughput target trial emulation for Alzheimer’s disease drug repurposing with real-world data
Moving towards best practice when using inverse probability of treatment weighting (IPTW) using the propensity score to estimate causal treatment effects in observational studies
An empirical transition matrix for non-homogeneous markov chains based on censored observations
Nature Communications. 2025.
[PMC11961632](#)

71. Ann M. Bruno, Chengxi Zang, Zhengxing Xu, Fei Wang, Mark G. Weiner, Nick Guthe, Megan Fitzgerald, Rainu Kaushal, Thomas W. Carton, Torri D. Metz, et al.. Association between acquiring SARS-CoV-2 during pregnancy and post-acute sequelae of SARS-CoV-2 infection: RECOVER electronic health record cohort analysis
Coronavirus disease 2019
COVID-19 treatment guidelines
A clinical case definition of post COVID-19 condition by a Delphi consensus
Long COVID, a comprehensive systematic scoping review
Risk factors associated with post-acute sequelae of SARS-CoV-2 in an EHR cohort: a national COVID cohort collaborative (N3C) analysis as part of the NIH RECOVER program
Researching COVID to enhance recovery (RECOVER) pregnancy study: rationale,

objectives and design Association of SARS-CoV-2 infection with serious maternal morbidity and mortality from obstetric complications Clinical presentation of coronavirus disease 2019 (COVID-19) in pregnant and recently pregnant people Data-driven analysis to understand long COVID using electronic health records from the RECOVER initiative Changing the research landscape: the New York City clinical data research Network Identification of delivery encounters using international classification of diseases, Tenth revision, diagnosis and procedure codes Validation of ICD-10-CM diagnosis codes for gestational age at birth ACOG Practice Bulletin, No. 233. American college of obstetricians and gynecologists Post-acute COVID-19 syndrome Comorbidity measures for use with administrative data Long COVID risk and pre-COVID vaccination in an EHR-based cohort study from the RECOVER program Reductions in stillbirths and preterm birth in COVID-19-vaccinated women: a multicenter cohort study of vaccination uptake and perinatal outcomes Symptom prevalence, duration, and risk of hospital admission in individuals infected with SARS-CoV-2 during periods of omicron and delta variant dominance: a prospective observational study from the ZOE COVID Study ACOG Practice Bulletin, No. 196. American College of Obstetrician and Gynecologists Adjusting for multiple testing--when and how? The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study Development of a definition of postacute sequelae of SARS-CoV-2 infection Data-driven identification of post-acute SARS-CoV-2 infection subphenotypes Symptoms and risk factors for

long COVID in non-hospitalized adults Long covid-mechanisms, risk factors, and management A statewide population-based approach to examining Long COVID symptom prevalence and predictors in Michigan Comparative analysis of long-term self-reported COVID-19 symptoms among pregnant women Clinical characteristics and outcomes of hospitalized women giving birth with and without COVID-19 Disease severity and perinatal outcomes of pregnant patients with coronavirus disease 2019 (COVID-19) Cardiovascular complications of pregnancy-associated COVID-19 infections Clinical risk factors of adverse outcomes among women with COVID-19 in the pregnancy and postpartum period: a sequential, prospective meta-analysis The immune system in pregnancy: a unique complexity Understanding the immune system in fetal protection and maternal infections during pregnancy The influence of COVID-19 infection-associated immune response on the female reproductive system Moving towards best practice when using inverse probability of treatment weighting (IPTW) using the propensity score to estimate causal treatment effects in observational studies eClinicalMedicine. 2024. [PMC11137338](#)

72. Torri D Metz, Harrison T Reeder, Rebecca G Clifton, Valerie Flaherman, Leyna V Aragon, Leah Castro Baucom, Carmen J Beamon, Alexis Braverman, Jeanette Brown, Megan Carmilani, Tingyi Cao, Ann Chang, Maged M Costantine, Jodie A. Dionne, Kelly S Gibson, Rachel S Gross, Estefania Guerreros, Mounira Habli, Rachel Hess, Leah Hillier, Sally Hodder, M Camille Hoffman, Matthew K Hoffman, Weixing Huang, Brenna L Hughes, Xiaolin Jia, Minal Kale, Stuart D Katz, Victoria Laleau, Hector Mendez-Figueroa, Grace A McComsey, Igho Ofotokun, Megumi J Okumura, Luis D Pacheco, Anna Palatnik, Kristy T S Palomares, Samuel Parry,

Christian M Pettker, Beth A Plunkett, Athena Poppas, Patrick Ramsey, Uma M Reddy, Dwight J Rouse, George R Saade, Grecio J Sandoval, Frank Sciurba, Hyagriv N Simhan, Daniel W Skupski, Amber Sowles, John M Thorp, Alan TN Tita, Samantha Wiegand, Steven J Weiner, Lynn M Yee, Leora I Horwitz, Andrea S Foulkes, Vanessa L Jacoby, et al.. Long COVID Following Acquisition of the Omicron Variant of SARS-CoV-2 During Pregnancy Compared with Outside of Pregnancy Pathophysiological, immunological, and inflammatory features of long COVID Multiomic immune clockworks of pregnancy Maternal immunological adaptation during normal pregnancy Female Sex Is a Risk Factor Associated with Long-Term Post-COVID Related-Symptoms but Not with COVID-19 Symptoms: The LONG-COVID-EXP-CM Multicenter Study Sex Differences in Long COVID Association between acquiring SARS-CoV-2 during pregnancy and post-acute sequelae of SARS-CoV-2 infection: RECOVER electronic health record cohort analysis Post-Acute Sequelae of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) After Infection During Pregnancy Post-Viral Fatigue Following SARS-CoV-2 Infection during Pregnancy: A Longitudinal Comparative Study RECOVER Initiative. Researching COVID to enhance recovery (RECOVER) pregnancy study: Rationale, objectives and design Researching COVID to Enhance Recovery (RECOVER) adult study protocol: Rationale, objectives, and design 2024 Update of the RECOVER-Adult Long COVID Research Index Using full matching to estimate causal effects in nonexperimental studies: examining the relationship between adolescent marijuana use and adult outcomes mice: Multivariate Imputation by Chained Equations in R MatchThem:: Matching and Weighting after Multiple Imputation Long COVID after SARS-CoV-2 during Pregnancy in the United States Acute and

post-acute sequelae of SARS-CoV-2 infection: a review of risk factors and social determinants Pre-existing conditions associated with post-acute sequelae of COVID-19 Prevalence and risk factors of post-acute sequelae of COVID-19 among United States Veterans Obstetrics and gynecology. 2026. [PMC12915694](#)

73. Yinghui Wei, Elsie MF Horne, Rochelle Knight, Genevieve Cezard, Alex J Walker, Louis Fisher, Rachel Denholm, Kurt Taylor, Venexia Walker, Stephanie Riley, Dylan M Williams, Robert Willans, Simon Davy, Sebastian Bacon, Ben Goldacre, Amir Mehrkar, Spiros Denaxas, Felix Greaves, Richard J Silverwood, Aziz Sheikh, Nish Chaturvedi, Angela M Wood, John Macleod, Claire Steves, Jonathan Sterne, et al.. Patient characteristics associated with clinically coded long COVID: an OpenSAFELY study using electronic health records Long COVID: aiming for a consensus High-dimensional characterization of post-acute sequelae of COVID-19 Post-COVID syndrome (long COVID) A clinical case definition of post-COVID-19 condition by a Delphi consensus COVID-19 rapid guideline: managing the long-term effects of COVID-19 Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records in situ using opensafely Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Attributes and predictors of long COVID Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK Primary care post-COVID syndrome diagnosis and referral coding [Preprint] Long COVID: the NHS plan for 2021/22 Changes to COVID-19 testing in England from 1 April Nutrition for a healthy life - WHO recommendations Eurostat - An overview of the 3 NUTS and 2 LAU layers in the UK Association of COVID-19 with major arterial and venous thrombotic diseases: a population-wide cohort study of 48

million adults in England and Wales On the C-statistics for evaluating overall adequacy of risk prediction procedures with censored survival data opensafely/long-covid-risk-factors-and-prediction Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK Comparative effectiveness of BNT162b2 versus mRNA-1273 covid-19 vaccine boosting in England: matched cohort study in OpenSAFELY-TPP OpenSAFELY: representativeness of electronic health record platform OpenSAFELY-TPP data compared to the population of England Spatial distribution of clinical computer systems in primary care in England in 2016 and implications for primary care electronic medical record databases: a cross-sectional population study Long covid symptoms and diagnosis in primary care: a cohort study using structured and unstructured data in the Health Improvement Network primary care database The table 2 fallacy: presenting and interpreting confounder and modifier coefficients Symptoms and risk factors for long COVID in non-hospitalized adults Long-term effects of COVID-19 on workers in health and social services in Germany Prevalence and risk factors of post-COVID-19 condition in adults and children at 6 and 12 months after hospital discharge: a prospective, cohort study in Moscow (stopCOVID) ONS: Short report on Long COVID Risk factors for post-COVID-19 condition in previously hospitalised children using the ISARIC global follow-up protocol: a prospective cohort study Long COVID in hospitalized and non-hospitalized patients in a large cohort in Northwest Spain, a prospective cohort study Prognostic factors for post-COVID-19 syndrome: a systematic review and meta-analysis The NHS England OpenSAFELY COVID-19 service - privacy notice Data Security and Protection Toolkit - NHS Digital ISB1523: Anonymisation Standard for Publishing

Health and Social Care Data Health Research Authority BJGP
Open. 2025. [PMC12820521](#)

74. Yun Lu, Arnstein Lindaas, Hector S. Izurieta, Myrna Cozen, Mikhail Menis, Xiangyu Shi, Whitney R. Steele, Michael Wernecke, Yoganand Chillarige, Jeffrey A. Kelman, Richard A. Forshee, et al.. Lessons Learned From Characterizing Long COVID Among US Medicare Beneficiaries “COVID-19 Mortality Overview,” Post-COVID Conditions Among Adult COVID-19 Survivors Aged 18–64 and ≥65 Years—United States, March 2020–November 2021 Persistent Symptoms in Patients After Acute COVID-19 Short-Term and Long-Term Rates of Postacute Sequelae of SARS-CoV-2 Infection: A Systematic Review Post-Acute COVID-19 Syndrome Sequelae in Adults at 6 Months After COVID-19 Infection Symptoms and Risk Factors for Long COVID in Non-Hospitalized Adults High-Dimensional Characterization of Post-Acute Sequelae of COVID-19 “Long COVID or Post-COVID Conditions,” “HHS Releases Long COVID Report Providing Insights and Opportunities to Support Patient Community,” RECOVER Consortium Coding Long COVID: Characterizing a New Disease Through an ICD-10 Lens Symptoms and Functional Impairment Assessed 8 Months After Mild COVID-19 Among Health Care Workers Sequelae in Adults at 12 Months After Mild-to-Moderate Coronavirus Disease 2019 (COVID-19) Post-Acute and Long-COVID-19 Symptoms in Patients With Mild Diseases: A Systematic Review Post-COVID Syndrome in Non-Hospitalised Patients With COVID-19: A Longitudinal Prospective Cohort Study Risk of Persistent and New Clinical Sequelae Among Adults Aged 65 Years and Older During the Post-Acute Phase of SARS-CoV-2 Infection: Retrospective Cohort Study Rates and Factors Associated With Documentation of Diagnostic Codes for Long COVID in the National Veterans Affairs Health Care System “Nearly One in

Five American Adults Who Have Had COVID-19 Still Have 'Long COVID'," Long COVID: A Clinical Update "Guidance for Industry: Real-World Data: Assessing Electronic Health Records and Medical Claims Data to Support Regulatory Decision Making for Drug and Biological Products," 2024 How Common Is Long COVID? Why Studies Give Different Answers An Intersectional Analysis of Long COVID Prevalence Persistent COVID-19 Symptoms in a Community Study of 606,434 People in England Acute and Post-Acute Sequelae of SARS-CoV-2 Infection: A Review of Risk Factors and Social Determinants Prevalence and Correlates of Long COVID Symptoms Among US Adults Is Myopathy Part of Long-Covid? Myopathy as a Cause of Fatigue in Long-Term Post-COVID-19 Symptoms: Evidence of Skeletal Muscle Histopathology Pharmacoepidemiology and Drug Safety. 2025. [PMC11753895](#)

75. Alasdair D. Henderson, Ben FC. Butler-Cole, John Tazare, Laurie A. Tomlinson, Michael Marks, Mark Jit, Andrew Briggs, Liang-Yu Lin, Oliver Carlile, Chris Bates, John Parry, Sebastian CJ. Bacon, Iain Dillingham, William A. Dennison, Ruth E. Costello, Yinghui Wei, Alex J. Walker, William Hulme, Ben Goldacre, Amir Mehrkar, Brian MacKenna, AlexAmeliaAmirAndreaAndrewBenBenBrianCarolineCarolineCatherineChr WalkerGreenMehrkarSchafferBrownGoldacreButler-ColeMacKennaMortonWaltersStablesCunninghamWoodAndrewsEvansHickman, Alex Walker, Amelia Green, Amir Mehrkar, Andrea Schaffer, Andrew Brown, Ben Goldacre, Ben Butler-Cole, Brian MacKenna, Caroline Morton, Caroline Walters, Catherine Stables, Christine Cunningham, Christopher Wood, Colm Andrews, David Evans, George Hickman, Helen Curtis, Henry Drysdale, Iain Dillingham, Jessica Morley, Jon Massey, Linda Nab, Lisa Hopcroft, Louis Fisher, Lucy Bridges, Milan Wiedemann, Nicholas DeVito, Orla Macdonald, Peter Inglesby,

Rebecca Smith, Richard Croker, Robin Park, Rose Higgins, Sebastian Bacon, Simon Davy, Steven Maude, Thomas O'Dwyer, Tom Ward, Victoria Speed, William Hulme, Liam Hart, Pete Stokes, Krishnan Bhaskaran, Ruth Costello, Thomas Cowling, Ian Douglas, Rosalind Eggo, Stephen Evans, Harriet Forbes, Richard Grieve, Daniel Grint, Emily Herrett, Sinead Langan, Viyaasan Mahalingasivam, Kathryn Mansfield, Rohini Mathur, Helen McDonald, Edward Parker, Christopher Rentsch, Anna Schultze, Liam Smeeth, John Tazare, Laurie Tomlinson, Jemma Walker, Elizabeth Williamson, Kevin Wing, Angel Wong, Bang Zheng, Christopher Bates, Jonathan Cockburn, John Parry, Frank Hester, Sam Harper, Shaun O'Hanlon, Alex Eavis, Richard Jarvis, Dima Avramov, Paul Griffiths, Aaron Fowles, Nasreen Parkes, Rafael Perera, David Harrison, Kamlesh Khunti, Jonathan Sterne, Jennifer Quint, Emily Herrett, Rosalind M. Eggo, et al.. Clinical coding of long COVID in primary care 2020–2023 in a cohort of 19 million adults: an OpenSAFELY analysis Long COVID: major findings, mechanisms and recommendations Excess risk and clusters of symptoms after COVID-19 in a large Norwegian cohort Why we need a deeper understanding of the pathophysiology of long COVID The immunology of long COVID Innate immune cell activation causes lung fibrosis in a humanized model of long COVID SARS-CoV-2 airway infection results in the development of somatosensory abnormalities in a hamster model Viral persistence, reactivation, and mechanisms of long COVID How “long covid” is shedding light on postviral syndromes The prevalence of stigma in a UK community survey of people with lived experience of long COVID Short-term and long-term impacts of COVID-19 on economic vulnerability: a population-based longitudinal study (COVIDENCE UK) Conceptualising the episodic nature of disability among adults living with Long COVID: a qualitative

study Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK - Office for National Statistics Attributes and predictors of long COVID Persistent COVID-19 symptoms in a community study of 606,434 people in England Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Post-COVID-19 condition and persisting symptoms in English schoolchildren: repeated surveys to March 2022 The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study Long covid outcomes at one year after mild SARS-CoV-2 infection: nationwide cohort study Persistence of somatic symptoms after COVID-19 in The Netherlands: an observational cohort study Post-acute effects of SARS-CoV-2 infection in individuals not requiring hospital admission: a Danish population-based cohort study Long COVID: 3 years in Data resource profile: clinical practice research datalink (CPRD) Promises and pitfalls of electronic health record analysis The Danish civil registration system as a tool in epidemiology Big data from electronic health records for early and late translational cardiovascular research: challenges and potential Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records in situ using OpenSAFELY A retrospective cohort analysis leveraging augmented intelligence to characterize long COVID in the electronic health record: a precision medicine framework Identifying who has long COVID in the USA: a machine learning approach using N3C data Coding long COVID: characterizing a new disease through an ICD-10 lens Data-driven analysis to understand long COVID using electronic health records from the RECOVER initiative Next-generation

phenotyping of electronic health records Avoidable flaws in observational analyses: an application to statins and cancer Patient online management information (POMI) Patient and public involvement within epidemiological studies of long COVID in the UK Protocol for an OpenSAFELY cohort study collecting patient-reported outcome measures using the TPP Airmid smartphone application and linked big data to quantify the health and economic costs of long COVID (OpenPROMPT) SNOMED CT Graphical depiction of longitudinal study designs in health care databases OpenCodelists: NICE managing the long-term effects of COVID-19 OpenCodelists: referral and signposting for long COVID OpenCodelists: assessment instruments and outcome measures for long covid Comparative effectiveness of two- and three-dose COVID-19 vaccination schedules involving AZD1222 and BNT162b2 in people with kidney disease: a linked OpenSAFELY and UK Renal Registry cohort study Comparative effectiveness of BNT162b2 versus mRNA-1273 covid-19 vaccine boosting in England: matched cohort study in OpenSAFELY-TPP OpenCodelists: ethnicity (SNOMED) OpenCodelists: high risk from COVID-19 code The inverse care law The 'five safes' framework and applying it to OpenSAFELY | Bennett Institute for applied data science Development and validation of an algorithm to accurately identify atopic eczema patients in primary care electronic health records from the UK CODE-EHR best-practice framework for the use of structured electronic health-care records in clinical research Importance of accurately identifying disease in studies using electronic health records The long COVID evidence gap: comparing self-reporting and clinical coding of long COVID using longitudinal study data linked to healthcare records Long Covid symptoms and diagnosis in primary care: a cohort study using structured and unstructured data in the Health

Improvement Network primary care database Potential pitfalls in the use of real-world data for studying long COVID Early experiences of the Your COVID Recovery® digital programme for individuals with long COVID Public use of the “Your COVID Recovery” website designed to help individuals manage their COVID-19 Recovery: observational study Deriving and validating a risk prediction model for long COVID-19: protocol for an observational cohort study using linked Scottish data Differences in clinical presentation with long COVID after community and hospital infection and associations with all-cause mortality: English sentinel network database study Long COVID and significant activity limitation among adults, by age — United States, June 1–13, 2022, to June 7–19, 2023 Trajectory of long covid symptoms after covid-19 vaccination: community based cohort study Protective effect of COVID-19 vaccination against long COVID syndrome: a systematic review and meta-analysis Risk factors associated with Post–COVID-19 condition: a systematic review and meta-analysis Effect of covid-19 vaccination on long covid: systematic review Long COVID in a highly vaccinated population infected during a SARS-CoV-2 Omicron wave - Australia, 2022 The effectiveness of COVID-19 vaccines to prevent long COVID symptoms: staggered cohort study of data from the UK, Spain, and Estonia Efficacy of first dose of covid-19 vaccine versus no vaccination on symptoms of patients with long covid: target trial emulation based on ComPaRe e-cohort A prospective cohort study assessing the relationship between long-COVID symptom incidence in COVID-19 patients and COVID-19 vaccination Association between BNT162b2 vaccination and reported incidence of post-COVID-19 symptoms: cross-sectional study 2020-21, Israel Vaccination status and long COVID symptoms in patients discharged from hospital The impact of COVID

vaccination on symptoms of long COVID: an international survey of people with lived experience of long COVID Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study Six-month sequelae of post-vaccination SARS-CoV-2 infection: a retrospective cohort study of 10,024 breakthrough infections Post-acute sequelae after severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection by viral variant and vaccination status: a multicenter cross-sectional study Severe fatigue and persistent symptoms at three months following SARS-CoV-2 infections during the pre-delta, delta, and omicron time periods: a multicenter prospective cohort study Heterologous infection and vaccination shapes immunity against SARS-CoV-2 variants Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Prevalence and risk factors for long COVID among adults in Scotland using electronic health records: a national, retrospective, observational cohort study Emerging evidence on heterologous COVID-19 vaccine schedules—to mix or not to mix? Homologous and heterologous covid-19 booster vaccinations Effectiveness of heterologous and homologous covid-19 vaccine regimens: living systematic review with network meta-analysis Effectiveness of COVID-19 booster vaccines against COVID-19-related symptoms, hospitalization and death in England Trends and clinical characteristics of COVID-19 vaccine recipients: a federated analysis of 57.9 million patients' primary care records in situ using OpenSAFELY Factors associated with COVID-19-related death using OpenSAFELY eClinicalMedicine. 2024. [PMC11127160](#)

76. Kin Wah Fung, Fitsum Baye, Seo H. Baik, Zhaonian Zheng, Clement J. McDonald, Philippa Dodd, Philippa Dodd, Philippa Dodd, Philippa Dodd, et al.. Prevalence and characteristics of

long COVID in elderly patients: An observational cohort study of over 2 million adults in the US High-dimensional characterization of post-acute sequelae of COVID-19 Long-term Health Consequences of COVID-19 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records Long-Term Symptoms Among Adults Tested for SARS-CoV-2—United States, January 2020–April 2021. More than 50 long-term effects of COVID-19: a systematic review and meta-analysis. Persistent Symptoms in Patients After Acute COVID-19 Frequency, signs and symptoms, and criteria adopted for long COVID-19: A systematic review. "LONG COVID": an insight. Long COVID, a comprehensive systematic scoping review. Short-term and Long-term Rates of Postacute Sequelae of SARS-CoV-2 Infection: A Systematic Review Challenges in defining Long COVID: Striking differences across literature, Electronic Health Records, and patient-reported information. COVID Symptoms, Symptom Clusters, and Predictors for Becoming a Long-Hauler: Looking for Clarity in the Haze of the Pandemic. Four-Month Clinical Status of a Cohort of Patients After Hospitalization for COVID-19 Population-Based Estimates of Post-acute Sequelae of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection (PASC) Prevalence and Characteristics. Post-acute effects of SARS-CoV-2 infection in individuals not requiring hospital admission: a Danish population-based cohort study Post-Acute Sequelae of SARS-CoV-2 Infection Among Adults Aged $\geq$ 18 Years—Long Beach, California, April 1–December 10, 2020. Sequelae in Adults at 6 Months After COVID-19 Infection Point Prevalence Estimates of Activity-Limiting Long-Term Symptoms among U.S. Adults $\geq$ 1 Month After Reported

SARS-CoV-2 Infection, November 1, 2021. Attributes and predictors of long COVID Long-COVID: An evolving problem with an extensive impact Risk of persistent and new clinical sequelae among adults aged 65 years and older during the post-acute phase of SARS-CoV-2 infection: retrospective cohort study Health-related quality of life, functional decline, and long-term mortality in older patients following hospitalisation due to COVID-19. Post-infectious disease syndrome. Long COVID or Post-acute Sequelae of COVID-19 (PASC): An Overview of Biological Factors That May Contribute to Persistent Symptoms. Taking pandemic sequelae seriously: from the Russian influenza to COVID-19 long-haulers Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: A systematic review and meta-analysis. Long term outcomes in survivors of epidemic Influenza A (H7N9) virus infection. Postacute COVID-19: An Overview and Approach to Classification. The Charlson comorbidity index is adapted to predict costs of chronic disease in primary care patients. A general introduction to adjustment for multiple comparisons Application of the Logistic Function to Bio-Assay Using Propensity Scores to Help Design Observational Studies: Application to the Tobacco Litigation. The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis. Identifying who has long COVID in the USA: a machine learning approach using N3C data Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records in situ using OpenSAFELY. Coding Long COVID: Characterizing a new disease through an ICD-10 lens. Post-COVID Syndrome: An Insight on Its

Pathogenesis. Neuroinvasion of SARS-CoV-2 in human and mouse brain The cognitive consequences of the COVID-19 epidemic: collateral damage COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-Up: JACC State-of-the-Art Review Coagulopathy and Antiphospholipid Antibodies in Patients with Covid-19 Pulmonary function and radiological features 4 months after COVID-19: first results from the national prospective observational Swiss COVID-19 lung study Effect of common maintenance drugs on the risk and severity of COVID-19 in elderly patients. Decision Letter 0 Decision Letter 1 Decision Letter 2 Decision Letter 3 PLOS Medicine. 2023. [PMC10150975](https://doi.org/10.1371/journal.pmed.1015097)

77. Xiaoli Jia, Jingjing Wang, Xiaomeng Cui, Ting Li, Yan Tian, Rui Lu, Yindi Tian, Xin Shen, Shuangsoo Dang, Wenjun Wang, et al.. Trends in long COVID among US adults, 2022–2024 A clinical case definition of post-COVID-19 condition by a Delphi consensus Global prevalence of long COVID, its subtypes and risk factors: an updated systematic review and meta-analysis Long COVID in adults: United States, 2022 Prevalence of and factors associated with long COVID among US adults: a nationwide survey From emergency response to long-term management: the many faces of the endemic state of COVID-19 Long COVID and significant activity limitation among adults, by age - United States, June 1-13, 2022, to June 7-19, 2023 The prevalence and long-term health effects of long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records The Behavioral Risk Factor Surveillance System Determinants of recovery from post-COVID-19 dyspnoea: analysis of UK prospective cohorts of hospitalised COVID-19 patients and community-based

controls Effect of covid-19 vaccination on long covid: systematic review Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: a systematic review Protective effect of COVID-19 vaccination against long COVID syndrome: a systematic review and meta-analysis COVID-19 booster dose vaccination coverage and factors associated with booster vaccination among adults, United States, march 2022 Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Acute and Postacute COVID-19 outcomes among immunologically naive adults during Delta vs omicron waves Association of Treatment with Nirmatrelvir and the risk of post-COVID-19 condition Attributes and predictors of long COVID Integrated immune, hormonal, and transcriptomic profiling reveals sex-specific dysregulation in long COVID patients with ME/CFS Sex differences in long COVID Gender and the pandemic: associations between caregiving, working from home, personal and career outcomes for women and men Sex differences in symptomatology and immune profiles of long COVID Association between county-level risk groups and COVID-19 outcomes in the United States: a socioecological study A systematic review of racial/ethnic and socioeconomic disparities in COVID-19 Associations of psychological wellbeing with COVID-19 hospitalization and mortality in adults aged 50 years or older from 25 European countries and Israel Social determinants of recovery from ongoing symptoms following COVID-19 in two UK longitudinal studies: a prospective cohort study Impact of fatigue as the primary determinant of functional limitations among patients with post-COVID-19 syndrome: a cross-sectional observational study Association of Post-COVID-19 condition symptoms and employment status Frontiers in Public Health. 2026. [PMC13136101](#)

78. Kimberly H. Nguyen, Yingjun Bao, Julie Mortazavi, Jennifer D. Allen, Patricia O. Chocano-Bedoya, Laura Corlin. Prevalence and Factors Associated with Long COVID Symptoms among U.S. Adults, 2022 Statement on the Fifteenth Meeting of the IHR (2005) Emergency Committee on the COVID-19 Pandemic WHO Chief Declares End to COVID-19 as a Global Health Emergency COVID Data Tracker Long COVID or Post-COVID Conditions A Clinical Case Definition of Post COVID-19 Condition by a Delphi Consensus 2021 Long COVID and mental health correlates: A new chronic condition fits existing patterns Impact of Long COVID on health and quality of life Measuring the impact of COVID-19 on the quality of life of the survivors, partners and family members: A cross-sectional international online survey What Are the Implications of Long COVID for Employment and Health Coverage? Racial/Ethnic Disparities in Post-acute Sequelae of SARS-CoV-2 Infection in New York: An EHR-Based Cohort Study from the RECOVER Program People with Certain Medical Conditions Mental health symptoms and association with COVID-19 vaccination receipt and intention to vaccinate among adults, United States Disparities in COVID-19 Vaccination Status, Intent, and Perceived Access for Noninstitutionalized Adults, by Disability Status—National Immunization Survey Adult COVID Module, United States, 30 May–26 June 2021 Food Insecurity and COVID-19 Vaccination Status and Vaccination Hesitancy in the United States Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: A systematic review Trajectory of long covid symptoms after covid-19 vaccination: Community based cohort study 2022 BRFSS Survey Data and Documentation Census Regions and Divisions of the United States Prevalence of COVID-19 and long COVID—Results from the 2022 Behavioral Risk Factor Surveillance System, 50 States

Associations Between Long COVID and Healthcare Access: A Cross-Sectional Analysis of the Behavioral Risk Factor Surveillance System (BRFSS) 2022 Disparities in national and state estimates of COVID-19 vaccination receipt and intent to vaccinate by race/ethnicity, income, and age group among adults ≥ 18 years, United States COVID-19 Vaccination and Intent for Vaccination of Adults with Reported Medical Conditions Sex differences in sequelae from COVID-19 infection and in long COVID syndrome: A review Long COVID risk—A signal to address sex hormones and women's health Prevalence and Correlates of Long COVID Symptoms Among US Adults Attributes and predictors of long COVID Distinguishing features of long COVID identified through immune profiling Characterising long COVID: A living systematic review Long COVID Through a Public Health Lens: An Umbrella Review End of the Federal COVID-19 Public Health Emergency (PHE) Declaration Long COVID in Adults: United States, 2022 Long COVID and Significant Activity Limitation Among Adults, by Age—United States, June 1–13, 2022, to June 7–19, 2023 Guidance on “Long COVID” as a Disability Under the ADA, Section 504, and Section 1557 Vaccines. 2024. [PMC10820629](#)

79. Claire E. Hastie, David J. Lowe, Andrew McAuley, Nicholas L. Mills, Andrew J. Winter, Corri Black, Janet T. Scott, Catherine A. O'Donnell, David N. Blane, Susan Browne, Tracy R. Ibbotson, Jill P. Pell, et al.. True prevalence of long-COVID in a nationwide, population cohort study The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study Physical and mental health

3 months after SARS-CoV-2 infection (long COVID) among adolescents in England (CLOcK): a national matched cohort study Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Post-covid medical complaints following infection with SARS-CoV-2 Omicron vs Delta variants Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study What is the recovery rate and risk of long-term consequences following a diagnosis of COVID-19? A harmonised, global longitudinal observational study protocol Nature Communications. 2023. [PMC10689486](#)

80. Lauren E. Wisk, Michelle L'Hommedieu, Kate Diaz Roldan, Imtiaz Ebna Mannan, Erica S. Spatz, Robert A. Weinstein, Arjun K. Venkatesh, Michael Gottlieb, Ryan Huebinger, Kristin L. Rising, Juan Carlos C. Montoy, Kari A. Stephens, Robert M. Rodriguez, Mandy J. Hill, Kelli N. O'Laughlin, Nicole L. Gentile, Ahamed H. Idris, Shu-Xia Li, Michelle Santangelo, Efrat R. Kean, Samuel A. McDonald, Kristyn Gatling, Joann G. Elmore, et al.. Variability in Long COVID Definitions and Validation of Published Prevalence Rates A clinical case definition of post-COVID-19 condition by a Delphi consensus Development of a definition of postacute sequelae of SARS-CoV-2 infection Prevalence of long-term symptoms varies when using different post-COVID-19 definitions in positively and negatively tested adults: the PRIME post-COVID study Definition of post-COVID-19 condition among published research studies Toward a universal definition of post-COVID-19 condition—how do we proceed? Long COVID definitions and models of care: a scoping review Identification and diagnosis of long COVID-19: a scoping review Toward unbiased evaluation of postacute sequelae of SARS-CoV-2 infection: challenges and solutions for the long haul ahead

Understanding the experience of long COVID symptoms in hospitalized and non-hospitalized individuals: a random, cross-sectional survey study

Systematic review of the prevalence of long COVID

Using self-reports or claims to assess disease prevalence: it's complicated

Study protocol for the Innovative Support for Patients With SARS-CoV-2 Infections Registry (INSPIRE): a longitudinal study of the medium and long-term sequelae of SARS-CoV-2 infection

The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies

Racial, ethnic, and sex disparities in the incidence and cognitive symptomology of long COVID-19

Risk predictors and symptom features of long COVID within a broad primary care patient population including both tested and untested patients

Attributes and predictors of long COVID

Factors associated with post-acute sequelae of SARS-CoV-2 (PASC) after diagnosis of symptomatic COVID-19 in the inpatient and outpatient setting in a diverse cohort

Post-COVID syndrome in non-hospitalised patients with COVID-19: a longitudinal prospective cohort study

Follow-up of adults with noncritical COVID-19 two months after symptom onset

Post-COVID-19 syndrome in outpatients: a cohort study

Onset, duration and unresolved symptoms, including smell and taste changes, in mild COVID-19 infection: a cohort study in Israeli patients

Prevalence and associated factors of post-COVID-19 syndrome in a Brazilian cohort after 3 and 6 months of hospital discharge

Post COVID-19 condition of the Omicron variant of SARS-CoV-2

Post-COVID-19 symptoms 6 months after acute infection among hospitalized and non-hospitalized patients

Post-COVID-19 syndrome and quality of life impairment in severe COVID-19 Mexican patients

Predictors of post-COVID-19 and the impact of persistent symptoms in non-hospitalized patients 12 months after

COVID-19, with a focus on work ability Mechanisms, biomarkers and targets for adult-onset Still's disease Differentiation of prior SARS-CoV-2 infection and postacute sequelae by standard clinical laboratory measurements in the RECOVER cohort Severe fatigue and persistent symptoms at 3 months following severe acute respiratory syndrome coronavirus 2 infections during the pre-Delta, Delta, and Omicron time periods: a multicenter prospective cohort study Evidence mounts that about 7% of US adults have had long COVID Association of initial SARS-CoV-2 test positivity with patient-reported well-being 3 months after a symptomatic illness Long COVID clinical phenotypes up to 6 months after infection identified by latent class analysis of self-reported symptoms Agreement between reported questionnaire data and medical records on diagnosis and COVID-19 symptoms at onset Benefits and drawbacks of electronic health record systems Ethnic and racial differences in self-reported symptoms, health status, activity level, and missed work at 3 and 6 months following SARS-CoV-2 infection JAMA Network Open. 2025. [PMC12344537](#)

81. J Coste, C Delpierre, S Tebeka, C Lemogne, O Robineau, O Steichen, M Rushyizekera, M van den Akker, T T Makovski, et al.. Difference in long COVID prevalence due to different definitions and long COVID related risk factors The European Journal of Public Health. 2024. [PMC11516473](#)
82. Karen Jeffrey, Lana Woolford, Rishma Maini, Siddharth Basetti, Ashleigh Batchelor, David Weatherill, Chris White, Vicky Hammersley, Tristan Millington, Calum Macdonald, Jennifer K. Quint, Robin Kerr, Steven Kerr, Syed Ahmar Shah, Igor Rudan, Adeniyi Francis Fagbamigbe, Colin R. Simpson, Srinivasa Vittal Katikireddi, Chris Robertson, Lewis Ritchie, Aziz Sheikh, Luke Daines, et al.. Prevalence and risk factors

for long COVID among adults in Scotland using electronic health records: a national, retrospective, observational cohort study Long COVID: major findings, mechanisms and recommendations A clinical case definition of post-COVID-19 condition by a Delphi consensus Long-Term sequelae of COVID-19: a systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms Risks and burdens of incident diabetes in long COVID: a cohort study Association of COVID-19 with major arterial and venous thrombotic diseases: a population-wide cohort study of 48 million adults in England and Wales Preparing for the long-haul: autonomic complications of COVID-19 Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK: 30 March 2023 Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study Persistent COVID-19 symptoms in a community study of 606,434 people in England Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records in situ using OpenSAFELY Identifying who has long COVID in the USA: a machine learning approach using N3C data Predictive attributes for developing long COVID—a study using machine learning and real-world data from primary care physicians in Germany Finding the ‘right’ GP: a qualitative study of the experiences of people with long-COVID Incidence, co-occurrence, and evolution of long-COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19 Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Data-driven identification of post-acute SARS-CoV-2 infection subphenotypes PASClex: a comprehensive post-acute sequelae of COVID-19 (PASC) symptom lexicon derived from

electronic health record clinical notes Deriving and validating a risk prediction model for long COVID-19: protocol for an observational cohort study using linked Scottish data The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies Cohort profile: early pandemic evaluation and enhanced surveillance of COVID-19 (EAVE II) database Grouping of codes for conditions (RCGs) GP consultation rates for sequelae after acute covid-19 in patients managed in the community or hospital in the UK: population based study Long Covid symptoms and diagnosis in primary care: a cohort study using structured and unstructured data in the Health Improvement Network primary care database Controlling the false discovery rate: a practical and powerful approach to multiple testing Post-covid medical complaints following infection with SARS-CoV-2 Omicron vs Delta variants eClinicalMedicine. 2024. [PMC11016856](#)

83. Alaleh Azhir, Jonas Hgel, Jiazi Tian, Jingya Cheng, Ingrid V. Bassett, Douglas S. Bell, Elmer V. Bernstam, Maha R. Farhat, Darren W. Henderson, Emily S. Lau, Michele Morris, Yevgeniy R. Semenov, Virginia A. Triant, Shyam Visweswaran, Zachary H. Strasser, Jeffrey G. Klann, Shawn N. Murphy, Hossein Estiri, et al.. Precision Phenotyping for Curating Research Cohorts of Patients with Post-Acute Sequelae of COVID-19 (PASC) as a Diagnosis of Exclusion Precision phenotyping for curating research cohorts of patients with unexplained post-acute sequelae of COVID-19 medRxiv. 2024. [PMC11065031](#)
84. Skyler Resendez, Steven H Brown, Hugo Sebastian Ruiz Ayala, Prahalad Rangan, Jonathan Nebeker, Diane Montella, Peter L Elkin, et al.. Defining the Subtypes of Long COVID and Risk Factors for Prolonged Disease: Population-Based

Case-Crossover Study Confronting our next national health disaster—long-haul covid Persistent symptoms in patients after acute COVID-19 COVID-19 and the gastrointestinal tract: more than meets the eye Gastrointestinal symptoms associated with COVID-19: impact on the gut microbiome Characterizing long COVID in an international cohort: 7 months of symptoms and their impact SARS-CoV-2 is associated with changes in brain structure in UK Biobank Real-time tracking of self-reported symptoms to predict potential COVID-19 Attributes and predictors of long COVID Epidemiology, symptomatology, and risk factors for long COVID symptoms: population-based, multicenter study Long COVID: major findings, mechanisms and recommendations Characterization of autonomic symptom burden in long COVID: a global survey of 2,314 adults Risk of arrhythmias following COVID-19: nationwide self-controlled case series and matched cohort study Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study Sequelae in adults at 6 months after COVID-19 infection Long COVID in the United States The costs of long COVID Muscle abnormalities worsen after post-exertional malaise in long COVID Addressing the long-term effects of COVID-19 Overview of VA data, information systems, National Databases and Research uses COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines A randomized trial of long-term oxygen for COPD with moderate desaturation Socioeconomic status and its relationship to chronic respiratory disease Socioeconomic status and hypertension: a meta-analysis Impact of socioeconomic status on diabetes and cardiovascular risk factors: results of a large French survey Chest CT findings of COVID-19 pneumonia by duration of symptoms Pulmonary vascular endothelialitis, thrombosis,

and angiogenesis in Covid-19 Persistent 129Xe MRI pulmonary and CT vascular abnormalities in symptomatic individuals with post-acute COVID-19 syndrome Cardiac complications more common after COVID-19 than vaccination Gastrointestinal post-acute COVID-19 syndrome Ocular findings among patients surviving COVID-19 What is long COVID? National healthcare quality and disparities report chartbooks Optimizing data on race and ethnicity for veterans affairs patients COVID-19 epidemiological update—22 December 2023 JMIR Public Health and Surveillance. 2024. [PMC11094603](#)

85. Agneta Cederström, George Frederick Mkoma, Thomas Benfield, Charles Agyemang, Marie Nørredam, Mikael Rostila. Long COVID and its risk factors in migrants: a nationwide register study from Sweden COVID-19: WHO declares end of global health emergency Long COVID: 3 years in Clustering analysis identified three long COVID phenotypes and their association with general health status and working ability Five cluster classifications of long COVID and their background factors: a cross-sectional study in Japan Clinical outcomes and risk factors for COVID-19 among migrant populations in high-income countries: a systematic review Risk of SARS-CoV-2 infection in migrants and ethnic minorities compared with the general population in the European WHO region during the first year of the pandemic: a systematic review Inequitable impact of infection: social gradients in severe COVID-19 outcomes among all confirmed SARS-CoV-2 cases during the first pandemic wave in Sweden Disparities in coronavirus disease 2019 mortality by country of birth in Stockholm, Sweden: a total-population-based cohort study Risk of long COVID and associated symptoms after acute SARS-COV-2 infection in ethnic minorities: a nationwide register-linked cohort study in Denmark Racial,

ethnic, and sex disparities in the incidence and cognitive symptomology of long COVID-19 Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records COVID-19 illness severity and 2-year prevalence of physical symptoms: an observational study in Iceland Risk factors associated with post-COVID-19 condition: a systematic review and meta-analysis Symptoms and risk factors for long COVID in non-hospitalized adults Female gender is associated with long COVID syndrome: a prospective cohort study Sex differences in sequelae from COVID-19 infection and in long COVID syndrome: a review Healthy migrant effect in the Swedish context: a register-based, longitudinal cohort study Inequalities in COVID-19 severe morbidity and mortality by country of birth in Sweden Socioeconomic and demographic risk factors in COVID-19 hospitalization among immigrants and ethnic minorities Ethnic minority status as social determinant for COVID-19 infection, hospitalisation, severity, ICU admission and deaths in the early phase of the pandemic: a meta-analysis COVID-19 vaccine effectiveness against post-COVID-19 condition among 589 722 individuals in Sweden: population based cohort study Sociodemographic characteristics and COVID-19 testing rates: spatiotemporal patterns and impact of test accessibility in Sweden Health literacy and e-health literacy among Arabic-speaking migrants in Sweden: a cross-sectional study Mitigating the impacts of the COVID-19 pandemic on vulnerable populations: lessons for improving health and social equity BMC Medicine. 2025. [PMC11776292](#)

86. Kristen Nishimi, Thomas C Neylan, Daniel Bertenthal, Karen H Seal, Aoife O'Donovan. Association of Psychiatric Disorders with Clinical Diagnosis of Long COVID in US Veterans Long COVID after breakthrough SARS-CoV-2 infection Obesity Among Those with Mental Disorders: A National Institute of

Mental Health Meeting Report The immunology of long COVID Persistence of somatic symptoms after COVID-19 in the Netherlands: An observational cohort study Recovery and symptom trajectories up to two years after SARS-CoV-2 infection: Population based, longitudinal cohort study Access to Medical Care Among Persons With Psychotic and Major Affective Disorders Post-COVID Conditions Among Adult COVID-19 Survivors Aged 18-64 and ≥ 65 Years—United States, March 2020–November 2021 Effect of covid-19 vaccination on long covid: Systematic review A comprehensive characterization of patients diagnosed with post-COVID-19 condition in Sweden 16 months after the introduction of the International Classification of Diseases Tenth Revision diagnosis code (U09.9): A population-based cohort study Pathophysiology and mechanism of long COVID: A comprehensive review Global Prevalence of Post-Coronavirus Disease 2019 (COVID-19) Condition or Long COVID: A Meta-Analysis and Systematic Review Unexplained post-acute infection syndromes Association of Psychiatric Illness and Obesity, Physical Inactivity, and Smoking among a National Sample of Veterans Long COVID: Major findings, mechanisms and recommendations Age Differences in Posttraumatic Stress Disorder, Psychiatric Disorders, and Healthcare Service Use Among Veterans in Veterans Affairs Primary Care Clinics Evidence for a Dysregulated Immune System in the Etiology of Psychiatric Disorders Serious Mental Illness Diagnosis and COVID-19 Vaccine Uptake in the Veterans Health Administration Post COVID-19 condition diagnosis: A population-based cohort study of occurrence, associated factors, and healthcare use by severity of acute infection Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Rates and Factors Associated With Documentation of Diagnostic Codes

for Long COVID in the National Veterans Affairs Health Care System Prevalence of and factors associated with post-coronavirus disease 2019 (COVID-19) condition in the 12 months after the diagnosis of COVID-19 in adults followed in general practices in Germany Post-traumatic stress disorder (PTSD) as a systemic disorder: Pathways to cardiovascular disease Mental and neurological disorders and risk of COVID-19 susceptibility, illness severity and mortality: A systematic review, meta-analysis and call for action Use of the Postacute Sequelae of COVID-19 Diagnosis Code in Routine Clinical Practice in the US Association between Mental Disorders and Subsequent Medical Conditions Post-traumatic stress disorder and risk for hospitalization and death following COVID-19 infection Association of Psychiatric Disorders With Incidence of SARS-CoV-2 Breakthrough Infection Among Vaccinated Adults Elevated risk for autoimmune disorders in Iraq and Afghanistan veterans with posttraumatic stress disorder The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis A peripheral inflammatory signature discriminates bipolar from unipolar depression: A machine learning approach The Nature and Dimensions of Social Support Among Individuals with Severe Mental Illnesses Bringing the war back home: Mental health disorders among 103,788 US veterans returning from Iraq and Afghanistan seen at Department of Veterans Affairs facilities Psychiatric disorders and subsequent risk of cardiovascular disease: A longitudinal matched cohort study across three countries A clinical case definition of post-COVID-19 condition by a Delphi consensus Symptoms and risk factors for long COVID in non-hospitalized adults Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Long COVID burden and risk factors in

10 UK longitudinal studies and electronic health records Risk Factors Associated With Post-COVID-19 Condition: A Systematic Review and Meta-analysis Mental disorders and risk of COVID-19-related mortality, hospitalisation, and intensive care unit admission: A systematic review and meta-analysis Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: A systematic review and meta-analysis Psychiatric, Neurophysical and Neurocognitive Sequelae of Post-Acute COVID-19 Syndrome: A Systematic Review Immune response to vaccination in adults with mental disorders: A systematic review Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Inflammation-related biomarkers in major psychiatric disorders: A cross-disorder assessment of reproducibility and specificity in 43 meta-analyses A systematic review and meta-analysis of long term physical and mental sequelae of COVID-19 pandemic: Call for research priority and action Potential pitfalls in the use of real-world data for studying long COVID Peripheral cytokine levels across psychiatric disorders: A systematic review and network meta-analysis A modified poisson regression approach to prospective studies with binary data Psychological medicine. 2024. [PMC11345858](#)

87. Veer Vekaria, Rohith Kumar Thiruvalluru, Zoe Verzani, Sajjad Abedian, Mark Olfson, Braja Gopal Patra, Yunyu Xiao, Katherine S. Salamon, Karin Hoth, Frank Blancero, Maxwell M. Hornig-Rohan, Teresa Akintonwa, Mahfuza Sabiha, Mark G. Weiner, Thomas W. Carton, Rainu Kaushal, Jyotishman Pathak, et al.. Schizophrenia, Bipolar, or Major Depressive Disorder and Postacute Sequelae of COVID-19 Estimated number of COVID-19 infections, hospitalizations, and deaths prevented among vaccinated persons in the US, December

2020 to September 2021 Development of a definition of postacute sequelae of SARS-CoV-2 infection Disentangling the postacute sequelae of SARS-CoV-2: E unibus pluram (from one, many) Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review SARS-CoV-2 reservoir in post-acute sequelae of COVID-19 (PASC) COVID-19-related mortality risk in people with severe mental illness: a systematic and critical review Severe mental illness, race/ethnicity, multimorbidity and mortality following COVID-19 infection: nationally representative cohort study Heightened COVID-19 mortality in people with severe mental illness persists after vaccination: a cohort study of greater Manchester residents PCORnet 2020: current state, accomplishments, and future directions Launching PCORnet, a national patient-centered clinical research network The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies A new method of classifying prognostic comorbidity in longitudinal studies: development and validation Concordance of rural-urban self-identity and zip code-derived rural-urban commuting area (RUCA) designation Incubation period of COVID-19 caused by unique SARS-CoV-2 strains: a systematic review and meta-analysis SARS-CoV-2 viral variants-tackling a moving target Distinguishing admissions specifically for COVID-19 from incidental SARS-CoV-2 admissions: national retrospective electronic health record study Post-COVID-19 depressive symptoms: epidemiology, pathophysiology, and pharmacological treatment Generalisable long COVID subtypes: findings from the NIH N3C and RECOVER programmes COVID-19, oxidative stress, and neuroinflammation in the depression route Mental illness surveillance among adults in the United States Demographic

risk factors for COVID-19 infection, severity, ICU admission and death: a meta-analysis of 59 studies Prognostic factors for severity and mortality in patients infected with COVID-19: a systematic review One-year adverse outcomes among US adults with post-COVID-19 condition vs those without COVID-19 in a large commercial insurance database Assessing the risk and costs of COVID-19 in immunocompromised populations in a large United States commercial insurance health plan: the EPOCH-US Study Factors associated with post-acute sequelae of SARS-CoV-2 (PASC) after diagnosis of symptomatic COVID-19 in the inpatient and outpatient setting in a diverse cohort Risk factors associated with post-acute sequelae of SARS-CoV-2: an N3C and NIH RECOVER study Age, sex and previous comorbidities as risk factors not associated with SARS-CoV-2 infection for long COVID-19: a systematic review and meta-analysis How does multimorbidity affect middle-aged adults? A cross-sectional survey in the Singapore primary healthcare setting The relationships among age, chronic conditions, and healthcare costs Association between race and COVID-19 outcomes among 2.6 million children in England The association of race and COVID-19 mortality Disparities in COVID-19 outcomes by race, ethnicity, and socioeconomic status: a systematic-review and meta-analysis Extracting social support and social isolation information from clinical psychiatry notes: comparing a rule-based natural language processing system and a large language model Physical and mental health complications post-COVID-19: scoping review The psychological impact of preexisting mental and physical health conditions during the COVID-19 pandemic Perception, prevalence, and prediction of severe infection and post-acute sequelae of COVID-19 Long COVID or post-acute sequelae of COVID-19 (PASC): an overview of biological factors that may

contribute to persistent symptoms Disparities in COVID-19 infection, hospitalisation and death in people with schizophrenia, bipolar disorder, and major depressive disorder: a cohort study of the UK Biobank Evaluating telehealth adoption and related barriers among hospitals located in rural and urban areas Using effect size-or why the P value is not enough JAMA Network Open. 2025. [PMC12573031](#)

88. Xiaotong Zhang, Alfred Jerrod Anzalone, Daisy Dai, Gary Cochran, Ran Dai, Mark E Rupp, Adam BAdam MAlexisAlfred (Jerrod)AminAmitAmyAndreaAndrew EAndrewAndrew TAnitaAnjali ABenjaminBenjaminBrianBrijeshCalebCarolynCavinCharisseChristineChris ADiegoDonEilisElaineElizabethEmily CarlsonEmily REvanFarrukh MFedericoFredGeorgeGregHaroldHeidiHemalkumarHongfangHythemJ WJamiJaylynJeremy RichardJessicaJinJoelJoel HJoelJohannaJohnJomolJoni LJulie AJustinJustinKarenKatie RebeccaKellie MKenKenneth RKenrick DwainKimberlyKristinLavanceLee AllanLeonieLesleyLiliMariamMark MMarshallMaryMary MorrisonMatvey BMelissa AMeredithMeredithMichael GMicheleNabeelNasiaNicoleNohaOferPatricia APenny WungPeterPhilip R ORafaelRandeepRebeccaRenaRichard ARichard LRishiRobertRobert TSaijuSam GSamuelSandeepSatyanarayanaScottShawn TSokoStephanie SSteveTellen DTiffanyUmitUsmanValeryVigneshWarren AWenndyWillWillWilliamXiaohan Tanner WilcoxLeeGravesAnzaloneMannaSahaOlexZhouWilliamsSoutherlandGirvin CavinessMadlock- BrownSuverChuteDillonWuSchmittTakemotoHousmanGabrielEichmannMa HayangaPincavitchClarkHarperIslamGeGagnierSaltzSaltzLoombaBuseMat O'ConnorKurillaMorrisQureshiSafdarGarbariniSharafeldinSadanFrancisBu

CohenPatelMoffittZhuKamaleswaranHurleyMillerPyarajanMichaelBozzette
Adam B Wilcox, Adam M Lee, Alexis Graves, Alfred (Jerrod)
Anzalone, Amin Manna, Amit Saha, Amy Olex, Andrea Zhou,
Andrew E Williams, Andrew Southerland, Andrew T Girvin,
Anita Walden, Anjali A Sharathkumar, Benjamin Amor,
Benjamin Bates, Brian Hendricks, Brijesh Patel, Caleb
Alexander, Carolyn Bramante, Cavin Ward-Caviness, Charisse
Madlock-Brown, Christine Suver, Christopher Chute,
Christopher Dillon, Chunlei Wu, Clare Schmitt, Cliff
Takemoto, Dan Housman, Davera Gabriel, David A Eichmann,
Diego Mazzotti, Don Brown, Eilis Boudreau, Elaine Hill,
Elizabeth Zampino, Emily Carlson Marti, Emily R Pfaff, Evan
French, Farrukh M Koraishy, Federico Mariona, Fred Prior,
George Sokos, Greg Martin, Harold Lehmann, Heidi Spratt,
Hemalkumar Mehta, Hongfang Liu, Hythem Sidky, J W Awori
Hayanga, Jami Pincavitch, Jaylyn Clark, Jeremy Richard
Harper, Jessica Islam, Jin Ge, Joel Gagnier, Joel H Saltz, Joel
Saltz, Johanna Loomba, John Buse, Jomol Mathew, Joni L
Rutter, Julie A McMurry, Justin Guinney, Justin Starren, Karen
Crowley, Katie Rebecca Bradwell, Kellie M Walters, Ken
Wilkins, Kenneth R Gersing, Kenrick Dwain Cato, Kimberly
Murray, Kristin Kostka, Lavance Northington, Lee Allan Pyles,
Leonie Misquitta, Lesley Cottrell, Lili Portilla, Mariam Deacy,
Mark M Bissell, Marshall Clark, Mary Emmett, Mary
Morrison Saltz, Matvey B Palchuk, Melissa A Haendel,
Meredith Adams, Meredith Temple-O'Connor, Michael G
Kurilla, Michele Morris, Nabeel Qureshi, Nasia Safdar, Nicole
Garbarini, Noha Sharafeldin, Ofer Sadan, Patricia A Francis,
Penny Wung Burgoon, Peter Robinson, Philip R O Payne,
Rafael Fuentes, Randeep Jawa, Rebecca Erwin-Cohen, Rena
Patel, Richard A Moffitt, Richard L Zhu, Rishi Kamaleswaran,
Robert Hurley, Robert T Miller, Saiju Pyarajan, Sam G
Michael, Samuel Bozzette, Sandeep Mallipattu,

Satyanarayana Vedula, Scott Chapman, Shawn T O'Neil, Soko Setoguchi, Stephanie S Hong, Steve Johnson, Tellen D Bennett, Tiffany Callahan, Umit Topaloglu, Usman Sheikh, Valery Gordon, Vignesh Subbian, Warren A Kibbe, Wenndy Hernandez, Will Beasley, Will Cooper, William Hillegass, Xiaohan Tanner Zhang, et al.. Chronic Lung Disease as a Risk Factor for Long COVID in Patients Diagnosed With Coronavirus Disease 2019: A Retrospective Cohort Study Post-acute COVID-19 syndrome Post-acute COVID-19 syndrome (PCS) and health-related quality of life (HRQoL)—a systematic review and meta-analysis Prevalence and correlates of long COVID symptoms among US adults Cross-sectional study evaluating the impact of SARS-CoV-2 variants on long COVID outcomes in UK hospital survivors The role of multimorbidity and socio-economic characteristics as potential risk factors for long Covid: evidence from the multilevel analysis of the Survey of Health, Ageing and Retirement in Europe's corona surveys (2020-2021) Symptoms and risk factors for long COVID in non-hospitalized adults Long COVID in hospitalized and non-hospitalized patients in a large cohort in northwest Spain, a prospective cohort study Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Association between BNT162b2 vaccination and long COVID after infections not requiring hospitalization in health care workers Risk factors associated with post-acute sequelae of SARS-CoV-2 in an EHR cohort: a national COVID cohort collaborative (N3C) analysis as part of the NIH RECOVER program A longitudinal study of COVID-19 sequelae and immunity: baseline findings Development of a definition of postacute sequelae of SARS-CoV-2 infection Long covid—mechanisms, risk factors, and management Female sex is a risk factor associated with long-term post-COVID related-

symptoms but not with COVID-19 symptoms: the LONG-COVID-EXP-CM Multicenter Study Prevalence and risk factors of post-COVID-19 condition in adults and children at 6 and 12 months after hospital discharge: a prospective, cohort study in Moscow (StopCOVID) Predictive models of long COVID Long term predictors of breathlessness, exercise intolerance, chronic fatigue and well-being in hospitalized patients with COVID-19: a cohort study with 4 months median follow-up Risk factors associated with post-COVID-19 condition Air pollution and risk of chronic obstructed pulmonary disease: the modifying effect of genetic susceptibility and lifestyle Differences in rural and urban health information access and use Polyautoimmunity in rheumatological conditions Determinants of COVID-19 disease severity in patients with underlying rheumatic disease Obesity and inflammation: the linking mechanism and the complications Chronic inflammatory status observed in patients with type 2 diabetes induces modulation of cytochrome P450 expression and activity Associations of depression, anxiety, worry, perceived stress, and loneliness prior to infection with risk of post-COVID-19 conditions Long COVID: major findings, mechanisms and recommendations COVID-19 long-term sequelae: Omicron versus Alpha and Delta variants Risk factors for long coronavirus disease 2019 (long COVID) among healthcare personnel, Brazil, 2020-2022 COVID-19 in immunocompromised hosts: what we know so far Smokers are less likely than non-smokers to seek help for a lung cancer 'alarm' symptom A population-based study on social inequality and barriers to healthcare-seeking with lung cancer symptoms Help seeking behavior for problematic substance uses in north-west Ethiopia Effect of COVID-19 vaccines on reducing the risk of long COVID in the real world: a systematic review and meta-analysis Prevalence and

predictors of post-acute COVID-19 symptoms in Italian primary care patients Long COVID and significant activity limitation among adults, by age—United States, June 1–13, 2022, to June 7–19, 2023 Open Forum Infectious Diseases. 2024. [PMC11342242](#)

89. Yasin Abul, Daniel Harris, Preeti Chachlani, Kaley Hayes, Vincent Mor, Stefan Gravenstein. UTILIZATION OF THE LONG COVID DIAGNOSIS CODE IN US NURSING HOME RESIDENTS Innovation in Aging. 2024. [PMC11690510](#)

90. Ekaterina Pazukhina, Esteban Garcia-Gallo, Luis Felipe Reyes, Anders Benjamin Kildal, Waasila Jassat, Murray Dryden, Jan Cato Holter, Allegra Chatterjee, Kyle Gomez, Arne Søråas, Matteo Puntoni, Nicola Latronico, Fernando A Bozza, Michael Edelstein, Bronner P Gonçalves, Christiana Kartsonaki, Oksana Kruglova, Sérgio Gaião, Yock Ping Chow, Yash Doshi, Sara Isabel Duque Vallejo, Elsa D Ibáñez-Prada, Yuli V Fuentes, Claire Hastie, Margaret E O'Hara, Valeria Balan, Tigist Menkir, Laura Merson, Sadie Kelly, Barbara Wanjiru Citarella, Malcolm G Semple, Janet T Scott, Daniel Munblit, Louise Sigfrid, AdamJohn HAndreaValeriaIbrahimLeonardoAnnaFrankLucilleFernando ADaniloCaterinaGailAnnaDanielMugeAllegraYockBarbaraYashThomas MSarahMurrayAnneMichaelRachelRobYuli VAnneliesKyleBronner PLynseySérgioSilvioKatrinaMargaretaEwen MMadihaClaireLarsAquilesAntoniaJanElsa DJaneNinaWaasilaEdwinChristianaSadieAndersOksanaSulaimanNicolaJam JMilesSinnaduraiJoanneTigistLauraRobertaSarahBenCarolineDanielSriniva KEllenEkaterinaSimoneDaniel RMatteoJordiLilianaLuisSergioClark DSteffiJanet TMalcolm GLouiseGirishArneRenaudLanceNatalieCaroline AliAmuasiAnghebenBalanRichard

BanguraBastosBeltrameBloosBlumbergBozzaBuonsensoCaminitiCarsonCa
ChowCitarellaDoshiDrakeDuqueDrydenMargarita Dyrhol
RiiseEdelsteinEvansFowlerFuentesGillesenGomezGoncalvezGoodwinGaião
TrujilloHoCato
HolterIbañezIresonJamiesonJassatJesudasonKartsonakiKellyBenjamin
KildalKrugalovaLakohLatronicoLeeShen
LimLissaeurLoneLoweLunnManohanMcPeakeMenkirMersonMetaMooreM
ReyesRuiz
SaltanaRussellRyckaertScottSempleSigfridSindhvani
PulmSøraasTamisierTurtleWrightVika,
LaurentAmalYounesAliyaMartaBeatriceKévinAdamKazaliClotildeNathalieC
BenoîtEliseJeanJohannHuguesAdrienEyvind
WLaurèneCecileBenjaminDelphineClaudineJohnFirouzéWendy
SMarieJoaquínRomainJulesAlexandraHusnaMarineNicolasDelphineJoséGi
HélèneIsabelaPierre-
AdrienRaphaëlElisabethLilaOlivierSabellineKévinLaurenceCamileAnne-
SophieMaudeAuroreMarielleAxelleSonjaPolinaMarielleDaniloIngrid
GDenisRoarAndréEderCyrilJoseValentinePaulineNicolasThierryMarie-
ChristineGailMaire-LaureGuylaineFrançois-
XavierMinervaAnissaCatherineMeeraJean-
MarcCharlotteJulieLéoAntoineThibaultCatherineBernardMarie-
CharlotteYockBarbaraSaraGwenhaëlMarieGraham
SHuguesAndreaGrégorySandrineRoxaneStéphanieJuanJaimeElodieEricAna
PierreRomainEveIsabelleKarenChristellePierreElisaEmmanuelleDominique
LucVincentFaraAlphonsineFélixAnnemarie BChristl
ACélineNathalieThomasAmiel
AVincentFrançoisAlexandreSusannePaulJakeBertrandXavierAnneMichaelL
JKarineRaphaëlFrançois-XavierEglantine
FerrandNicolasCélineWilliamThomasTomAline-
MarieVictorErwanChristopheStéphanieValérieRostaneJean-
CharlesAmandineLindaSérgioCarrolEstebanNoeliaDenisValérieAnatoliyAle
AWilliamSegolèneFionaJérémieMartinRomainAnneLaurentThomasAliNadir
MJanetLarsRossMaximeDianaRupertHikomboAntoniaAlexandreIsabelleJan

SébastienSamreenPatrickMariachiaraMargauxMetteDanielleSalmaJulienC
 RARSèneOksanaGalynaMarieMarieNadhemMarieFabriceOlivierAntonioMar
 DanielAdrienVéroniqueAnthonyMarcQuentinFrançois-
 XavierOlivierMathieuSophieBrunoYvesClaireGeoffreyWeiBrunoAndreasGui
 ChristopheCarlosMilesLiemDominiqueMoïseGabrielGuillermoRafaelSophie
 EKennethACécileFranceAlexander
 JEmmanuelleAntoineMaykaLauraRobertaAgnèsAlison
 MVaninaMehdiIsabelleShona
 CSarahLinaLuciaClaraJulienJimmyDanielMarlèneSrinivasAlaminKarlFredri
 LWilnaPeterClaudiaPaulNadiaRachidaEricJustine
 PagesMassimoCarloNathalieAurélieJérémieBrunoChristelleWilliam
 AJean-
 FrançoisMiguelFlorentNathanVincentDanielThomasVincentVentzislavaFran
 GVíctorElseChristianMarieBlandineChristopheStanislasSarahDagAnne-
 SophieMatthieuLuisAntoniaLaurentPatrickKarineStephanieDavid
 LOlivierPaolaAmandaRobertoMélanieManuelAndreaPatrickCarineBenoîtCl
 DAleksanderNadiaCharlotteHélèneOlivierVanessaPierre-
 FrançoisBenjamineEgleArnaudMarionJanet
 TNicholasMalcolmGERicPabloCatherine
 AVictoriaNassimaJeanneLouiseDarioVegardSueLeneTomAgnèsAlbertEdoua
 CapucineOlivierNicolasVincentSimon-DjamelBenoîtEmma
 CMathewRyanSVadimPeterSJean-
 FrançoisNoémieKristianCécileTiffanyJeanneChristelleSarahLance
 CWAndersFrançoisTimothy
 MPieroSylvieNoémieCharlineAurélieFannySteveJiaMurrayPaulAurélieNata
 AbelAbrousAitTamlihatMohammed
 AlameenAlessiAlexAlexandreAliEnagnon
 AlidjnouAllavenaAllouAndrejakAnghebenAngoulvantAnsartArletArtaud-
 MacariBaptiste
 AssieAuchabieAumaitreAuvetAxelsenAzemarAzoulayBachBacheletBadrKen
 BaillieBani-
 SadrBarclayBartoliBaruchBasmaciBauerBedossaBegumBeluzeBenechBerg
 Bernal

SobrinoBertoliBessisBevilacquaBezulierBhavsarBisoffiBitkerBlotBodenesB
NeversBouadmaBouchaudBouchezBouillerBouilletBouisseBoureauBouscan
BesseyreBraconnierHjellegjerde
BrunvollBugaevaBuissonBuonsensoBustosButnaruBævre-
JensenCabieCaceresCadozAndres CalvacheCampanaCaraux-
PazCarlierCarmoiCarretCarsonCasanovaCastor-
AlexandreCatherineCervantes-
GonzalezChairChakveatzeChandChapplainCharpentierChasChenardCheret
ChowWanjiru
CitarellaClohiseyColinConnorCookeCordelCortegianiCorvaisierCouffin-
CadierguesCourtoisCousseLuis Cruz BermúdezCruz
RojoCurlierD'Ortenzioda Silva FilipeDa SilveiraDagensArne
DahlDaltonDe MontmollinDe Rosede
SilvaDeCastroDebardDebrayDecoursDefousDelacroixDelavigneDelmasDelo
Da
SilvaDidierDiehlDinotDiopDioufDjossouDochertyDonnellyDorivalDournonD
Dyrhol-RiiseEdelsteinMargrete EggesbøEl
SanharawiElharrarEllingjord-
DaleEloyEnderleEngelmannEnoufEpaulardEsperouEsposito-
FareseEtienneFabreFabreFairfieldFaureFavoryFerrandDevougeFerriereFic
BrunonGail SkeieGaiãoGambleGarcia-GalloGarcía
BarrioGarotGarraitGavrylovGaymardGeraudGerbaud
MorlaesGhosnGiganteGiordanoGirvanGissotGlybochkoGoehringerGomezG
HolterHorbyHouasHulotIjazImbertIppolitoIsnardStausland
IstreJaafarJaafouraJabotJacksonJaureguiberryJegoJenumJosephJourdainBal
JørgensenKafifKaguelidouKaliTrygve
KallebergKartsonakiKeatingKellyKennonKerroumiKhalilKhooKiland
GranerudBenjamin KildalKimmounFloerecke Kjetland
KjetlandKlenermanKloumann
BekkenKnightKpangonKruglovaKutsynaLachatreLacosteLafhejLagrangeLa
RichardLangloisLaouénanLaribiLariviereLasryLaunayLaureillardLavie-
BadieLawLeLe BihanLe BrisLe FalherLe FevreLe HingratLe
MaréchalLe MestreLe MoalLe MoingLe

NagardLeGacLeeLeemingLefebvreLefebvreLefèvreLelievreLemaignanLem
 MarchalLiegeonShen LimLinaLindLingasLion-
 DaolioLivrozetLoubetLouftiLouisLucetLumbreras
 BermejoLunnLuongLutonMachadoMachedaMaestro de la
 CalleMahieuMahyMailletMaitreMalvyMandaMandelbrotMankikianManuel
 BlondelMartinotMartynenkoMatteiMaulinMazzoniMcDonaldMcLeanMear-
 PassardMentréMentzerMercierMerckxMergeay-
 FabreMersonMetaMeybeckMeynertMeysonnierMezidiMicheletMooreMoore
 CelyMoroMouton
 PerrotMoyetMullaertMunblitMurrismurthyMustafaErik
 MüllerMüllerNdureNeantNekliudovNghiNguyenNoursadeghiNseirNyaman
 NygaardOlliaroOosthuyzenOpenshawMilena Orozco-
 ChamorroOtikuOuamaraOuissaOziolMaïder
 PagadoyPalmariniPalmieriPansuPapadopoulosPasquierPastenePaulPaxtonP
 JiménezPeelmanPeiffer-
 SmadjaPeignePerezPerpointPestrePetrov-SanchezOlav
 PettersenPeytavinPicardPiconePirothPiubelliPiusPlantierPoissyPokeerbuxP
 GonzálezQuist-
 PaulsenRabaudRafiqRammaertRappRebaudetRedlHenrik
 ReikvamResseguierRevestFelipe
 ReyesRicchiutoRichierRispalRissoRobertsRobertsonRobineauRodariRojekF
 AlbuquerqueRorizRosa-
 CalatravaRossaneseRossignolRoyRozeRussellRygh
 HoltenSaidaniSalmon GandonniereSalvatorSanchezSancho-
 ShimizuSandrineSartonSaviciuteScherpereelSchneiderScottSedillotSemple
 BalazoteShawShawSi
 MohammedSibiudeSigfridSinattiSkogenSmithBergendal
 SolbergSolomonSommetSottoSoumSpinuzzaSriskandanStablerSteinsvikSt
 Der WerfVanelVauchyVeislingerVuottoWebbWeiWhamHenri
 WickyWiedemannWrightYazdanpanahYelnikYonisZabbeZambonZucman,
 et al.. Long Covid: a global health issue – a prospective,
 cohort study set in four continents Variations in the financial
 impact of the COVID-19 pandemic across 5 continents: A

cross-sectional, individual level analysis Socioeconomic impacts of COVID-19 in low-income countries WHO dashboard Unexplained post-acute infection syndromes Immune determinants of chronic sequelae after respiratory viral infection Pneumonia as a cardiovascular disease Severe Pneumococcal Pneumonia Causes Acute Cardiac Toxicity and Subsequent Cardiac Remodeling Complication of Community-Acquired Pneumonia (Including Cardiac Complications) A clinical case definition of post COVID-19 condition by a Delphi consensus 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study True prevalence of long-COVID in a nationwide, population cohort study Determinants of recovery from post-COVID-19 dyspnoea: analysis of UK prospective cohorts of hospitalised COVID-19 patients and community-based controls Effect of covid-19 vaccination on long covid: systematic review Effect of COVID-19 Vaccines on Reducing the Risk of Long COVID in the Real World: A Systematic Review and Meta-Analysis Characterising long COVID: a living systematic review Long COVID: aiming for a consensus Long COVID in low-income and middle-income countries: the hidden public health crisis Clinical characteristics, risk factors and outcomes in patients with severe COVID-19 registered in the International Severe Acute Respiratory and Emerging Infection Consortium WHO clinical characterisation protocol: a prospective, multinational, multicentre, observational study Ten months of temporal variation in the clinical journey of hospitalised patients with COVID-19: An observational cohort Research priorities for Long Covid: refined through an international multi-stakeholder forum What is the recovery rate and risk of long-term consequences following a diagnosis of COVID-19? A harmonised, global longitudinal observational study protocol Research electronic data capture (REDCap)--a metadata-

driven methodology and workflow process for providing translational research informatics support ISARIC-COVID-19 dataset: A Prospective, Standardized, Global Dataset of Patients Hospitalized with COVID-19 Thrombotic and hemorrhagic complications of COVID-19 in adults hospitalized in high-income countries compared with those in adults hospitalized in low- and middle-income countries in an international registry Respiratory support in patients with severe COVID-19 in the International Severe Acute Respiratory and Emerging Infection (ISARIC) COVID-19 study: a prospective, multinational, observational study Major adverse cardiovascular events (MACE) in patients with severe COVID-19 registered in the ISARIC WHO clinical characterization protocol: A prospective, multinational, observational study Post covid-19 condition (long covid) World Bank country and lending groups 2024 COVID-19 long-term sequelae: Omicron versus Alpha and Delta variants A cohort study of post-COVID-19 condition across the Beta, Delta, and Omicron waves in South Africa: 6-month follow-up of hospitalized and nonhospitalized participants Long-term physical impairments in survivors of COVID-19-associated ARDS compared with classic ARDS: A two-center study Physical, cognitive and mental health outcomes in 1-year survivors of COVID-19-associated ARDS Core outcome measurement instruments for use in clinical and research settings for adults with post-COVID-19 condition: an international Delphi consensus study Long COVID: major findings, mechanisms and recommendations Author Correction: The immunology of long COVID A living mapping review for COVID-19 funded research projects: final (27 month) update BMJ Global Health. 2024. [PMC11552006](#)

91. Lars G. Fritsche, Weijia Jin, Andrew J. Admon, Bhramar Mukherjee. Characterizing and Predicting Post-Acute

Sequelae of SARS CoV-2 Infection (PASC) in a Large Academic Medical Center in the US
Characterizing and Predicting Post-Acute Sequelae of SARS CoV-2 infection (PASC) in a Large Academic Medical Center in the US
Bing COVID-19 Tracker
High-dimensional characterization of post-acute sequelae of COVID-19
Global Prevalence of Post COVID-19 Condition or Long COVID: A Meta-Analysis and Systematic Review
Long-COVID in children and adolescents: A systematic review and meta-analyses
Post-COVID Conditions: Information for Healthcare Providers
Public Health Recommendations
Long COVID or Post-COVID Conditions
Post-acute COVID-19 syndrome
Chronic COVID syndrome: Need for an appropriate medical terminology for long-COVID and COVID long-haulers
Long-Haul COVID Symptoms, complications and management of long COVID: A review
Assessment and characterisation of post-COVID-19 manifestations
6-month consequences of COVID-19 in patients discharged from hospital: A cohort study
Post Acute Coronavirus (COVID-19) Syndrome
Follow up of patients with severe coronavirus disease 2019 (COVID-19): Pulmonary and extrapulmonary disease sequelae
Neurological manifestations of long-COVID syndrome: A narrative review
Characterizing long COVID in an international cohort: 7 months of symptoms and their impact
Neurological and psychiatric risk trajectories after SARS-CoV-2 infection: An analysis of 2-year retrospective cohort studies including 1,284,437 patients
Mid and long-term neurological and neuropsychiatric manifestations of post-COVID-19 syndrome: A meta-analysis
Long-term cardiovascular outcomes in COVID-19 survivors among non-vaccinated population: A retrospective cohort study from the TriNetX US collaborative networks
Long-term neurologic outcomes of COVID-19
Trajectory of long covid symptoms after covid-19 vaccination: Community based

cohort study Long COVID after breakthrough SARS-CoV-2 infection Female gender is associated with long COVID syndrome: A prospective cohort study Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Factors Associated with Post-Acute Sequelae of SARS-CoV-2 (PASC) After Diagnosis of Symptomatic COVID-19 in the Inpatient and Outpatient Setting in a Diverse Cohort Attributes and predictors of long COVID Immunoglobulin signature predicts risk of post-acute COVID-19 syndrome Identifying who has long COVID in the USA: A machine learning approach using N3C data Multiple early factors anticipate post-acute COVID-19 sequelae Phenotype risk scores (PheRS) for pancreatic cancer using time-stamped electronic health record data: Discovery and validation in two large biobanks A Phenome-Wide Association Study (PheWAS) of COVID-19 Outcomes by Race Using the Electronic Health Records Data in Michigan Medicine Evolving phenotypes of non-hospitalized patients that indicate long COVID Evaluating and Caring for Patients with Post-COVID Conditions: Interim Guidance comorbidity: An R package for computing comorbidity scores Comorbidity measures for use with administrative data Mapping ICD-10 and ICD-10-CM Codes to Phecodes: Workflow Development and Initial Evaluation R PheWAS: Data analysis and plotting tools for phenome-wide association studies in the R environment MatchIt: Nonparametric Preprocessing for Parametric Causal Inference Regularization Paths for Generalized Linear Models via Coordinate Descent Ridge Regression: Biased Estimation for Nonorthogonal Problems Ridge regression and its application to medical data Rcompanion: Functions to Support Extension Education Program Evaluation {DescTools}: Tools for Descriptive Statistics {ROCNReg}: An {R} Package for Receiver Operating Characteristic Curve

Inference with and Without Covariates {brglm2}: Bias Reduction in Generalized Linear Models Risk factors for post-COVID-19 condition in previously hospitalised children using the ISARIC Global follow-up protocol: A prospective cohort study Coronavirus Disease (COVID-19) Caused by (SARS-CoV-2) Infections: A Real Challenge for Human Gut Microbiota Altered gut microbial metabolites could mediate the effects of risk factors in Covid-19 Gut-brain axis in the neurological comorbidity of COVID-19 Risk factors and abnormal cerebrospinal fluid associate with cognitive symptoms after mild COVID-19 A Systematic Review of Persistent Symptoms and Residual Abnormal Functioning following Acute COVID-19: Ongoing Symptomatic Phase vs. Post-COVID-19 Syndrome Long-COVID severe refractory cough: Discussion of a case with 6-week longitudinal cough characterization Symptoms Experienced at the Acute Phase of SARS-CoV-2 Infection as Risk Factor of Long-term Post-COVID Symptoms: The LONG-COVID-EXP-CM Multicenter Study Frequency, signs and symptoms, and criteria adopted for long COVID-19: A systematic review Long COVID in the Faroe Islands: A Longitudinal Study Among Nonhospitalized Patients Skeletal muscle alterations in patients with acute Covid-19 and post-acute sequelae of Covid-19 Long-term cardiovascular outcomes of COVID-19 A tutorial on sensitivity analyses in clinical trials: The what, why, when and how Sensitivity analysis: A review of recent advances Post-acute sequelae of COVID-19 in a non-hospitalized cohort: Results from the Arizona CoVHORT Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Persistent COVID-19 symptoms in a community study of 606,434 people in England Clinical characteristics with inflammation profiling of long COVID and association with 1-year recovery following hospitalisation in the UK: A

prospective observational study Management of post-acute covid-19 in primary care New ICD-10-CM code for Post-COVID Conditions, following the 2019 Novel Coronavirus (COVID-19) CDC Announces Approval of ICD-10 Code for Post-Acute Sequelae of COVID-19 Pediatric long-COVID: An overlooked phenomenon? A Multidisciplinary NHS COVID-19 Service to Manage Post-COVID-19 Syndrome in the Community Long COVID Household Pulse Survey PASClex: A comprehensive post-acute sequelae of COVID-19 (PASC) symptom lexicon derived from electronic health record clinical notes Selection Bias and Information Bias in Clinical Research Global Percentage of Asymptomatic SARS-CoV-2 Infections Among the Tested Population and Individuals With Confirmed COVID-19 Diagnosis: A Systematic Review and Meta-analysis Proportion of asymptomatic coronavirus disease 2019: A systematic review and meta-analysis Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status 2022 ACC Expert Consensus Decision Pathway on Cardiovascular Sequelae of COVID-19 in Adults: Myocarditis and Other Myocardial Involvement, Post-Acute Sequelae of SARS-CoV-2 Infection, and Return to Play: A Report of the American College of Cardiology Solution Set Oversight Committee A central role for amyloid fibrin microclots in long COVID/PASC: Origins and therapeutic implications Addressing the post-acute sequelae of SARS-CoV-2 infection: A multidisciplinary model of care Caring for People with Post-COVID Conditions Lack of Antinuclear Antibodies in Convalescent Coronavirus Disease 2019 Patients With Persistent Symptoms Short-term and Long-term Rates of Postacute Sequelae of SARS-CoV-2 Infection: A Systematic Review Use of COVID-19 Vaccines in the United States Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine Safety and Efficacy of the BNT162b2

mRNA Covid-19 Vaccine Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against Covid-19 Journal of Clinical Medicine. 2023. [PMC9967320](#)

92. Lauren L. O'Mahoney, Ash Routen, Clare Gillies, Winifred Ekezie, Anneka Welford, Alexa Zhang, Urvi Karamchandani, Nikita Simms-Williams, Shabana Cassambai, Ashkon Ardavani, Thomas J. Wilkinson, Grace Hawthorne, Ffion Curtis, Andrew P. Kingsnorth, Abdullah Almaqhawi, Thomas Ward, Daniel Ayoubkhani, Amitava Banerjee, Melanie Calvert, Roz Shafran, Terence Stephenson, Jonathan Sterne, Helen Ward, Rachael A. Evans, Francesco Zaccardi, Shaney Wright, Kamlesh Khunti, et al.. The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis Corrigendum to "The prevalence and long-term health effects of long Covid among hospitalised and non-hospitalised populations: a systematic review and meta-analysis" WHO Coronavirus (COVID-19) dashboard Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK: 6 May 2022 Nearly one in five American adults who have had COVID-19 still have "Long COVID" Characterising long COVID: a living systematic review Symptoms, complications and management of long COVID: a review Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review More than 50 Long-term effects of COVID-19: a systematic review and meta-analysis Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: a systematic review Multiorgan impairment in low-risk individuals with post-COVID-19 syndrome: a prospective, community-based study Long covid: damage to multiple organs presents in young, low risk patients Global prevalence of post COVID-19 condition or long COVID: a meta-analysis

and systematic review Study quality assessment tools Measuring inconsistency in meta-analyses Pulmonary function test and computed tomography features during follow-up after SARS, MERS and COVID-19: a systematic review and meta-analysis Development and validation of the symptom burden questionnaire for long covid (SBQ-LC): rasch analysis A core outcome set for post-COVID-19 condition in adults for use in clinical practice and research: an international Delphi consensus study Long COVID: an overview Characteristics and impact of long covid: findings from an online survey A dynamic review of the evidence around ongoing Covid 19 symptoms (often called Long Covid) Characterizing long COVID in an international cohort: 7 months of symptoms and their impact The role of poverty and racial discrimination in exacerbating the health consequences of COVID-19 The disproportionate impact of COVID-19 on racial and ethnic minorities in the United States Understanding and tracking the impact of long COVID in the United Kingdom Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records in situ using OpenSAFELY The occurrence of long COVID: a rapid review eClinicalMedicine. 2023. [PMC9714474](#)

93. Luisa Frallonardo, Francesco Vladimiro Segala, Kajal D. Chhaganlal, Mohmaoud Yelshazly, Roberta Novara, Sergio Cotugno, Giacomo Guido, Roberta Papagni, Agnese Colpani, Andrea De Vito, Mario Barbagallo, Giordano Madeddu, Sergio Babudieri, Peter Lochoro, Jerry Ictho, Giovanni Putoto, Nicola Veronese, Annalisa Saracino, Francesco Di Gennaro, et al.. Incidence and burden of long COVID in Africa: a systematic review and meta-analysis A clinical case definition of post-COVID-19 condition by a Delphi consensus Long COVID: Major findings, mechanisms and recommendations Prognostic factors for post-COVID-19 syndrome: A systematic review and

meta-analysis Long-term consequences of COVID-19 at 6 months and above: A systematic review and meta-analysis Characteristics and predictors of acute and chronic post-COVID syndrome: A systematic review and meta-analysis A cohort study of post-COVID-19 condition across the Beta, Delta, and Omicron waves in South Africa: 6-month follow-up of hospitalized and nonhospitalized participants Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID symptom study app: A prospective, community-based, nested, case-control study Post-COVID-19 condition 3 months after hospitalisation with SARS-CoV-2 in South Africa: A prospective cohort study Interventions for improving long COVID-19 symptomatology: A systematic review How to properly use the PRISMA statement Assessing the quality of studies in meta-analyses: Advantages and limitations of the Newcastle Ottawa Scale Assessing the quality of studies in meta-research: Review/guidelines on the most important quality assessment tools Incidence of long COVID-19 in people with previous SARS-Cov2 infection: a systematic review and meta-analysis of 120,970 patients Meta-analysis in clinical trials Bias in meta-analysis detected by a simple, graphical test Female gender is associated with long COVID syndrome: A prospective cohort study Long COVID risk: A signal to address sex hormones and women's health Female sex is a risk factor associated with long-term post-COVID related-symptoms but not with COVID-19 symptoms: The LONG-COVID-EXP-CM multicenter study Association of sex, age, and comorbidities with mortality in COVID-19 patients: A systematic review and meta-analysis Risk factors prediction, clinical outcomes, and mortality in COVID-19 patients A minimal common outcome measure set for COVID-19 clinical research Intensive care unit capacity in low-income countries: A systematic review Fatigue and

cognitive impairment in post-COVID-19 Syndrome: A systematic review and meta-analysis Neuropsychiatric post-acute sequelae of COVID-19: Prevalence, severity, and impact of vaccination Mental health in Africa Scientific Reports. 2023. [PMC10700349](#)

94. Ying Xu, Jia-Wen Xu, You Wu, Li-Juan Rong, Li Ye, Oscar H. Franco, Ching-Wen Chien, Xiao-Ru Feng, Jia-Yu Chen, Tao-Hsin Tung, et al.. Prevalence and prognosis of sarcopenia in acute COVID-19 and long COVID: a systematic review and meta-analysis Long COVID: a clinical update Long COVID: major findings, mechanisms and recommendations Metabolomic and immune alterations in long COVID patients with chronic fatigue syndrome Exploring the role of galectin-9 and artemin as biomarkers in long COVID with chronic fatigue syndrome: links to inflammation and cognitive function Understanding long COVID: a modern medical challenge Sarcopenia: revised European consensus on definition and diagnosis Identifying and managing malnutrition, frailty and sarcopenia in the community: a narrative review Comparative performance of current definitions of sarcopenia against the prospective incidence of falls among community-dwelling seniors age 65 and older Sarcopenia and impairment in cognitive and physical performance Frailty and sarcopenia: the potential role of an aged immune system Sarcopenia and mortality risk in frail older persons aged 80 years and older: results from ilSIRENTE study Associated factors of sarcopenia in community-dwelling older adults: a systematic review and meta-analysis Post-COVID-19 acute sarcopenia: physiopathology and management Cytokine storm COVID-19: a major cause of cachexia and sarcopenia? Covid-19-associated myopathy caused by type I interferonopathy The coupling between sarcopenia and COVID-19 is the real

problem Prevalence of sarcopenia in patients with COVID-19: a systematic review and meta-analysis Sarcopenia as a predictor of mortality among the critically ill in an intensive care unit: a systematic review and meta-analysis Coronavirus disease 2019 (Covid-19) outcomes in patients with sarcopenia: a meta-analysis and meta-regression A 24-step guide on how to design, conduct, and successfully publish a systematic review and meta-analysis in medical research The PRISMA 2020 statement: an updated guideline for reporting systematic reviews Frailty prevalence and impact on outcomes in patients with atrial fibrillation: a systematic review and meta-analysis of 1,187,000 patients Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis Meta-analysis in clinical trials Can meta-analyses be trusted? Measuring inconsistency in meta-analyses Bias in meta-analysis detected by a simple, graphical test Frequency and factors associated with sarcopenia prediction in adult and elderly patients hospitalized for COVID-19 Role of sarcopenia risk in predicting COVID-19 severity and length of hospital stay in older adults: a prospective cohort study Impact of COVID-19 in nutritional and functional status of survivors admitted in intensive care units during the first outbreak. Preliminary results of the NUTRICOID study Nutritional factors associated with mortality in hospitalized patients with COVID-19 Evaluation of muscle mass and stiffness with limb ultrasound in COVID-19 survivors Influence of reduced muscle mass and quality on ventilator weaning and complications during intensive care unit stay in COVID-19

patients Muscle strength and muscle mass as predictors of hospital length of stay in patients with moderate to severe COVID-19: a prospective observational study Reduced muscle mass as predictor of intensive care unit hospitalization in COVID-19 patients Skeletal muscle mass, sarcopenia and rehabilitation outcomes in post-acute COVID-19 patients Nutritional and functional impact of acute SARS-CoV-2 infection in hospitalized patients Body composition and risk factors associated with sarcopenia in post-COVID patients after moderate or severe COVID-19 infections The impact of body composition on mortality of COVID-19 hospitalized patients: a prospective study on abdominal fat, obesity paradox and sarcopenia Low thoracic skeletal muscle index is associated with negative outcomes in 244 patients with respiratory COVID-19 Sarcopenia is a frequent disease in SARS-COV-2 infection Grip strength as a predictor of disease severity in hospitalized COVID-19 patients Prognostic implication of baseline sarcopenia for length of hospital stay and survival in patients with coronavirus disease 2019 Thoracic sarcopenia as a predictive factor of SARS-COV2 evolution Long term follow-up of sarcopenia and malnutrition after hospitalization for COVID-19 in conventional or intensive care units COVID-19 in older patients: assessment of post-COVID-19 sarcopenia The role of SARC-F scale in predicting progression risk of COVID-19 in elderly patients: a prospective cohort study in Wuhan Sarcopenia as potential biological substrate of long COVID-19 syndrome: prevalence, clinical features, and risk factors The relationship between frailty, nutritional status, co-morbidity, CT-body composition and systemic inflammation in patients with COVID-19 Relation between body composition, systemic inflammatory response, and clinical outcomes in patients admitted to an urban teaching hospital with COVID-19 Impact of sarcopenia

in SARS-CoV-2 patients during two different epidemic waves
Low thoracic skeletal muscle area is not associated with
negative outcomes in patients with COVID-19 Prediction of
abdominal CT body composition parameters by thoracic
measurements as a new approach to detect sarcopenia in a
COVID-19 cohort Low muscle mass in COVID-19 critically-ill
patients: prognostic significance and surrogate markers for
assessment Impact of COVID-19 on nutritional status during
the first wave of the pandemic Identification of factors
impairing exercise capacity after severe COVID-19 pulmonary
infection: a 3-month follow-up of prospective COVulnerability
cohort Prevalence of risk of malnutrition and risk of
sarcopenia in a reference hospital for COVID-19: relationship
with mortality Neutrophil-to-lymphocyte ratio is not
associated with risk of sarcopenia in elderly COVID-19
patients Visceral to subcutaneous fat ratio predicts short-term
mortality in patients with Covid 19. A multicenter study The
prognostic value of pneumonia severity score and pectoralis
muscle area on chest CT in adult COVID-19 patients Poor
nutritional status, risk of sarcopenia and nutrition related
complaints are prevalent in COVID-19 patients during and
after hospital admission Clinical outcomes and prevalence of
sarcopenia in patients with moderate to severe COVID-19
Myosteatorsis predicting risk of transition to severe COVID-19
infection Long-term outcomes in critically ill patients who
survived COVID-19: the NUTRICOVID observational cohort
study Incidence and risk factors of sarcopenia in hospitalized
survivors of COVID-19; a retrospective cohort study Risk
factors associated with acute sarcopenia in patients
hospitalized with COVID-19 Persistent symptoms in patients
after acute COVID-19 Key pathophysiological role of skeletal
muscle disturbance in post COVID and Myalgic
Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS):

accumulated evidence Which vertebral level should be used to calculate sarcopenia in covid-19 patients? A systematic review and meta-analysis Skeletal muscle alterations in patients with acute Covid-19 and post-acute sequelae of Covid-19 Sarcopenia during COVID-19 lockdown restrictions: long-term health effects of short-term muscle loss Characterizing long COVID in an international cohort: 7 months of symptoms and their impact The long-term health outcomes, pathophysiological mechanisms and multidisciplinary management of long COVID Function over mass: a meta-analysis on the importance of skeletal muscle quality in COVID-19 patients Physiotherapy in the intensive care unit: an evidence-based, expert driven, practical statement and rehabilitation recommendations Feasibility of exercise testing in patients who are critically ill: a prospective, observational multicenter study Effect of neuromuscular electrical stimulation on the recovery of people with COVID-19 admitted to the intensive care unit: a narrative review Muscle abnormalities worsen after post-exertional malaise in long COVID Nutritional supplements to support resistance exercise in countering the sarcopenia of aging Synergistic effect of bodyweight resistance exercise and protein supplementation on skeletal muscle in sarcopenic or dynapenic older adults Health maintenance through home-based interventions for community-dwelling older people with sarcopenia during and after the COVID-19 pandemic: a systematic review and meta-analysis Sarcopenia and COVID-19 outcomes *Annals of Medicine*. 2025. [PMC12931308](#)

95. G Onder, M Floridia, M Giuliano, A Silenzi, L Castriotta, F Pricci, T Grisetti, T Grassi, D Tiple, M Villa, S Brusaferro, et al.. Long-COVID in Italy: a country perspective *The European Journal of Public Health*. 2023. [PMC10595608](#)

96. F McGuire, U Fallon, P Kavanagh, P Carney. Characterising Long COVID in an Irish cohort: a study of symptom type, severity and determinants *The European Journal of Public Health*. 2024. [PMC11517363](#)
97. César Fernández-de-las-Peñas, Kin Israel Notarte, Princess Juneire Peligro, Jacqueline Veronica Velasco, Miguel Joaquín Ocampo, Brandon Michael Henry, Lars Arendt-Nielsen, Juan Torres-Macho, Gustavo Plaza-Manzano, et al.. Long-COVID Symptoms in Individuals Infected with Different SARS-CoV-2 Variants of Concern: A Systematic Review of the Literature
Emergence of SARS-CoV-2 variants in the world: How could this happen? Emerging SARS-CoV-2 Variants of Concern (VOCs): An Impending Global Crisis Viral loads and profile of the patients infected with SARS-CoV-2 Delta, Alpha, or R.1 variants in Tokyo The Delta SARS-CoV-2 variant has a higher viral load than the Beta and the historical variants in nasopharyngeal samples from newly diagnosed COVID-19 patients Omicron SARS-CoV-2 variant: A new chapter in the COVID-19 pandemic Clinical outcomes in patients infected with different SARS-CoV-2 variants at one hospital during three phases of the COVID-19 epidemic in Marseille, France Clinical and virological characteristics of SARS-CoV-2 B.1.617.2 (Delta) variant: A prospective cohort study Long COVID: Current definition A clinical case definition of post-COVID-19 condition by a Delphi consensus Prevalence of Post-COVID-19 symptoms in hospitalized and non-hospitalized COVID-19 survivors: A systematic review and meta-analysis Long-term sequelae of COVID-19: A systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms Global prevalence of post COVID-19 condition or long COVID: A meta-analysis and systematic review Post-acute COVID-19 syndrome (PCS) and health-related quality of life (HRQoL)—A systematic review and meta-analysis

COVID-19: Two million people in the UK are estimated to be experiencing long COVID, says ONS Long COVID risk with omicron may be half that of delta COVID-19: Long COVID risk is lower with omicron than delta, researchers find Global surveillance, research, and collaboration needed to improve understanding and management of long COVID Long COVID: Mechanisms, risk factors, and management Prognostic factors for post-COVID-19 syndrome: A systematic review and meta-analysis Effect of COVID-19 vaccines on reducing the risk of long COVID in the real world: A systematic review and meta-analysis The PRISMA 2020 statement: An updated guideline for reporting systematic reviews The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Post COVID-19 condition of the Omicron variant of SARS-CoV-2 Association between BNT162b2 vaccination and long COVID after infections not requiring hospitalization in health care workers Associated-onset symptoms and post-COVID-19 symptoms in hospitalized COVID-19 survivors infected with Wuhan, Alpha or Delta SARS-CoV-2 variant Long COVID following Omicron wave in Eastern India—A retrospective cohort study The prevalence of SARS-CoV-2 infection and long COVID in US adults during the BA.5 surge, June–July 2022 Post-COVID-19 symptoms 2 years after SARS-CoV-2 infection among hospitalized vs non-hospitalized patients Long-term sequelae following previous coronavirus epidemics Burden of post-COVID-19 syndrome and implications for healthcare service planning: A population-based cohort study Exploring the recovery curve for long-term post-COVID dyspnea and fatigue A paradigm for post-COVID-19 fatigue syndrome analogous to ME/CFS Endothelial dysfunction and altered endothelial biomarkers in patients with post-

COVID-19 syndrome and chronic fatigue syndrome (ME/CFS) Milder disease with Omicron: Is it the virus or the pre-existing immunity? Adaptive immunity to SARS-CoV-2 and COVID-19 Analysis of online search trends suggests that SARS-CoV-2 Omicron (B.1.1.529) variant causes different symptoms Can symptoms of anosmia and dysgeusia be diagnostic for COVID-19? Centers for Disease Control and Prevention, National Center for Immunization and Respiratory Diseases (NCIRD) Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: A systematic review Post-acute COVID-19 syndrome after reinfection and vaccine breakthrough by the SARS-CoV-2 Gamma variant in Brazil Newcastle-Ottawa Scale: Comparing reviewers' to authors' assessments Development and validation of the long COVID symptom and impact tools, a set of patient-reported instruments constructed from patients' lived experience Viruses. 2022. [PMC9785120](#)

98. Fei Wang, Chengxi Zang, Haoyang Li, Dhru Khullar, Yongkang Zhang, Stephenson Strobel, Yong Chen, Marc Sala, Payal Patel, Alejandro Comellas, Andrew Wylam, Mark Weiner, Christopher Forrest, Thomas Carton, Rainu Kaushal, et al.. Real-World Effectiveness of Nirmatrelvir in Protecting Long COVID for Outpatient Adult Patients - A Large-Scale Observational Cohort Study from the RECOVER Initiative Long covid—mechanisms, risk factors, and management Long COVID: major findings, mechanisms and recommendations Grants Full Approval to Paxlovid, COVID-19 Antiviral Treatment Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19 Association of nirmatrelvir for acute SARS-CoV-2 infection with subsequent Long COVID symptoms in an observational cohort study Nirmatrelvir and Molnupiravir and Post-COVID-19 Condition in Older Patients

Studies Investigate Whether Antivirals Like Paxlovid May Prevent Long COVID Association of Treatment With Nirmatrelvir and the Risk of Post-COVID-19 Condition Nirmatrelvir for Vaccinated or Unvaccinated Adult Outpatients with Covid-19 Data-driven identification of post-acute SARS-CoV-2 infection subphenotypes Target Trial Emulation: A Framework for Causal Inference From Observational Data Data-driven analysis to understand long COVID using electronic health records from the RECOVER initiative Use of Stabilized Inverse Propensity Scores as Weights to Directly Estimate Relative Risk and Its Confidence Intervals Moving towards best practice when using inverse probability of treatment weighting (IPTW) using the propensity score to estimate causal treatment effects in observational studies High-throughput target trial emulation for Alzheimer's disease drug repurposing with real-world data Estimated Global Proportions of Individuals With Persistent Fatigue, Cognitive, and Respiratory Symptom Clusters Following Symptomatic COVID-19 in 2020 and 2021 Negat Control Outcomes Research Square. 2024. [PMC11213188](#)

99. Yan Xie, Taeyoung Choi, Ziyad Al-Aly. Molnupiravir and risk of post-acute sequelae of covid-19: cohort study High-dimensional characterization of post-acute sequelae of COVID-19 Long COVID after breakthrough SARS-CoV-2 infection Long COVID: major findings, mechanisms and recommendations Long COVID: long-term health outcomes and implications for policy and research The immunology and immunopathology of COVID-19 Association of Treatment With Nirmatrelvir and the Risk of Post-COVID-19 Condition Mechanism of molnupiravir-induced SARS-CoV-2 mutagenesis Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized Patients Molnupiravir and risk of hospital admission or death in adults with covid-19: emulation of a

randomized target trial using electronic health records Long-COVID treatments: why the world is still waiting Making Neighborhood-Disadvantage Metrics Accessible - The Neighborhood Atlas Diabetes after SARS-CoV-2 infection Risk of persistent and new clinical sequelae among adults aged 65 years and older during the post-acute phase of SARS-CoV-2 infection: retrospective cohort study Post-COVID Conditions Among Adult COVID-19 Survivors Aged 18-64 and ≥ 65 Years — United States, March 2020–November 2021 Risk of clinical sequelae after the acute phase of SARS-CoV-2 infection: retrospective cohort study Post-covid syndrome in individuals admitted to hospital with covid-19: retrospective cohort study Persistent Symptoms in Patients After Acute COVID-19 Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19 Long-term cardiovascular outcomes of COVID-19 Kidney Outcomes in Long COVID Risks of mental health outcomes in people with covid-19: cohort study Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19 Characterizing long COVID in an international cohort: 7 months of symptoms and their impact 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records Risks and burdens of incident diabetes in long COVID: a cohort study Long-term neurologic outcomes of COVID-19 Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Diabetes after SARS-CoV-2 infection Long-term gastrointestinal outcomes of COVID-19 Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic

COVID-19 in 2020 and 2021 Estimated Global Proportions of Individuals With Persistent Fatigue, Cognitive, and Respiratory Symptom Clusters Following Symptomatic COVID-19 in 2020 and 2021 Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019 Comparative evaluation of clinical manifestations and risk of death in patients admitted to hospital with covid-19 and seasonal influenza: cohort study Acute Kidney Injury in a National Cohort of Hospitalized US Veterans with COVID-19 Long-term cardiovascular outcomes of COVID-19 Multiple imputation of discrete and continuous data by fully conditional specification Molnupiravir plus usual care versus usual care alone as early treatment for adults with COVID-19 at increased risk of adverse outcomes (PANORAMIC): an open-label, platform-adaptive randomised controlled trial Acute and postacute sequelae associated with SARS-CoV-2 reinfection The BMJ. 2023. [PMC10126525](https://pubmed.ncbi.nlm.nih.gov/40126525/)

100. Ka Chun Chong, Yuchen Wei, Katherine Min Jia, Christopher Boyer, Guozhang Lin, Huwen Wang, Conglu Li, Chi Tim Hung, Xiaoting Jiang, Carrie Ho Kwan Yam, Tsz Yu Chow, Yawen Wang, Shi Zhao, Kehang Li, Aimin Yang, Chris Ka Pun Mok, David SC Hui, Eng Kiong Yeoh, Zihao Guo, et al.. SARS-CoV-2 rebound and post-acute mortality and hospitalization among patients admitted with COVID-19: cohort study The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis SARS-CoV-2 rebound with and without use of COVID-19 oral antivirals ACTIV-2/A5401 study team. Symptom and viral rebound in untreated SARS-CoV-2 infection EPIC-HR trial investigators. Nirmatrelvir-ritonavir and viral load rebound in Covid-19 Incidence of viral rebound after treatment with nirmatrelvir-ritonavir and molnupiravir

Viral burden rebound in hospitalised patients with COVID-19 receiving oral antivirals in Hong Kong: A population-wide retrospective cohort study Evaluation of SARS-CoV-2 RNA rebound after nirmatrelvir/ritonavir treatment in randomized, double-blind, placebo-controlled trials - United States and International Sites, 2021-2022 SARS-CoV-2 virologic rebound with nirmatrelvir-ritonavir therapy: An observational study Symptoms, viral loads, and rebound among COVID-19 outpatients treated with nirmatrelvir/ritonavir compared with propensity score-matched untreated individuals Wellcome Sanger Institute COVID-19 Surveillance Team; COVID-19 Infection Survey Group; COVID-19 Genomics UK (COG-UK) Consortium; Barrett J, Fraser C, Bonsall D, Walker AS, Lythgoe K. Prevalence of persistent SARS-CoV-2 in a large community surveillance study Changes in all-cause and cause-specific excess mortality before and after the Omicron outbreak of COVID-19 in Hong Kong Long COVID: Major findings, mechanisms and recommendations Association of SARS-CoV-2 infection and persistence with long COVID ATACCC study investigators. Onset and window of SARS-CoV-2 infectiousness and temporal correlation with symptom onset: A prospective, longitudinal, community cohort study A compromised specific humoral immune response against the SARS-CoV-2 receptor-binding domain is related to viral persistence and periodic shedding in the gastrointestinal tract SARS-CoV-2 infection and persistence in the human body and brain at autopsy COVID-19 and cardiac arrhythmias: A contemporary review Drug interactions with nirmatrelvir-ritonavir in older adults using multiple medications Viral kinetics of SARS-CoV-2 following onset of COVID-19 in symptomatic patients infected with the ancestral strain and omicron BA.2 in Hong Kong: a retrospective observational study Comorbidities prolonged viral shedding of

patients infected with SARS-CoV-2 omicron variant in Shanghai: A multi-center, retrospective, observational study Household transmission study team. Persistent SARS-CoV-2 RNA shedding without evidence of infectiousness: A cohort study of individuals with COVID-19 Dynamics of SARS-CoV-2 shedding in the respiratory tract depends on the severity of disease in COVID-19 patients COVID-19 rebound after VV116 vs nirmatrelvir-ritonavir treatment: A randomized clinical trial: A randomized clinical trial Quantifying the impact of immune history and variant on SARS-CoV-2 viral kinetics and infection rebound: A retrospective cohort study Real-world effectiveness of early molnupiravir or nirmatrelvir-ritonavir in hospitalised patients with COVID-19 without supplemental oxygen requirement on admission during Hong Kong's omicron BA.2 wave: a retrospective cohort study Long-term post-acute sequelae of COVID-19 infection: a retrospective, multi-database cohort study in Hong Kong and the UK Temporal dynamics in viral shedding and transmissibility of COVID-19 SARS-CoV-2 viral load and shedding kinetics Modelling upper respiratory viral load dynamics of SARS-CoV-2 COVID-19 vaccination modified the effect of nirmatrelvir-ritonavir on post-acute mortality and rehospitalization: a retrospective cohort study Effectiveness of nirmatrelvir/ritonavir and molnupiravir on post-COVID-19 outcomes among outpatients: a target trial emulation investigation Propensity score methods for confounding control in nonexperimental research Using the standardized difference to compare the prevalence of a binary variable between two groups in observational research Test for additive interaction in proportional hazards models Nature Communications. 2025. [PMC12304394](#)

101. Naomi J. Patel, Xiaosong Wang, Miao Lin, Emily N. Kowalski, Claire E. Cook, Kathleen M.M. Vanni, Krishan Guzzo, Grace

Qian, Katarina J. Bade, Alene Saavedra, Rathnam Venkat, Shruthi Srivatsan, Zachary K. Williams, Jennifer S. Hanberg, Yumeko Kawano, Abigail E. Schiff, Jeffrey A. Sparks, Zachary S. Wallace, et al.. Factors Associated with an Electronic Health Record-Based Definition of Post-Acute Sequelae of COVID-19 in Patients with Systemic Autoimmune Rheumatic Disease Clinical characteristics and outcomes of patients with coronavirus disease 2019 (COVID-19) and rheumatic disease: a comparative cohort study from a US 'hot spot' COVID-19 Outcomes in Patients With Systemic Autoimmune Rheumatic Diseases Compared to the General Population: A US Multicenter, Comparative Cohort Study Coronavirus disease 2019 outcomes among patients with rheumatic diseases 6 months into the pandemic Coronavirus Disease 2019 Outcomes Among Recipients of Anti-CD20 Monoclonal Antibodies for Immune-Mediated Diseases: A Comparative Cohort Study Characteristics associated with hospitalisation for COVID-19 in people with rheumatic disease: data from the COVID-19 Global Rheumatology Alliance physician-reported registry Vaccine effectiveness against SARS-CoV-2 infection and severe outcomes among individuals with immune-mediated inflammatory diseases tested between March 1 and Nov 22, 2021, in Ontario, Canada: a population-based analysis Immunogenicity, breakthrough infection, and underlying disease flare after SARS-CoV-2 vaccination among individuals with systemic autoimmune rheumatic diseases SARS-CoV-2 Breakthrough Infections Among Vaccinated Individuals with Rheumatic Disease: Results From the COVID-19 Global Rheumatology Alliance Provider Registry Evaluation of COVID-19 vaccine breakthrough infections among immunocompromised patients fully vaccinated with BNT162b2 Effect of Immunosuppression on the Immunogenicity of mRNA Vaccines to SARS-CoV-2 : A

Prospective Cohort Study Impact of disease-modifying antirheumatic drugs on vaccine immunogenicity in patients with inflammatory rheumatic and musculoskeletal diseases Methotrexate hampers immunogenicity to BNT162b2 mRNA COVID-19 vaccine in immune-mediated inflammatory disease Humoral and Cellular Immune Responses to SARS-CoV-2 Infection and Vaccination in Autoimmune Disease Patients With B Cell Depletion Immunogenicity and safety of the BNT162b2 mRNA COVID-19 vaccine in adult patients with autoimmune inflammatory rheumatic diseases and in the general population: a multicentre study Autoantibodies related to systemic autoimmune rheumatic diseases in severely ill patients with COVID-19 Defining Post-COVID Symptoms (Post-Acute COVID, Long COVID, Persistent Post-COVID): An Integrative Classification COVID-19 and antiphospholipid antibodies Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Impact of vaccination on postacute sequelae of SARS CoV-2 infection in patients with rheumatic diseases Long COVID after breakthrough SARS-CoV-2 infection Factors associated with COVID-19 breakthrough infection among vaccinated patients with rheumatic diseases: A cohort study DMARD disruption, rheumatic disease flare, and prolonged COVID-19 symptom duration after acute COVID-19 among patients with rheumatic disease: A prospective study Prolonged COVID-19 symptom duration in people with systemic autoimmune rheumatic diseases: results from the COVID-19 Global Rheumatology Alliance Vaccine Survey Post-COVID-19 condition in patients with autoimmune rheumatic diseases: the COVID-19 Vaccination in Autoimmune Diseases (COVAD) study Symptoms and risk factors for long COVID in non-hospitalized adults Racial/Ethnic Disparities in Post-acute Sequelae of SARS-CoV-2 Infection in New York: an EHR-

Based Cohort Study from the RECOVER Program Post-acute COVID-19 syndrome Long covid in persons with self-reported arthritis - symptoms, associated factors and functional limitations High risk of autoimmune diseases after COVID-19 Long COVID and Significant Activity Limitation Among Adults, by Age - United States, June 1-13, 2022, to June 7-19, 2023 The Journal of rheumatology. 2024. [PMC11065568](#)

102. Ludovica Segneri, Nandor Babina, Teresa Hammerschmidt, Andrea Fronzetti Colladon, Peter A. Gloor, Steve Zimmerman, Michal Ptaszynski, Michal Ptaszynski, Michal Ptaszynski, et al.. Too much focus on your health might be bad for your health: Reddit user's communication style predicts their Long COVID likelihood Too much focus on your health might be bad for your health: Reddit user's communication style predicts their Long COVID likelihood. Too much focus on your health might be bad for your health: Reddit user's communication style predicts their Long COVID likelihood. Early clues regarding the pathogenesis of long-COVID Long COVID: major findings, mechanisms and recommendations Defining long COVID: Going back to the start Long COVID: mechanisms, risk factors and recovery The Conundrum of 'Long-COVID-19': A Narrative Review Representation of long COVID syndrome in the awareness of the population is revealed by Google Trends analysis Long-Term COVID: Case Report and Methodological Proposals for Return to Work Long COVID Through a Public Health Lens: An Umbrella Review Number of initial symptoms is more related to long COVID-19 than acute severity of infection: a prospective cohort of hospitalized patients Characteristics and predictors of Long COVID among diagnosed cases of COVID-19 Using crowdsourced medicine to manage uncertainty on Reddit: The case of COVID-19 long-haulers Mining long-COVID symptoms from Reddit: characterizing post-COVID syndrome from

patient reports Loneliness during the COVID-19 Pandemic: A Comparison of Urban and Rural Areas Living with "long COVID": A systematic review and meta-synthesis of qualitative evidence Anti-social behaviour and economic decision-making: Panel experimental evidence in the wake of COVID-19 Reduced social distancing early in the COVID-19 pandemic is associated with antisocial behaviors in an online United States sample Self-reported COVID-19 symptoms on Twitter: an analysis and a research resource Within and between-day variation and associations of symptoms in Long Covid: Intensive longitudinal study Mental Health Discourse on reddit: Self-Disclosure, Social Support, and Anonymity Medical sentiment analysis using social media: towards building a patient assisted system Methods in predictive techniques for mental health status on social media: a critical review Positive Emotional Style Predicts Resistance to Illness After Experimental Exposure to Rhinovirus or Influenza A Virus Social media use and mental health during the COVID-19 pandemic: Moderator role of disaster stressor and mediator role of negative affect Understanding and measuring psychological stress using social media Natural emotion vocabularies as windows on distress and well-being Linguistic styles: Language use as an individual difference Perceptual and memory biases for health-related information in hypochondriacal individuals Memory bias in health anxiety is related to the emotional valence of health-related words Memory bias in patients with hypochondriasis and somatoform pain disorder Adjustment Disorder in the Face of COVID-19 Outbreak: The Impact of Death Anxiety, Media Exposure, Fear of Contagion and Hypochondriasis Symptoms Speaking under pressure: Low linguistic complexity is linked to high physiological and emotional stress reactivity Increase in Linguistic Complexity in Older Adults During COVID-19 Is

a cover letter still needed? Using LIWC to predict application success Mental health problems and social media exposure during COVID-19 outbreak Characterizing Audience Engagement and Assessing Its Impact on Social Media Disclosures of Mental Illnesses What drives unverified information sharing and cyberchondria during the COVID-19 pandemic? Trust in social media and COVID-19 beliefs and behaviours Post-COVID-19 syndrome among symptomatic COVID-19 patients: A prospective cohort study in a tertiary care center of Bangladesh Multimodal mental health analysis in social media Social Networking Sites, Depression, and Anxiety: A Systematic Review The development of adolescents' loneliness during the COVID-19 pandemic: The role of peer status and contact with friends Crisis response during the COVID-19 pandemic: Changes in social contact and social participation of older Americans On analyzing antisocial behaviors amid covid-19 pandemic Centrality in social networks conceptual clarification A faster algorithm for betweenness centrality The length of words reflects their conceptual complexity Justices and Legal Clarity: Analyzing the Complexity of U.S. Supreme Court Opinions Speech and language processing: An introduction to natural language processing, computational linguistics, and speech recognition The role of topic, interviewee and question in predicting rich interview data in the field of health research The Psychological Meaning of Words: LIWC and Computerized Text Analysis Methods The mt-86 β aphasia battery: A subset of normative data in relation to age and level of school education Effects of age and education on the lexico-semantic content of connected speech in adults Adult spoken discourse: the influences of age and education Linguistic Skills of Adult Native Speakers, as a Function of Age and Level of Education A comparative study of stemming algorithms Psychological

Aspects of Natural Language Use: Our Words, Our Selves A Comparison Analysis of Modal Auxiliary Verbs in Technical and General English Language Use and Personality during Crises: Analyses of Mayor Rudolph Giuliani's Press Conferences The language of depression Language use of depressed and depression-vulnerable college students Word use in the poetry of suicidal and nonsuicidal poets Linguistic bases of social perception Gender Differences in Language Use: An Analysis of 14,000 Text Samples The relationship of sex and intelligence to choice of words: A normative study of verbal behavior The sounds of social life: A psychometric analysis of students' daily social environments and natural conversations Male/female language differences and effects in same-sex and mixed-sex dyads: The gender-linked language effect Sentiment Analysis in Social Media and Its Application: Systematic Literature Review A multinational, multicentre study on the psychological outcomes and associated physical symptoms amongst healthcare workers during COVID-19 outbreak The relationship between repressive and defensive coping styles and monocyte, eosinophile, and serum glucose levels: support for the opioid peptide hypothesis of repression Lost in transmission? Self- and other-annotation of emotional words Linguistic markers of processing the first months of the pandemic COVID-19: a psycholinguistic analysis of Italian university students' diaries Examining the Impact of COVID-19 Lockdown in Wuhan and Lombardy: A Psycholinguistic Analysis on Weibo and Twitter The kernel of truth in text-based personality assessment: A meta-analysis of the relations between the Big Five and the Linguistic Inquiry and Word Count (LIWC) Personality in its natural habitat: Manifestations and implicit folk theories of personality in daily life Linguistic predictors of adaptive bereavement Revolutionary leaders: Long-term success as a function of

changes in conceptual complexity Exercise and well-being during COVID 19 -time to boost your immunity Pre-existing conditions associated with post-acute sequelae of COVID-19 Characteristics and impact of Long Covid: Findings from an online survey Author response to previous submission Decision Letter 0 Author response to Decision Letter 0 Decision Letter 1 Author response to Decision Letter 1 Decision Letter 2 Acceptance letter PLOS ONE. 2024. [PMC11302924](#)

103. Arinjita Bhattacharyya, Anand Seth, Shesh Rai. The effects of long COVID-19, its severity, and the need for immediate attention: Analysis of clinical trials and Twitter data The impact of nutrition on COVID-19 susceptibility and long-term consequences Global prevalence of post-acute sequelae of COVID-19 (PASC) or long COVID: A Meta-analysis and systematic review Long-term health consequences of COVID-19 Text mining infrastructure in R ISIS's COVID-19 messaging on twitter: An analysis of tweet sentiment and emotions More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Case report and systematic review suggest that children may experience similar long-term effects to adults after clinical COVID-19 Being a public health statistician during a global Pandemic COVID-19-related medical research: a meta-research and critical appraisal Long-term consequences of COVID-19: research needs The Challenge of Studying Long COVID: An Updated Review Frontiers in Big Data. 2022. [PMC9797990](#)
104. Lauren G. Staples, Olav Nielssen, Blake F. Dear, Madelyne A. Bisby, Alana Fisher, Rony Kayrouz, Nickolai Titov, et al.. Prevalence and Predictors of Long COVID in Patients Accessing a National Digital Mental Health Service Long COVID: Major findings, mechanisms and recommendations

Symptoms, complications and management of long COVID: A review
Attributes and predictors of long COVID
Differences in incidence, nature of symptoms, and duration of long COVID among hospitalised migrant and non-migrant patients in the Netherlands: A retrospective cohort study
Symptoms and risk factors for long COVID in non-hospitalized adults
Associations of Depression, Anxiety, Worry, Perceived Stress, and Loneliness Prior to Infection with Risk of Post-COVID-19 Conditions
Department of Health and Aged Care. Clinical Features of COVID-19 Disease. 2023
The complex association of barriers and interest in internet-delivered cognitive behavior therapy for depression and anxiety: Informing e-health policies through exploratory path analysis
Telehealth use in Australian primary healthcare during COVID-19: A cross-sectional descriptive survey
User characteristics and outcomes from a national digital mental health service: An observational study of registrants of the Australian MindSpot Clinic
A Comparison of Self-Referral and Referral via Primary Care Providers, through Two Similar Digital Mental Health Services in Western Australia
Rapid report 2: Symptoms of anxiety and depression during the first 12 weeks of the Coronavirus (COVID-19) pandemic in Australia
Rapid Report 3: Mental health symptoms, characteristics, and regional variation, for users of an Australian digital mental health service during the first 8 months of COVID-19
Rapid report: Early demand, profiles and concerns of mental health users during the coronavirus (COVID-19) pandemic
The PHQ-9: Validity of a brief depression severity measure
A brief measure for assessing generalized anxiety disorder: The GAD-7
Short screening scales to monitor population prevalences and trends in non-specific psychological distress
Long COVID and mental health correlates: A new chronic condition fits existing patterns
The high mental health burden

of “Long COVID” and its association with on-going physical and respiratory symptoms in all adults discharged from hospital The Effects of COVID-19 Lockdown on Social Connectedness and Psychological Distress in U.S. Adults with Chronic Diseases Comorbid Chronic Diseases are Strongly Correlated with Disease Severity among COVID-19 Patients: A Systematic Review and Meta-Analysis Cross-sectional study evaluating the impact of SARS-CoV-2 variants on Long COVID outcomes in UK hospital survivors What do we mean by long-COVID? A scoping review of the cognitive sequelae of SARS-CoV-2 infection Identity and COVID-19 in Canada: Gender, ethnicity, and minority status Factors Associated With Long COVID Symptoms in an Online Cohort Study Racial Disparities-Associated COVID-19 Mortality among Minority Populations in the US Depression and Generalized Anxiety as Long-Term Mental Health Consequences of COVID-19 in Iraqi Kurdistan International Journal of Environmental Research and Public Health. 2023. [PMC10531183](#)

105. David G. Blanchflower, Alex Bryson, Kanchan Thapa, Kanchan Thapa, Kanchan Thapa, Kanchan Thapa, Kanchan Thapa, et al.. Long COVID in the United States Post COVID-19 Condition (Long COVID) Challenges in defining long COVID: striking differences across literature, electronic health records, and patient-reported information COVID and Mental Health in America Long COVID: major findings, mechanisms and recommendations The prevalence and long-term health effects of Long COVID among hospitalised and non-hospitalised populations: a systematic review and meta-analysis 'Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19' Symptoms and risk factors for long COVID in non-hospitalized adults The prevalence of SARS-CoV-2 infection and long COVID in US adults during the

BA.5 surge', June-July 2022 Long COVID, Cognitive impairment, and the stalled decline in disability rates Prevalence and correlates of long COVID symptoms among US adults 'Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK' Trajectory of long COVID symptoms after COVID-19 vaccination: community-based cohort study Risk of Long COVID in People Infected with Severe Acute Respiratory Syndrome Coronavirus 2 After 2 Doses of Coronavirus Disease 2019 Vaccine: Community-Based, Matched Cohort Study' The psychological gains from COVID-19 vaccination The epidemiology of long COVID in US adults Global prevalence of post COVID-19 condition or long COVID: a meta-analysis and systematic review Wellbeing rankings Unhappiness and age The female happiness paradox Associations of depression, anxiety, worry, perceived stress, and loneliness prior to infection with risk of post-COVID-19 conditions Decision Letter 0 Author response to Decision Letter 0 Decision Letter 1 Author response to Decision Letter 1 Decision Letter 2 Author response to Decision Letter 2 Decision Letter 3 Acceptance letter PLOS ONE. 2023. [PMC10621843](#)

106. Christopher Kricorian. P-1908. The Relationship Between Long COVID and PHQ-4 Measures of Emotional Health Open Forum Infectious Diseases. 2025. [PMC11778124](#)
107. Sergio Padilla, Christian Ledesma, Javier García-Abellán, José Alberto García, Marta Fernández-González, Alba de la Rica, Antonio Galiana, Félix Gutiérrez, Mar Masiá, et al.. Long COVID across SARS-CoV-2 variants, lineages, and sublineages The evolution of SARS-CoV-2 Sub-lineages of the SARS-CoV-2 Omicron variants: Characteristics and prevention Comparative analysis of the risks of hospitalisation and death associated with SARS-CoV-2 omicron (B.1.1.529) and delta

(B.1.617.2) variants in England: a cohort study Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 SARS-CoV-2 omicron (B.1.1.529)-related COVID-19 sequelae in vaccinated and unvaccinated patients with cancer: results from the OnCovid registry Post-Acute Sequelae After Severe Acute Respiratory Syndrome Coronavirus 2 Infection by Viral Variant and Vaccination Status: A Multicenter Cross-Sectional Study Mortality and burden of post-COVID-19 syndrome have reduced with time across SARS-CoV-2 variants in haematology patients Potential pitfalls in the use of real-world data for studying long COVID Characteristics and predictors of Long COVID among diagnosed cases of COVID-19 Symptoms and functional impairment assessed 8 months after mild COVID-19 among health care workers Post-acute COVID-19 syndrome Persistent COVID-19 symptoms in a community study of 606,434 people in England Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection ONS website, statistical bulletin, Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK: 30 March 2023 Symptoms compatible with long-COVID in healthcare workers with and without SARS-CoV-2 infection—results of a prospective multicenter cohort Severity, predictors and clinical correlates of post-COVID syndrome (PCS) in Germany: a prospective, multi-centre, population-based cohort study COVID-19 long-term sequelae: Omicron versus Alpha and Delta variants Acute and postacute sequelae associated with SARS-CoV-2 reinfection Comorbidities, multimorbidity and COVID-19 Long-term symptom recovery and health-related quality of life in patients with mild-to-moderate-severe community-acquired pneumonia Recovery and symptom trajectories up to two

years after SARS-CoV-2 infection: population-based, longitudinal cohort study Course of post covid-19 disease symptoms over time in the ComPaRe long covid prospective e-cohort Post-acute sequelae of covid-19 six to 12 months after infection: population based study Association of SARS-CoV-2 infection and persistence with long COVID Omicron variants of SARS-CoV-2 and long COVID Long COVID: major findings, mechanisms and recommendations Analysis of anti-SARS-CoV-2 Omicron-neutralizing antibody titers in different vaccinated and unvaccinated convalescent plasma sources Effect of covid-19 vaccination on long covid: systematic review Preemptive interleukin-6 blockade in patients with COVID-19 Robust long-term immunity to SARS-CoV-2 in patients recovered from severe COVID-19 after interleukin-6 blockade T-cell immunity against SARS-CoV-2 measured by an interferon- γ release assay is strongly associated with patient outcomes in vaccinated persons hospitalized with Delta or Omicron variants Antibody Response to SARS-CoV-2 is Associated with Long-term Clinical Outcome in Patients with COVID-19: a Longitudinal Study iScience. 2024. [PMC10995878](#)

108. Jay K. Varma, Chengxi Zang, Thomas W. Carton, Jason P. Block, Dhruv J. Khullar, Yongkang Zhang, Mark G. Weiner, Russell L. Rothman, Edward J. Schenck, Zhenxing Xu, Kristin Lyman, Jiang Bian, Jie Xu, Elizabeth A. Shenkman, Christine Maughan, Leah Castro-Baucom, Lisa O'Brien, Fei Wang, Rainu Kaushal, Roberto Scendoni, Roberto Scendoni, Roberto Scendoni, Roberto Scendoni, et al.. Excess burden of respiratory and abdominal conditions following COVID-19 infections during the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER program Excess burden of respiratory and abdominal conditions following COVID-19 infections during

the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER Program Excess burden of respiratory and abdominal conditions following COVID-19 infections during the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER program. Excess burden of respiratory and abdominal conditions following COVID-19 infections during the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER program. Characterising long COVID: a living systematic review Global Prevalence of Post COVID-19 Condition or Long COVID: A Meta-Analysis and Systematic Review The origins and potential future of SARS-CoV-2 variants of concern in the evolving COVID-19 pandemic COVID-19 due to the B.1.617.2 (Delta) variant compared to B.1.1.7 (Alpha) variant of SARS-CoV-2: a prospective observational cohort study Pathophysiology and mechanism of long COVID: a comprehensive review PCORnet® 2020: current state, accomplishments, and future directions Changing the research landscape: the New York City Clinical Data Research Network OneFlorida Clinical Research Consortium: Linking a Clinical and Translational Science Institute With a Community-Based Distributive Medical Education Model Emergence and expansion of SARS-CoV-2 B.1.526 after identification in New York Data-driven analysis to understand long COVID using electronic health records from the RECOVER initiative An Empirical Transition Matrix for Non-Homogeneous Markov Chains Based on Censored Observations Use of Stabilized Inverse Propensity Scores as Weights to Directly Estimate Relative Risk and Its Confidence Intervals Data-driven identification of post-acute SARS-CoV-2 infection subphenotypes Making Neighborhood-Disadvantage Metrics Accessible—The Neighborhood Atlas Gastrointestinal

Manifestations of SARS-CoV-2 Infection and Virus Load in Fecal Samples From a Hong Kong Cohort: Systematic Review and Meta-analysis Gastrointestinal Involvement in SARS-CoV-2 Infection A core outcome set for post-COVID-19 condition in adults for use in clinical practice and research: an international Delphi consensus study An online breathing and wellbeing programme (ENO Breathe) for people with persistent symptoms following COVID-19: a parallel-group, single-blind, randomised controlled trial Pulmonary Embolism and Deep Vein Thrombosis in COVID-19: A Systematic Review and Meta-Analysis Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19 COVID-19 and Influenza Co-infection: A Systematic Review and Meta-Analysis Leading Pathogens Involved in Co-Infection and Super-Infection with COVID-19: Forensic Medicine Considerations after a Systematic Review and Meta-Analysis COVID-19 bacteremic co-infection is a major risk factor for mortality, ICU admission, and mechanical ventilation Co-infection of SARS-CoV-2 with other respiratory viruses and performance of lower respiratory tract samples for the diagnosis of COVID-19 Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: A systematic review Decision Letter 0 Author response to Decision Letter 0 Decision Letter 1 Author response to Decision Letter 1 Decision Letter 2 Acceptance letter PLOS ONE. 2024. [PMC11156291](https://doi.org/10.1371/journal.pone.0281291)

109. Jay K. Varma, Chengxi Zang, Thomas W. Carton, Jason P. Block, Dhruv J. Khullar, Yongkang Zhang, Mark G. Weiner, Russell L. Rothman, Edward J. Schenck, Zhenxing Xu, Kristin Lyman, Jiang Bian, Jie Xu, Elizabeth A. Shenkman, Christine Maughan, Leah Castro-Baucom, Lisa O'Brien, Fei Wang, Rainu Kaushal, et al.. Excess burden of respiratory and

abdominal conditions following COVID-19 infections during the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER Program

Excess burden of respiratory and abdominal conditions following COVID-19 infections during the ancestral and Delta variant periods in the United States: An EHR-based cohort study from the RECOVER program

Characterising long COVID: a living systematic review. Global Prevalence of Post COVID-19 Condition or Long COVID: A Meta-Analysis and Systematic Review. The origins and potential future of SARS-CoV-2 variants of concern in the evolving COVID-19 pandemic. COVID-19 due to the B.1.617.2 (Delta) variant compared to B.1.1.7 (Alpha) variant of SARS-CoV-2: a prospective observational cohort study. Pathophysiology and mechanism of long COVID: a comprehensive review. PCORnet® 2020: current state, accomplishments, and future directions. Changing the research landscape: the New York City Clinical Data Research Network. OneFlorida Clinical Research Consortium: Linking a Clinical and Translational Science Institute With a Community-Based Distributive Medical Education Model. Emergence and expansion of SARS-CoV-2 B.1.526 after identification in New York. An Empirical Transition Matrix for Non-Homogeneous Markov Chains Based on Censored Observations. Use of Stabilized Inverse Propensity Scores as Weights to Directly Estimate Relative Risk and Its Confidence Intervals. Making Neighborhood-Disadvantage Metrics Accessible—The Neighborhood Atlas. Gastrointestinal Manifestations of SARS-CoV-2 Infection and Virus Load in Fecal Samples From a Hong Kong Cohort: Systematic Review and Meta-analysis. Gastrointestinal Involvement in SARS-CoV-2 Infection. A core outcome set for post-COVID-19 condition in adults for use in clinical practice and research: an international Delphi

consensus study. An online breathing and wellbeing programme (ENO Breathe) for people with persistent symptoms following COVID-19: a parallel-group, single-blind, randomised controlled trial. Pulmonary Embolism and Deep Vein Thrombosis in COVID-19: A Systematic Review and Meta-Analysis. Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19. Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: A systematic review. medRxiv. 2023. [PMC9980238](#)

110. Liang En Wee, Muhammad Ismail Bin Abdul Malek, Yong Yi Tan, Jue Tao Lim, Wei Chong Tan, Jinghao Nicholas Ngiam, Matilda Lee, Elise Kiat Yee Vong, Calvin J. Chiew, Russell Jingxian Li, Iain Bee Huat Tan, David Chien Lye, Kelvin Bryan Tan, et al.. Postacute Sequelae Following Omicron COVID-19 in Patients With Cancer The consequences of SARS-CoV-2 within-host persistence Targeting the SARS-CoV-2 reservoir in long COVID Identification of risk factors of long COVID and predictive modeling in the RECOVER EHR cohorts Higher prevalence of long COVID observed in cancer survivors: insights from a US nationwide survey How methodological pitfalls have created widespread misunderstanding about long COVID Prevalence and risk factors for long COVID among cancer patients: a systematic review and meta-analysis Prevalence and impact of COVID-19 sequelae on treatment and survival of patients with cancer who recovered from SARS-CoV-2 infection: evidence from the OnCovid retrospective, multicentre registry study COVID-19 severity and survival over time in patients with hematologic malignancies: a population-based registry study Long COVID in cancer: a matched cohort study of 1-year mortality and long COVID prevalence among patients with cancer who

survived an initial severe SARS-CoV-2 infection
Booster vaccination and post-COVID-19 condition during the Omicron variant-dominant wave: a large population-based study
SARS-CoV-2 Omicron (B.1.1.529)-related COVID-19 sequelae in vaccinated and unvaccinated patients with cancer: results from the OnCovid registry
Post-acute sequelae of COVID-19 in cancer patients: two cohorts in UK and Hong Kong
Factors associated with delayed diagnosis of symptomatic adult COVID-19 cases presenting to primary care: a population-wide study during transition from Delta to Omicron BA.1 in Singapore
Access to COVID-19 vaccination by socio-economic status in older Singaporean adults: a population-based cohort study
Long-term real-world protection afforded by third mRNA doses against symptomatic severe acute respiratory syndrome coronavirus 2 infections, coronavirus disease 19-related emergency attendances and hospitalizations amongst older Singaporeans during an Omicron XBB wave
Long-term cardiovascular, cerebrovascular, and thrombotic complications after SARS-CoV-2-Omicron infection: a retrospective cohort study
Long-term neuropsychiatric sequelae of Delta versus Omicron SARS-CoV-2 infection
Autoimmune sequelae after Delta or Omicron variant SARS-CoV-2 infection in a highly vaccinated cohort
Nirmatrelvir/ritonavir treatment and risk for postacute sequelae of COVID-19 in older Singaporeans
Prevalence and characteristics of long COVID in elderly patients: an observational cohort study of over 2 million adults in the US
COVID-19 severity and waning immunity after up to 4 mRNA vaccine doses in 73 608 patients with cancer and 621 475 matched controls in Singapore: a nationwide cohort study
Addressing extreme propensity scores via the overlap weights
Remdesivir and risk of long-term multi-systemic sequelae following COVID-19 hospitalization
Enhanced infection

prevention measures including universal N95 usage and daily testing: the impact on SARS-CoV-2 transmission in cohorted hospital cubicles through successive Delta and Omicron waves Data quality at the Singapore Cancer Registry: an overview of comparability, completeness, validity and timeliness Comparison of medical and mental health sequelae following hospitalization for COVID-19, influenza, and sepsis Long-term outcomes following hospital admission for COVID-19 versus seasonal influenza: a cohort study Long-term multisystemic sequelae post-hospitalisation for Omicron COVID-19 vs influenza: a retrospective cohort study Long-term sequelae post-hospitalization for respiratory syncytial virus versus Omicron SARS-CoV-2 or influenza in adults and children: a retrospective cohort study Decreased influenza incidence under COVID-19 control measures, Singapore Long-lasting symptoms after an acute COVID-19 infection and factors associated with their resolution The immunology of long COVID Post-acute sequelae following Omicron COVID-19 in transplant recipients: a population-based cohort study Mortality and burden of post-COVID-19 syndrome have reduced with time across SARS-CoV-2 variants in haematology patients SARS-CoV-2 shedding and evolution in patients who were immunocompromised during the Omicron period: a multicentre, prospective analysis Humoral and cellular immune responses against SARS-CoV-2 post-vaccination in immunocompetent and immunocompromised cancer populations Real-world assessment of immunogenicity in immunocompromised individuals following SARS-CoV-2 mRNA vaccination: a one-year follow-up of the prospective clinical trial COVAXID Cardiovascular events 1 year after respiratory syncytial virus infection in adults Vaccine hesitancy in cancer patients: a rapid review Hesitancy to receive the booster doses of COVID-19 vaccine among cancer

patients in China: a multicenter cross-sectional survey—four PLADs, China, 2022 Factors influencing COVID-19 vaccine hesitancy among patients with serious chronic illnesses during the initial Australian vaccine rollout: a multi-centre qualitative analysis using the health belief model Behaviors, perceptions, and impact of the COVID-19 pandemic and vaccination on oncology patients in New Mexico with substantial representation of racial minorities and rural residents Characterizing the clinical and economic burden of COVID-19 among individuals with immunocompromising conditions in Ontario, Canada—a matched, population-based observational study Association of nirmatrelvir-ritonavir with post-acute sequelae and mortality among patients who are immunocompromised with COVID-19 in Hong Kong: a retrospective cohort study JAMA Network Open. 2026. [PMC13040405](#)

111. Hongkun Xu, Tingting Lu, Yajie Liu, Jingqi Yang, Simeng Ren, Baojin Han, Honghao Lai, Long Ge, Jie Liu, et al.. Prevalence and risk factors for long COVID among cancer patients: a systematic review and meta-analysis Follow up of patients with severe coronavirus disease 2019 (COVID-19): Pulmonary and extrapulmonary disease sequelae Persistence of symptoms and quality of life at 35 days after hospitalization for COVID-19 infection Long-term impact of COVID-19: A systematic review of the literature and meta-analysis 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study More than 100 persistent symptoms of SARS-coV-2 (Long COVID): A scoping review Long-term cardiovascular outcomes of COVID-19 A clinical case definition of post-COVID-19 condition by a Delphi consensus Definition of post-COVID-19 condition among published research studies COVID-19 mortality in patients with cancer on chemotherapy or other anticancer treatments: a

prospective cohort study COVID-19 in patients with thoracic Malignancies (TERAVOLT): first results of an international, registry-based, cohort study COVID-19 prevalence and mortality in patients with cancer and the effect of primary tumour subtype and patient demographics: a prospective cohort study Clinical impact of COVID-19 on patients with cancer (CCC19): a cohort study One-year adverse outcomes among US adults with post-COVID-19 condition vs those without COVID-19 in a large commercial insurance database Prevalence and impact of COVID-19 sequelae on treatment and survival of patients with cancer who recovered from SARS-CoV-2 infection: evidence from the OnCovid retrospective, multicentre registry study The effect of clinical decision making for initiation of systemic anticancer treatments in response to the COVID-19 pandemic in England: a retrospective analysis The PRISMA 2020 statement: an updated guideline for reporting systematic reviews Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data Undue reliance on I^2 in assessing heterogeneity may mislead A basic introduction to fixed-effect and random-effects models for meta-analysis Conducting meta-analyses in R with the metafor package Long COVID in cancer: A matched cohort study of 1-year mortality and long COVID prevalence among patients with cancer who survived an initial severe SARS-coV-2 infection Long COVID in cancer patients: preponderance of symptoms in majority of patients over long time period Characterization and predictive risk scoring of long COVID in a south Indian cohort after breakthrough COVID infection; a prospective single centre study 16 months follow up of patients' Behavior and mild COVID-19 patterns in a large cohort of cancer patients during the pandemic Long-term effects of COVID-19

on cancer patients: the experience from Guy's Cancer Centre
One-year mortality and consequences of COVID-19 in cancer patients: A cohort study
Long-COVID in patients with cancer previously treated with early anti-SARS-coV-2 therapies in an out-of-hospital setting: A single-center experience
Persistence of long-term COVID-19 sequelae in patients with cancer: An analysis from the OnCovid registry
Mortality and burden of post-COVID-19 syndrome have reduced with time across SARS-CoV-2 variants in haematology patients
Assessment of SARS-coV-2 infection according to previous metabolic status and its association with mortality and post-acute COVID-19
Long-lasting symptoms after an acute COVID-19 infection and factors associated with their resolution
Acute and post-acute COVID-19 severity and mortality in patients with hematologic Malignancies: A population-based registry study
One-year temporal changes in long COVID prevalence and characteristics: A systematic review and meta-analysis
Long-term sequelae of COVID-19: A systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms
Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study
Physical phenotype of blood cells is altered in COVID-19
Gut microbiota dynamics in a prospective cohort of patients with post-acute COVID-19 syndrome
A one-year follow-up study of systematic impact of long COVID symptoms among patients post SARS-CoV-2 omicron variants infection in Shanghai, China
Post-discharge persistent symptoms and health-related quality of life after hospitalization for COVID-19
Risk factors associated with post-COVID-19 condition: A systematic review and meta-analysis
Long COVID: systemic inflammation and obesity as therapeutic targets
Diabetes and COVID19: a bidirectional relationship
Post COVID-19 syndrome ("Long COVID") and diabetes: challenges in diagnosis and

management Smoking increases the risk of post-acute COVID-19 syndrome: Results from a French community-based survey The effect of smoking on COVID-19 severity: A systematic review and meta-analysis Increased expression of ACE2, the SARS-CoV-2 entry receptor, in alveolar and bronchial epithelium of smokers and COPD subjects Angiotensin-converting enzyme 2 expression in COPD and IPF fibroblasts: the forgotten cell in COVID-19 The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice The isolated effect of age on the risk of COVID-19 severe outcomes: a systematic review with meta-analysis Sex differences in sequelae from COVID-19 infection and in long COVID syndrome: a review Symptoms and risk factors for long COVID in non-hospitalized adults *Frontiers in Oncology*. 2025. [PMC11774732](#)

112. Carolyn T. Bramante, Thomas G. Stewart, David R. Boulware, Matthew W. McCarthy, Yue Gao, Russell L. Rothman, Ahmad Mourad, Florence Thicklin, Jonathan B. Cohen, Idania T. Garcia del Sol, Nirav S. Shah, Manisha Mehta, Orlando Quintero Cardona, Jake Scott, Adit A. Ginde, Mario Castro, Dushyantha Jayaweera, Mark Sulkowski, Nina Gentile, Kathleen McTigue, G. Michael Felker, Sean Collins, Sarah E. Dunsmore, Stacey J. Adam, Christopher J. Lindsell, Adrian F. Hernandez, Susanna Naggie, et al.. Metformin on the Presence of COVID-19 Symptoms Over 6 Months: The ACTIV-6 Randomized Clinical Trial *medRxiv*. 2025. [PMC12363699](#)
113. Steve Johnson, Carolyn Bramante, Til Stürmer, John Buse, Talia Wiggen, Jared Huling, Andrew Toler, et al.. 595 Is PHASTR faster? A target trial emulation case study in the N3C *Journal of Clinical and Translational Science*. 2025. [PMC12038486](#)

114. Anna Puigdemívol-Sánchez, Antonio Arévalo-Genicio, M^a Carmen García-Arqué, Marta Gragea-Nocete, Celia Lozano-Paz, Vanessa Moro-Casasola, Cristina Pérez-Díaz, Roger Valls-Foix, Ramon Roca-Puig, Maria Llistosella, et al.. Long COVID and Reduced Thrombosis in Antihistamine-Treated Patients: An Observational Study in the Metropolitan Area of Barcelona
The risk of Long COVID symptoms: A systematic review and meta-analysis of controlled studies
Post-acute COVID-19 syndrome
Long COVID: Major findings, mechanisms and recommendations
Long COVID Mechanisms, Microvascular Effects, and Evaluation Based on Incidence
Post COVID-19 Condition (Long COVID)
Long COVID Syndrome Prevalence in 2025 in an Integral Healthcare Consortium in the Metropolitan Area of Barcelona: Persistent and Transient Symptoms
Risk of acute myocardial infarction and ischaemic stroke following COVID-19 in Sweden: A self-controlled case series and matched cohort study
Association of COVID-19 With Major Arterial and Venous Thrombotic Diseases: A Population-Wide Cohort Study of 48 Million Adults in England and Wales
Effect of antiplatelet therapy after COVID-19 diagnosis: A systematic review with meta-analysis and trial sequential analysis
Alleviation of COVID-19 Symptoms and Reduction in Healthcare Utilization Among High-risk Patients Treated With Nirmatrelvir/Ritonavir (NMV/R): A Phase 3 Randomized Trial
Oral Nirmatrelvir-Ritonavir as Postexposure Prophylaxis for COVID-19
Nirmatrelvir-Ritonavir and Symptoms in Adults with Postacute Sequelae of SARS-CoV-2 Infection: The STOP-PASC Randomized Clinical Trial
A SARS-CoV-2 Protein Interaction Map Reveals Targets for Drug-Repurposing
Antihistamines and azithromycin as a treatment for COVID-19 on primary health care—A retrospective observational study in elderly patients
Clinical Characteristics, Frailty, and Mortality of Residents with

COVID-19 in Nursing Homes of a Region of Madrid
Antihistamines as an early treatment for COVID-19
COVID-19 in Relation to Chronic Antihistamine Prescription
COVID-19 Pandemic Waves and 2024–2025 Winter Season in Relation to
Angiotensin-Converting Enzyme Inhibitors, Angiotensin
Receptor Blockers and Amantadine CoVariants. Institute of
Social and Preventive Medicine University of Bern,
Switzerland & SIB Swiss Institute of 375 Bioinformatics
OpenEpi: Open Source Epidemiologic Statistics for Public
Health, Versión Multiple Hypotheses Testing with Weights
Exact Calculations of Average Power for the Benjamini-
Hochberg Procedure JASP Linear regression. Collinearity. On
'An Introduction to Statistical Learning' Effect of famotidine
on cognitive and behavioral dysfunctions induced in post-
COVID-19 infection: A randomized, double-blind, and placebo-
controlled study Network Pharmacology Study to Elucidate
the Key Targets of Underlying Antihistamines against
COVID-19 Disease-Associated Variants in GRIN1, GRIN2A and
GRIN2B genes: Insights into NMDA Receptor Structure,
Function, and Pathophysiology Effects of H1- and H2-
antihistamines on platelet-activating factor and bradykinin-
induced inflammatory responses in human skin How to handle
off-label prescriptions of rupatadine, a second-generation
antihistamine and PAF antagonist: A review Oral famotidine
versus placebo in non-hospitalised patients with COVID-19: A
randomised, double-blind, data-intensive, phase 2 clinical trial
Pandemic H1N1 influenza infection and vascular thrombosis
Acute Cardiovascular Events Associated With Influenza in
Hospitalized Adults: A Cross-sectional Study Thromboembolic
Events in Patients with Influenza: A Scoping Review
Generalitat de Catalunya Generalitat de Catalunya The
seasonal distribution of immune thrombotic
thrombocytopenic purpura is influenced by geography:

Epidemiologic findings from a multi-center analysis of 719 disease episodes Seasonal variations in hospitalizations of heart failure patients: A United States nationwide analysis Association between air pollution-cold wave sequential events and ischaemic stroke incidence among elderly adults in Tianjin, China: A retrospective cohort study Impact of Estrogen on Purinergic Signaling in Microvascular Disease Comparability of clinical trials and spontaneous reporting data regarding COVID-19 vaccine safety Update on Thromboembolic Events After Vaccination Against COVID-19 Thrombocytopenic, thromboembolic and haemorrhagic events following second dose with BNT162b2 and ChAdOx1: Self-controlled case series analysis of the English national sentinel cohort The Role of Sterile Inflammation in Thrombosis: Consequences for Cardiovascular Disease and COVID-19 Diabetic cardiomyopathy and COVID-19: Intersecting pathways and amplified cardiovascular risk Viruses. 2026. [PMC12945107](#)

115. Freddy Frost, José Miguel Rivera-Caravaca, Gregory Y. H. Lip. The use of oral anticoagulation at the time of acute COVID-19 infection and subsequent development of long-COVID/post-acute sequelae of SARS-CoV-2 infection Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19 Postdischarge thromboembolic outcomes and mortality of hospitalized patients with COVID-19: the CORE-19 registry Sustained prothrombotic changes in COVID-19 patients 4 months after hospital discharge Serum metabolic profile in patients with Long-Covid (PASC) syndrome: clinical implications Triple antithrombotic therapy after ACS and PCI in patients on chronic oral anticoagulation: update Long COVID: the nature of thrombotic sequelae determines the necessity of early anticoagulation The association of

preinfection daily oral anticoagulation use and all-Cause in hospital mortality from novel coronavirus 2019 at 21 days: a retrospective cohort study Rivaroxaban versus no anticoagulation for post-discharge thromboprophylaxis after hospitalisation for COVID-19 (MICHELLE): an open-label, multicentre, randomised, controlled trial Vascular thromboembolic events following COVID-19 hospital discharge: incidence and risk factors Journal of Thrombosis and Thrombolysis. 2025. [PMC12043746](#)

116. Matthew Brown, John Gerrard, Lynne McKinlay, John Marquess, Teneika Sparrow, Ross Andrews. Ongoing symptoms and functional impairment 12 weeks after testing positive for SARS-CoV-2 or influenza in Australia: an observational cohort study A clinical case definition of post-COVID-19 condition by a Delphi consensus Long COVID: long-term health outcomes and implications for policy and research Post-COVID syndrome. The never ending challenge Using condition specific patient reported outcome measures for long Covid Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: A systematic review The US now has a research plan for long COVID-is it enough The prevalence and long-term health effects of long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis Association between BNT162b2 vaccination and reported incidence of post-COVID-19 symptoms: cross-sectional study 2020-21 Taking pandemic sequelae seriously: from the Russian influenza to COVID-19 long-haulers Long COVID in Australia - a review of the literature Effect of COVID-19 vaccination on long Covid: systematic review COVID-19 vaccine Rollout List of Notifiable conditions Queensland weekly influenza surveillance report The strengthening the reporting of observational studies in

epidemiology (STROBE) statement: guidelines for reporting observational studies Version 2 Can the post-COVID-19 functional status scale discriminate between patients with different levels of fatigue, quality of life and functional performance? Chapter 4. how to interpret SEIFA score distributions Influenza vaccination coverage - by jurisdiction Coronavirus (COVID-19) case numbers and Statistics Causation or confounding: why controls are critical for characterizing long COVID Prevalence of ongoing symptoms following Coronavirus (COVID-19) infection in the UK: 1 September 2022 Epidemiologic and economic Modelling of optimal COVID-19 policy: public health and social measures, masks and vaccines in Victoria, Australia Long COVID: major findings, mechanisms and recommendations Incidence, Co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19 Risk of long COVID associated with Delta versus Omicron variants of SARS-Cov-2 BMJ Public Health. 2023. [PMC11812696](https://doi.org/10.1136/bmjph-2023-100496)

117. Jianpeng Cai, Ke Lin, Haocheng Zhang, Quanlin Xue, Kun Zhu, Guanmin Yuan, Yuhan Sun, Feng Zhu, Jingwen Ai, Sen Wang, Wenhong Zhang, et al.. A one-year follow-up study of systematic impact of long COVID symptoms among patients post SARS-CoV-2 omicron variants infection in Shanghai, China Correction A clinical case definition of post-COVID-19 condition by a delphi consensus Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Outcomes among confirmed cases and a matched comparison group in the long-COVID in Scotland study Six-month sequelae of post-vaccination SARS-CoV-2 infection: a retrospective cohort study of 10,024 breakthrough infections Risk of long COVID in people

infected with severe acute respiratory syndrome coronavirus 2 after 2 doses of a coronavirus disease 2019 vaccine: community-based, matched cohort study Prevalence and correlates of long COVID symptoms among US adults Symptoms and risk factors for long COVID in non-hospitalized adults The GAD-7 scale was accurate for diagnosing generalised anxiety disorder A brief measure for assessing generalized anxiety disorder: the GAD-7 EQ-5D: a measure of health status from the EuroQol group The PHQ-9: validity of a brief depression severity measure Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care The posttraumatic stress disorder checklist for DSM-5 (PCL-5): development and initial psychometric evaluation A new equation to estimate glomerular filtration rate Some characteristics of clinical sequelae of COVID-19 survivors from Wuhan, China: a multi-center longitudinal study Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study Two-year health outcomes in hospitalized COVID-19 survivors in China 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study Follow-up of patients with COVID-19 by the delta variant after hospital discharge in Guangzhou, guandong, China Long COVID following omicron wave in eastern India-a retrospective cohort study Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Long COVID in a prospective cohort of home-isolated patients Comparative analysis of the risks of hospitalisation and death associated with SARS-CoV-2 omicron (B.1.1.529) and delta (B.1.617.2) variants in England: a cohort study Clinical severity of, and effectiveness of mRNA vaccines against, COVID-19 from omicron, delta, and alpha SARS-CoV-2 variants in the United States: prospective observational study Post-acute COVID-19 syndrome (PCS) and health-related

quality of life (HRQoL)-a systematic review and meta-analysis
 Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments
 Association between BNT162b2 vaccination and long COVID after infections not requiring hospitalization in health care workers
 Vaccinating adolescents against SARS-CoV-2 in England: a risk-benefit analysis
 Multiorgan impairment in low-risk individuals with post-COVID-19 syndrome: a prospective, community-based study
 Multi-organ impairment and long COVID: a 1-year prospective, longitudinal cohort study
 SARS-CoV-2 omicron variant: recent progress and future perspectives
 Clinical status of patients 1 year after hospital discharge following recovery from COVID-19: a prospective cohort study
 Post-COVID-19 functional status six-months after hospitalization
 Medium-term effects of SARS-CoV-2 infection on multiple vital organs, exercise capacity, cognition, quality of life and mental health, post-hospital discharge
 Emerging
 Microbes & Infections. 2023. [PMC10281439](https://pubmed.ncbi.nlm.nih.gov/3681439/)

118. Hui Zhang, Peng Yang, Xiaoying Gu, Ying Sun, Rongling Zhang, Daitao Zhang, Jiaojiao Zhang, Yeming Wang, Chunna Ma, Min Liu, Jiaxin Ma, Aili Li, Yingying Wang, Xiao Ma, Xiaojing Cui, Yimin Wang, Zhibo Liu, Wei Wang, Zhi Zheng, Yong Li, Jin Wu, Quanyi Wang, Bin Cao, et al.. Health outcomes one year after Omicron infection among 12,789 adults: a community-based cross-sectional study
 COVID-19 cases | WHO COVID-19 dashboard. datadot 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study
 1-year outcomes in hospital survivors with COVID-19: a longitudinal cohort study
 Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study
 3-year outcomes of discharged survivors of COVID-19 following the SARS-CoV-2 omicron (B.1.1.529) wave in 2022 in China: a longitudinal

cohort study A clinical case definition of post-COVID-19 condition by a Delphi consensus Development of a definition of postacute sequelae of SARS-CoV-2 infection Persistence of somatic symptoms after COVID-19 in The Netherlands: an observational cohort study Lung function trajectories in mild COVID-19 with two-year T follow-up Lung-function trajectories in COVID-19 survivors after discharge: a two-year longitudinal cohort study Kidney outcomes in long COVID Long-term cardiovascular outcomes of COVID-19 Long-term gastrointestinal outcomes of COVID-19 Long-term neurologic outcomes of COVID-19 Risks and burdens of incident diabetes in long COVID: a cohort study Risks of mental health outcomes in people with covid-19: cohort study High-dimensional characterization of post-acute sequelae of COVID-19 Three-year outcomes of post-acute sequelae of COVID-19 Recovery and symptom trajectories up to two years after SARS-CoV-2 infection: population based, longitudinal cohort study Cognitive and psychiatric symptom trajectories 2-3 years after hospital admission for COVID-19: a longitudinal, prospective cohort study in the UK Long COVID science, research and policy Estimating the transmission dynamics of SARS-CoV-2 Omicron BF.7 in Beijing after adjustment of the zero-COVID policy in November-December 2022 Characterisation of SARS-CoV-2 variants in Beijing during 2022: an epidemiological and phylogenetic analysis Long-term prognosis at 1.5 years after infection with wild-type strain of SARS-CoV-2 and alpha, delta, as well as omicron variants Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 Post-covid medical complaints following infection with SARS-CoV-2 Omicron vs Delta variants China center for disease control and prevention Long COVID in a highly vaccinated but largely unexposed Australian population following the 2022 SARS-

CoV-2 Omicron wave: a cross-sectional survey Long-term health impacts of COVID-19 among 242,712 adults in England Postacute sequelae of SARS-CoV-2 infection in the pre-delta, delta, and omicron eras Prevalence and determinants of long-term post-COVID conditions in the United States: 2022 behavioral risk factor surveillance system Prevalence and risk factors for persistent symptoms after COVID-19: a systematic review and meta-analysis Long COVID: major findings, mechanisms and recommendations Acute and postacute sequelae associated with SARS-CoV-2 reinfection Early biological markers of post-acute sequelae of SARS-CoV-2 infection Confronting COVID-19-associated cough and the post-COVID syndrome: role of viral neurotropism, neuroinflammation, and neuroimmune responses 3-month, 6-month, 9-month, and 12-month respiratory outcomes in patients following COVID-19-related hospitalisation: a prospective study Long-term radiological and pulmonary function abnormalities at 3 years after COVID-19 hospitalisation: a longitudinal cohort study Normal lung function and mortality in world trade center responders and national health and nutrition examination survey III participants Comparative single-cell analysis reveals IFN- γ as a driver of respiratory sequelae after acute COVID-19 SARS-CoV-2 infection and lung regeneration Immuno-proteomic profiling reveals aberrant immune cell regulation in the airways of individuals with ongoing post-COVID-19 respiratory disease Immune signatures underlying post-acute COVID-19 lung sequelae Monocyte and macrophage function in respiratory viral infections Postacute sequelae of COVID-19 at 2 years Neurological and psychiatric risk trajectories after SARS-CoV-2 infection: an analysis of 2-year retrospective cohort studies including 1 284 437 patients Taking pandemic sequelae seriously: from the Russian influenza to COVID-19

long-haulers Long-term outcomes of pandemic 2009 influenza A(H1N1)-associated severe ARDS Narcolepsy onset is seasonal and increased following the 2009 H1N1 pandemic in China - PubMed Long-term risk of Parkinson disease following influenza and other infections Cerebral micro-structural changes in COVID-19 patients - an MRI-based 3-month follow-up study SARS-CoV-2 is associated with changes in brain structure in UK Biobank Brain 18F-FDG PET imaging in outpatients with post-COVID-19 conditions: findings and associations with clinical characteristics Mild respiratory COVID can cause multi-lineage neural cell and myelin dysregulation Blood-brain barrier disruption and sustained systemic inflammation in individuals with long COVID-associated cognitive impairment Para-infectious brain injury in COVID-19 persists at follow-up despite attenuated cytokine and autoantibody responses Virus-induced acute respiratory distress syndrome causes cardiomyopathy through eliciting inflammatory responses in the heart Cardiovascular effects of the post-COVID-19 condition Monocyte migration profiles define disease severity in acute COVID-19 and unique features of long COVID Elevated circulating monocytes and monocyte activation in COVID-19 convalescent individuals Persistence of SARS CoV-2 S1 protein in CD16+ monocytes in post-acute sequelae of COVID-19 (PASC) up to 15 Months post-infection Long-term multiple metabolic abnormalities among healthy and high-risk people following nonsevere COVID-19 Risks and burdens of incident dyslipidaemia in long COVID: a cohort study Altered lipid profile in COVID-19 patients and metabolic reprogramming Fibrin drives thromboinflammation and neuropathology in COVID-19 COVID-19 and its implications for thrombosis and anticoagulation High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective

cohort study Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study Persistent endotheliopathy in the pathogenesis of long COVID syndrome Long COVID: pathophysiological factors and abnormalities of coagulation Differentiation of prior SARS-CoV-2 infection and postacute sequelae by standard clinical laboratory measurements in the RECOVER cohort The Lancet Regional Health: Western Pacific. 2025. [PMC11992576](#)

119. Liane S Canas, Erika Molteni, Jie Deng, Carole H Sudre, Benjamin Murray, Eric Kerfoot, Michela Antonelli, Khaled Rjoob, Joan Capdevila Pujol, Lorenzo Polidori, Anna May, Marc F Österdahl, Ronan Whiston, Nathan J Cheetham, Vicky Bowyer, Tim D Spector, Alexander Hammers, Emma L Duncan, Sebastien Ourselin, Claire J Steves, Marc Modat, et al.. Profiling post-COVID-19 condition across different variants of SARS-CoV-2: a prospective longitudinal study in unvaccinated wild-type, unvaccinated alpha-variant, and vaccinated delta-variant populations Coronavirus (COVID-19) Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2 COVID-19 rapid guideline: managing the long-term effects of COVID-19 Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK: 22 June 2022 Recommendations for the recognition, diagnosis, and management of long COVID: a Delphi study Impact of COVID-19 vaccination on the risk of developing long-COVID and on existing long-COVID symptoms: a systematic review UK marks one year since deploying world's first COVID-19 vaccine Proposed subtypes of post-COVID-19 syndrome (or long-COVID) and their respective potential therapies COVID-19 trajectories among 57 million adults in England: a cohort study using electronic health records Emerging potential mechanisms and predispositions to the

neurological manifestations of COVID-19 Axes of prognosis: identifying subtypes of COVID-19 outcomes Rapid implementation of mobile technology for real-time epidemiology of COVID-19 Safety and efficacy of the BNT162b2 mRNA COVID-19 vaccine Variants of concern (VOC) and under investigation (VUI) and any other variant by weeks and days Symptom clusters in COVID-19: a potential clinical prediction tool from the COVID Symptom Study app Multivariate time series clustering based on common principal component analysis Accessible data curation and analytics for international-scale citizen science datasets Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation (PHOSP-COVID): a UK multicentre, prospective cohort study Characterising patterns of COVID-19 and long COVID symptoms: evidence from nine UK longitudinal studies High-dimensional characterization of post-acute sequelae of COVID-19 SARS-CoV-2 is associated with changes in brain structure in UK Biobank Prospective evaluation of autonomic dysfunction in post-acute sequela of COVID-19 Clues to long COVID Symptoms and risk factors for long COVID in non-hospitalized adults Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study Do vaccines protect from long COVID? Long COVID: to investigate immunological mechanisms and sex/gender related aspects as fundamental steps for tailored therapy Early detection of COVID-19 in the UK using self-reported symptoms: a large-scale, prospective, epidemiological surveillance study The Lancet. Digital Health. 2023. [PMC10187990](https://doi.org/10.1016/S2666-5264(23)00090-0)

120. Fernando Luis Scolari, Julia Spinardi, Mariana Motta Dias da Silva, Geraldine Trott, Cristina de Oliveira Rodrigues, Marciane Maria Rover, Emanuel Maltempi de Souza, Josélia

Larger Manfio, Nathan Iori Camargo, Ana Paula de Souza, Denise de Souza, Raíne Fogliati De Carli, Emelyn de Souza Roldão, Duane Mocellin, Aline Paula Miozzo, Gabrielle Nunes da Silva, Jennifer Menna Barreto de Souza, Rosa da Rosa Minho dos Santos, Carolina Rothmann Itaquí, Gabriela Soares Rech, Vivian Menezes Irineu, Saionara Cristina Francisco, Odilson Silvestre, Precil Diego Miranda de Menezes Neves, Lucas Tramujas, Vandack Nobre, Sidiclei Machado Carvalho, Carlos Delmar do Amaral Ferreira, Jaqueline Carvalho de Oliveira, Carla Adriane Royer, Rafael Messias Luiz, Valter Antonio Baura, Daniela Fiori Gradia, Ana Paula Carneiro Brandalize, Hellen Abreu Pereira, Carolina Gracia Poitevin, Caroline Cabral Robinson, Bruna Brandao Barreto, Paulo R. Schvartzman, Milena Marcolino, Ana Carolina Peçanha Antonio, Carisi Anne Polanczyk, Juçara Gasparetto Maccari, Luiz Antonio Nasi, Srinivas Rao Valluri, Viviane Wal Julião, Florence Lefebvre d'Hellencourt, Moe H. Kyaw, Graciela Del Carmen Morales Castillo, Maicon Falavigna, Regis Goulart Rosa, et al.. Impact of long COVID phenotypes on quality of life following symptomatic omicron infection in Brazil: a machine learning analysis Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Attributes and predictors of long COVID Association between acute disease severity and one-year quality of life among post-hospitalisation COVID-19 patients: coalition VII prospective cohort study Studying the post-COVID-19 condition: research challenges, strategies, and importance of core outcome set development Clinical phenotypes and quality of life to define post-COVID-19 syndrome: a cluster analysis of the multinational, prospective ORCHESTRA cohort Characterizing long COVID in an international cohort: 7 months of symptoms and their impact A distinct symptom

pattern emerges for COVID-19 long-haul: a nationwide study
Characterization of post-COVID syndromes by symptom
cluster and time period up to 12 months post-infection: a
systematic review and meta-analysis Identification of distinct
long COVID clinical phenotypes through cluster analysis of
self-reported symptoms Five cluster classifications of long
COVID and their background factors: a cross-sectional study
in Japan A machine learning approach identifies distinct
early-symptom cluster phenotypes which correlate with
hospitalization, failure to return to activities, and prolonged
COVID-19 symptoms Symptom cluster analysis of long
COVID-19 in patients discharged from the temporary
COVID-19 hospital in Mexico City COVID symptoms, symptom
clusters, and predictors for becoming a long-hauler looking
for clarity in the haze of the pandemic Global lineage
evolution pattern of SARS-CoV-2 in Africa, America, Europe,
and asia: a comparative analysis of variant clusters and their
relevance across continents Epidemiological surveillance
reveals the rise and establishment of the Omicron SARS-
CoV-2 variant in Brazil BNT162b2 against COVID-19 in Brazil
using a test-negative design: study protocol and statistical
analysis plan Long-term health-related quality of life and
outcomes after hospitalization for COVID-19 in brazil: Post-
COVID Brazil 1 study protocol Health-related quality of life
and long-term outcomes after mildly symptomatic COVID-19:
the Post-COVID Brazil study 2 protocol Association between
vaccination and persistent COVID-19-related symptoms
among patients with mild Omicron infection: a prospective
cohort study Brazilian valuation of EQ-5D-3L health States
The minimum clinically important difference for EQ-5D index:
a critical review EQ-5D Brazilian population norms NbClust:
an R package for determining the relevant number of clusters
in a data set Hierarchical grouping to optimize an objective

function Long-term reduced functional capacity and quality of life in hospitalized COVID-19 patients Recovery and symptom trajectories up to two years after SARS-CoV-2 infection: population-based, longitudinal cohort study Persistent COVID-19 symptoms 1 year after hospital discharge: a prospective multicenter study Review of organ damage from COVID and long COVID: a disease with a spectrum of pathology Understanding autoimmune response after SARS-CoV-2 infection and the pathogenesis/mechanisms of long COVID Circular RNA: a promising new star of vaccine A nationwide questionnaire study of post-acute symptoms and health problems after SARS-CoV-2 infection in Denmark Presence of COVID-19 self-reported symptoms at 12 months in patients discharged from hospital in 2020–2021: a Spanish cross-sectional study Symptom burden and immune dynamics 6 to 18 months following mild SARS-CoV-2 infection: a case-control study Post-COVID-19 syndrome: physical capacity, fatigue and quality of life Suggestions on home quarantine and recovery of novel coronavirus patients Post-acute sequelae of COVID-19 six to 12 months after infection: population-based study Cognitive dysfunction and health-related quality of life among older Chinese BMC Infectious Diseases. 2025. [PMC12595840](#)

121. Kristine M. Erlandson, Linda N. Geng, Caitlin A. Selvaggi, Tanayott Thaweethai, Peter Chen, Nathan B. Erdmann, Jason D. Goldman, Timothy J. Henrich, Mady Hornig, Elizabeth W. Karlson, Stuart D. Katz, C. Kim, Sushma K. Cribbs, Adeyinka O. Laiyemo, Rebecca Letts, Janet Y. Lin, Jai Marathe, Sairam Parthasarathy, Thomas F. Patterson, Brittany D. Taylor, Elizabeth R. Duffy, Monika Haack, Boris Julg, Gabrielle Maranga, Carla Hernandez, Nora Singer, Jenny Han, Priscilla Pemu, Hassan Brim, Hassan Ashktorab, Alexander W. Charney, Juan Wisnivesky, Jenny Lin, Helen Y. Chu, Minjoung

Go, Upinder Singh, Emily B. Levitan, Paul A. Goepfert, Janko Ž. Nikolich, Harvey Hsu, Michael J. Peluso, J. Daniel Kelly, Megumi Okumura, Valerie J Flaherman, John G. Quigley, Jerry A. Krishnan, Mary Beth Scholand, Rachel Hess, Torri D. Metz, Maged M. Costantine, Dwight J Rouse, Barbara S. Taylor, Mark P. Goldberg, Gailen D. Marshall, Jeremy Wood, David Warren, Leora Horwitz, Andrea S. Foulkes, Grace A McComsey, et al.. Differentiation of Prior SARS-CoV-2 Infection and Postacute Sequelae by Standard Clinical Laboratory Measurements in the RECOVER Cohort Estimated Global Proportions of Individuals With Persistent Fatigue, Cognitive, and Respiratory Symptom Clusters Following Symptomatic COVID-19 in 2020 and 2021 Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status Generalisable long COVID subtypes: findings from the NIH N3C and RECOVER programmes Characterizing long COVID in an international cohort: 7 months of symptoms and their impact 1-year outcomes in hospital survivors with COVID-19: a longitudinal cohort study Post-acute COVID-19 syndrome Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection More than 50 long-term effects of COVID-19: a systematic review and meta-analysis Multidisciplinary Center Care for Long COVID Syndrome-A Retrospective Cohort Study "Living like an empty gas tank with a leak": Mixed methods study on post-acute sequelae of COVID-19 Persistence, Magnitude, and Patterns of Postacute Symptoms and Quality of Life Following Onset of SARS-CoV-2 Infection: Cohort Description and Approaches for Measurement Long COVID: major findings, mechanisms and recommendations Pathogenic mechanisms of post-acute sequelae of SARS-CoV-2 infection (PASC) Long COVID or Post-acute Sequelae of COVID-19 (PASC): An Overview of Biological Factors That May Contribute to

Persistent Symptoms Immune mechanisms underlying COVID-19 pathology and post-acute sequelae of SARS-CoV-2 infection (PASC) Force RMPT. Viral persistence, reactivation, and mechanisms of long COVID Inflammatory and vascular biomarkers in post-COVID-19 syndrome: A systematic review and meta-analysis of over 20 biomarkers Biomarkers in long COVID-19: A systematic review Laboratory Findings and Biomarkers in Long COVID: What Do We Know So Far? Insights into Epidemiology, Pathogenesis, Therapeutic Perspectives and Challenges Increased interleukin-6 is associated with long COVID-19: a systematic review and meta-analysis Circulating anti-nuclear autoantibodies in COVID-19 survivors predict long COVID symptoms Autoimmunity is a hallmark of post-COVID syndrome Distinguishing features of long COVID identified through immune profiling Chronic viral coinfections differentially affect the likelihood of developing long COVID Herpesvirus infections and post-COVID-19 manifestations: a pilot observational study A Longitudinal Study of COVID-19 Sequelae and Immunity Researching COVID to Enhance Recovery (RECOVER) adult study protocol: Rationale, objectives, and design Mechanisms of COVID-19 pathogenesis in diabetes COVID-19 Vaccination Prior to SARS-CoV-2 Infection Reduced Risk of Subsequent Diabetes Mellitus: A Real-World Investigation Using U.S. Electronic Health Records Risks and burdens of incident diabetes in long COVID: a cohort study Post COVID-19 Syndrome ("Long COVID") and Diabetes: Challenges in Diagnosis and Management Inflammatory and Prothrombotic Biomarkers Contribute to the Persistence of Sequelae in Recovered COVID-19 Patients Chronic Fatigue, Depression and Anxiety Symptoms in Long COVID Are Strongly Predicted by Neuroimmune and Neuro-Oxidative Pathways Which Are

Caused by the Inflammation during Acute Infection Clinical and pulmonary function analysis in long-COVID revealed that long-term pulmonary dysfunction is associated with vascular inflammation pathways and metabolic syndrome TEG(®), Microclot and Platelet Mapping for Guiding Early Management of Severe COVID-19 Coagulopathy SARS-CoV-2 Initiates Programmed Cell Death in Platelets A central role for amyloid fibrin microclots in long COVID/PASC: origins and therapeutic implications Muscle abnormalities worsen after post-exertional malaise in long COVID Plasmapheresis to remove amyloid fibrin(ogen) particles for treating the post-COVID-19 condition Long-term perturbation of the peripheral immune system months after SARS-CoV-2 infection Persisting Platelet Activation and Hyperactivity in COVID-19 Survivors Persistent complement dysregulation with signs of thromboinflammation in active Long Covid The Occurrence of Hyperactivated Platelets and Fibrinoid Microclots in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) Platelet Storage Pool Deficiency and Elevated Inflammatory Biomarkers Are Prevalent in Postural Orthostatic Tachycardia Syndrome Long-term effect of COVID-19 infection on kidney function among COVID-19 patients followed in post-COVID-19 recovery clinics in British Columbia, Canada Long-term clinical and biochemical residue after COVID-19 recovery Characteristics of Renal Function in Patients Diagnosed With COVID-19: An Observational Study The impact of SARS-CoV-2 infection on renal function in patients with biopsy-proven kidney diseases Long-term effect of COVID-19 infection on kidney function among COVID-19 patients followed in post-COVID-19 recovery clinics in British Columbia, Canada Is the kidney a target of SARS-CoV-2? The Association of COVID-19 With Acute Kidney Injury Independent of Severity of Illness: A Multicenter Cohort

Study Urinary Albumin-to-Creatinine Ratio in Normal Range, Cardiovascular Health, and All-Cause Mortality Elevated urine albumin creatinine ratio increases cardiovascular mortality in coronary artery disease patients with or without type 2 diabetes mellitus: a multicenter retrospective study Association between kidney function and albuminuria with cardiovascular events in HIV-infected persons Urinary albumin-to-creatinine ratio is associated with endothelial dysfunction in HIV-infected patients receiving antiretroviral therapy Urine Albumin-to-Creatinine Ratio: A Marker of Early Endothelial Dysfunction in Youth Cardiac Involvement Due to COVID-19: Insights from Imaging and Histopathology Knockout of angiotensin converting enzyme-2 receptor leads to morphological aberrations in rodent olfactory centers and dysfunctions associated with sense of smell Association between chemosensory dysfunctions and inflammatory biomarkers in patients with SARS-CoV-2 infection: a systematic review and meta-analysis COVID-19 anosmia and gustatory symptoms as a prognosis factor: a subanalysis of the HOPE COVID-19 (Health Outcome Predictive Evaluation for COVID-19) registry Annals of internal medicine. 2024. [PMC11408082](#)

122. Ruud P.H. Raijmakers, Lise Lund Berven, Stephan P. Keijmel, Chaturaka Rodrigo, Vegard B.B. Wyller, Ben Z. Katz, Dedra Buchwald, Rachael A. Evans, Patrick Gérardin, Hans Knoop, Maria Prins, Knut Stavem, Tonje Stiansen-Sonerud, Renee Taylor, Braulio Mark Valencia Arroyo, Knut-Arne Wensaas, Joel Pradeepkumar Selvakumar, Cees van den Wijngaard, Andrew R. Lloyd, Carolina X. Sandler, et al.. Immunological associations in post-infective fatigue syndromes including Long COVID—a systematic review and meta-analysis Unexplained post-acute infection syndromes The international collaborative on fatigue following infection (COFFI)

Postinfectious fatigue: prospective cohort study in primary care A prospective observational study of post-COVID-19 chronic fatigue syndrome following the first pandemic wave in Germany and biomarkers associated with symptom severity Long COVID and post-infective fatigue syndrome: a review Characterization of fatigue states in medicine and psychiatry by structured interview Genetic associations of fatigue and other symptom domains of the acute sickness response to infection A validated scale for assessing the severity of acute infectious mononucleosis Post-COVID-19 fatigue: a systematic review Prevalence and characteristics associated with post-COVID-19 condition among nonhospitalized adolescents and young adults The chronic fatigue syndrome: a comprehensive approach to its definition and study. International Chronic Fatigue Syndrome Study Group A report--chronic fatigue syndrome: guidelines for research Myalgic encephalomyelitis: International consensus criteria Predictions and associations of fatigue syndromes and mood disorders that occur after infectious mononucleosis The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America Multidisciplinaire LCI-Richtlijn Q-koorts-vermoeidheidssyndroom (QVS) [Revision of guideline on Q fever fatigue syndrome (QFS)] How does post COVID differ from other post-viral conditions in childhood and adolescence (0-20 years old)? A systematic review Fatigue and symptom-based clusters in post COVID-19 patients: a multicentre, prospective, observational cohort study Acute sickness behaviour: an immune system-to-brain communication? Biological basis of the behavior of sick animals Cytokine-induced sickness behaviour: mechanisms and implications Long-term persistence of *Coxiella burnetii* in the host after

primary Q fever Q fever: persistence of antigenic non-viable cell residues of Coxiella burnetii in the host--implications for post Q fever infection fatigue syndrome and other chronic sequelae The natural history of acute Q fever: a prospective Australian cohort Prolonged illness after infectious mononucleosis is associated with altered immunity but not with increased viral load Measurement of circulating viral antigens post-SARS-CoV-2 infection in a multicohort study Persistence of spike protein at the skull-meninges-brain axis may contribute to the neurological sequelae of COVID-19 Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline Identifying the PECO: a framework for formulating good questions to explore the association of environmental and other exposures with health outcomes Covidence systematic review software Development and reliability assessment of a new quality appraisal tool for cross-sectional studies using biomarker data (BIOCROSS) Early biomarkers for prediction of severe manifestations of dengue fever: a systematic review and a meta-analysis Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range Tracking post-infectious fatigue in clinic using routine lab tests Predictors for Developing Severe Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Following Infectious Mononucleosis Risks for developing myalgic Encephalomyelitis/Chronic fatigue syndrome in college students following infectious mononucleosis: a prospective cohort Study Predictors of chronic fatigue in adolescents six months after acute Epstein-Barr virus infection: a prospective cohort study Atopy and elevation of IgE, IgG3, and IgG4 May be risk factors for post COVID-19 condition in children and adolescents Long-term symptom

severity and clinical biomarkers in post-COVID-19/chronic fatigue syndrome: results from a prospective observational cohort Inflammatory markers and pulmonary function in adolescents and young adults 6 months after mild COVID-19 Bulk RNA sequencing for analysis of post COVID-19 condition in adolescents and young adults CCL19 as a chemokine risk factor for posttreatment Lyme disease syndrome: a prospective clinical cohort Study Longitudinal transcriptome analysis reveals a sustained differential gene expression signature in patients treated for acute Lyme disease Serological and virological investigation of the role of the herpesviruses EBV, CMV and HHV-6 in post-infective fatigue syndrome Gene expression correlates of postinfective fatigue syndrome after infectious mononucleosis Serum cytokine levels in postinfective fatigue syndrome Serum of Post-COVID-19 syndrome patients with or without ME/CFS differentially affects endothelial cell function in vitro Peripheral blood gene expression in postinfective fatigue syndrome following from three different triggering infections Endothelial dysfunction and altered endothelial biomarkers in patients with post-COVID-19 syndrome and chronic fatigue syndrome (ME/CFS) Normal salivary cortisol and NK cell function in adolescents with chronic fatigue syndrome following infectious mononucleosis Clinical symptoms and markers of disease mechanisms in adolescent chronic fatigue following Epstein-Barr virus infection: an exploratory cross-sectional study Chronic fatigue syndrome: an emerging sequela in COVID-19 survivors? Preliminary evidence of mitochondrial dysfunction associated with post-infective fatigue after acute infection with Epstein Barr virus Postinfective fatigue syndrome is not associated with altered cytokine production Neurological involvement among non-hospitalized adolescents and young adults 6 months after

acute COVID-19 Metabolomic and immune alterations in long COVID patients with chronic fatigue syndrome Diverse immunological dysregulation, chronic inflammation, and impaired erythropoiesis in long COVID patients with chronic fatigue syndrome Determinants of persistent symptoms after treatment for Lyme borreliosis: a prospective observational cohort study Development of a fatigue scale The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection A clinical case definition of post-COVID-19 condition by a Delphi consensus Researching COVID to Enhance Recovery (RECOVER) adult study protocol: rationale, objectives, and design Post-exertional malaise in long COVID: subjective reporting versus objective assessment Immunological fingerprint in coronavirus disease-19 convalescents with and without post-COVID syndrome Distinguishing features of long COVID identified through immune profiling Post-acute sequelae of COVID-19 is characterized by diminished peripheral CD8(+)beta7 integrin(+) T cells and anti-SARS-CoV-2 IgA response Improvement of immune dysregulation in individuals with long COVID at 24-months following SARS-CoV-2 infection Biomarkers in long COVID-19: a systematic review Neuroinflammation in post-acute sequelae of COVID-19 (PASC) as assessed by [(11)C]PBR28 PET correlates with vascular disease measures Muscle abnormalities worsen after post-exertional malaise in long COVID A review of cytokine-based pathophysiology of long COVID symptoms The kynurenine pathway relates to post-acute COVID-19 objective cognitive impairment and PASC Prolonged indoleamine 2,3-dioxygenase-2 activity and associated cellular stress in post-acute sequelae of SARS-CoV-2 infection Complement dysregulation is a prevalent and therapeutically amenable feature of long COVID Persistent complement dysregulation

with signs of thromboinflammation in active Long Covid A research agenda for post-COVID-19 fatigue Multi-omics examination of Q fever fatigue syndrome identifies similarities with chronic fatigue syndrome eBioMedicine. 2025. [PMC12596658](#)

123. Elisa Gentilotti, Carolina Alvarez Garavito, Anna Górska, Roy Gusinow, Lorenzo Maria Canziani, Pasquale De Nardo, Alessandro Visentin, Maria Giulia Caponcello, Michela Di Chiara, Aline-Marie Florence, Gerolf de Boer, Salvatore Cataudella, Gabriel Levy Hara, Adriana Tami, Maddalena Giannella, Cédric Laouénan, Jan Hasenauer, Jesús Rodríguez-Baño, Evelina Tacconelli, et al.. Persistent Low-Grade Inflammation and Post-COVID Condition: Evidence from the ORCHESTRA Cohort Post-COVID Syndrome: An Insight on Its Pathogenesis Long COVID or Post-acute Sequelae of COVID-19 (PASC): An Overview of Biological Factors That May Contribute to Persistent Symptoms A Clinical Case Definition of Post COVID-19 Condition by a Delphi Consensus National Academies of Sciences, Engineering, and Medicine Committee on Examining the Working Definition for Long Covid. Long Covid Defined Post-acute COVID-19 syndrome (PCS) and health-related quality of life (HRQoL)—A systematic review and meta-analysis How methodological pitfalls have created widespread misunderstanding about long COVID Chemosensory assessment and impact on quality of life in neurosensorial cluster of the post COVID 19 syndrome Persistent capillary rarefaction in long COVID syndrome Longitudinal Lung Function Assessment of Patients Hospitalized with COVID-19 Using 1H and 129Xe Lung MRI Laboratory abnormalities in patients with COVID-2019 infection The critical role of laboratory medicine during coronavirus disease 2019 (COVID-19) and other viral outbreaks Blood lactate concentration in COVID-19: A

systematic literature review Procalcitonin as a predictive marker in COVID-19: A systematic review and meta-analysis Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China Serum ferritin as a predictive biomarker in COVID-19. A systematic review, meta-analysis and meta-regression analysis A review of cytokine-based pathophysiology of Long COVID symptoms Biomarker-Based Risk Assessment Strategy for Long COVID: Leveraging Spike Protein and Proinflammatory Mediators to Inform Broader Postinfection Sequelae Sustained vascular inflammatory effects of SARS-CoV-2 spike protein on human endothelial cells Inflammatory profiles are associated with long COVID up to 6 months after COVID-19 onset: A prospective cohort study of individuals with mild to critical COVID-19 Post-COVID-19 syndrome, low-grade inflammation and inflammatory markers: A cross-sectional study Clinical phenotypes and quality of life to define post-COVID-19 syndrome: A cluster analysis of the multinational, prospective ORCHESTRA cohort The RAND 36-Item Health Survey 1.0 SF-36 health survey update Follow-Up of COVID-19 Long Term Sequelae Study Protocol. Prospective Multinational Cohort Study on Characteristics and Determinants of Post-COVID-19 Long-Term Sequelae Challenges of data sharing in European COVID-19 projects: A learning opportunity for advancing pandemic preparedness and response Harmonization and standardization of data for a pan-European cohort on SARS-CoV-2 pandemic SciPy 1.0: Fundamental algorithms for scientific computing in Python When and how should multiple imputation be used for handling missing data in randomised clinical trials—a practical guide with flowcharts Variable selection for generalized linear mixed models by L1-penalized estimation Comparison of Intracranial Pressure Measurements Before and After Hypertonic Saline or

Mannitol Treatment in Children with Severe Traumatic Brain Injury
Molecular states during acute COVID-19 reveal distinct etiologies of long-term sequelae
Regularization Paths for Generalized Linear Models via Coordinate Descent
A Minimal Common Outcome Measure Set for COVID-19 Clinical Research
Biomarkers in long COVID-19: A systematic review
Inflammatory and vascular biomarkers in post-COVID-19 syndrome: A systematic review and meta-analysis of over 20 biomarkers
Laboratory Findings and Biomarkers in Long COVID: What Do We Know So Far? Insights into Epidemiology, Pathogenesis, Therapeutic Perspectives and Challenges
Differentiation of Prior SARS-CoV-2 Infection and Postacute Sequelae by Standard Clinical Laboratory Measurements in the RECOVER Cohort
Evaluation of the Hematological Patterns from Up to 985 Days of Long COVID: A Cross-Sectional Study
An umbrella review of meta-analyses on diagnostic accuracy of C-reactive protein
Exploring the Pathophysiology of Long COVID: The Central Role of Low-Grade Inflammation and Multisystem Involvement
C Reactive Protein
Chronic inflammation in the etiology of disease across the life span
Low-grade inflammation is negatively associated with physical Health-Related Quality of Life in healthy individuals: Results from The Danish Blood Donor Study (DBDS)
Chronic inflammatory systemic diseases: An evolutionary trade-off between acutely beneficial but chronically harmful programs
Persistent Autoimmune Activation and Proinflammatory State in Post-Coronavirus Disease 2019 Syndrome
Problems of Pathogenesis and Pathogenetic Therapy of COVID-19 from the Perspective of the General Theory of Pathological Systems (General Pathological Processes)
Handling missing data in clinical research
Biomedicines. 2025. [PMC12839268](https://pubmed.ncbi.nlm.nih.gov/4012839268/)

124. Jonika Tannous, Alan P Pan, Thomas Potter, Abdulaziz T Bako, Katharine Dlouhy, Ashley Drews, Henry Dirk Sostman, Farhaan S Vahidy, et al.. Real-world effectiveness of COVID-19 vaccines and anti-SARS-CoV-2 monoclonal antibodies against postacute sequelae of SARS-CoV-2: analysis of a COVID-19 observational registry for a diverse US metropolitan population COVID-19 map. Johns Hopkins Coronavirus Resource Center SARS-cov-2 pathogenesis COVID-19 treatments and medications. Centers for Disease Control and Prevention Scientists set out to connect the dots on long COVID Do vaccines protect against long COVID? what the data say Rapid response to drive COVID-19 research in a learning health care system: rationale and design of the Houston Methodist COVID-19 surveillance and outcomes registry (CURATOR) (Preprint) Monoclonal antibody therapy Long-term symptoms among adults tested for SARS-cov-2 - united states, january 2020-april 2021 A new method of classifying prognostic comorbidity in longitudinal studies: development and validation A minimal common outcome measure set for COVID-19 clinical research Neighborhood socioeconomic disadvantage is associated with multimorbidity in a geographically-defined community Area deprivation and widening inequalities in US mortality, 1969-1998 Making neighborhood-disadvantage metrics accessible - the neighborhood atlas TriNetX Understanding long COVID: a modern medical challenge Incidence, co-occurrence, and evolution of long-COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19 Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments Short-term and long-term rates of postacute sequelae of SARS-cov-2 infection: a systematic review Attributes and predictors of long COVID Risk factors and disease profile of post-vaccination SARS-cov-2 infection

in UK users of the COVID symptom study APP: a prospective, community-based, nested, case-control study Efficacy of COVID-19 vaccination on the symptoms of patients with long COVID: a target trial emulation using data from the compare e-cohort in France Markers of immune activation and inflammation in individuals with post-acute sequelae of SARS-cov-2 infection Diverse functional autoantibodies in patients with COVID-19 Neuro-COVID long-haulers exhibit broad dysfunction in T cell memory generation and responses to vaccination Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation (PHOSP-COVID): a UK multicentre, prospective cohort study Sex differences in immune responses BMJ Open. 2023. [PMC10083521](#)

125. Melsew Setegn Alie, Getayeneh Antehunegn Tesema, Gossa Fetene Abebe, Desalegn Girma. The prolonged health sequelae “of the COVID-19 pandemic” in sub-Saharan Africa: a systematic review and meta-analysis How and why patients made long COVID Why the patient-made term 'long COVID' IS needed COVID-19 symptoms: longitudinal evolution and persistence in outpatient settings Long COVID in the Faroe Islands: a longitudinal study among nonhospitalized patients Confronting the pathophysiology of long COVID Outcomes of cardiovascular magnetic resonance imaging in patients recently recovered from coronavirus disease 2019 (COVID-19) COVID symptoms, symptom clusters, and predictors for becoming a long-hauler looking for clarity in the haze of the pandemic Multiorgan impairment in low-risk individuals with post-COVID-19 syndrome: a prospective, community-based study Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Persistent symptoms after COVID-19: qualitative study of 114 “long Covid” patients and draft quality principles for services Finding the 'right' GP: a qualitative study of the experiences of

people with long-COVID Impact of the burden of COVID-19 in Italy: results of disability-adjusted life years (DALYs) and productivity loss Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine Challenges in defining long COVID: striking differences across literature, electronic health records, and patient-reported information Post-COVID-19 condition 3 months after hospitalisation with SARS-CoV-2 in South Africa: a prospective cohort study The PRISMA 2020 statement: an updated guideline for reporting systematic reviews Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement Undue reliance on I^2 in assessing heterogeneity may mislead Assessing heterogeneity in meta-analysis: Q statistic or I^2 index? Post-discharge symptoms among hospitalized covid-19 patients in Nigeria: a single-center study A cohort study of post-COVID-19 condition across the Beta, Delta, and omicron waves in South Africa: 6-month follow-up of hospitalized and nonhospitalized participants Assessing the quality of studies in meta-research: review/guidelines on the most important quality assessment tools Incidence of long COVID-19 in people with previous SARS-Cov2 infection: a systematic review and meta-analysis of 120,970 patients Incidence and burden of long COVID in Africa: a systematic review and meta-analysis Long-COVID in children and adolescents: a systematic review and meta-analyses A systematic review and meta-analysis of long COVID symptoms Mid and long-term neurological and neuropsychiatric manifestations of post-COVID-19 syndrome: a meta-analysis Trends in COVID-19 admissions and deaths among people living with HIV in South Africa: analysis of national surveillance data Epidemiology of severe COVID-19 from South Africa Methods for early characterisation of the severity and dynamics of SARS-CoV-2 variants: a population-based time series analysis in South

Africa Characteristics and predictors of acute and chronic post-COVID syndrome: a systematic review and meta-analysis
The neurological manifestations of post-acute sequelae of SARS-CoV-2 infection
COVID-19: dealing with a potential risk factor for chronic neurological disorders
Neurological manifestations and neuro-invasive mechanisms of the severe acute respiratory syndrome coronavirus type 2
Neuroinvasion, neurotropic, and neuroinflammatory events of SARS-CoV-2: understanding the neurological manifestations in COVID-19 patients
Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic
Patient outcomes after hospitalisation with COVID-19 and implications for follow-up: results from a prospective UK cohort
COVID-19 after effects: concerns for singers
Prevalence of post-COVID-19 symptoms in hospitalized and non-hospitalized COVID-19 survivors: a systematic review and meta-analysis
Prevalence of potential respiratory symptoms in survivors of hospital admission after coronavirus disease 2019 (COVID-19): a systematic review and meta-analysis
Persistence of long COVID symptoms in COVID-19 survivors worldwide and its potential pathogenesis- a systematic review and meta-analysis
Cardiovascular sequelae in uncomplicated COVID-19 survivors
COVID-19 cardiac injury: implications for long-term surveillance and outcomes in survivors
Long-term sequelae of COVID-19: a systematic review and meta-analysis of one-year follow-up studies on post-COVID symptoms
Long-term neurological sequelae among severe COVID-19 patients: a systematic review and meta-analysis
Pooled prevalence of long COVID-19 symptoms at 12 months and above follow-up period: a systematic review and meta-analysis
Long-term consequences of COVID-19 at 6 months and above: a systematic review and

meta-analysis Mental morbidities and chronic fatigue in severe acute respiratory syndrome survivors: long-term follow-up Post-viral fatigue and COVID-19: lessons from past epidemics Clinical sequelae of the novel coronavirus: does COVID-19 infection predispose patients to cancer? Attributes and predictors of long COVID Determinants of post-acute COVID-19 syndrome among hospitalized severe COVID-19 patients: a 2-year follow-up study Predictive models of long COVID Risk predictors and symptom features of long COVID within a broad primary care patient population including both tested and untested patients Risk factors associated with post- COVID-19 condition: a systematic review and meta-analysis. JAMA Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Long COVID: long-term effects? The isolated effect of age on the risk of COVID-19 severe outcomes: a systematic review with meta-analysis Characteristics and predictors of long COVID among diagnosed cases of COVID-19 Clinical characteristics and factors associated with long COVID in Zambia, august 2020 to January 2023 Age, sex and previous comorbidities as risk factors not associated with SARS-CoV-2 infection for long COVID-19: a systematic review and meta-analysis Long COVID in hospitalized COVID-19 patients: a retrospective cohort study Ventilator-associated lung injury: a search for better therapeutic targets Prediction of the development of pulmonary fibrosis using serial thin-section CT and clinical features in patients discharged after treatment for COVID-19 pneumonia *Frontiers in Public Health*. 2025. [PMC11803863](#)

126. Valdez Jaures Njio Heugno, Odette Linda Kamdem, Estelle Geraldine Essangui Same, Elysée Claude Bika Lele, Yves Mbama Biloa, Clarisse Ayina Ayina, Jessica Guyot, Bienvenu Bongue, Samuel Honore Mandengue, Carole Else Eboumbou Moukoko, et al.. Factors associated with long COVID in sub-

Saharan Africa: a scoping review WHO Declares COVID-19 a Pandemic Coronavirus disease 2019 (COVID-19): current status and future perspectives 2019 novel coronavirus (2019-nCoV) outbreak: A new challenge A pneumonia outbreak associated with a new coronavirus of probable bat origin An in-depth statistical analysis of the COVID-19 pandemic's initial spread in the WHO African region Relatively low mortality rate of COVID-19 in Cameroon: what if age was the main factor? Symptoms and Functional Impairment Assessed 8 Months After Mild COVID-19 Among Health Care Workers Post-acute COVID-19 syndrome A cohort study of post-COVID-19 condition across the Beta, Delta, and Omicron waves in South Africa: 6-month follow-up of hospitalized and nonhospitalized participants High-dimensional characterization of post-acute sequelae of COVID-19 A clinical case definition of post-COVID-19 condition by a Delphi consensus The persistence of SARS-CoV-2 in tissues and its association with long COVID symptoms: a cross-sectional cohort study in China Why we need a deeper understanding of the pathophysiology of long COVID COVID-19 pandemic: The African paradox A review of co-morbidity between infectious and chronic disease in Sub Saharan Africa: TB and Diabetes Mellitus, HIV and Metabolic Syndrome, and the impact of globalization Burden of non-communicable diseases in sub-Saharan Africa, 1990-2017: results from the Global Burden of Disease Study 2017 Ambient air pollution and health in Sub-Saharan Africa: Current evidence, perspectives and a call to action Impact of poor disease surveillance system on COVID-19 response in africa: Time to rethink and rebuilt Coronavirus Disease 2019 Diagnostics: Key to Africa's Recovery Effets du changement climatique sur la pauvreté et les inégalités en Afrique subsaharienne Impact of Climate Change on Poverty and Inequality in Sub-Saharan Africa The

Demography of Sub-Saharan Africa in the 21st Century. Transformations Since 2000, Outlook to 2050 Connaissances et perceptions de la maladie à coronavirus (COVID-19) dans la commune de Ouagadougou The Pandemic City: Urban Issues in the Time of COVID-19 Croyances, perceptions et attitudes des populations face à la pandémie du Covid-19 en Afrique de l'Ouest: une étude qualitative à Abidjan, Côte d'Ivoire Scoping studies: towards a methodological framework Scoping studies: advancing the methodology Qualitative research synthesis: methodological guidance for systematic reviewers utilizing meta-aggregation The PRISMA 2020 statement: An updated guideline for reporting systematic reviews Clinical presentation and persistent symptoms in patients at a post-COVID-19 clinic in Ghana Two months follow-up of patients with non-critical COVID-19 in Cape Town, South Africa Determinants of post-acute COVID-19 syndrome among hospitalized severe COVID-19 patients: A 2-year follow-up study Clinical characteristics and factors associated with long COVID among post-acute COVID-19 clinic patients in Zambia, August 2020 to January 2023: A cross-sectional and longitudinal study design Post-COVID-19 pulmonary complications among recovered COVID-19 patients: a cross-sectional study from Addis Ababa, Ethiopia Post-COVID-19 condition 3 months after hospitalisation with SARS-CoV-2 in South Africa: a prospective cohort study Persistent Symptoms among Frontline Health Workers Post-Acute COVID-19 Infection Proteomics of fibrin amyloid microclots in long COVID/post-acute sequelae of COVID-19 (PASC) shows many entrapped pro-inflammatory molecules that may also contribute to a failed fibrinolytic system Prevalence of symptoms, comorbidities, fibrin amyloid microclots and platelet pathology in individuals with Long COVID/Post-Acute

Sequelae of COVID-19 (PASC) Two-month follow-up of persons with SARS-CoV-2 infection-Zambia, September 2020: a cohort study Post-Discharge Symptoms among Hospitalized COVID-19 Patients in Nigeria: A Single-Center Study « Long COVID »: persistent COVID-19 symptoms in survivors managed in Lagos State, Nigeria Excessive eating and weight gain:A rare post-acute COVID-19 syndrome Determinants of COVID-19-Related Length of Hospital Stays and Long COVID in Ghana: A Cross-Sectional Analysis Age, Sex and Previous Comorbidities as Risk Factors Not Associated with SARS-CoV-2 Infection for Long COVID-19: A Systematic Review and Meta-Analysis Prognostic Factors for Post-COVID-19 Syndrome: A Systematic Review and Meta-Analysis Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Sex differences in immune responses that underlie COVID-19 disease outcomes Incidence and risk factors of post COVID-19 syndrome: a Tunisian cohort study Incidence of New-Onset Hypertension Post-COVID-19: Comparison With Influenza Risk of long COVID and associated symptoms after acute SARS-COV-2 infection in ethnic minorities: A nationwide register-linked cohort study in Denmark Risk factors and clinical presentations of long COVID in Africa: A scoping review [The impact of carbohydrate metabolism disorders on the early and long-term clinical outcomes of patients with COVID-19 according to the AKTIV and AKTIV 2 registries] COVID-19 and Cardiovascular diseases. Scoping review study Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province Multidisciplinary Management Strategies for Long COVID: A Narrative Review Rehabilitation Research Framework for Patients With COVID-19 Defined by Cochrane Rehabilitation and the World Health Organization Rehabilitation Programme SPACO+: a mixed methods

protocol to assessing the effectiveness of an educative intervention in patients with Long Covid Immunological Interactions between Intestinal Helminth Infections and Tuberculosis Abstract 15637: Detection and Characterization of Fibrin/Amyloid Microclots in Patients With Post-Acute Sequelae of Covid-19 BMC Infectious Diseases. 2026. [PMC13067598](#)

127. John P. Powers, Tomas J. McIntee, Abhishek Bhatia, Charisse R. Madlock-Brown, Jaime Seltzer, Anisha Sekar, Nita Jain, Mady Hornig, Elle Seibert, Peter J. Leese, Melissa Haendel, Richard Moffitt, Emily R. Pfaff, Christopher G. Chute, Christopher G. Chute, et al.. Identifying commonalities and differences between EHR representations of PASC and ME/CFS in the RECOVER EHR cohort A comparative review of systemic and neurological symptomatology in 12 outbreaks collectively described as chronic fatigue syndrome, epidemic neuromyasthenia, and myalgic encephalomyelitis Unexplained post-acute infection syndromes ME/CFS and Long COVID share similar symptoms and biological abnormalities: road map to the literature Epidemiology, clinical presentation, pathophysiology, and management of long COVID: an update ME/CFS and Post-Exertional Malaise among Patients with Long COVID Long COVID: major findings, mechanisms and recommendations Chronic inflammation, neuroglial dysfunction, and plasmatogen deficiency as a new pathobiological hypothesis addressing the overlap between post-COVID-19 symptoms and myalgic encephalomyelitis/chronic fatigue syndrome Post-COVID syndrome: pulmonary complications COVID-19 Induced Postural Orthostatic Tachycardia Syndrome (POTS): A Review A narrative review on the validity of electronic health record-based research in epidemiology Generalisable long COVID subtypes: findings from the NIH N3C and RECOVER

programmes Coding long COVID: characterizing a new disease through an ICD-10 lens The National COVID Cohort Collaborative (N3C): Rationale, design, infrastructure, and deployment Correction to: Methods for identifying 30 chronic conditions: application to administrative data Combining billing codes, clinical notes, and medications from electronic health records provides superior phenotyping performance Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data Updating and validating the Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using data from 6 countries Identifying who has long COVID in the USA: a machine learning approach using N3C data De-black-boxing health AI: demonstrating reproducible machine learning computable phenotypes using the N3C-RECOVER Long COVID model in the All of Us data repository Index for rating diagnostic tests Fast unfolding of communities in large networks Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Myalgic encephalomyelitis: International Consensus Criteria Myalgic encephalomyelitis/chronic fatigue syndrome Estimating prevalence, demographics, and costs of ME/CFS using large scale medical claims data and machine learning Review of case definitions for myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) Establishing a consensus on ME/CFS exclusionary illnesses Association between fatigue, peripheral serotonin, and L-carnitine in hypothyroidism and in chronic fatigue syndrome Higher prevalence of “low T3 syndrome” in patients with chronic fatigue syndrome: A case-control study Myalgic encephalomyelitis/chronic fatigue syndrome: essentials of diagnosis and management Serum 25-hydroxy vitamin D levels in chronic fatigue syndrome: a retrospective survey Chronic fatigue syndrome (CFS): Suggestions for a

nutritional treatment in the therapeutic approach Chronic fatigue syndrome: a personalized integrative medicine approach Review of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: an evidence-based approach to diagnosis and management by clinicians Racial and Ethnic Disparities in the Quality of Health Care Healthcare utilization is a collider: an introduction to collider bias in EHR data reuse Differences in symptoms among Black and White patients with ME/CFS A systematic review of chronic fatigue, its syndromes and ethnicity: prevalence, severity, co-morbidity and coping Chronic fatigue syndrome: sociodemographic subtypes in a community-based sample: Sociodemographic subtypes in A community-based sample Racial/ethnic disparities in post-acute sequelae of SARS-CoV-2 infection in New York: An EHR-based cohort study from the RECOVER program Detecting orthostatic intolerance in Long COVID in a clinic setting Prevalence of orthostatic intolerance in long covid clinic patients and healthy volunteers: A multicenter study Communications Medicine. 2025. [PMC11986062](#)

128. Hengrui Liu, Zewen Xu, Ilayda Karsidag, Panpan Wang, Jielsing Weng. Immune cell communication networks and memory CD8+ T cell signatures sustaining chronic inflammation in COVID-19 and Long COVID Correction: Immune cell communication networks and memory CD8+ T cell signatures sustaining chronic inflammation in COVID-19 and Long COVID Weekly epidemiological update on COVID-19-13 June 2023 Safety and efficacy of the BNT162b2 mRNA covid-19 vaccine COVID-19 vaccines vs variants, determining how much immunity is enough The evolution of SARS-coV-2 Disease and healthcare burden of COVID-19 in the United States Impact of sex and metabolic comorbidities on coronavirus disease 2019 (COVID-19) mortality risk across age groups: 66-646 inpatients across 613 U.S. Hospitals

COVID-19 in patients with pulmonary hypertension: A national prospective cohort study
COVID-19 as a trigger for type 1 diabetes
Comprehensive analysis of scRNA-Seq and bulk RNA-Seq reveals ubiquitin promotes pulmonary fibrosis in chronic pulmonary diseases
Breathing face down
Acute respiratory distress syndrome
The trinity of COVID-19: immunity, inflammation and intervention
Long COVID symptoms in exposed and infected children, adolescents and their parents one year after SARS-CoV-2 infection: A prospective observational cohort study
Temporal dynamics in viral shedding and transmissibility of COVID-19
Clinical and immunological features of severe and moderate coronavirus disease 2019
COVID-19: consider cytokine storm syndromes and immunosuppression
Single-cell landscape of immunological responses in patients with COVID-19
COVID-19 immune features revealed by a large-scale single-cell transcriptome atlas
Reduction and functional exhaustion of T cells in patients with coronavirus disease 2019 (COVID-19)
A single-cell atlas of the peripheral immune response in patients with severe COVID-19
Immune cell profiling of COVID-19 patients in the recovery stage by single-cell sequencing
CD8⁺ T cells contribute to survival in patients with COVID-19 and hematologic cancer
Machine learning applications in genetics and genomics
A unified approach to interpreting model predictions
From local explanations to global understanding with explainable AI for trees & Zeng, J. A review on deep learning techniques for the diagnosis of novel coronavirus (COVID-19)
Diverse functional autoantibodies in patients with COVID-19
Longitudinal analyses reveal immunological misfiring in severe COVID-19
Large-scale multi-omic analysis of COVID-19 severity
Integrated analysis of multimodal single-cell data
Current best practices in single-cell RNA-seq analysis: a tutorial

Comprehensive integration of single-cell data
Evaluation of efficiency prediction algorithms and development of ensemble model for CRISPR/Cas9 gRNA selection
Single-cell Mayo Map (scMayoMap): an easy-to-use tool for cell type annotation in single-cell RNA-sequencing data analysis
Pan-cancer single-cell landscape of tumor-infiltrating T cells
Fully-automated and ultra-fast cell-type identification using specific marker combinations from single-cell transcriptomic data
Single-cell transcriptomics of human T cells reveals tissue and activation signatures in health and disease
clusterProfiler: an R package for comparing biological themes among gene clusters
Inference and analysis of cell-cell communication using CellChat
Towards a systems understanding of MHC class I and MHC class II antigen presentation
MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data
Building predictive models in R using the caret package
Deep immune profiling of COVID-19 patients reveals distinct immunotypes with therapeutic implications
T cell responses in patients with COVID-19
Cytokine profiles in the detection of severe lung involvement in hospitalized patients with COVID-19: The IL-8/IL-32 axis
The differential immune responses to COVID-19 in peripheral and lung revealed by single-cell RNA sequencing
Complex immune dysregulation in COVID-19 patients with severe respiratory failure
Magnitude and dynamics of the T-cell response to SARS-coV-2 infection at both individual and population levels
Ribosomal protein S26 serves as a checkpoint of T-cell survival and homeostasis in a p53-dependent manner
Deficiency of ribosomal protein S26, which is mutated in a subset of patients with Diamond Blackfan anemia, impairs erythroid differentiation
Frontiers in Immunology. 2025.
[PMC12585964](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC12585964/)

129. Snehal M Pinto Pereira, Manjula D Nugawela, Natalia K Rojas, Roz Shafran, Kelsey McOwat, Ruth Simmons, Tamsin Ford, Isobel Heyman, Shamez N Ladhani, Emily Y Cheung, Lana Fox-Smith, Emma Dalrymple, Terence Stephenson, et al.. Post-COVID-19 condition at 6 months and COVID-19 vaccination in non-hospitalised children and young people WHO/Europe | Child and adolescent health - COVID-19 and Children Persistent symptoms following SARS-CoV-2 infection amongst children and young people: a meta-analysis of controlled and uncontrolled studies Post-acute COVID-19 outcomes in children with mild and asymptomatic disease Illness duration and symptom profile in symptomatic UK school-aged children tested for SARS-CoV-2 Risk factors for post-COVID-19 condition in previously hospitalised children using the ISARIC global follow-up protocol: a prospective cohort study Healthcare use in 700 000 children and adolescents for six months after covid-19: before and after register based cohort study Long COVID symptoms and duration in SARS-CoV-2 positive children - a nationwide cohort study Long COVID symptoms in SARS-CoV-2-positive adolescents and matched controls (LongCOVIDKidsDK): a national, cross-sectional study Vaccinations in the UK | coronavirus in the UK Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID symptom study APP: a prospective, community-based, nested, case-control study Trajectory of long covid symptoms after covid-19 vaccination: community based cohort study Long COVID and the mental and physical health of children and young people: national matched cohort study protocol (the clock study) ISARIC global COVID-19 paediatric follow-up Development of the EQ-5D-Y: a child-friendly version of the EQ-5D Psychometric properties of the strengths and difficulties questionnaire The Warwick-Edinburgh mental

well-being scale (WEMWBS): development and UK validation
 Development of a fatigue scale
 Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK statistical bulletins
 Physical and mental health 3 months after SARS-CoV-2 infection (long COVID) among adolescents in England (clock): a national matched cohort study
 Long COVID (post-COVID-19 condition) in children: a modified Delphi process
 Developing a model for predicting impairing physical symptoms in children 3 months after a SARS-CoV-2 PCR-test: the clock study
 Strengthening the reporting of observational studies in epidemiology (STROBE): explanation and elaboration
 Difficult questions about long COVID in children
 Archives of Disease in Childhood. 2023. [PMC10086284](#)

130. Yong-Bo Zheng, Na Zeng, Kai Yuan, Shan-Shan Tian, Ying-Bo Yang, Nan Gao, Xuan Chen, An-Yi Zhang, Alexandra L. Kondratiuk, Pei-Pei Shi, Fang Zhang, Jie Sun, Jing-Li Yue, Xiao Lin, Le Shi, Ajit Lalvani, Jie Shi, Yan-Ping Bao, Lin Lu, et al..
 Prevalence and risk factor for long COVID in children and adolescents: A meta-analysis and systematic review
 The lasting misery of coronavirus long-haulers
 A clinical case definition of post-COVID-19 condition by a Delphi consensus
 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study
 The phenotype and prediction of long-term physical, mental and cognitive COVID-19 sequelae
 20 months after recovery, a community-based cohort study in China
 Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: a systematic review
 More than 50 long-term effects of COVID-19: a systematic review and meta-analysis
 A systematic review and meta-analysis of long term physical and mental sequelae of COVID-19 pandemic: call for research priority and action
 Estimated global proportions of individuals with persistent

fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 A clinical case definition of post-COVID-19 condition by a Delphi consensus Mental and neurological disorders and risk of COVID-19 susceptibility, illness severity and mortality: a systematic review, meta-analysis and call for action Neurological issues in children with COVID-19 Epidemiology of COVID-19 infection in young children under five years: a systematic review and meta-analysis 'Long COVID': Symptom persistence in children hospitalised for COVID-19 6-month multidisciplinary follow-up and outcomes of patients with paediatric inflammatory multisystem syndrome (PIMS-TS) at a UK tertiary paediatric hospital: a retrospective cohort study Physical and mental health 3 months after SARS-CoV-2 infection (long COVID) among adolescents in England (CLOcK): a national matched cohort study Clinical characteristics of 58 children with a pediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 Post-COVID-19 conditions among children 90 days after SARS-CoV-2 infection Risk factors for long covid in previously hospitalised children using the ISARIC Global follow-up protocol: a prospective cohort study Comparison of long-term complications of COVID-19 illness among a diverse sample of children by MIS-C status Persistent symptoms following SARS-CoV-2 infection amongst children and young people: a meta-analysis of controlled and uncontrolled studies Long-COVID in children and adolescents: a systematic review and meta-analyses Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement Long COVID in children: observations from a designated pediatric clinic Pediatric long-

COVID: an overlooked phenomenon? Clinical characteristics, activity levels and mental health problems in children with long coronavirus disease: a survey of 510 children Intriguing new faces of Covid-19: persisting clinical symptoms and cardiac effects in children Pulmonary function and long-term respiratory symptoms in children and adolescents after COVID-19 Clinical characteristics of paediatric COVID-19 patients followed for up to 13 months Long-term symptoms after SARS-CoV-2 infection in children and adolescents Smell status in children infected with SARS-CoV-2 Post-acute COVID-19 outcomes in children with mild and asymptomatic disease Persistent symptoms in Swedish children after hospitalisation due to COVID-19 Mental Health in Children in the Context of COVID-19: Focus on Discharged Children Loneliness, internalizing symptoms, and inflammatory markers in adolescent COVID-19 survivors Cardiac magnetic resonance follow-up of children after pediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 with initial cardiac involvement Comparison of mental health outcomes in seropositive and seronegative adolescents during the COVID19 pandemic Long COVID in a prospective cohort of home-isolated patients Pulmonary function and persistent clinical symptoms in children and their parents 12 months after mild SARS-CoV-2 infection Post-COVID condition in adults and children living in the same household in Italy: a prospective cohort study using the ISARIC global follow-up protocol The prevalence, characteristics and risk factors of persistent symptoms in non-hospitalized and hospitalized children with SARS-CoV-2 infection followed-up for up to 12 months: a prospective, cohort study in Rome, Italy Sequelae of COVID-19 in hospitalized children: a 4-months follow-up The respiratory consequences of COVID-19 lasted for a median of 4 months in a cohort of children aged 2–18 years of

age Patients diagnosed with multisystem inflammatory syndrome in children have persistent neurologic, sleep, and psychiatric symptoms after hospitalization Persistent cough and asthma-like symptoms post COVID-19 hospitalization in children Persistent symptoms and decreased health-related quality of life after symptomatic pediatric COVID-19: a prospective study in a Latin American tertiary hospital Extended coagulation profile of children with Long Covid: a prospective study Neurological and psychological effects of long COVID in a young population: a cross-sectional study Pulmonary dysfunction after pediatric COVID-19 Long-term persistence of anti-sars-cov-2 antibodies in a pediatric population Multisystem inflammatory syndrome in children: salient echocardiogram findings in the acute phase and longitudinal follow-up A non-randomized comparative study of olfactory and gustatory functions in children who recovered from COVID-19 (1-year follow-up) Predictive model for long COVID in children 3 months after a SARS-CoV-2 PCR test Pulmonary function tests in the follow-up of children with COVID-19 Long-term pulmonary sequelae in adolescents post-SARS-CoV-2 infection Short-term outcomes in children recovered from multisystem inflammatory syndrome associated with SARS-CoV-2 infection Prevalence and risk factors of post-COVID-19 condition in adults and children at 6 and 12 months after hospital discharge: a prospective, cohort study in Moscow (StopCOVID) Multisystem inflammatory syndrome in children: clinical presentation, management, and short- and long-term outcomes Evolution of echocardiographic and cardiac magnetic resonance imaging abnormalities during follow-up in patients with multisystem inflammatory syndrome in children 6-month multidisciplinary follow-up and outcomes of patients with paediatric inflammatory multisystem syndrome (PIMS-TS) at a UK

tertiary paediatric hospital: a retrospective cohort study
2019-nCoV epidemic: address mental health care to empower
society Prevalence of and risk factors associated with mental
health symptoms among the general population in China
during the coronavirus disease 2019 pandemic The impact of
quarantine on mental health status among general population
in China during the COVID-19 pandemic Mental health status
of late-middle-aged adults in China during the coronavirus
disease 2019 pandemic A systematic review and meta-
analysis on prevalence of and risk factors associated with
depression, anxiety and insomnia in infectious diseases,
including COVID-19: a call to action Mental health
considerations for children quarantined because of COVID-19
Prevalence of posttraumatic stress disorder after infectious
disease pandemics in the twenty-first century, including
COVID-19: a meta-analysis and systematic review
Multisystem inflammatory syndrome in U.S. children and
adolescents. Persistent symptoms following SARS-CoV-2
infection amongst children and young people: a meta-analysis
of controlled and uncontrolled studies Journal of Infection and
Public Health. 2023. [PMC9990879](#)

131. Rachel S. Gross, Tanayott Thaweethai, Lawrence C. Kleinman, Jessica N. Snowden, Erika B. Rosenzweig, Joshua D. Milner, Kelan G. Tantisira, Kyung E. Rhee, Terry L. Jernigan, Patricia A. Kinser, Amy L. Salisbury, David Warburton, Sindhu Mohandas, John C. Wood, Jane W. Newburger, Dongngan T. Truong, Valerie J. Flaherman, Torri D. Metz, Elizabeth W. Karlson, Lori B. Chibnik, Deepti B. Pant, Aparna Krishnamoorthy, Richard Gallagher, Michelle F. Lamendola-Essel, Denise C. Hasson, Stuart D. Katz, Shonna Yin, Benard P. Dreyer, Megan Carmilani, K. Coombs, Megan L. Fitzgerald, Nick Güthe, Mady Hornig, Rebecca J. Letts, Aimee K. Peddie, Brittany D. Taylor, Andrea S. Foulkes, Melissa S. Stockwell,

Venkataraman Balaraman, Amanda Bogie, Hulya Bukulmez, Allen J. Dozor, Daniel Eckrich, Amy J. Elliott, Danielle N. Evans, Jonathan S. Farkas, E. Vincent S. Faustino, Laura Fischer, Sunanda Gaur, Ashraf S. Harahsheh, Uzma N. Hasan, Daniel S. Hsia, Gredia Huerta-Montañez, Kathy D. Hummel, Matt P. Kadish, David C. Kaelber, Sankaran Krishnan, Jessica S. Kosut, Jerry Larrabee, Peter Paul C. Lim, Ian C. Michelow, Carlos R. Oliveira, Hengameh Raissy, Zaira Rosario-Pabon, Judith L. Ross, Alice I. Sato, Michelle D. Stevenson, Maria M. Talavera-Barber, Ronald J. Teufel, Kathryn E. Weakley, Emily Zimmerman, Marie-Abele C. Bind, James Chan, Zoe Guan, Richard E. Morse, Harrison T. Reeder, Natascha Akshoomoff, Judy L. Aschner, Rakesh Bhattacharjee, Lesley A. Cottrell, Kelly Cowan, Viren A. D'Sa, Alexander G. Fiks, Maria L. Gennaro, Katherine Irby, Manaswitha Khare, Jeremy Landeo Guttierrez, Russell J. McCulloh, Shalu Narang, Manette Ness-Cochinwala, Sheila Nolan, Paul Palumbo, Julie Ryu, Juan C. Salazar, Rangaraj Selvarangan, Cheryl R. Stein, Alan Werzberger, William T. Zempsky, Robin Aupperle, Fiona C. Baker, Marie T. Banich, Deanna M. Barch, Arielle Baskin-Sommers, James M. Bjork, Susan Y. Bookheimer, Sandra A. Brown, BJ Casey, Linda Chang, Duncan B. Clark, Anders M. Dale, Mirella Dapretto, Thomas M. Ernst, Damien A. Fair, Sarah W. Feldstein Ewing, John J. Foxe, Edward G. Freedman, Naomi P. Friedman, Hugh Garavan, Dylan G. Gee, Raul Gonzalez, Kevin M. Gray, Mary M. Heitzeg, Megan M. Herting, Joanna Jacobus, Angela R. Laird, Christine L. Larson, Krista M. Lisdahl, Monica Luciana, Beatriz Luna, Pamela A.F. Madden, Erin C. McGlade, Eva M. Müller-Oehring, Bonnie J. Nagel, Michael C. Neale, Martin P. Paulus, Alexandra S. Potter, Perry F. Renshaw, Elizabeth R. Sowell, Lindsay M. Squeglia, Susan Tapert, Lucina Q. Uddin, Sylia Wilson, Deborah A. Yurgelun-Todd, et al.. Characterizing Long COVID

in Children and Adolescents Uncovering Long COVID in Children. 2024 Update of the RECOVER-Adult Long COVID Research Index JAMA. 2024. [PMC11339705](#)

132. Rachel S. Gross, Tanayott Thaweethai, Erika B. Rosenzweig, James Chan, Lori B. Chibnik, Mine S. Cicek, Amy J. Elliott, Valerie J. Flaherman, Andrea S. Foulkes, Margot Gage Witvliet, Richard Gallagher, Maria Laura Gennaro, Terry L. Jernigan, Elizabeth W. Karlson, Stuart D. Katz, Patricia A. Kinser, Lawrence C. Kleinman, Michelle F. Lamendola-Essel, Joshua D. Milner, Sindhu Mohandas, Praveen C. Mudumbi, Jane W. Newburger, Kyung E. Rhee, Amy L. Salisbury, Jessica N. Snowden, Cheryl R. Stein, Melissa S. Stockwell, Kelan G. Tantisira, Moriah E. Thomason, Dongngan T. Truong, David Warburton, John C. Wood, Shifa Ahmed, Almary Akerlundh, Akram N. Alshawabkeh, Brett R. Anderson, Judy L. Aschner, Andrew M. Atz, Robin L. Aupperle, Fiona C. Baker, Venkataraman Balaraman, Dithi Banerjee, Deanna M. Barch, Arielle Baskin-Sommers, Sultana Bhuiyan, Marie-Abele C. Bind, Amanda L. Bogie, Tamara Bradford, Natalie C. Buchbinder, Elliott Bueler, Hülya Bükülmez, B. J. Casey, Linda Chang, Maryanne Chrisant, Duncan B. Clark, Rebecca G. Clifton, Katharine N. Clouser, Lesley Cottrell, Kelly Cowan, Viren D'Sa, Mirella Dapretto, Soham Dasgupta, Walter Dehority, Audrey Dionne, Kirsten B. Dummer, Matthew D. Elias, Shari Esquenazi-Karonika, Danielle N. Evans, E. Vincent S. Faustino, Alexander G. Fiks, Daniel Forsha, John J. Foxe, Naomi P. Friedman, Greta Fry, Sunanda Gaur, Dylan G. Gee, Kevin M. Gray, Stephanie Handler, Ashraf S. Harahsheh, Keren Hasbani, Andrew C. Heath, Camden Hebson, Mary M. Heitzeg, Christina M. Hester, Sophia Hill, Laura Hobart-Porter, Travis K. F. Hong, Carol R. Horowitz, Daniel S. Hsia, Matthew Huentelman, Kathy D. Hummel, Katherine Irby, Joanna Jacobus, Vanessa L. Jacoby, Pei-Ni Jone, David C.

Kaelber, Tyler J. Kasmarcak, Matthew J. Kluko, Jessica S. Kosut, Angela R. Laird, Jeremy Landeo-Gutierrez, Sean M. Lang, Christine L. Larson, Peter Paul C. Lim, Krista M. Lisdahl, Brian W. McCrindle, Russell J. McCulloh, Kimberly McHugh, Alan L. Mendelsohn, Torri D. Metz, Julie Miller, Elizabeth C. Mitchell, Lerraughn M. Morgan, Eva M. Müller-Oehring, Erica R. Nahin, Michael C. Neale, Manette Ness-Cochinwala, Sheila M. Nolan, Carlos R. Oliveira, Onyekachukwu Osakwe, Matthew E. Oster, R. Mark Payne, Michael A. Portman, Hengameh Raissy, Isabelle G. Randall, Suchitra Rao, Harrison T. Reeder, Johana M. Rosas, Mark W. Russell, Arash A. Sabati, Yamuna Sanil, Alice I. Sato, Michael S. Schechter, Rangaraj Selvarangan, S. Kristen Sexson Tejtelt, Divya Shakti, Kavita Sharma, Lindsay M. Squeglia, Shubika Srivastava, Michelle D. Stevenson, Jacqueline Szmuszkovicz, Maria M. Talavera-Barber, Ronald J. Teufel, Deepika Thacker, Felicia Trachtenberg, Mmekom M. Udosen, Megan R. Warner, Sara E. Watson, Alan Werzberger, Jordan C. Weyer, Marion J. Wood, H. Shonna Yin, William T. Zempsky, Emily Zimmerman, Benard P. Dreyer, Seyed Aria Nejadghaderi, Seyed Aria Nejadghaderi, et al.. Researching COVID to enhance recovery (RECOVER) pediatric study protocol: Rationale, objectives and design Researching COVID to enhance recovery (RECOVER) pediatric study protocol: Rationale, objectives and design. Researching COVID to enhance recovery (RECOVER) pediatric study protocol: Rationale, objectives and design. A clinical case definition of post-COVID-19 condition by a Delphi consensus Long COVID: major findings, mechanisms and recommendations How common is Long COVID in children and adolescents? Post-COVID-19 symptoms and conditions among children and adolescents—United States, March 1, 2020-January 31, 2022 Long COVID symptoms and duration in SARS-CoV-2 positive children—a

nationwide cohort study Severity of acute COVID-19 in children <18 years old March 2020 to December 2021 Long-COVID in children and adolescents: a systematic review and meta-analyses Epidemiological and clinical characteristics of non-severe and severe pediatric and adult COVID-19 patients across different geographical regions in the early phase of pandemic: a systematic review and meta-analysis of observational studies Post-COVID-19 conditions among children 90 days after SARS-CoV-2 infection Short-term and long-term rates of Postacute Sequelae of SARS-CoV-2 Infection: A systematic review Multisystem Inflammatory Syndrome in Children in New York State Physical and mental health 3 months after SARS-CoV-2 infection (long COVID) among adolescents in England (CLOCK): a national matched cohort study Prevalence of select new symptoms and conditions among persons aged younger than 20 years and 20 years or older at 31 to 150 days after testing positive or negative for SARS-CoV-2 Clinical characteristics, activity levels and mental health problems in children with long coronavirus disease: a survey of 510 children Long COVID symptoms in SARS-CoV-2-positive adolescents and matched controls (LongCOVIDKidsDK): a national, cross-sectional study Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Long COVID symptoms in exposed and infected children, adolescents and their parents one year after SARS-CoV-2 infection: A prospective observational cohort study Post-COVID-19-associated morbidity in children, adolescents, and adults: A matched cohort study including more than 157,000 individuals with COVID-19 in Germany Pediatric Long-COVID: A review of the definition, epidemiology, presentation, and pathophysiology Clinical features and burden of Postacute

Sequelae of SARS-CoV-2 Infection in children and adolescents
Long-term COVID 19 sequelae in adolescents: the overlap with orthostatic intolerance and ME/CFS
Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description
The conception of the ABCD study: From substance use to a broad NIH collaboration
The NHLBI Study on Long-term Outcomes after the Multisystem Inflammatory Syndrome In Children (MUSIC): Design and objectives
Approaches to improving health literacy: lessons from the field
Recruiting the ABCD sample: Design considerations and procedures
Measuring PROMIS®
Global Health in Early Childhood
Development of the PROMIS ® pediatric global health (PGH-7) measure
Development of physical and mental health summary scores from the patient-reported outcomes measurement information system (PROMIS) global items
Children's Somatization Inventory: psychometric properties of the revised form (CSI-24)
The "All of Us" Research Program
Identifying children with special health care needs: development and evaluation of a short screening instrument
An evaluation of the linguistic and cultural validity of the Spanish language version of the children with special health care needs screener
The development, preliminary validation, and clinical application of the quick parenting assessment
Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome
Postural tachycardia in children and adolescents: what is abnormal?
The Beighton Score as a measure of generalised joint hypermobility
Revision of a parent-completed development screening tool: Ages and Stages Questionnaires
Validation of the modified checklist for Autism in toddlers, revised with follow-up (M-CHAT-R/F)
Cognition assessment using the NIH Toolbox
Development of

six PROMIS pediatrics proxy-report item banks Psychometric Evaluation of the PROMIS® Pediatric Psychological and Physical Stress Experiences Measures Development and Evaluation of the PROMIS(®) Pediatric Positive Affect Item Bank, Child-Report and Parent-Proxy Editions Assessment of symptoms of DSM-IV anxiety and depression in children: a revised child anxiety and depression scale The Strengths and Difficulties Questionnaire: a research note Development of three web-based computerized versions of the Kiddie Schedule for Affective Disorders and Schizophrenia Child Psychiatric Diagnostic Interview: Preliminary Validity Data A modified poisson regression approach to prospective studies with binary data Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection Understanding pediatric long COVID using a tree-based scan statistic approach: an EHR-based cohort study from the RECOVER Program Algorithms for hierarchical clustering: an overview Consensus clustering: a resampling-based method for class discovery and visualization of gene expression microarray data A sharper Bonferroni procedure for multiple tests of significance Statistical power of multiplicity adjustment strategies for correlated binary endpoints Multiple imputation using chained equations: Issues and guidance for practice Correcting for noncompliance and dependent censoring in an AIDS Clinical Trial with inverse probability of censoring weighted (IPCW) log-rank tests Pattern-mixture models for multivariate incomplete data with covariates Decision Letter 0 Author response to Decision Letter 0 Decision Letter 1 PLOS ONE. 2024. [PMC11075869](https://doi.org/10.1371/journal.pone.0311075)

133. Rachel Gross, Tanayott Thaweethai, Erika B. Rosenzweig, James Chan, Lori B. Chibnik, Mine S. Cicek, Amy J. Elliott, Valerie J. Flaherman, Andrea S. Foulkes, Margot Gage Witvliet, Richard Gallagher, Maria Laura Gennaro, Terry L.

Jernigan, Elizabeth W. Karlson, Stuart D. Katz, Patricia A. Kinser, Lawrence C. Kleinman, Michelle F. Lamendola-Essel, Joshua D. Milner, Sindhu Mohandas, Praveen C. Mudumbi, Jane W. Newburger, Kyung E. Rhee, Amy L. Salisbury, Jessica N. Snowden, Cheryl R. Stein, Melissa S. Stockwell, Kelan G. Tantisira, Moriah E. Thomason, Dongngan T. Truong, David Warburton, John C. Wood, Shifa Ahmed, Almary Akerlundh, Akram N. Alshawabkeh, Brett R. Anderson, Judy L. Aschner, Andrew M. Atz, Robin L. Aupperle, Fiona C. Baker, Venkataraman Balaraman, Dithi Banerjee, Deanna M. Barch, Arielle Baskin-Sommers, Sultana Bhuiyan, Marie-Abele C. Bind, Amanda L. Bogie, Natalie C. Buchbinder, Elliott Bueler, Hülya Bükülmez, B.J. Casey, Linda Chang, Duncan B. Clark, Rebecca G. Clifton, Katharine N. Clouser, Lesley Cottrell, Kelly Cowan, Viren D'Sa, Mirella Dapretto, Soham Dasgupta, Walter Dehority, Kirsten B. Dummer, Matthew D. Elias, Shari Esquenazi-Karonika, Danielle N. Evans, E. Vincent S. Faustino, Alexander G. Fiks, Daniel Forsha, John J. Foxe, Naomi P. Friedman, Greta Fry, Sunanda Gaur, Dylan G. Gee, Kevin M. Gray, Ashraf S. Harahsheh, Andrew C. Heath, Mary M. Heitzeg, Christina M. Hester, Sophia Hill, Laura Hobart-Porter, Travis K.F. Hong, Carol R. Horowitz, Daniel S. Hsia, Matthew Huentelman, Kathy D. Hummel, William G. Iacono, Katherine Irby, Joanna Jacobus, Vanessa L. Jacoby, Pei-Ni Jone, David C. Kaelber, Tyler J. Kasmarcak, Matthew J. Kluko, Jessica S. Kosut, Angela R. Laird, Jeremy Landeo-Gutierrez, Sean M. Lang, Christine L. Larson, Peter Paul C. Lim, Krista M. Lisdahl, Brian W. McCrindle, Russell J. McCulloh, Alan L. Mendelsohn, Torri D. Metz, Lerraughn M. Morgan, Eva M. Müller-Oehring, Erica R. Nahin, Michael C. Neale, Manette Ness-Cochinwala, Sheila M. Nolan, Carlos R. Oliveira, Matthew E. Oster, R. Mark Payne, Hengameh Raissy, Isabelle G. Randall, Suchitra Rao, Harrison T. Reeder, Johana M.

Rosas, Mark W. Russell, Arash A. Sabati, Yamuna Sanil, Alice I. Sato, Michael S. Schechter, Rangaraj Selvarangan, Divya Shakti, Kavita Sharma, Lindsay M. Squeglia, Michelle D. Stevenson, Jacqueline Szmuszkovicz, Maria M. Talavera-Barber, Ronald J. Teufel, Deepika Thacker, Mmekom M. Udosen, Megan R. Warner, Sara E. Watson, Alan Werzberger, Jordan C. Weyer, Marion J. Wood, H. Shonna Yin, William T. Zempsky, Emily Zimmerman, Benard P. Dreyer, et al.. Researching COVID to enhance recovery (RECOVER) pediatric study protocol: Rationale, objectives and design Researching COVID to Enhance Recovery (RECOVER) adult study protocol: Rationale, objectives, and design Long COVID: major findings, mechanisms and recommendations. How common is Long COVID in children and adolescents? Post-COVID-19 symptoms and conditions among children and adolescents - United States, March 1, 2020-January 31, 2022. Long COVID symptoms and duration in SARS-CoV-2 positive children - a nationwide cohort study. Severity of acute COVID-19 in children <18 years old March 2020 to December 2021. Long-COVID in children and adolescents: a systematic review and meta-analyses. Epidemiological and clinical characteristics of non-severe and severe pediatric and adult COVID-19 patients across different geographical regions in the early phase of pandemic: a systematic review and meta-analysis of observational studies. Post-COVID-19 conditions among children 90 days after SARS-CoV-2 infection. Short-term and long-term rates of Postacute Sequelae of SARS-CoV-2 Infection: A systematic review. Multisystem Inflammatory Syndrome in Children in New York State. Physical and mental health 3 months after SARS-CoV-2 infection (long COVID) among adolescents in England (CLOcK): a national matched cohort study. Prevalence of select new symptoms and conditions among persons aged

younger than 20 years and 20 years or older at 31 to 150 days after testing positive or negative for SARS-CoV-2. Clinical characteristics, activity levels and mental health problems in children with long coronavirus disease: a survey of 510 children. Long COVID symptoms in SARS-CoV-2-positive adolescents and matched controls (LongCOVIDKidsDK): a national, cross-sectional study. Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021. Long COVID symptoms in exposed and infected children, adolescents and their parents one year after SARS-CoV-2 infection: A prospective observational cohort study. Post-COVID-19-associated morbidity in children, adolescents, and adults: A matched cohort study including more than 157,000 individuals with COVID-19 in Germany. Pediatric Long-COVID: A review of the definition, epidemiology, presentation, and pathophysiology. Clinical features and burden of Postacute Sequelae of SARS-CoV-2 Infection in children and adolescents. Long-term COVID 19 sequelae in adolescents: the overlap with orthostatic intolerance and ME/CFS. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. The conception of the ABCD study: From substance use to a broad NIH collaboration. The NHLBI Study on Long-term Outcomes after the Multisystem Inflammatory Syndrome In Children (MUSIC): Design and objectives. Approaches to improving health literacy: lessons from the field. Recruiting the ABCD sample: Design considerations and procedures. Measuring PROMIS® Global Health in Early Childhood. Development of the PROMIS ® pediatric global health (PGH-7) measure. Development of physical and mental health summary scores from the patient-reported outcomes measurement information

system (PROMIS) global items. Children's Somatization Inventory: psychometric properties of the revised form (CSI-24). The "All of Us" Research Program. Identifying children with special health care needs: development and evaluation of a short screening instrument. An evaluation of the linguistic and cultural validity of the Spanish language version of the children with special health care needs screener. The development, preliminary validation, and clinical application of the quick parenting assessment. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. Postural tachycardia in children and adolescents: what is abnormal? The Beighton Score as a measure of generalised joint hypermobility. Revision of a parent-completed development screening tool: Ages and Stages Questionnaires. Validation of the modified checklist for Autism in toddlers, revised with follow-up (M-CHAT-R/F). Cognition assessment using the NIH Toolbox. Development of six PROMIS pediatrics proxy-report item banks. Psychometric Evaluation of the PROMIS® Pediatric Psychological and Physical Stress Experiences Measures. Development and Evaluation of the PROMIS(®) Pediatric Positive Affect Item Bank, Child-Report and Parent-Proxy Editions. Assessment of symptoms of DSM-IV anxiety and depression in children: a revised child anxiety and depression scale. The Strengths and Difficulties Questionnaire: a research note. Beery VMI: The Beery-Buktenica Developmental test of visual-motor integration with supplemental developmental tests of visual perception and motor coordination: Stepping stones age norms from birth to age six. P. Algorithms for hierarchical clustering: an overview. Consensus clustering: a resampling-based method for class discovery and visualization of gene expression microarray data. medRxiv. 2023. [PMC10197716](https://doi.org/10.1101/2023.05.15.23281116)

134. Tanayott Thaweethai, Sarah E. Donohue, Jeffrey N. Martin, Mady Hornig, Jarrod M. Mosier, Daniel J. Shinnick, Hassan Ashktorab, Ornina Atieh, Andra Blomkalns, Hassan Brim, Yu Chen, Melissa M. Cortez, Nathan B. Erdmann, Valerie Flaherman, Paul Goepfert, Jason D. Goldman, Naomi M. Hamburg, Jenny E. Han, James R. Heath, Vanessa Jacoby, Sarah E. Jolley, J. Daniel Kelly, Sara W. Kelly, C. Kim, Jerry A. Krishnan, Rebecca Letts, Emily B. Levitan, Matthew E. Modes, Grace A. McComsey, Torri D. Metz, Janet M. Mullington, Igho Ofotokun, Megumi J. Okumura, Claudia Castillo Paredes, Thomas F. Patterson, Michael J. Peluso, Rebecca Reece, Zaki A. Sherif, Hyagriv N. Simhan, Christopher Simmons, Upinder Singh, Barbara S. Taylor, Brittany D. Taylor, Joel D. Trinity, Andrea B. Troxel, Paul J. Utz, Andrew J. Vasey, Elisheva Weinberger, Zanthia Wiley, Juan Wisnivesky, Lynn M. Yee, Leora Horwitz, Andrea S. Foulkes, Bruce D. Levy, et al.. Long COVID trajectories in the prospectively followed RECOVER-Adult US cohort Long COVID and significant activity limitation among adults, by age-United States, June 1-13, 2022, to June 7-19, 2023 True prevalence of long-COVID in a nationwide population cohort study Researching COVID to enhance recovery (RECOVER) adult study protocol: rationale, objectives, and design Development of a definition of postacute sequelae of SARS-CoV-2 infection Mechanisms of long COVID and the path toward therapeutics FlexMix version 2: finite mixtures with concomitant variables and varying and constant parameters Long-term prognosis at 1.5 years after infection with wild-type strain of SARS-CoV-2 and alpha, delta, as well as Omicron variants Health outcomes in people 2 years after surviving hospitalisation with COVID-19: a longitudinal cohort study Two- and 3-year outcomes in convalescent individuals with COVID-19: a prospective cohort study Trajectories of

health-related quality of life and their predictors in adult COVID-19 survivors: a longitudinal analysis of the Biobanque quebecoise de la COVID-19 (BQC-19) Examining the trajectory of health-related quality of life among coronavirus disease patients Postacute sequelae of SARS-CoV-2 infection in the pre-delta, delta, and Omicron eras Three-year outcomes of post-acute sequelae of COVID-19 The effect of non-random loss to follow-up on group mean estimates in a longitudinal study Multiple imputation of missing data for multilevel models: simulations and recommendations Strategies for multiple imputation in longitudinal studies A framework for multiple imputation in cluster analysis Nature Communications. 2025. [PMC12623977](#)

135. Jiunn Wang, Henry Goodfellow, Sarah Walker, Ann Blandford, Paul Pfeffer, John R Hurst, David Sunkersing, Katherine Bradbury, Chris Robson, William Henley, Manuel Gomes, et al.. Trajectories of functional limitations, health-related quality of life and societal costs in individuals with long COVID: a population-based longitudinal cohort study COVID-19 rapid guideline: managing the long-term effects of covid-19 In the Wake of the Pandemic: Preparing for Long COVID Policy Brief Characterizing long COVID in an international cohort: 7 months of symptoms and their impact Impact of Long COVID on productivity and informal caregiving Long covid-mechanisms, risk factors, and management Impact of fatigue as the primary determinant of functional limitations among patients with post-COVID-19 syndrome: a cross-sectional observational study Development, deployment and evaluation of digitally enabled, remote, supported rehabilitation for people with long COVID-19 (Living With COVID-19 Recovery): protocol for a mixed-methods study Long COVID Through a Public Health Lens: An Umbrella Review The Work and Social Adjustment Scale: a

simple measure of impairment in functioning Optimizing the clinical utility of four proposed criteria for a persistent and impairing grief disorder by emphasizing core, rather than associated symptoms Cancer-related fatigue: definitions and clinical subtypes Valuing health-related quality of life: An EQ-5D-5L value set for England Unit Costs of Health and Social Care 2020. Personal Social Services Research Unit, University of Kent, Canterbury National tariff Average hourly pay Social dysfunctioning after mild to moderate first-ever stroke at vocational age The work and social adjustment scale: reliability, sensitivity and value Systematic review of health state utility values in metastatic non-small cell lung cancer with a focus on previously treated patients Health state utility values by cancer stage: a systematic literature review Recovery and symptom trajectories up to two years after SARS-CoV-2 infection: population based, longitudinal cohort study Profiling post-COVID-19 condition across different variants of SARS-CoV-2: a prospective longitudinal study in unvaccinated wild-type, unvaccinated alpha-variant, and vaccinated delta-variant populations Determinants of the onset and prognosis of the post-COVID-19 condition: a 2-year prospective observational cohort study Post-acute sequelae of covid-19 six to 12 months after infection: population based study Course of post COVID-19 disease symptoms over time in the ComPaRe long COVID prospective e-cohort Returning to work and the impact of post COVID-19 condition: A systematic review Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records Post COVID-19 condition, work ability and occupational changes in a population-based cohort Long Term Impact of COVID-19 Infection on Sleep and Mental Health Long-term outcomes after NeuroCOVID: A 6-month follow-up study on 60 patients The impact of COVID-19 critical illness on new disability,

functional outcomes and return to work at 6 months: a prospective cohort study Living With Covid Recovery Survey [Conferencepresentation]. NHS England Long COVID Meeting. London, UK When and how should multiple imputation be used for handling missing data in randomised clinical trials - a practical guide with flowcharts Review Process File BMJ Open. 2024. [PMC11574431](#)

136. Clemence Servier, Raphael Porcher, Isabelle Pane, Philippe Ravaud, Viet-Thi Tran. Trajectories of the evolution of post-COVID-19 condition, up to two years after symptoms onset Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021 Post-COVID-19 symptoms 2 years after SARS-CoV-2 infection among hospitalized vs nonhospitalized patients Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records A clinical case definition of post-COVID-19 condition by a Delphi consensus Course of post-COVID-19 disease symptoms over time in the ComPaRe long COVID prospective e-cohort Lifelines Corona Research Initiative. Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study 1-year outcomes in hospital survivors with COVID-19: a longitudinal cohort study Characterising long COVID: a living systematic review Respiratory and psychophysical sequelae among patients with COVID-19 four months after hospital discharge High-dimensional characterization of post-acute sequelae of COVID-19 Six-month Follow-up Chest CT findings after Severe COVID-19 Pneumonia Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens: prospective cohort study Efficacy of first dose of Covid-19 vaccine versus no vaccination on symptoms of patients with long covid: target trial emulation based on

ComPaRe e-cohort Prolonged SARS-CoV-2 RNA shedding: not a rare phenomenon Pathological sequelae of long-haul COVID Long COVID: major findings, mechanisms and recommendations Association of self-reported COVID-19 infection and SARS-CoV-2 serology test results with persistent physical symptoms among French adults during the COVID-19 pandemic CoviCare Study Team. Prevalence of symptoms more than seven months after diagnosis of symptomatic COVID-19 in an outpatient setting Symptoms and risk factors for long COVID in non-hospitalized adults Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation (PHOSP-COVID): a UK multicentre, prospective cohort study COVID-19 symptoms: longitudinal evolution and persistence in outpatient settings The GRoLTS-checklist: guidelines for reporting on latent trajectory studies Collaborative Open platform E-cohorts for research acceleration in trials and epidemiology Evidence of previous SARS-CoV-2 infection in seronegative patients with long COVID Development and validation of the long covid symptom and impact tools, a set of patient-reported instruments constructed from patients' lived experience Living risk prediction algorithm (QCOVID) for risk of hospital admission and mortality from coronavirus 19 in adults: national derivation and validation cohort study Estimation of Extended Mixed Models Using Latent Classes and Latent Processes: the R package lcmm Attributes and predictors of long COVID A patient-centric characterization of systemic recovery from SARS-CoV-2 infection Unexplained post-acute infection syndromes Post-acute sequelae after SARS-CoV-2 infection by viral variant and vaccination status: a multicenter cross-sectional study International Journal of Infectious Diseases. 2023. [PMC10176960](https://pubmed.ncbi.nlm.nih.gov/40176960/)