A Report On The Calculation Of The Excited States Of Z_T_azobenzene At The DFT/def2-SVP Level

hgt20

Abstract

The calculation of excited states for the system 'Z_T_azobenzene' is presented, accompanied by automated analysis and image generation provided by the Digichem software package. The calculation was performed using the ORCA software package at the DFT/def2-SVP level of theory. The total self-consistent field (SCF) energy of the system was found to be -15565.18 eV after 1 step. The highest-occupied molecular orbital (HOMO) and lowest-unoccupied molecular orbital (LUMO) were calculated to be -5.61 and -2.15 eV respectively, corresponding to a HOMO-LUMO band gap of 3.47 eV. The permanent dipole moment (PDM) was calculated to be 3.48 D. In total, 2 excited states were calculated with singlet multiplicity. The most intense absorption peaks were calculated to be at 327 and 563 nm. The lowest energy singlet excited state (S_1) was calculated to be 2.17 eV (571 nm).

Table 1: *Summary of overall calculation metadata.*

Date[a] (Duration[b])	CPUs (Memory)	Success (Converged)	Computational package	Level of theory	Solvent (model)	Calculations	Wavefunction	Multiplicity	T[c] / K	P[d] / atm
N/A (3 m, 19 s)	1 (4 GB)	True (N/A)	ORCA (5.0.3)	DFT/def2-SVP	Gas-phase (N/A)	Excited States	restricted	1 (singlet)	N/A	N/A

[a]: The date and time at which the calculation was completed. [b]: Total combined duration in real-time (wall-time) for all components of the calculation. [c]: Temperature used for thermochemistry analysis. [d]: Pressure used for thermochemistry analysis.

Summary Of Results

SCF Energy

Table 2: *Summary of SCF energy properties.*

No. of steps	1
Final energy	-15565.1826 eV
Final energy	-1,501,812 kJ·mol^{-1}

Geometry

Table 3: *Summary of geometry properties.*

Formula	$C_{12}H_{10}N_2$
SMILES	c1ccc(/N=N\c2ccccc2)cc1
Exact mass	182.2260 g·mol^{-1}
Molar mass	182.2212 g·mol^{-1}
Alignment method	Minimal
X extension	7.58 Å
Y extension	6.02 Å
Z extension	5.89 Å
Linearity ratio	0.21
Planarity ratio	0.02

Molecular Orbitals

Table 4: *Summary of HOMO & LUMO properties.*

$E_{HOMO,LUMO}$	3.47 eV
E_{HOMO}	-5.61 eV
E_{LUMO}	-2.15 eV

Permanent Dipole Moment

Table 5: *Summary of the permanent dipole moment properties.*

Total	3.48 D
X axis angle	83.47 °
XY plane angle	57.78 °

Excited States

Table 6: *Summary of the calculated excited states. E_x: The energy of excited state x. λ_x: The wavelength of a photon of equivalent energy to excited state x. f_x: The oscillator strength of the excited state transition x. ΔE_{xy}: The difference in energy between the lowest excited states of multiplicity x and y.*

No. calculated singlets	2
E_{S_1}	2.17 eV
λ_{S_1} **(colour, CIE)**	571 nm (Yellow, (0.45, 0.55))
f_{S_1}	< 0.01
Simulated Absorption Peaks	327 and 563 nm

Methodology

Metadata

The calculation of the excited states was performed using the **ORCA (5.0.3)** program, the **DFT** method and the **def2-SVP** basis set. It was completed after a total duration of **3 m, 19 s** and **finished successfully**. The base multiplicity of the system under study was **1 (singlet)**. Finally, a **restricted wavefunction** was used, resulting in a single set of doubly occupied orbitals. The full calculation metadata is tabulated in table 1.

Analysis

The report presented here was generated using the Digichem software package. This toolset relies upon a number of third-party applications and libraries which should be cited appropriately in derivative works. In particular, the calculation results described within were parsed by the cclib library.[1] Scientific constants which were used, among other things, for the interconversion of scientific units were provided by SciPy.[2] Commission internationale de l'éclairage (CIE) coordinates, along with visual representations of the equivalent colour, were calculated using the Colour Science library.[3] Three-dimensional

plots of atom positions and calculated densities, including molecular orbitals, were rendered using Visual Molecular Dynamics (VMD)[4] and the Tachyon ray-tracer.[5] Finally, two-dimensional graphs were plotted using the MatPlotlib library,[6] while this report itself was prepared using the Mako template library[7] and the Weasyprint library[8], the latter of which was responsible for generation of the PDF file.

Discussion

Total SCF Energy

The total energy of the system was calculated at the **self-consistent field (SCF)** level, corresponding to the energy calculated by the density-functional theory (DFT) method, with a value of -15565.18 eV, corresponding to -1,501,812 $KJmol^{-1}$. A plot of the total SCF electron density is shown in figure 1.

A: *B:*
C: *D:*

Figure 1: *Plot of the total SCF electron density, plotted with an isovalue of 0.02. A: In the X/Y plane, B: In the X/Z plane, C: In the Z/Y plane, D: 45° to the axes.*

Geometry



Figure 2: *Labelled structure of Z_T_azobenzene.*

The **empirical formula** of the studied system was $C_{12}H_{10}N_2$, corresponding to a **molecular mass** of 182.22 $gmol^{-1}$ and an **exact mass**, considering only specific atomic isotopes, of 182.23 $gmol^{-1}$. The molecular structure, with atom labelling, is shown in figure 2. The molecular geometry was aligned to the cartesian (X, Y and Z) axes by the **Minimal (MIN)** method, and the resulting atomic position are displayed in figure 3. Using this method, the **extent of the molecular system** in the X, Y and Z axes (L_X, L_Y and L_Z, corresponding to the molecular width, length and height respectively) was determined to be 7.58, 6.02 and 5.89 Å respectively. These extensions give rise to a **molecular linearity ratio** ($1-(L_Y/L_X)$) and **planarity ratio** ($1-(L_X/L_Y)$) of 0.21 and 0.02 respectively.

A: *B:*
C: *D:*

Figure 3: *The molecular structure, aligned using the Minimal (MIN) method. A: In the X/Y plane, B: In the X/Z plane, C: In the Z/Y plane, D: 45° to the axes.*

Permanent Dipole Moment

The calculated **permanent dipole moment** was 3.48 D, with a vector (x,y,z) of -0.56, -2.46, -2.40 D. The origin point of the dipole vector was (-2.08, 4.52, -13.74). The angle between the dipole moment vector and the x-axis was 83.47 °, while the angle between the dipole moment and the xy-plane was 57.78 °. A plot of the permanent dipole moment is shown in figure 4.

A: *B:*
C: *D:*

Figure 4: *The permanent dipole moment (red arrow) plotted against the aligned molecular geometry with a scale of 1 Å = 1.0 D. A: In the X/Y plane, B: In the X/Z plane, C: In the Z/Y plane, D: 45° to the axes.*

Molecular Orbitals

In total, 246 doubly occupied molecular orbitals were calculated, divided into 48 occupied orbitals and 198 unoccupied (or virtual) orbitals. The calculated energies of the **HOMO and LUMO** were -5.61 and -2.15 eV respectively, corresponding to a **HOMO-LUMO band gap** of 3.47 eV (figure 8). Plots of the orbital density for the HOMO and LUMO are shown in figures 5-6 respectively, while the orbital overlap between the HOMO and LUMO is shown in figure 7.



Figure 8: *Graph of the calculated molecular orbital energies in close proximity to the HOMO-LUMO gap. Solid lines: occupied orbitals, dashed lines: virtual orbitals.*

A: *B:*
C: *D:*

Figure 5: *Orbital density plots of the HOMO, plotted with isovalue: 0.02. A: In the X/Y plane, B: In the X/Z plane, C: In the Z/Y plane, D: 45° to the axes.*

A: *B:*
C: *D:*

Figure 6: *Orbital density plots of the LUMO, plotted with isovalue: 0.02. A: In the X/Y plane, B: In the X/Z plane, C: In the Z/Y plane, D: 45° to the axes.*

A: *B:*
C: *D:*

Figure 7: *Orbital density plots of the HOMO (blue) and LUMO (red), plotted simultaneously with isovalue: 0.02. A: In the X/Y plane, B: In the X/Z plane, C: In the Z/Y plane, D: 45° to the axes.*

Excited States

In total, the energies of two singlet electronic excited states were calculated, which are shown in figure 9. The energy of the

lowest **singlet excited state (S_1)** was 2.17 eV, corresponding to absorption by a photon with a wavelength of 571 nm, a yellow 'color' ■ and CIE coordinates of (0.45, 0.55). A complete table of the calculated excited state properties is available in table 7. In addition, an electronic transition spectrum was simulated using a gaussian function with full-width at half maximum (FWHM) of 0.40 eV, from which the **two most intense peaks** were found at 327 and 563 nm. The full simulated absorption spectrum is shown in figure 10.



Figure 9: *Graph of the calculated excited states. f: oscillator strength of the relevant ground to excited state transition.*



Figure 10: *Graph of the simulated absorption spectrum. Excited states are shown as vertical black lines, while peaks simulated with a gaussian function with FWHM: 0.40 eV are shown as a blue line. Peaks can be found at: 327 and 563 nm.*

Tables Of Results

Atom Coordinates

Table 8: *Coordinates of the atoms of the system under study, as aligned to the cartesian axes by the Minimal method. Atoms that are chemically equivalent have been assigned the same group number.s*

Index	Element	Group	X Coord /Å	Y Coord /Å	Z Coord /Å
1	C	C_1	0.2919900	4.1728700	-9.4811900
2	C	C_2	1.4167700	3.3467700	-9.5420600
3	C	C_1	1.8519100	2.6749800	-8.3972600
4	C	C_3	1.1590000	2.8217400	-7.1915800
5	C	C_4	0.0113400	3.6387000	-7.1258700
6	C	C_3	-0.4032400	4.3225600	-8.2772200
7	H	H_3	-1.2716700	4.9685000	-8.2412000
8	H	H_1	-0.0403100	4.6994700	-10.3664900
9	H	H_2	1.9552400	3.2331200	-10.4740400
10	H	H_1	2.7300100	2.0437800	-8.4425500
11	H	H_3	1.5275100	2.3045500	-6.3163100
12	N	N_5	-0.7173400	3.8422000	-5.9156900
13	N	N_5	-1.6800000	3.1557000	-5.4918900
14	C	C_4	-2.2626300	2.0167500	-6.1245400
15	C	C_3	-3.3742500	1.4135800	-5.5198500
16	C	C_1	-3.9932500	0.3089800	-6.1131300
17	C	C_2	-3.5135800	-0.1956000	-7.3241600
18	C	C_1	-2.4187500	0.4082600	-7.9469800
19	C	C_3	-1.7967700	1.5127200	-7.3565300
20	H	H_3	-0.9624100	1.9721400	-7.8685400
21	H	H_3	-3.7571600	1.7982300	-4.5827600
22	H	H_1	-4.8464900	-0.1542500	-5.6347300
23	H	H_2	-3.9946500	-1.0494700	-7.7834200
24	H	H_1	-2.0545700	0.0235700	-8.8908600

Molecular Orbitals

Table 9: *Energies of the calculated molecular orbitals.*

Level	Label	Symmetry	Energy /eV
64	LUMO+15	A	4.0772
63	LUMO+14	A	4.0692
62	LUMO+13	A	3.9671
61	LUMO+12	A	3.9094
60	LUMO+11	A	3.8139
59	LUMO+10	A	3.1688
58	LUMO+9	A	3.0710
57	LUMO+8	A	2.7169
56	LUMO+7	A	2.6876
55	LUMO+6	A	1.7507
54	LUMO+5	A	1.6564
53	LUMO+4	A	0.5293
52	LUMO+3	A	-0.2474
51	LUMO+2	A	-0.3078
50	LUMO+1	A	-0.7471
49	**LUMO**	**A**	**-2.1480**
48	**HOMO**	**A**	**-5.6147**
47	HOMO-1	A	-7.0090
46	HOMO-2	A	-7.0474
45	HOMO-3	A	-7.2180
44	HOMO-4	A	-7.3842
43	HOMO-5	A	-9.2627
42	HOMO-6	A	-9.4953
41	HOMO-7	A	-9.9882
40	HOMO-8	A	-10.0427
39	HOMO-9	A	-10.1725
38	HOMO-10	A	-10.2385
37	HOMO-11	A	-10.6754
36	HOMO-12	A	-11.1649
35	HOMO-13	A	-11.6769
34	HOMO-14	A	-11.8967
33	HOMO-15	A	-12.1182

Excited States

Table 7: *Energies and other properties of the calculated excited states.*

Number	Symbol	Symmetry	Energy /eV	Wavelength /nm	Colour (CIE x,y)	Oscillator Strength	Transitions (Probability)
1	S_1	Singlet	2.1730	570.58	Yellow (0.45, 0.55)	0.0011	
2	S_2	Singlet	3.7782	328.16	Ultraviolet (0.00, 0.00)	0.0005	

References

1. N. M. O'boyle, A. L. Tenderholt and K. M. Langner, *Journal of Computational Chemistry,* 2008, **29,** 839--845
2. P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. Jarrod Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. Carey, İ. Polat, Y. Feng, E. W. Moore, J. Vand erPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt and S. 1. 0. Contributors, *Nature Methods,* 2020, **17,** 261--272
3. T. Mansencal, M. Mauderer, M. Parsons, N. Shaw, K. Wheatley, S. Cooper, J. D. Vandenberg, L. Canavan, K. Crowson, O. Lev, K. Leinweber, S. Sharma, T. J. Sobotka, D. Moritz, M. Pppp, C. Rane, P. Eswaramoorthy, J. Mertic, B. Pearlstine, M. Leonhardt, O. Niemitalo, M. Szymanski and M. Schambach, Colour 0.3.15, Zenodo, 2020
4. W. Humphrey, A. Dalke and K. Schulten, *Journal of Molecular Graphics,* 1996, **14,** 33--38
5. J. Stone, Masters Thesis, Computer Science Department, University of Missouri-Rolla, 1998
6. J. D. Hunter, *Computing in Science & Engineering,* 2007, **9,** 90--95
7. M. Bayer, https://www.makotemplates.org, (accessed May 2020)
8. K. Community, https://weasyprint.org, (accessed May 2020)