

# **IE 400**

## **Project**



## **Group 49**

## **Report**

Hammad Khan Musakhel

21801175

December 30, 2021

In this project, we have worked on the optimization problem for Dr. Who's basic regimen for chemotherapy sessions. There are seven drugs which are to be used for cancer patients.

However, these seven drugs have an impact on the patient's Q-score value, which indicates the quality of life for the patients. A higher Q-score depicts a better We have been provided with an equation which estimates the Q-score:

$$\begin{aligned} Q(\mathbf{p}, \mathbf{y}, \mathbf{x}) = & -5p_1 - 0.5p_2 - 12p_3 - 8p_4 - 5p_5 - 5p_6 - 1p_7 - 3p_8 - 2p_9 \\ & - 5y_1 - 6y_2 - 4y_3 - 4y_4 - 8y_5 - 6y_6 - 7y_7 \\ & + 0.28x_1 + 0.30x_2 + 0.25x_3 + 0.17x_4 + 0.31x_5 + 0.246x_6 + 0.40x_7 \end{aligned}$$

This equation is used in all the parts of the project to predict the amount of dosage of each drug to be used. The primary focus of this project is the optimization of different drug dosages in order to maximize the Q-score for patients and minimize the cost of addition and removal of drugs in the base regimen.

## **Part A**

In the first part of the project, we are tasked to maximize the Q-score of the patient with four drugs provide. In this part of the project, there is no specific Q-score to achieve, rather maximize the Q-score with certain drugs. A base regimen is provided to us in which the four drugs are included: Melphalan (X1), Decitabine (X3), Pentostatin (X4), and Thiotepa (X7). A unit of an acceptable chemotherapy regimen is 100 centiliters (cl) and must include one or more of these drugs. Thus, we are initially provided with 20 cl of X1, 30 cl of X3, 15 cl of X4, and 35 cl of X7.

The goal was to maximize the Q-score from the above-mentioned equation by using these drugs as the only drugs in the regimen. We could alter the dosage of these four drugs in order to maximize the Q-score. For this, we programmed a mathematical optimization solver with CPLEX in order to calculate the maximized Q-score. Moreover, to start with the program, we outlined our necessary decision variables and constraints:

## **Decision Variables**

$x_i$  = dosage of drug  $j$  in the regimen, and  $x = 1, 2, 3, 4, 5, 6, 7$

$p_i = (p_1, p_2, \dots, p_9)$  patient's characteristics which are (1,1,0,0,1,1,0,0,0)

$y_i = (y_1, y_2, \dots, y_7)$  the presence of drug 1 to 7 which is 1 if it is present and 0 if not

### Objective Function:

$P\_value = -5 \cdot 1 - 0.5 \cdot 1 - 12 \cdot 0 - 8 \cdot 0 - 5 \cdot 1 - 1 \cdot 0 - 3 \cdot 0 - 2 \cdot 0 \rightarrow$  which is -15.5

$Y\_value = -5 \cdot 1 - 6 \cdot 0 - 4 \cdot 1 - 4 \cdot 1 - 8 \cdot 0 - 6 \cdot 0 - 7 \cdot 1 \rightarrow$  which is -20

$Q\_score = P\_value + Y\_value + 0.28 \cdot X_1 + 0.30 \cdot X_2 + 0.25 \cdot X_3 + 0.17 \cdot X_4 + 0.31 \cdot X_5 + 0.246 \cdot X_6 + 0.40 \cdot X_7$

### max Q-score

Subject to:

The lower and upper bounds for the four drugs' dosage and equating the others to zero:

$80 \geq X_1 \geq 20$

$100 \geq X_3 \geq 20$

$100 \geq X_4 \geq 10$

$50 \geq X_7 \geq 20$

$X_2, X_5, X_6 = 0$

The constraint of total maximum quantity:

$X_1 + X_3 + X_4 + X_7 = 100$

### Feasible Solution:

Objective value/Q-score value: **-3.200003**

Values of decision variables: **[20, 20, 10, 50]** for **[X1, X3, X4, X7]**

It can be concluded that the best value for Q-score could be -3.2 with drugs X1, X3, X4, X7 used in the base regimen only. The best combination for these four drugs to attain the maximum Q-score value **X1 = 20 cl, X3 = 20 cl, X4 = 10 cl, X7 = 50 cl.**

## Part b

### Decision Variables

$x_i$  = Amount of dosage of drug i

$y_i$  = 1 if drug i is included, 0 otherwise

$c_i$  = 1 if drug i is included, 0 otherwise

$a_i$  = Amount of dosage changed of drug i

### Objective Function

$$\min (25c_1 + 1x_1) + (50c_2 + 2x_2) + (10c_3 + 1x_3) + (25c_4 + 3x_4) + (20c_5 + 2x_5) + \\ (30c_6 + 1x_6) + (40 + 1x_7)$$

s.t.

$$Q(\mathbf{p}, \mathbf{y}, \mathbf{x}) = -5p_1 - 0.5p_2 - 12p_3 - 8p_4 - 5p_5 - 5p_6 - 1p_7 - 3p_8 - 2p_9 \\ - 5y_1 - 6y_2 - 4y_3 - 4y_4 - 8y_5 - 6y_6 - 7y_7 \\ + 0.28x_1 + 0.30x_2 + 0.25x_3 + 0.17x_4 + 0.31x_5 + 0.246x_6 + 0.40x_7$$

$$Q(\mathbf{p}, \mathbf{x}, \mathbf{y}) = 25$$

$$c_1 = 1 - y_1$$

$$c_2 = y_2$$

$$c_3 = 1 - y_3$$

$$c_4 = 1 - y_4$$

$$c_5 = 1 - y_5$$

$$c_6 = y_6$$

$$c_7 = 1 - y_7$$

$$add_i + remove_i \leq 1 \text{ for } i = 1, 2, 3, 4, 5, 6, 7$$

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 = 200$$

$$20 \leq x_1 \leq 80$$

$$10 \leq x_2 \leq 50$$

$$20 \leq x_3 \leq 100$$

$$10 \leq x_4 \leq 100$$

$$10 \leq x_5 \leq 70$$

$$10 \leq x_6 \leq 70$$

$$20 \leq x_7 \leq 50$$

### **Feasible Solution**

After minimizing the cost, we get the following variable dosages:

$$x_1 = 71$$

$$x_2 = 0$$

$$x_3 = 100$$

$$x_4 = 0$$

$$x_5 = 0$$

$$x_6 = 0$$

$$x_7 = 29$$

Quality of life = 25

Using these variables, the cost is minimized cost is 206

## Part C

In this part, we are tasked to add further constraints to our model in part B. In this part, we primarily focus on further improvising the base regimen's content by introducing additional constraints. In B part, our focus was to reach the Q-score of 25 (for our patient number 49) by altering the base regimen's content in terms of all the drugs, that is we could administer a base regimen with all drugs included and with maximum quantity of 100 cl. Moreover, we were tasked to minimize the cost of addition/removal of those drugs along with the unit cost of change in dosage.

In part c, the additional constraints we added to our LP/IP model are as follows:

1) Melphalan and Oxaliplatin dosage:

$$X1 + X2 \geq 30$$

$$X1 + X2 \leq 70$$

2) Either Epirubicin included or Decitabine should be less than 25 cl:

$$w = \{ 0 \text{ if Epirubicin is included (Y5 = 1) , 1 otherwise (Y5 = 0) } \}$$

$$Y3 \leq w$$

$$X5(1-Y3) < 25$$

3) If both Pentostatin and Lomoustine included then at least one of Thiotepa and Epirubicin also be included:

$$z = \{ 1 \text{ if } (Y4 + Y6 = 2), 0 \text{ otherwise } \}$$

$$Y4 + Y6 = 2z$$

$$Y5 + Y7 \geq z$$

## Feasible Solution

After minimizing the cost, we get the following variable dosages:

$$x1 = 68$$

$$x2 = 0$$

$$x3 = 0$$

$$\mathbf{x4 = 0}$$

$$\mathbf{x5 = 0}$$

$$\mathbf{x6 = 90}$$

$$\mathbf{x7 = 43}$$

$$\text{Quality of life} = 25$$

Using these variables, the cost is minimized and it is 286.8

