


The Five Color Theorem


Danny Hammer

August 26, 2021


- Recall that a **Graph** is a mathematical structure consisting of a set of vertices V and edges E .




- Recall that a **Graph** is a mathematical structure consisting of a set of vertices V and edges E .
 - A **Vertex** (or **Node**) represents an entity in the graph.




- Recall that a **Graph** is a mathematical structure consisting of a set of vertices V and edges E .
 - A **Vertex** (or **Node**) represents an entity in the graph.
 - An **Edge** represents connection between vertices.




- A **Planar Graph** is a graph that can be drawn on a two dimensional surface such that no edges overlap. [3]





- A **Planar Graph** is a graph that can be drawn on a two dimensional surface such that no edges overlap. [3]
 - The area enclosed by a set of edges is called a **Face**.





- A **Planar Graph** is a graph that can be drawn on a two dimensional surface such that no edges overlap. [3]
 - The area enclosed by a set of edges is called a **Face**.
 - If a graph cannot be drawn this way, it is not planar.




- Geographical maps can be translated into planar graphs.
- Nations are nodes and their borders are edges.



- Geographical maps can be translated into planar graphs.
- Nations are nodes and their borders are edges.
 - This translation allows us to use graph theory when constructing maps. [1]




- The **chromatic number** $\chi(G)$ of a graph G is the minimum number of colors needed to assign a color to every vertex in G such that no two vertices sharing an edge are the same color.



Theorem (Euler's Formula)


Let G be a connected planar graph with size e and order v . Any planar depiction of G has $e - v + 2$ faces.



Theorem (Euler's Formula)

Let G be a connected planar graph with size e and order v . Any planar depiction of G has $e - v + 2$ faces.

- e is the number of edges (size)
- v is the number of vertices (order)



Corollary (1)

If G is a connected planar graph with $e > 1$, then $e \leq 3v - 6$.

Corollary (2)

Every planar graph G has a vertex of degree ≤ 5 .

- Derived from Euler's formula
- Basis for the Five Color Theorem

Theorem (The Five Color Theorem)

Every planar graph G has $\chi(G) \leq 5$.

- Trivially true for any G with order ≤ 5 .
- Will use induction on the order of the graph v for the non-trivial cases. [2]

- Inductive Hypothesis: Assume $\chi(G) \leq 5$ for every planar graph G of order v and consider a planar graph H of order $v + 1$.

- Inductive Hypothesis: Assume $\chi(G) \leq 5$ for every planar graph G of order v and consider a planar graph H of order $v + 1$.
- From Corollary 3 we know H contains a vertex x with degree ≤ 5 .

- Inductive Hypothesis: Assume $\chi(G) \leq 5$ for every planar graph G of order v and consider a planar graph H of order $v + 1$.
- From Corollary 3 we know H contains a vertex x with degree ≤ 5 .
- Now look at J , which is $H - x$ and all edges incident with it.

- Inductive Hypothesis: Assume $\chi(G) \leq 5$ for every planar graph G of order v and consider a planar graph H of order $v + 1$.
- From Corollary 3 we know H contains a vertex x with degree ≤ 5 .
- Now look at J , which is $H - x$ and all edges incident with it.
- We know J is planar and of order v , so $\chi(J) \leq 5$.

- Inductive Hypothesis: Assume $\chi(G) \leq 5$ for every planar graph G of order v and consider a planar graph H of order $v + 1$.
- From Corollary 3 we know H contains a vertex x with degree ≤ 5 .
- Now look at J , which is $H - x$ and all edges incident with it.
- We know J is planar and of order v , so $\chi(J) \leq 5$.
- Suppose J has been colored with at most 5 colors, and now we must attempt to rejoin x into J .




Figure: Planar Depiction of a subgraph of H

Case 1: J is colored using < 5 colors.

Case 1: J is colored using < 5 colors.

- Reintroduce x with an unused color.

Case 1: J is colored using < 5 colors.

- Reintroduce x with an unused color.

Case 2: J is colored using 5 colors but x has degree < 5 .

Case 1: J is colored using < 5 colors.

- Reintroduce x with an unused color.

Case 2: J is colored using 5 colors but x has degree < 5 .

- Again, we can reintroduce x with the color not used by any of its neighbors.

Case 1: J is colored using < 5 colors.

- Reintroduce x with an unused color.

Case 2: J is colored using 5 colors but x has degree < 5 .

- Again, we can reintroduce x with the color not used by any of its neighbors.

Case 3: J is colored using 5 colors and x has degree 5.

Case 1: J is colored using < 5 colors.


- Reintroduce x with an unused color.

Case 2: J is colored using 5 colors but x has degree < 5 .

- Again, we can reintroduce x with the color not used by any of its neighbors.

Case 3: J is colored using 5 colors and x has degree 5.


- Identify the five nodes (clockwise) adjacent to x as a, b, c, d , and e and the five available colors as 1, 2, 3, 4, and 5.
- If any two of these vertices are the same color, x can use the remaining fifth color.



- Consider (WLOG) two non-adjacent vertices a and c (colored 1 and 3).

- Consider (WLOG) two non-adjacent vertices a and c (colored 1 and 3).
- If there is no walk connecting a and c consisting of only colors 1 and 3, simply alternate the coloring of a and its adjacent nodes.

- Consider (WLOG) two non-adjacent vertices a and c (colored 1 and 3).
- If there is no walk connecting a and c consisting of only colors 1 and 3, simply alternate the coloring of a and its adjacent nodes.
- Thus, color 1 is now available for x .




- If there *is* a walk connecting a and c consisting of only colors 1 and 3...

- If there *is* a walk connecting a and c consisting of only colors 1 and 3...
- Consider two other non-adjacent vertices b and d (colored 2 and 4).

- If there *is* a walk connecting a and c consisting of only colors 1 and 3...
- Consider two other non-adjacent vertices b and d (colored 2 and 4).
- Since J is planar, there cannot exist a walk connecting b and d consisting only of colors 2 and 4.

- If there *is* a walk connecting a and c consisting of only colors 1 and 3...
- Consider two other non-adjacent vertices b and d (colored 2 and 4).
- Since J is planar, there cannot exist a walk connecting b and d consisting only of colors 2 and 4.
- So alternate the coloring of b and its adjacent vertices.

- If there *is* a walk connecting a and c consisting of only colors 1 and 3...
- Consider two other non-adjacent vertices b and d (colored 2 and 4).
- Since J is planar, there cannot exist a walk connecting b and d consisting only of colors 2 and 4.
- So alternate the coloring of b and its adjacent vertices.
- Thus, color 2 is now available for x .



- Therefore if G is a planar graph of order v such that $\chi(G) \leq 5$, then $\chi(H) \leq 5$ for a planar graph H of order $v + 1$.

- Therefore if G is a planar graph of order v such that $\chi(G) \leq 5$, then $\chi(H) \leq 5$ for a planar graph H of order $v + 1$.
- Thus we can see that any planar graph can be colored with at most five colors.

- Strictest coloring bounds provable without computers.

- Strictest coloring bounds provable without computers.
- There is an improved Four Color Conjecture, but can only be proven by computational brute-force.

- Strictest coloring bounds provable without computers.
- There is an improved Four Color Conjecture, but can only be proven by computational brute-force.
- All map projections can be colored with ≤ 5 colors.

References I



Moti Ben-Ari.

The five-color theorem.
2021.



Richard J. Trudeau.

Introduction to Graph Theory.

Dover Books on Mathematics. Dover Publications, 2nd
edition, 1994.



Matthew Wahab.

5-coloring planar maps.
website, 2003.