

Buffon's Needle

Danny Hammer, Julia Januchowski

October 12th, 2021

- If you dropped a needle onto a sheet of ruled paper, what is the probability that the needle will land across a line?

- If you dropped a needle onto a sheet of ruled paper, what is the probability that the needle will land across a line?
- It would of course depend on the length l of your needle and the distance d between lines.

- If you dropped a needle onto a sheet of ruled paper, what is the probability that the needle will land across a line?
- It would of course depend on the length l of your needle and the distance d between lines.
- Hint: This probability is related to π !

Theorem (Buffon's Needle)

If a needle of length l is dropped on a paper with equally spaced lines of distance $d \geq l$, then the probability P that the needle lands in such a way that it rests on a line is

$$P = \frac{2l}{\pi d}$$

- Let n be the number of needles dropped and h be the number of needles that land on a line.

- Let n be the number of needles dropped and h be the number of needles that land on a line.
- Recall that l is the length of our needle and d is the distance between lines, and $l \leq d$

- Let n be the number of needles dropped and h be the number of needles that land on a line.
- Recall that l is the length of our needle and d is the distance between lines, and $l \leq d$

$$\frac{h}{n} \approx \frac{2l}{\pi d}$$

- Let n be the number of needles dropped and h be the number of needles that land on a line.
- Recall that l is the length of our needle and d is the distance between lines, and $l \leq d$

$$\frac{h}{n} \approx \frac{2l}{\pi d} \rightarrow \pi \approx 2 \frac{ln}{dh}$$


- Let n be the number of needles dropped and h be the number of needles that land on a line.
- Recall that l is the length of our needle and d is the distance between lines, and $l \leq d$

$$\frac{h}{n} \approx \frac{2l}{\pi d} \rightarrow \pi \approx 2 \frac{ln}{dh}$$

- Since l and d are constants, we can estimate π by dropping needles! [1]

- Consider an arbitrary needle dropped onto a plane with evenly ruled horizontal lines of distance d apart.

- Consider an arbitrary needle dropped onto a plane with evenly ruled horizontal lines of distance d apart.
- We can then find the angle the needle makes with the horizontal (α) and the height of the needle using trigonometry



We will then use probability to say the probability that a needle crosses a line is:

$$P = \frac{l \sin(\alpha)}{d}$$

We can then say that the average value of a function from a to b is :

$$f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$$

We can then say that the average value of a function from a to b is :

$$f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$$

In our case:

- $f(x) = P = \frac{l \sin(\alpha)}{d}$

We can then say that the average value of a function from a to b is :

$$f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$$

In our case:

- $f(x) = P = \frac{l \sin(\alpha)}{d}$
- $a = 0$

We can then say that the average value of a function from a to b is :

$$f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$$

In our case:

- $f(x) = P = \frac{l \sin(\alpha)}{d}$
- $a = 0$
- $b = \frac{\pi}{2}$

Therefore, the average probability that a given needle of length l crosses a line at any α is:

$$P = \frac{2}{\pi} \int_0^{\frac{\pi}{2}} \frac{l \sin(\alpha)}{d} d\alpha$$

$$P = \frac{2}{\pi} \frac{l}{d} [-\cos \alpha]_0^{\frac{\pi}{2}}$$

$$P = \frac{2}{\pi} \frac{l}{d}$$



Additional solution: E. Barbier (1860) “Buffon’s noodle”

- Drop any size or shaped needle
- Focused on the probability of the number of times a needle crossed a line
 - utilizes limits
- Circular needle where the diameter = d




Figure: Example of “noodles” dropped on a sheet of paper

- In 1901, Lazzarini experimented with 3,408 needles of length 5 and distance of 6. He had 1,808 successes.

- In 1901, Lazzarini experimented with 3,408 needles of length 5 and distance of 6. He had 1,808 successes.
- $2 \frac{5 \cdot 3408}{6 \cdot 1808} \approx 3.1415929203539825$

- In 1901, Lazzarini experimented with 3,408 needles of length 5 and distance of 6. He had 1,808 successes.
- $2 \frac{5 \cdot 3408}{6 \cdot 1808} \approx 3.1415929203539825$
- Lazzarini's "estimation" was accurate to six decimal places with a suspiciously low number of needles.

- In 1901, Lazzarini experimented with 3,408 needles of length 5 and distance of 6. He had 1,808 successes.
- $2 \frac{5 \cdot 3408}{6 \cdot 1808} \approx 3.1415929203539825$
- Lazzarini's "estimation" was accurate to six decimal places with a suspiciously low number of needles.
- But why 3,408?

- Consider the well-known approximation $\frac{355}{113} = 3.14159292035$

- Consider the well-known approximation $\frac{355}{113} = 3.14159292035$

$$\frac{355}{133} = \frac{5n}{6h} \rightarrow h = \frac{113n}{213}$$

- Consider the well-known approximation $\frac{355}{113} = 3.14159292035$

$$\frac{355}{133} = \frac{5n}{6h} \rightarrow h = \frac{113n}{213}$$

- So long as n is a multiple of 213, h will be an integer and the formula will estimate π to six decimal places.

- Consider the well-known approximation $\frac{355}{113} = 3.14159292035$

$$\frac{355}{133} = \frac{5n}{6h} \rightarrow h = \frac{113n}{213}$$

- So long as n is a multiple of 213, h will be an integer and the formula will estimate π to six decimal places.
- All one needs to do is drop 213 needles at a time until the number of successes h is a multiple of 113.

- Consider the well-known approximation $\frac{355}{113} = 3.14159292035$

$$\frac{355}{133} = \frac{5n}{6h} \rightarrow h = \frac{113n}{213}$$

- So long as n is a multiple of 213, h will be an integer and the formula will estimate π to six decimal places.
- All one needs to do is drop 213 needles at a time until the number of successes h is a multiple of 113.
- Note $3,408 = 16 \cdot 213$ [2]

Significance

- Estimate π by simply dropping sticks
- Monte Carlo Simulation

References I



Martin Aigner and Gnter M. Ziegler.

Proofs from THE BOOK.

Springer Publishing Company, Incorporated, 4th edition, 2009.



Lee Badger.

Lazzarini's lucky approximation of π .

Mathematics Magazine, 67(2):83–91, 1994.