

# PODS Lab 4: Probability Distributions, Central Tendency, and Dispersion

Hamza Alshamy

Center for Data Science, NYU  
ha2486@nyu.edu

02/14/25

- 1 **Probability Distributions**
- 2 **Central Tendency and Ergodicity**
- 3 **Dispersion**
- 4 **Quiz + Discussion**
- 5 **Pain Relief** (Questions?)

## What Are Probability Distributions?

- ▶ **Probability distributions** describe how probabilities are assigned to different possible outcomes of a random variable.
- ▶ They are **governed by parameters**, which control their shape and properties.

## Why Are They Relevant?

- 1 **Describing Random Variables:** Distributions characterize the behavior of random variables, assigning probabilities to their possible values.
- 2 **Probabilistic Modeling:** If we have data, we can use distributions to model uncertainty and make predictions.

## Example of Probability Distributions

### Continuous: Normal Distribution

#### Normal Distribution Equation:

$$P(\mathcal{X} = x; \mu, \sigma) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

## Example of Probability Distributions

### Continuous: Normal Distribution

#### Normal Distribution Equation:


$$P(\mathcal{X} = x; \mu, \sigma) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- ▶ **In English:** "The probability of the random variable  $\mathcal{X}$  taking on the value  $x$  given the parameters, the mean  $\mu$  and standard deviation  $\sigma$ "
- ▶ **Logic:** Many natural phenomena (e.g., heights, test scores) follow a normal distribution.
- ▶ **Parameters:**
  - 1  $\mu$  (**mean**): Determines the center of the distribution.
  - 2  $\sigma$  (**Standard Deviation**): Controls the spread of the distribution.

# Example of Probability Distributions

## Continuous: Normal Distribution

- Notice how the shape of the normal distributions differs based on the **parameters** ( $\mu$ ,  $\sigma$ )



## Example of Probability Distributions

### Discrete: Poisson Distribution

#### Poisson Distribution Equation:

$$P(\mathcal{X} = k; \lambda) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad k = 0, 1, 2, \dots$$

- ▶ **In English:** "The probability of the random variable  $\mathcal{X}$  taking on the value  $k$  given the rate parameter  $\lambda$ ."
- ▶ **Logic:** Models the number of events occurring in a fixed interval of time or space when events happen independently at a constant average rate.
- ▶ **Parameter:**
  - ①  **$\lambda$  (Rate Parameter):** Represents the expected number of occurrences in the given interval.
- ▶ **Example:** Model the number of earthquakes that will occur given a time interval.

# Random Variables and Probability Distributions

## Example: Measuring IQ

- ▶ Random variables can follow specific probability distributions.
- ▶ **Example:** Let  $\mathfrak{X}$  represent IQ scores.

$$\mathfrak{X} \sim \mathcal{N}(\mu = 100, \sigma = 15)$$

- ▶ **In English:** "The random variable  $\mathfrak{X}$  follows a normal distribution with mean  $\mu = 100$  and standard deviation  $\sigma = 15$ ."
- ▶ **Interpretation:**
  - 1 Most IQ scores will be close to 100, with variation controlled by  $\sigma = 15$ .
  - 2 About 68% of IQ scores fall within one standard deviation ( $85 \leq \mathfrak{X} \leq 115$ ).
  - 3 Approximately 95% of IQ scores lie within two standard deviations ( $70 \leq \mathfrak{X} \leq 130$ ).



## Central Tendency, Typical Value, Average

- ▶ **Definition:** Measures of central tendency describe a typical or central value of a dataset.
- ▶ **Relevance:**
  - ▶ If you know nothing about the data, the best guess for an unknown value is often the *average*.
  - ▶ Helps summarize data with a single representative value.
- ▶ **Main Measures of Central Tendency:**
  - 1 **(Arithmetic) Mean:** The average of all values.
  - 2 **Median:** The middle value when data is ordered.
  - 3 **Mode:** The most frequently occurring value.

## (1) The Arithmetic Mean

### Best used when:

- 1 (Approximately) Normal or symmetrical data
- 2 No (or few) extreme values – Mean is not very robust
- 3 Not heavily skewed – Mean will be dragged towards the tail

### Example:

$$\text{Set} = \{9, 3, 300, 8, 7, 10, 8, 5\}$$

- Calculation of the Mean:

## (1) The Arithmetic Mean

### Best used when:

- 1 (Approximately) Normal or symmetrical data
- 2 No (or few) extreme values – Mean is not very robust
- 3 Not heavily skewed – Mean will be dragged towards the tail

### Example:

$$\text{Set} = \{9, 3, 300, 8, 7, 10, 8, 5\}$$

- Calculation of the Mean:

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i = \frac{9 + 3 + 300 + 8 + 7 + 10 + 8 + 5}{8} = \frac{350}{8} = \boxed{43.75}$$

- The mean is dragged towards the outlier (300), making it not representative of most values in the set.

## (2) The Median

- ▶ **Definition:** The middle value when data is ordered.
- ▶ **Best used when:**
  - 1 Extreme values – Median is more robust than the mean.
  - 2 Data is skewed (not symmetrical).

### Example:

$$\begin{aligned}\text{Set} &= \{9, 3, 300, 8, 7, 10, 8, 5\} \\ &= \{3, 5, 7, 8, 8, 9, 10, 300\} \quad (\text{Ordered})\end{aligned}$$

- ▶ Calculation of the Median:

$$\text{Median} = \frac{8 + 8}{2} = \boxed{8}$$

- ▶ Notice how the median is not affected by the extreme value (300), unlike the mean.
- ▶ Representative of the set!

### (3) The Mode

- ▶ **Definition:** The most frequently occurring value in a dataset.
- ▶ **Best used when:**
  - 1 With categorical data.
  - 2 When numbers are used as labels.

#### Example: Diagnosing Disorders in DSM-IV

Disorders = {Anxiety, Depression}

Disorder Codes = {300.00, 311.00}

- ▶ **Incorrect Approach (Mean):**

$$\frac{300.00 + 311.00}{2} = 305.5 \quad (\text{Code for Opioid Abuse})$$

### (3) The Mode

- ▶ **Definition:** The most frequently occurring value in a dataset.
- ▶ **Best used when:**
  - 1 With categorical data.
  - 2 When numbers are used as labels.

#### Example: Diagnosing Disorders in DSM-IV

Disorders = {Anxiety, Depression}

Disorder Codes = {300.00, 311.00}

- ▶ **Incorrect Approach (Mean):**

$$\frac{300.00 + 311.00}{2} = 305.5 \quad (\text{Code for Opioid Abuse})$$

- ▶ The mean does not represent the state of the patients.
- ▶ **Correct Approach (Mode):**
  - ▶ With categorical data, the most frequent value (mode) is used to represent the typical case.

## Summary of Measures of Central Tendency

### Given Set:

$$\begin{aligned}\text{Set} &= \{9, 3, 300, 8, 7, 10, 8, 5\} \\ &= \{3, 5, 7, 8, 8, 9, 10, 300\} \quad (\text{Ordered})\end{aligned}$$

### Summary Table:


	Mean	Median	Mode
Value	43.75	8	8

- ▶ Measures of central tendency allow us to summarize data with a single representative value.
- ▶ The **median is more robust than the mean** because it is not affected by extreme values.
- ▶ The **mode is best suited for categorical data**, where taking an average does not make sense.

## Note: The Mean and Ergodicity

### Law of Large Numbers (LLN):

- ▶ Given enough samples, the sample mean  $\bar{x}$  converges to the expected value  $\mathbb{E}[X]$ .
- ▶ **However, this only holds if the system is ergodic!**
  - ▶ Otherwise, the mean is not very meaningful.





## Note: The Mean and Ergodicity (cont.)

### What is Ergodicity?

- ▶ The average of the results obtained from a large number of independent random samples converge to the true value
- ▶ The measure of an individual (one coin flip) over time is predictive of the ensemble average (1000 coin flips)

### Assumptions:

- 1 **Stationary:** Statistical Properties are constant over time
  - ▶ The mean, variance, and probability distribution of the system remain consistent.
- 2 **State Space Convergence:** The System Must Visit All Possible States
  - ▶ Over time, air molecules will spread everywhere in the room.
- 3 **Non-Determinism:** Future events are independent of past events.

# Ergodicity: Examples

## Ergodic: Roulette (Assume it's fair)

- ▶ **Total pockets:** 37 → Red: 18, Black: 18, Green: 1
- ▶  $P(\text{red}) = \frac{18}{37}$ ,  $P(\text{black}) = \frac{18}{37}$ ,  $P(\text{green}) = \frac{1}{37}$
- ▶ Results are about the same if you spin 1000 roulettes at the same time or soon one roulette 1000 times

## Non-ergodic: Russian Roulette

- ▶ Total chambers in a revolver: 6 and 5 are empty
- ▶ **60 people playing at the same time:** Survival rate is  $\frac{5}{6} \approx 83.4\%$
- ▶ **One person playing 60 times:** Survival rate is  $(\frac{5}{6})^{60} \approx 1.7\%$

## A Second Way to Characterize Data: Dispersion

- ▶ **Central Tendency:** If I don't know anything about the data, which single value best represents a typical value?
- ▶ **Dispersion:** How much the data deviates from the center.
- ▶ **Main Measures of Dispersion:**
  - ① **Standard Deviation ( $\sigma$ )** – Least robust (sensitive to outliers).
  - ② **Mean Absolute Deviation (MAD)** – More robust than standard deviation.
  - ③ **Median Absolute Deviation (MeAD)** – Most robust (resistant to extreme values).

## (1) Standard Deviation ( $\sigma$ )

### ► Arriving at the Standard Deviation (SD):

$$\frac{1}{n} \sum_i (x_i - \bar{x}) \quad (\text{Always} = 0)$$

$$\frac{1}{n} \sum_i (x_i - \bar{x})^2 \quad (\text{Better, but large deviations has large effects})$$

$$\sigma = \sqrt{\frac{1}{n} \sum_i (x_i - \bar{x})^2} \quad (\text{Taking the square root mitigates that it})$$

### ► Interpretation:

- **Low SD:** Data points are clustered close to the mean
- **High SD:** Data points are more dispersed and further away from the mean

### ► Issue: Influenced by outliers because of squaring!

## Effect of an Outlier on Standard Deviation

### Example: Without an Outlier

$$s = \{5, 6, 7, 8, 9\}, \quad \text{Mean} = 7$$

► **Squared deviations:**

$$\{(5-7)^2, (6-7)^2, (7-7)^2, (8-7)^2, (9-7)^2\} = \{4, 1, 0, 1, 4\}$$

► **Variance:**  $\frac{4+1+0+1+4}{5} = 2 \implies \sigma = \sqrt{2} \approx \boxed{1.41}$

### Example: With an Outlier (Changing 9 to 99)

$$s = \{5, 6, 7, 8, 99\}, \quad \text{Mean} = 25$$

► **Squared deviations:**

$$\{(5-25)^2, (6-25)^2, \dots, (99-25)^2\} = \{400, 361, 324, 289, 5476\}$$

► **Variance:**

$$\frac{400+361+324+289+5476}{5} = 1470 \implies \sigma = \sqrt{1470} \approx \boxed{38.36}$$

## (2) Mean Absolute Deviation (MAD)

- ▶ **Logic:** Squaring values in standard deviation **magnifies deviations**, making it sensitive to outliers.
- ▶ **Instead:** Use absolute values to measure dispersion **without over-emphasizing large deviations**.

### Formula:

$$\text{MAD} = \frac{1}{n} \sum_{i=1}^n |x_i - \bar{x}|$$

- ▶ **More robust** than standard deviation because outliers have less influence.
- ▶ Unlike standard deviation, MAD does not disproportionately weight large deviations.

### (3) Median Absolute Deviation (MeAD)

- ▶ **Logic:** In Mean Absolute Deviation (MAD), we use the **mean**, which is sensitive to outliers.
- ▶ **Instead**, use the **median**, which is **more robust** to extreme values.

#### Formula:

$$\text{MeAD} = \text{Median}(|x_i - \text{Median}(X)|)$$

- ▶ **Most robust** measure of dispersion – outliers have **minimal influence**.
- ▶ Often used in **non-parametric statistics** where distributions may be skewed.

## Summary of Measures of Dispersion and Central Tendency

### Central Tendency

- 1 Mean
- 2 Median
- 3 Mode

#### Example Set:

$$s = \{3, 6, 7, 8, 8, 10, 12, 25\}$$

#### Computed Central Tendencies:

Measure	Value
Mean	9.88
Median	8
Mode	8

### Dispersion

- 1 Standard Deviation
- 2 Mean Absolute Deviation
- 3 Median Absolute Deviation

#### Computed Dispersion Measures:

Measure	Value
Standard Deviation	6.67
Mean Absolute Deviation	4.59
Median Absolute Deviation	2



## Quiz

# Quiz Discussion

# Any questions?