

PODS Lab 3: Probability Theory

Hamza Alshamy

Center for Data Science, NYU
ha2486@nyu.edu

02/07/25

- 1 Basics of Probability
- 2 Conditional Probability
- 3 Random Variables
- 4 Quiz + Discussion

Motivation: Why Do We Need Probability?

- ▶ **Goal:** Reason quantitatively about uncertain phenomena
- ▶ **Questions:**

Motivation: Why Do We Need Probability?

- ▶ **Goal:** Reason quantitatively about uncertain phenomena
- ▶ **Questions:**
 - ▶ Will it rain tomorrow?
 - ▶ Will the value of a stock drop tomorrow?
 - ▶ Who will win the 2026 FIFA world cup?

Main Concepts in Probability

Example: Selecting a Random Locker

Scenario: A gym has 6 lockers labeled 1 through 6. You randomly choose one to store your backpack.

- ▶ **Sample Space (Ω):** The set of all possible outcomes.
 - ▶ The sample space is:

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$

- ▶ **Event:** A subset of the sample space.
 - ▶ The event "Choosing an even-numbered locker" is:


$$B = \{2, 4, 6\}$$

- ▶ **Probability (Frequentist):** The long-run relative frequency of an event occurring.
 - ▶ Since 3 out of the 6 lockers are even-numbered, the probability is:

$$P(B) = \frac{\text{\# of outcomes in } B}{\text{\# of all possible outcomes}} = \frac{3}{6} = 0.5 \quad (\text{or } 50\%)$$


Basic Probability Rules: Union (\cup "Or")

General Union Rule:



$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$


Mutually Exclusive Case:



$$P(A \cup B) = P(A) + P(B) \quad (\text{if } A \cap B = 0)$$


Mutually Exclusive Events

- ▶ **Goal:** Spin the gears
- ▶ **Sample Space:** $\Omega = \{1, 2, \dots, 10\}$
- ▶ **Event:** The number at which the gear stops
- ▶ $P(8 \text{ and } 3) = P(8 \cap 3) = ?$



Mutually Exclusive Events

- ▶ **Goal:** Spin the gears
- ▶ **Sample Space:** $\Omega = \{1, 2, \dots, 10\}$
- ▶ **Event:** The number at which the gear stops
- ▶ $P(8 \text{ and } 3) = P(8 \cap 3) = 0$
 - ▶ **Mutually exclusive:** Cannot happen at the same time!



Conditional Probability: Updating Beliefs

Key Idea: Conditional probability helps us revise our beliefs when new information is available.

Conditional Probability Formula



$$P(B \mid A) = \frac{P(A \cap B)}{P(A)}$$

In words: What is the probability B Given that A has happened? how does this affect our belief in B ?

Conditional Probability: Where does the formula come from?

Remember!


$$P(B) = \frac{\text{\# of outcomes in } B}{\text{\# of all outcomes}} \Rightarrow P(B | A) = \frac{P(B \cap A)}{P(A)}$$



- We're **restricting** the sample space to only cases where event A has occurred.
- In a sense, we are **zooming in** on the sample space!


Relationship Between Independence and Mutual Exclusivity

- **Question:** If two events A and B are mutually exclusive, does that mean they are independent?



Relationship Between Independence and Mutual Exclusivity

- **Question:** If two events A and B are mutually exclusive, does that mean they are independent?



- **No!** If two events A and B are mutually exclusive, they are **NOT** independent.
- **Why?**
 - In this case, if we know event A happened, we know for a fact that B cannot happen. We are **gaining** information by knowing that event A happened.

Independence: Gaining No Extra Information

Definition: Two events A and B are **independent** if knowing that A occurred **does not** provide any information about whether B occurred.

1. Without Independence:

$$P(B | A) = \frac{P(A \cap B)}{P(A)}$$

Rewriting:

$$P(A \cap B) = P(A) \cdot P(B | A)$$

2. With Independence:

$$P(B | A) = P(B)$$

Rewriting:

$$P(A \cap B) = P(A) \cdot P(B)$$

Summary of Probability

(1) Addition Rule

General: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

Mutually Exclusive: $P(A \cup B) = P(A) + P(B)$

(2) Multiplication Rule

General: $P(A \cap B) = P(A) \cdot P(B | A)$

Independent: $P(A \cap B) = P(A) \cdot P(B)$

(3) Conditional Probability

General: $P(B | A) = \frac{P(A \cap B)}{P(A)}$

Independent: $P(B | A) = P(B)$

What is a Random Variable (RV)?

- ▶ **Random Variable (RV):** A *function* that *maps* all elements of a sample space to the real numbers
 - ▶ **Mathematically:** $\Omega \rightarrow \mathbb{R}$
 - ▶ **RV Notation:** $\mathcal{X}, X, \tilde{x}$
- ▶ **Intuition:** Sometimes, the sample space is not represented in numbers
 - ▶ The role of RVs is to map all those elements in the sample space Ω (that are, at times, not numbers) into real numbers \mathbb{R}
- ▶ **Relevance:**
 - 1 We will use RVs to describe probability distributions (next week!)
 - 2 We use RVs to deal with probabilities

Random Variables as Functions?

Example: Flipping a Coin ($\Omega \rightarrow \mathbb{R}$)

- ▶ **Scenario:** You flip a coin and want to reason about the outcome quantitatively.
- ▶ **Sample Space:**

Random Variables as Functions?

Example: Flipping a Coin ($\Omega \rightarrow \mathbb{R}$)

- ▶ **Scenario:** You flip a coin and want to reason about the outcome quantitatively.
- ▶ **Sample Space:** $\Omega = \{\text{Heads, Tails}\}$
- ▶ Define a Random Variable \mathcal{X} that "transforms" the elements in the sample space to real numbers:

Random Variables as Functions?

Example: Flipping a Coin ($\Omega \rightarrow \mathbb{R}$)

- ▶ **Scenario:** You flip a coin and want to reason about the outcome quantitatively.
- ▶ **Sample Space:** $\Omega = \{\text{Heads, Tails}\}$
- ▶ Define a Random Variable \mathfrak{X} that "transforms" the elements in the sample space to real numbers:

$$\mathfrak{X}(\omega) := \begin{cases} 0, & \text{if } \omega = \text{Heads} \\ 1, & \text{if } \omega = \text{Tails} \end{cases}$$

- ▶ **In words:** The RV \mathfrak{X} takes on the value 0 if the outcome is Heads and 1 if Tails
 - ▶ There you go! We have successfully mapped outcomes from the sample space Ω to real numbers \mathbb{R}

Random Variables and Probability

Example: Flipping a Coin

- ▶ **Question:** What is the probability of observing the outcome Heads (0)?


$$P_{\mathfrak{X}}(0) = P(\mathfrak{X} = 0) = \frac{\text{\# of outcomes in our event}}{\text{\# of all outcomes}} = \frac{1}{2} = 0.5$$

- ▶ **In words:** "What is the probability of the random variable \mathfrak{X} taking the value 0?"
 - ▶ **NOT:** "What is the probability of getting heads?"
- ▶ **Key Idea:** The way we deal with probability it through RVs.
 - ▶ We ask "what is the probability of the random variable taking the value(s)?"

Two Types of Random Variables (RVs)


(1) Discrete

- ▶ Countable finite numbers of values
- ▶ **Example:** Goals scored in a soccer match



(2) Continuous

- ▶ Can take an infinite number of values, intervals
- ▶ **Example:** Marathon time



Quiz + Discussion