Taylor Series

Hamza Kamal

January 29, 2024

Contents

1	\mathbf{Pre}	requisites:	2
	1.1	Taylor Series Approximation:	6
		Linear Approximation:	
		1.2.1 Example:	4
	1.3	Absolute Value Equations:	
		1.3.1 Example:	•
2	Bin	omial Series:	4
	2.1	Examples:	ļ
		2.1.1 Example 1:	ļ
		2.1.2 Example 2:	ļ
		2.1.3 Example 3:	

1 Prerequisites:

1.1 Taylor Series Approximation:

The approximated expansion for the Taylor Series of the function f(x) centered at x = a is given by the following formula:

Taylor Series Approximate Expansion
$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a^3 + \dots)$$

Where each constant is represented by an increasing derivative of the function f(x) at some value a.

1.2 Linear Approximation:

Linear approximation, also known as linearization or tangent line approximation, is a method used in calculus to estimate the value of a function near a specific point by using the equation of the tangent line at that point. The linear approximation provides a good approximation of the function's behavior in the vicinity of the chosen point. The linear approximation is

based on the idea that a sufficiently smooth and well-behaved function can be approximated by a linear function (a straight line) near a particular point. The equation of the tangent line to the graph of the function at a given point (a, f(a)) is given by:

Linear Approximation
$$L(x) = f(a) + f'(a)(x - a)$$

Here:

- L(x) is the linear approximation to the function,
- f(a) is the value of the function at the point x = a,
- f'(a) is the derivative of the function evaluated at x = a,
- x is the variable.

1.2.1 Example:

Find the value of $\sqrt{4.3}$ using the function $f(x) = \sqrt{x}$.

In this case the closest whole number that the function f(x) outputs is $f(4) = \sqrt{4}$ which equals 2. So, our a = 4, f(a) = 2 and $f'(a) = \frac{1}{2\sqrt{4}}$ which is $\frac{1}{4}$.

Therefore, our equation of the tangent line is $L(x) = f(a) + f'(a)(x-a) = 2 + \frac{1}{4}(x-4)$. Plugging in L(4.3) we get that $\sqrt{4.3} \approx 2.075$.

1.3 Absolute Value Equations:

Absolute Value Function

$$|x| = \begin{cases} x & \text{if } x \ge 0 \\ -x & \text{if } x < 0 \end{cases}$$

The absolute value function |x| returns the non-negative distance of x from zero on the number line.

Absolute Value Equation

$$|f(x)| = g(x)$$

Absolute value equations relate the absolute value of an expression f(x) to another expression or constant g(x). Solving Procedure:

- 1. Solve f(x) = g(x)
- 2. Solve f(x) = -g(x)

Solving each case may yield two sets of solutions, and validity is checked by substitution into the original equation.

1.3.1 Example:

Solve |2x - 5| = 7.

- 1. Isolate the Absolute Value Expression: 2x 5 = 7 or 2x 5 = -7
- 2. Apply the Absolute Value Property: 2x = 12 or 2x = -2
- 3. Solve for the Variable: Solutions: x = 6 and x = -1.
- 4. Check for Extraneous Solutions: Both solutions are valid.

2 Binomial Series:

Binomial Definition: The term "binomial" is derived from the Latin words "bi" which means two and "nomen" which means name. Which shows that binomials involve expressions with two terms, for example: (a + b).

Here are some examples of binomials:

$$(1+x)^{0} = 1$$
$$(1+x)^{1} = 1+x$$
$$(1+x)^{2} = 1+2x+x^{2}$$
$$(1+x)^{3} = 1+3x+3x^{2}+x^{3}$$
$$(1+x)^{4} = 1+4x+6x^{2}+4x^{3}+x^{4}$$

This pattern is derived from Pascals Triangle which lists out the coefficients for each term.




Figure 1: Pascals Triangle

However, if we wanted to calculate larger and larger powers of (1+x) you can see that this would become difficult and cumbersome. For example if you wanted to calculate $(1+x)^n$ where n was a very large value like 8 it would not be wise to write out 8 rows of Pascals Triangle and then find the coefficients to write the answer.

We have a name for the terms that appear in the Pascals Triangle which are also the powers (1+x). We call them *binomial coefficients*.

Binomial Coefficients

$$\binom{p}{k} = \frac{p!}{k!(p-k)!}$$

Note: The numbers $\binom{p}{0}$, $\binom{p}{1}$, $\binom{p}{2}$, ..., $\binom{p}{p}$ form the "pth" row on the Pascals Triangle.

More generally, if p is any real number and k >= 1 is an integer, then:

Binomial Coefficients

$$\binom{p}{k} = \frac{p(p-1)(p-2)\dots(p-k+1)}{k!}$$

The Special Symbol $\binom{p}{k}$ is pronounced "p choose k".

2.1 Examples:

2.1.1 Example 1:

If you had a class of 30 students and you had to pick a team of 3 students. This would be "30 choose 3" or $\binom{30}{3}$.

$$\binom{30}{3} = \frac{30!}{3!(27)!}$$

You can solve this equation to get the answer to the problem.

2.1.2 Example 2:

What if we wanted to find the 8th row of Pascals Triangle. This would be the same as doing:

$${8 \choose 0} = 1, {8 \choose 1} = \frac{8!}{1!(7)!}, {8 \choose 2} = \frac{8!}{2!(6)!}, \dots, {8 \choose n} = \frac{8!}{n!(8-n)!}$$
 Solving this you can calculate $(1+x)^8$

which equals $1 + 8x + 28x^2 + 56x^3 + 70x^4 + 56x^5 + 28x^6 + 8x^7 + x^8$.

2.1.3 Example 3:

Find $\binom{-3}{2}$ and $\binom{\frac{1}{2}}{3}$.

$$\binom{3}{-2} = \frac{-3(-3-1)}{2!}$$

Because k=2 there are only 2 factors on the top -3(-3-1).

$$\binom{3}{-2} = \frac{-3(-3-1)}{2!}$$
$$= \frac{(-3)(-4)}{2}$$
$$= 6$$

$$\binom{\frac{1}{2}}{3} = \frac{\frac{1}{2}(\frac{1}{2} - 1)(\frac{1}{2} - 2)}{3!}$$

Because k=3 there are only 3 factors on the top $\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2)$

Binomial Series

$$\sum_{k=0}^{\infty} \binom{p}{k} x^k = 1 + \sum_{k=1}^{\infty} \frac{p(p-1)(p-2)\dots(p-k+1)}{k!} x^k$$