



Faculty of Engineering and Architectural Science

Department of Computer and Electrical Engineering

Course Number	ELE532
Course Title	Signal and Systems I
Semester/Year	Fall 2025
Instructor	Dr.Beheshti

Report Title	Fourier Series Analysis using Matlab
--------------	---

Section No.	12
Report Submission Date	Nov 9 2025

Name	Student ID	Signature*
Hamza Malik	501112545	H.M
Kattoji Aneesh	501233584	A.K

*By signing above, you attest that you have contributed to this submission and confirm that all work you have contributed to this submission is your own work. Any suspicion of copying or plagiarism in this work will result in an investigation of Academic Misconduct and may result in a “0” on the work, an “F” in the course, or possibly more severe penalties, as well as a Disciplinary Notice on your academic record under the Student Code of Academic Conduct, which can be found online at: <http://www.ryerson.ca/senate/policies/pol60.pdf>.

A.1

$$D_n = \frac{1}{T} \int_0^T x_1(t) e^{-jn\omega_0 t} dt \quad x_1(t) = \cos \frac{3\pi}{10} t + \frac{1}{2} \cos \frac{\pi}{10} t$$

$$3\omega_0$$

general integral

$$\omega_0 = \frac{\pi}{10} \quad T = \frac{2\pi}{\pi/10} = 20$$

$$D_n = \frac{1}{T} \int_0^T \left[\frac{1}{2} e^{j3\omega_0 t} + \frac{1}{2} e^{-j3\omega_0 t} + \frac{1}{4} e^{j\omega_0 t} + \frac{1}{4} e^{-j\omega_0 t} \right] e^{-jn\omega_0 t} dt$$

$$D_n = \frac{1}{T} \left[\frac{1}{2} \int_0^T e^{j(3-n)\omega_0 t} dt + \frac{1}{2} \int_0^T e^{j(-3-n)\omega_0 t} dt + \frac{1}{4} \int_0^T e^{j(1-n)\omega_0 t} dt + \frac{1}{4} \int_0^T e^{j(-1-n)\omega_0 t} dt \right]$$

for each integral

$$\int_0^T e^{jk\omega_0 t} dt = T \text{ if } k=0 \quad \text{or} \quad \int_0^T e^{jk\omega_0 t} dt = \frac{e^{jk\omega_0 T} - 1}{jk\omega_0} = 0 \text{ if } k \neq 0$$

where k is some int

so n must be such that added w/ can't n

$$k-n=0$$

$$D_n = \frac{1}{T} \left[\frac{1}{2}(T) \delta_{n,3} + \frac{1}{2}(T) \delta_{n,-3} + \frac{1}{4}(T) \delta_{n,1} + \frac{1}{4}(T) \delta_{n,-1} \right]$$

$$D_{\pm 3} = \frac{1}{2} \quad D_{\pm 1} = \frac{1}{4}$$

A1: Calculation for the value of D_n for function $x_1(t)$

4.2 general integral

$$D_n = \frac{1}{T} \int_{-T/2}^{T/2} x_2(t) e^{-jn\omega_0 t} dt$$

$$D_n = \frac{1}{T} \int_{-5}^5 e^{-jn\omega_0 t} dt = \frac{1}{T} \frac{e^{-jn\omega_0 (5)} - e^{jn\omega_0 (5)}}{-jn\omega_0} = \frac{2}{T\omega_0} \sin(n\omega_0 \cdot 5), n \neq 0$$

DC term

$$D_0 = \frac{1}{T} \int_{-5}^5 1 dt = \frac{10}{T}$$

Simplify

$$D_n = \frac{2}{Tn(2\pi/\pi)} \sin(n\omega_0 \cdot 5) = \frac{1}{\pi n} \sin(n\omega_0 \cdot 5), n \neq 0$$

for $x_2(t)$

$$T=20 \quad \omega_0 = \frac{2\pi}{20} = \frac{\pi}{10} \quad 5n\omega_0 = \frac{\pi n}{2}$$

$$D_n = \frac{10}{20} = \frac{1}{2}, \quad n=0 \quad D_n = \frac{1}{\pi n} \sin\left(\frac{\pi n}{2}\right) \quad n \neq 0$$

for $x_3(t)$

$$T=40 \quad \omega_0 = \frac{2\pi}{40} = \frac{\pi}{20}$$

$$5n\omega_0 = \frac{\pi n}{4}$$

$$D_n = \frac{10}{40} = \frac{1}{4}, \quad n=0 \quad D_n = \frac{1}{\pi n} \sin\left(\frac{\pi n}{4}\right) \quad n \neq 0$$

A2: Calculations for the value of D_n for function $x_2(t)$, $x_3(t)$, subbed in $n=0$ to the existing equation and found that $n=1/2$ for $x_2(t)$ and $n=1/4$ for $x_3(t)$

```

% A.3
function Dn = compute_Dn(n_values)
% compute_Dn - Returns the Exponential Fourier Series coefficients Dn
% for the signal:
%   x1(t) = cos(3πt/10) + 0.5*cos(πt/10)
%
% Usage:
%   Dn = compute_Dn(n_values)
%
% Example:
%   Dn = compute_Dn(-5:5)
% Initialize Dn as zeros
Dn = zeros(size(n_values));
% Loop through the user-specified n values
for k = 1:length(n_values)
    n = n_values(k);
    if abs(n) == 1
        Dn(k) = 1/4;           % From 0.5*cos(πt/10)
    elseif abs(n) == 3
        Dn(k) = 1/2;           % From cos(3πt/10)
    else
        Dn(k) = 0;            % All other harmonics are zero
    end
end
end



```



A3: Code for A3 that computes d_n and returns the fourier coefficients

```

% A.4 - Fourier Coefficients and Reconstruction
% --- x1(t) ---
% a) Compute and plot |Dn| and Dn for n = -5:5
figure(1);
n = (-5:5);
D_func = Dn(1);           % this selects the Dn definition
D_n = D_func(n);% Evaluate coefficients for chosen n
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% b) Increase harmonic range
figure(4);
n = (-20:20);
D_func = Dn(1);
D_n = D_func(n);
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% c) Increase to n = -50:50
figure(7);
n = (-50:50);
D_func = Dn(1);
D_n = D_func(n);
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% d) increases n = -500:500
figure(10);
n = (-500:500);
D_func = Dn(1);
D_n = D_func(n);
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');


```








A4: Code for $x_1(t)$ that computes d_n and returns the fourier coefficients and its respective graphs (graph a, graph b, graph c, graph d)

```
% --- x2(t) ---
% a) Compute coefficients for x2(t)
figure(2);
n = (-5:5);
D_func = Dn(2);
D_n = D_func(n);
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% b) n = -20:20
figure(5);
n = (-20:20);
D_func = Dn(2);
D_n = D_func(n);
D_n(n==0)=1/2; % we took the limit at n -> 0 and it is 1/2
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% c) n = -50:50
figure(8);
n = (-50:50);
D_func = Dn(2);
D_n = D_func(n);
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% d) n = -500:500
figure(11);
n = (-500:500);
D_func = Dn(2);
D_n = D_func(n);
D_n(n==0)=1/2;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel(|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel(\angle D_n [rad]);
%
```











A4: Code for $x_2(t)$ that computes d_n and returns the fourier coefficients and its respective graphs (graph a, graph b, graph c, graph d)

```
% a) Compute coefficients for x3(t)
figure(3);
n = (-5:5);
D_func = Dn(3);
D_n = D_func(n);
D_n(n==0)=1/4;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% b) n = -20:20
figure(6);
n = (-20:20);
D_func = Dn(3);
D_n = D_func(n);
D_n(n==0)=1/4;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% c) n = -50:50
figure(9);
n = (-50:50);
D_func = Dn(3);
D_n = D_func(n);
D_n(n==0)=1/4;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
% d) n = -500:500
figure(12);
n = (-500:500);
D_func = Dn(3);
D_n = D_func(n);
D_n(n==0)=1/4;
subplot(1,2,1); stem(n,abs(D_n),'.k'); xlabel('n'); ylabel('|D_n|');
subplot(1,2,2); stem(n,angle(D_n),'.k'); xlabel('n'); ylabel('\angle D_n [rad]');
```











A4: Code for $x_3(t)$ that computes d_n and returns the fourier coefficients and its respective graphs (graph a, graph b, graph c, graph d)



```
% A5(Dn): Reconstruct signal from Fourier coefficients
function [D] = A5(Dn)
    n = -200:200;
    D = Dn;
    t = -300:300;
    w = pi * 0.1;
    x = zeros(size(t));           % Initialize reconstructed signal
    % Compute partial Fourier sum
    for i = 1:length(n)
        x = x + D(i) * exp(ij * n(i) * w * t);
    end
    % Plot reconstructed signal
    figure;
    plot(t, real(x), 'k');
    xlabel('t (sec)');
    ylabel('x(t)');
    axis([-300 300 -1 2]);
    title('Reconstructed Signal from Fourier Coefficients');
    grid on;
end
```



A5: *Code for A5 that computes dn and reconstructs original time-domain signal*















A6: Graphs for the reconstructed time-domain signals $x_1(t)$, $x_2(t)$ and $x_3(t)$ with the fourier coefficients generated in Problem A4

B1.

$$x_1(t) = \cos\left(\frac{3\pi t}{10}\right) + \frac{1}{2} \cos\left(\frac{\pi t}{10}\right)$$

$$\omega_{o1} = \frac{3\pi}{10} \quad \omega_{o2} = \frac{\pi}{10}$$

$$\omega_o = \frac{6\pi F}{LCM} = \frac{\pi}{10} \text{ rad/s}$$

$x_2(t)$

$$T=20$$

$$\omega_o = \frac{2\pi}{T} = \frac{\pi}{10} \text{ rad/s}$$

$x_3(t)$

$$T=40s$$

$$\omega_o = \frac{2\pi}{T} = \frac{\pi}{20} \text{ rad/s}$$

B1: Calculation for B2 values.

B2:

The main differences between $x_1(t)$ and $x_2(t)$ are how the sin functions in $x_1(t)$ and sin functions in $x_2(t)$ are used. Additionally, $x_1(t)$ contains four distinct Fourier series coefficients, while $x_2(t)$ contains infinite fourier series as n grows to infinity. We also note that $x_2(t)$ has a DC value signal, a_0 , which is 0.5, while $x_1(t)$ does not have a DC value signal. Otherwise, the two signals share a fundamental frequency.

B3:

Since the period of $x_3(t)$ is greater, it will be inversely proportional to its fundamental frequency, which means that $x_3(t)$ has a smaller fundamental frequency compared to $x_2(t)$. Additionally, a shorter period will expand the change values of fourier coefficients. And vice versa; a greater period will compress the change in values of the fourier coefficients, as seen in signal $x_3(t)$.

B4:

$$\begin{aligned}
 & \text{By, } \frac{1}{20} \int_{-10}^{10} x_4(t) e^{-jn(\frac{\pi}{10})t} dt \\
 &= -\frac{1}{20} \int_{-10}^{-5} e^{-jn(\frac{\pi}{10})t} + \frac{1}{20} \int_{-5}^{5} e^{-jn(\frac{\pi}{10})t} - \frac{1}{20} \int_{5}^{10} e^{-jn(\frac{\pi}{10})t} \\
 &= -\frac{1}{20} \left[\frac{-10e^{-jn(\frac{\pi}{10})t}}{jn\pi} \right]_{-10}^5 + \frac{1}{20} \left[\frac{-10e^{-jn(\frac{\pi}{10})t}}{jn\pi} \right]_{-5}^5 - \frac{1}{20} \left[\frac{-10e^{-jn(\frac{\pi}{10})t}}{jn\pi} \right]_{5}^{10} \\
 &= \cancel{-\frac{1}{2} \frac{e^{jn(\frac{\pi}{2})t}}{jn\pi}} - \cancel{\frac{1}{2} \frac{e^{jn\pi t}}{jn\pi}} - \cancel{\frac{1}{2} \frac{e^{-jn(\frac{\pi}{2})t}}{jn\pi}} + \cancel{\frac{1}{2} \frac{e^{jn(\frac{\pi}{2})t}}{jn\pi}} \\
 &\quad + \cancel{\frac{1}{2} \frac{e^{-jn\pi t}}{jn\pi}} - \cancel{\frac{1}{2} \frac{e^{jn\pi t}}{jn\pi}} \\
 &= 0
 \end{aligned}$$

∴ D_n of $x_4(t)$ is 0.

B5:

It is known that $x_1(t)$ has a finite number of fourier coefficients, while for $x_2(t)$, there are infinite numbers of fourier coefficients. Thus, increasing the number of fourier coefficients will increase the accuracy of the reconstructed signal for $x_2(t)$, but $x_1(t)$ will stay the same.

B6:

Referencing Problem B5, we only need four fourier coefficients to perfectly reconstruct $x_1(t)$. For $x_2(t)$ and $x_3(t)$, we will need an infinite number of fourier coefficients for a perfect reconstruction.

B7:

For signals that require a large amount of finite fourier coefficients, it is not recommended to be used to reconstruct a fourier series signal due to space and time limitations. Such as signals $x_2(t)$ and $x_3(t)$, they require an infinite number of fourier coefficients, so unless there are constraints set on the number of fourier coefficients required, it will use a lot of space and time. For signals such as $x_1(t)$, it is possible since it only requires four fourier coefficients to replicate a signal.