





VERIFY PATIENT

14-09-2021 12:20 AM

If you wish to confirm the validity of this report, call 051-2120740.

ISO 9001: 2015 Certified

LAB. REPORT

TRACK ONLINE



Flight No

http://report.thebiogenelab.com/report.aspx?id=49-215355

Patient's Name: MOHSIN TAHIR Age / Sex: 59 Years / Male

Lab No. : 21133584 Sample Collection Date : 13-09-2021 12:14 AM

Panel: Cc Ali Sabar Report Date: 13-09-2021 07:40 AM

NIC #: Ref By: BRITISH AIRWAYS

Flight Date

Passport: 539336456 Phone:

Destination: LONDON Ticket No: KIGPWX

http://report.thebiogenelab.com User: 49-215355 Password: 2569

BA 260

2019-nCoV (Corona) Virus PCR

Specimen: Nasopharyngeal & Oropharyngeal Swab

Result: Not Detected

METHOD (TaqMan®Real Time RT-PCR Viral Load)

The 2019-n CoV (Corona) virus quantitation uses a TaqMan® Real Time Polymerase Chain Reaction (RT-PCR) technology to detect and quantify COVID -19 RNA. Under optimal PCR and can detect less than 100 copies of target template. The test is performed on Sa Cycler 96, onditions genesig 2019-n CoV detection kits have very high priming efficiencies of >90% Sacace Biotechnologies, Italy. Correlation of these results with clinical features and other laboratory findings is highly recommended.

Comments:

- 1. Clinical Correlation is highly recommended in case of Positive / Negative Result
- 2. One or more negative results do not rule out the possibility of COVID-19 virus infection
- 3. A number of factors could lead to a negative result in an infected individual including, Poor Quality of specimen, Containing little patient materialthe specimen was not handled and shipped appropriately, the specimen is collected late or very early in the infection, technical reasons inherent in the test, reasons inherent in the test, e.g Virus Mutations in primer probe target region.
- 4. If a negative result is obtaining from a patient with high index of suspicion for SARS-CoV 2 virus infection, particularly when only upper respiratory tract specimens were collected, additional specimens, including from the lower respiratory tract if possible, should be collected and tested.

5. The viral genes targeted can vary from lab to lab (The viral genes targeted so far include the N,E, S and RdRP genes)

Muhammad Noman Sa Biotechnologist Ph.D Biotechnology (IP)

Electronically generated report. No signature(s) required.