1. The probability of a leap year selected at random contain 53					
Sunday is:	/1 \	((1) =0 (0.6 =		
(a) 53/366					
•			A marble is drawn at		
random. The prob	-	-			
* *		(c) 0/5	· ·		
-	•		0.85. What is the		
probability that it			(1)		
			(d) none of these		
_	-		cted from the numbers		
(1, 2, 3,,15)	_		4.00		
	• •	(c) 2/15	* *		
5. What are the t					
		(c) 8			
-	-	e number selec	ted at random from the		
numbers (1,2,3,					
(a) 12/35	(b) 11/35	5 (c) 13/3	5 (d) none of these		
7. The sum of the	_				
) 0 (d) no			
8. The following	probabilities	are given; cho	ose the correct answer		
for that which is r	ot possible.				
(a) 0.15	(b) 2/7	(c) 7/5	(d) none of these.		
			han the probability of		
getting at least tw	o heads, is:				
(a) 1/4	(b) 3/8	(c) $\frac{1}{2}$	(d) 1/8		
10. À letter is ch					
♦ ASSASSINATIO	N�. The pro	bability that th	e letter chosen has:		
			(d) none of these.		
(1)	() , ,	(-)			
11. A dice is throw	wn. Find the p	robability of g	etting an even number.		
(A) 2/3	_	(C) 5/6	_		
(* ') =/ '	(-)	(3) 3, 3	., =		
12. Two coins are thrown at the same time. Find the probability of					
12. Two coins are	thrown at the	e same time. F	ind the probability of		
12. Two coins are getting both head		e same time. F	ind the probability of		
	s.		ind the probability of		

13. Two dice are thrown simultaneously. The probability of getting a sum of 9 is:

(A) 1/10	(B) 3/10	(C)	1/9	(D) 4/9			
		ered from 1	to 100. Fi	nd the probability of			
	rime number. (B) 27/50	(C) 1,	/4	(D) 29/100			
of drawing blue balls in	a blue ball is on a bag is:	double that	of a red ba	palls .If the probability all, then the number of			
(A) 5	<mark>(B)</mark> 10	(C) 15	(D) 2	0			
taken out a non-defect	16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:						
(A) 143/150	O (B) 14	7/150	(C) 1/25	(D) 1/50			
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100							
18. What is (A) 1/7	-		n g 53 Mond /7	lays in a leap year? (D) 7/366			
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
equally like 1,2,312	ly to come to	rest pointing that	ng to one o t it will poir	arrow which is of the number of to an odd number is: D) 5/12			
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4							

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is: (A) 364/365 (B) 31/365 (D) 1/365 (D) 1/133225							
(A) 304/303	(b) 31/303	1/303	(D) 1/133223				
2. Then the pr	x is chosen at ra cobability that x ² < 2/5 (C) 3/5	< 2 is?	umbers -2, -1, 0 , 1,				
a marble is dra red is 2/3, the	24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
Then the prob		multiple of 3 and	0 natural numbers. I 4 is:				
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
	red by batsman ir ne standard devia	_	hes are 50, 70, 82,				
•		c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
	b) 13, 18		d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$	b) $\frac{1}{3}$	c) ¹ / ₄	d) $\frac{1}{6}$				
a) 8 t	ate between 0 an o) 7	d 3. The value of	E(X²) is) 9				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of the following values, which one is not possible in probability?					
	b) ∑ x d) P(x	P(x) = 3 () = -0.5			
	2 and E(z) = 4 b) 6) Insufficient	data
34.The cova	ariance of two	independen	t random va	riable is	
a) 1	. b) 0	c) - 1	d) Undefined	
	= k ² – 8 then, b) 1			l) Insufficien	t data
, ,	0.5 and x = 4, b) 0.5	, ,		2	
37.In a disc is always?	rete probabilit	y distributio	n, the sum c	of all probabi	ilities
•	b) Infinite	c) 1	d)	Undefined	
38.If the pr variance.	obability of hit	tting the targ	jet is 0.4, fir	nd mean and	I
	b) 0.6, (0.24	c) 0.4, 0.1	6 d) 0.6	5, 0.16
39. If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6					
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 					

c) 5

d) 7

a) 3

b) 4

		and varia and varia		•				
42. \a) E(
	43.Mean of a random variable X is given by a) E(X) b) E(X2) c) E(X2) - (E(X))2 d) (E(X))2							
	44.Mean of a constant 'a' is a) 0							
45. V	45.Variance of a constant 'a' is . a) 0							
46.Find the mean and variance of X?								
г 	Х	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		

47. Find the expectation of a random variable X?

b) 3, 4/3

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5	,	c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

c) 2, 2/3

d) 3, 2/3

a) 2, 4/3



d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}