Selections of **Inspirations**

Han-bai Wang Typeset by LaTeX

2022 Spring

Abstract

The significance of probability theory is to describe the regularity shown by a large number of random factors.

— Kolmogorov

Contents

1 Set Theory **3**

2 Analysis **4**

3 Algebra **41**

4 Topology **46**

5 Complex Analysis **49**

6 Probability **53**

7 Stochastic Processes **117**

8 Functional Analysis **149**

9 Partial Differential Equations **155**

10 Econometrics **159**

11 Mathemetical Statistics **161**

12 Semiparametric Model **162**

References **169**

§ 1 Set Theory

QUESTION 1.1 *Suppose $\{A_i\}_{i\in I}$ and $\{B_j\}_{j\in I}$ are two set sequences. Then*

$$\left(\bigcap_{i\in I} A_i\right) \cap \left(\bigcap_{j\in I} B_j\right) \subset \bigcap_{i\in J}(A_i \cap B_i),$$
$$\left(\bigcup_{i\in I} A_i\right) \cup \left(\bigcup_{j\in I} B_j\right) \subset \bigcup_{i\in J}(A_i \cup B_i).$$

Proof. We start from the first inequality.

$$\left(\bigcap_{i\in I} A_i\right) \cap \left(\bigcap_{j\in I} B_j\right) = \bigcap_{i\in I}\left(A_i \cap \bigcap_{j\in I} B_j\right) \subset \bigcap_{i\in J}(A_i \cap B_i).$$

The second inequality follows from

$$\left(\bigcup_{i\in I} A_i\right) \cup \left(\bigcup_{j\in I} B_j\right) = \bigcup_{i\in I}\left(A_i \cap \bigcap_{j\in I} B_j\right)$$

□

QUESTION 1.2 *Suppose $\{A_n\}$ is a set sequence. Then*

$$\left(\bigcup_{n=1}^{m} A_n\right)' = \bigcup_{n=1}^{m} A_n', \quad \overline{\bigcup_{n=1}^{m} A_n} = \bigcup_{n=1}^{m} \overline{A_n}.$$

QUESTION 1.3 *Suppose A is a countable set, then $\mathscr{A} := \{F \subset A : F \text{ is finite}\}$ is countable.*

Proof. Wirte $A = \{a_m\}$. We use $|F|$ to denote the cardinal number of F. Then we have the following decomposition:

$$\mathscr{A} = \sum_{n=0}^{\infty}\{F \in \mathscr{A} : |F| = n\} =: \sum_{n=0}^{\infty} \mathscr{A}_n.$$

So we only need to verify $\mathscr{A}_n$ is at most countable for each $n \geq 0$. However, we know for each $n \geq 0$,

$$\varphi_n : \mathscr{A}_n \to \mathbb{N}^n : \{a_{m_k}\}_{k=1}^{n} \to (m_1, \cdots, m_k).$$

It is easy to verify φ_n is a one-to-one mapping. So $\mathscr{A}_n$ is countable and thus $\mathscr{A}$ is countable. □

§ 2 Analysis

QUESTION 2.1 *Prove the following inequality:*

$$|x \wedge C - y \wedge C| \leq |x - y|, \quad |x \vee C - y \vee C| \leq |x - y|,$$

where $x, y, C \in \mathbb{R}$.

Proof. 1. (a) When $x, y \leq C$, then $|x \wedge C - y \wedge C| = |x - y|$.

(b) When $x, y \geq C$, then $|x \wedge C - y \wedge C| = |C - C| = 0 \leq |x - y|$.

(c) Without loss of generality, we assume $x \leq C < y$, then $|x \wedge C - y \wedge C| = |x - C| \leq |x - y|$.

In conclusion, the first inequality holds.

2. (a) When $x, y \leq C$, then $|x \vee C - y \vee C| = |C - C| = 0 \leq |x - y|$.

(b) When $x, y \geq C$, then $|x \vee C - y \vee C| = |x - y|$.

(c) Without loss of generality, we assume $x \leq C < y$, then $|x \vee C - y \vee C| = |C - y| \leq |x - y|$.

In conclusion, the second inequality holds. □

QUESTION 2.2 *For any nonnegative (nonpositive) measurable function* $f(x) \in \mathscr{B}[a, \infty)$ *and* $t \geq a$*, we have*

$$\int_a^{b \wedge t} f(x)dx \leq (\geq) \int_a^b f(x \wedge t)dt.$$

Proof. If $a \leq b \leq t$, then the above two sides are actually equal. If $b > t$, then

$$\begin{aligned}\int_a^b f(x \wedge t)dt &= \int_a^t f(x)dt + f(t)(t - b) \\ &= \int_a^{b \wedge t} f(x)dt + f(t)(t - b) \\ &\geq (\leq) \int_a^{b \wedge t} f(x)dx.\end{aligned}$$

So the desired results follows. □

QUESTION 2.3 *Suppose* $f, g : U \to \mathbb{R}$ *and* $U \subset \mathbb{R}^n$ *is an open set. Assume exists* $C > 0$ *such that* $f(x) + g(x) \leq C$ *for each* $x \in U$*, then*

$$\sup_{x \in U} f(x) + \sup_{x \in U} g(x) \leq 2C.$$

Proof. The inequality $f(x) + g(x) \leq C, \forall x \in U$ yields

$$f(x) \leq C, \quad g(x) \leq C, \quad \forall x \in U.$$

So $\sup_{x\in U} f(x) \leq C$ and $\sup_{x\in U} g(x) \leq C$. This is what we desire. □

QUESTION 2.4 *Suppose $f : \mathbb{R} \to \mathbb{R}$ is generalized intergral, i.e.,*

$$\int_1^\infty f(x)dx = \lim_{\ell\to\infty} \int_1^\ell f(x)dx \in \mathbb{R}.$$

Use an example to show $f(x) \not\to 0, x \to \infty$ might happen.

Proof. Consider

$$f(x) = \frac{\sin x^2}{x}, \quad f'(x) = 2\cos x^2 - \frac{\sin x^2}{x^2}.$$

Notice $f' \in C^\infty[1, \infty)$, so $\forall \ell > 1, f' \in R[1, \ell]$. And

$$\begin{aligned}\int_1^\infty f'(x)dx &= \lim_{\ell\to\infty} \int_1^\ell f'(x)dx = \lim_{\ell\to\infty} [f(\ell) - f(1)] \\ &= \lim_{\ell\to\infty} \left(\frac{\sin \ell^2}{\ell} - \sin 1\right) = -\sin 1 \in \mathbb{R}.\end{aligned}$$

However

$$\lim_{x\to\infty} f'(x) = \lim_{x\to\infty} \left(2\cos x^2 - \frac{\sin x^2}{x^2}\right)$$

does not exist. □

QUESTION 2.5 *Suppose f is differentiable on $[a, b]$ (f has the left and right derivatives on endpoints). If $\forall \varepsilon > 0, \exists \delta > 0$, when $0 < |h| < \delta$, for all $x \in [a, b]$, we have*

$$\left|\frac{f(x + h) - f(x)}{h} - f'(x)\right| < \varepsilon,$$

then we say f is uniformly differentiable on $[a, b]$. Prove $f' \in C[a, b]$ iff f is uniformly differentiable.

Proof. For all $x \in [a, b]$, write

$$f_h(x) := \frac{f(x + h) - f(x)}{h},$$

then $f_h \in C[a, b]$.

1. If $f' \in C[a,b]$, then f' is uniformly continuous, then $\forall \varepsilon > 0, \exists \delta > 0, \forall \xi, x \in [a,b], |\xi - x| < \delta$, then $|f'(\xi) - f'(x)| < \varepsilon$. For all $x \in [a,b]$, when $0 < |h| < \delta$, by Lagrange theorem of mean, exist ξ between x and $x+h$,

$$\left| \frac{f(x+h) - f(x)}{h} - f'(x) \right| = |f'(\xi) - f'(x)| < \varepsilon,$$

which means f is uniformly differentiable.

2. If f is uniformly differentiable, then $\forall \varepsilon > 0, \exists \delta_1 > 0$, when $|h| < \delta_1, \forall x \in [a,b]$, we have

$$|f_h(x) - f'(x)| < \varepsilon/3.$$

Since f_h is continuous, $\forall \varepsilon > 0, \exists \delta > 0$, when $|\Delta| < \delta$, we have

$$|f_h(x+\Delta) - f_h(x)| < \varepsilon/3.$$

Finally, $\forall \varepsilon > 0, \exists \delta > 0$, when $|\Delta| < \delta$, we obtain

$$\begin{aligned} &|f'(x+\Delta) - f'(x)| \\ \leq &|f'(x+\Delta) - f_h(x+\Delta)| + |f_h(x+\Delta) - f_h(x)| + |f_h(x) - f'(x)| \\ < &\varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon, \end{aligned}$$

which implies $f' \in C[a,b]$. □

QUESTION 2.6 *Suppose $\sum_i a_i$ is a real series and $\sum_j A_j$ is a real series obtained by adding some brackets in some items of $\sum_i a_i$. If for every A_j, the addends of A_j are all positive or all negative, then prove $\sum_i a_i$ converges iff $\sum_j A_j$ converges, and when they converge, we have $\sum_i a_i = \sum_j A_j$.*

Proof. 1. Notice $\sum_{j=1}^n A_j, n \geq 1$ is just a sub-sequence of $\sum_{i=1}^n a_i, n \geq 1$, if $\sum_i a_i$ converges, then $\sum_j A_j$ must converges and $\sum_j A_j = \sum_i a_i$.

2. On the contrary, if $\sum_j A_j$ converges, then $\forall \varepsilon > 0, \exists N_1 \in \mathbb{N}, \forall n > N_1, p \in \mathbb{N}$, we have

$$\left| \sum_{j=n}^{n+p} A_j \right| < \varepsilon.$$

We write m_j to denote the number of items bracketed in $A_j, j \geq 1$ and write $a_i \in A_j$ to denote a_j is an addend of A_j. Consider $N_2 := \sum_{j=1}^{N_1} m_j$, then $\forall n > N_2, p \in \mathbb{N}, \exists \alpha, \beta \in \mathbb{N}$, s.t. $a_n \in A_\alpha$ and $a_{n+p} \in A_\beta$. Write

$n_\alpha := \min\{n : a_n \in A_\alpha\}$ and $n_\beta := \max\{n : a_n \in A_\beta\}$. Observe $\alpha > N_1$, then $\min\{|A_\alpha|, |A_\beta|, |\sum_{j=\alpha}^{\beta} A_j|\} < \varepsilon$. Whence

$$\begin{aligned}\left|\sum_{i=n}^{n+p} a_i\right| &= \left|\sum_{j=\alpha}^{\beta} A_j - \sum_{i=n_\alpha}^{n-1} a_i - \sum_{i=n+p+1}^{n_\beta} a_i\right| \\ &\le \left|\sum_{j=\alpha}^{\beta} A_j\right| + \left|\sum_{i=n_\alpha}^{n-1} a_i\right| + \left|\sum_{i=n+p+1}^{n_\beta} a_i\right| \\ &\le \left|\sum_{j=\alpha}^{\beta} A_j\right| + |A_\alpha| + |A_\beta| < 3\varepsilon,\end{aligned}$$

where the last $\le$ comes from all the addends of A_j are all positive or all negative. So this formula means $\sum_i a_i$ converges. Notice $\sum_{j=1}^{n} A_j, n \ge 1$ is just a sub-sequence of $\sum_{i=1}^{n} a_i, n \ge 1$, if $\sum_i a_i$ converges, then $\sum_j A_j$ must converges and $\sum_j A_j = \sum_i a_i$. □

QUESTION 2.7 *Suppose $f_0 : D \to \mathbb{R}$ is an increasing function, where D is a dense subset on $\mathbb{R}$. We define*

$$f(x) := \inf_{x<t\in D} f_0(t), \quad \forall x \in \mathbb{R}.$$

Prove the uniformly continuity of f_0 implies the uniformly continuity of f.

Proof. Since f_0 is uniformly continuous, then $\forall \varepsilon > 0, \exists \delta > 0$, for all $t_1, t_2 \in D$ and $|t_1 - t_2| < 3\delta$, we have

$$|f_0(t_1) - f_0(t_2)| < \varepsilon/2.$$

For all $x_1, x_2 \in \mathbb{R}$ and $|x_1 - x_2| < \delta$, we can find a sequence $\{t_1^{(n)}\}$ and $\{t_2^{(n)}\}$ in D, such that $t_j^{(n)} \searrow x_j, j = 1, 2$. Exist $N \in \mathbb{N}_+$, for all $n > N$, we have $|t_j^{(n)} - x_j| < \delta, j = 1, 2$. This implies $\forall n > N$,

$$|t_1^{(n)} - t_2^{(n)}| \le |t_1^{(n)} - x| + |x - y| + |y - t_2^{(n)}| < 3\delta.$$

Hence for all $n > N$,

$$|f_0(t_1^{(n)}) - f_0(t_2^{(n)})| < \varepsilon/2.$$

Let $n \to \infty$ in the above formula and observe the fact that

$$f(x) = \inf_{x<t\in D} f_0(t) = \lim_{n\to\infty} f_0(t_1^{(n)}),$$

then we will obtain

$$|f(x_1) - f(x_2)| \leq \varepsilon/2 < \varepsilon.$$

This provides the desired result. □

QUESTION 2.8 *Suppose* $\boldsymbol{x} = \boldsymbol{f}(\theta_1, \cdots, \theta_{n-1})$ *is a regular surface on* $\mathbb{R}^n$*. Then exist a coordinate transformation* $\boldsymbol{y} = \varphi(\boldsymbol{x})$*, such that the surface is* $y_n = 0$ *under the new coordinate.*

Proof. Since $\mathbf{x} = \mathbf{f}(\theta_1, \cdots, \theta_{n-1})$ is a regular surface, then

$$\begin{pmatrix} \partial_1 f_1 & \partial_2 f_1 & \cdots & \partial_{n-1} f_1 \\ \partial_1 f_2 & \partial_2 f_2 & \cdots & \partial_{n-1} f_2 \\ \partial_1 f_3 & \partial_2 f_3 & \cdots & \partial_{n-1} f_3 \\ \vdots & \vdots & \vdots & \vdots \\ \partial_1 f_{n-1} & \partial_2 f_{n-1} & \cdots & \partial_{n-1} f_{n-1} \\ \partial_1 f_n & \partial_2 f_n & \cdots & \partial_{n-1} f_n \end{pmatrix}_{n\times(n-1)}$$

has the rank of $n-1$. Write $\mathbf{x}' = (x_1, \cdots, x_{n-1})$ and $\mathbf{f}' = (f_1, \cdots, f_{n-1})$, then $\mathbf{x} = \mathbf{f}'(\theta_1, \cdots, \theta_{n-1})$. Suppose

$$\frac{\partial(f_1, \cdots, f_{n-1})}{\partial(\theta_1, \cdots, \theta_{n-1})} = \begin{pmatrix} \partial_1 f_1 & \partial_2 f_1 & \cdots & \partial_{n-1} f_1 \\ \partial_1 f_2 & \partial_2 f_2 & \cdots & \partial_{n-1} f_2 \\ \partial_1 f_3 & \partial_2 f_3 & \cdots & \partial_{n-1} f_3 \\ \vdots & \vdots & \vdots & \vdots \\ \partial_1 f_{n-1} & \partial_2 f_{n-1} & \cdots & \partial_{n-1} f_{n-1} \end{pmatrix}_{(n-1)\times(n-1)}$$

has the rank $n-1$. Then $\mathbf{f}^{-1}$ exists and $(\theta_1, \cdots, \theta_{n-1}) = \mathbf{f}^{-1}(\mathbf{x}')$. We can plug this into $x_n = f_n(\theta_1, \cdots, \theta_{n-1})$ to obtain

$$x_n = f_n(\mathbf{f}^{-1}(\mathbf{x}')) := h(x_1, \cdots, x_{n-1}).$$

So the surface has another expression: $x_n = h(x_1, \cdots, x_{n-1})$ with parameters $(x_1, \cdots, x_{n-1})$.

Consider the following mapping:

$$\varphi : \begin{cases} y_1 = x_1 \\ y_2 = x_2 \\ \quad \vdots \\ y_n = x_n - h(x_1, \cdots, x_{n-1}) \end{cases}$$

and

$$\det D\varphi(\mathbf{x}) = \det \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ -\partial_1 h & -\partial_2 h & \cdots & -\partial_{n-1} h & 1 \end{pmatrix} = 1 \neq 0.$$

Thus this is indeed a coordinate transformation and the surface in this new coordinate is

$$y_n = 0,$$

which is what we desire. □

QUESTION 2.9 *Suppose $\{f_n\}$ is a sequence of measurable functions on $(\Omega, \mathscr{F})$. Show*

$$\lim_{n\to\infty}\left(\sup_{k\leq n} f_k\right) = \sup_{n\in\mathbb{N}} f_n.$$

Proof. Since $\forall n \in \mathbb{N}, \sup_{k\leq n} f_k \leq \sup_n f_n$, let $n \to \infty$, we will obtain

$$\lim_{n\to\infty}\left(\sup_{k\leq n} f_k\right) \leq \sup_{n\in\mathbb{N}} f_n.$$

On the contrary, for all $x \in \Omega$, exists a sequence $\{n_\ell\}$, s.t. $f_{n_k}(x) \uparrow \sup_{n\in\mathbb{N}} f(x)$. Moreover

$$\sup_{k\leq n_k} f_k(x) \geq f_{n_k}(x).$$

Let $\ell \to \infty$ in the above inequality, we will obtain

$$\sup_{n\in\mathbb{N}} f_n(x) \geq \lim_{\ell\to\infty}\left(\sup_{k\leq n_\ell} f_k(x)\right) \geq \lim_{\ell\to\infty} f_{n_\ell}(x) = \sup_{n\in\mathbb{N}} f_n(x).$$

Because of the monotonicity of $\sup_{k\leq n} f_k$ with respect to $n \in \mathbb{N}$, then $\sup_{k\leq n} f_k$ and $\sup_{k\leq n_\ell} f_k$ share the same limit. Thus

$$\lim_{n\to\infty}\left(\sup_{k\leq n} f_k\right) = \sup_{n\in\mathbb{N}} f_n.$$

□

QUESTION 2.10 *Given function $f(x,y) : X \times Y \to \mathbb{R}$, where X, Y are two spaces. Then we have*

$$\sup_{y\in Y}\sup_{x\in X} f(x,y) = \sup_{x\in X}\sup_{y\in Y} f(x,y),$$
$$\inf_{y\in Y}\inf_{x\in X} f(x,y) = \inf_{x\in X}\inf_{y\in Y} f(x,y).$$

Proof. We just prove the first equality, since we can apply the first equality to $g(x,y) := -f(x,y)$. At first, for each $y \in Y$, we have

$$\sup_{x\in X} f(x,y) \le \sup_{x\in X}\left[\sup_{y\in Y} f(x,y)\right].$$

Let $\sup_{y\in Y}\cdot$ at both sides to obtain

$$\sup_{y\in Y}\sup_{x\in X} f(x,y) \le \sup_{x\in X}\sup_{y\in Y} f(x,y)$$

Then the symmetry between x and y guarantees the inverse inequality also holds. So we have completed the proof. □

QUESTION 2.11 *Suppose (E,d) is a metric space and $A \subset E$. Then the function $x \mapsto d(x,A)$ is continuous.*

Proof. For all $x, y, z \in E$, we have

$$d(x,z) \le d(x,y) + d(y,z).$$

Take $\inf_{z\in A}\cdot$ in both side of the above inequality:

$$\begin{aligned} d(x,A) &= \inf_{z\in A} d(x,z) \le \inf_{z\in A}[d(x,y) + d(y,z)] \\ &= d(x,y) + \inf_{z\in A} d(y,z) = d(x,y) + d(y,A), \end{aligned}$$

which implies

$$d(x,A) - d(y,A) \le d(x,y).$$

Interchange the positions of x and y, we will obtain

$$|d(x,A) - d(y,A)| \le d(x,y).$$

This is what we desire. □

QUESTION 2.12 *Suppose (E,d) is a metric space and $A, B, C \subset E$. Then we have the following triangle inequalities:*

$$\begin{aligned} d(x,A) &\le d(x,y) + d(y,A), \\ d(A,B) &\le d(A,y) + d(y,B). \end{aligned}$$

Moreover,

$$\begin{aligned} d(x,y) &\le d(x,A) + d(A,y), \\ d(x,B) &\le d(x,A) + d(A,B), \\ d(A,C) &\le d(A,B) + d(B,C). \end{aligned}$$

will not hold.

Proof. The previous **QUESTION** has shown $d(x, A) \leq d(x, y) + d(y, A)$ holds. We let $\inf_{x \in B} \cdot$ in this inequality to obtain

$$d(A, B) = \inf_{x \in B} d(A, x) \leq \inf_{x \in B} [d(A, y) + d(y, x)] = d(A, y) + d(y, B).$$

Next we use examples to show the last three inequalities will not hold. Consider $E = \mathbb{R}^2, A = B(0, 1), B = B(2/3, 1), C = B(7/3, 1)$ and $x = (-1, 0), y = (2/3, 0)$. Then

$$\begin{aligned} \frac{5}{3} &= d(x, y) > d(x, A) + d(A, y) = 0 + 0 = 0, \\ \frac{1}{3} &= d(x, B) > d(x, A) + d(A, B) = 0 + 0 = 0, \\ \frac{1}{3} &= d(A, C) > d(A, B) + d(B, C) = 0 + 0 = 0. \end{aligned}$$

□

QUESTION 2.13 *For any right (left) limiting function $f : \mathbb{R} \to \mathbb{R}$, the right (left) limit function $g : \mathbb{R} \to \mathbb{R}, x \mapsto f(x+)$ ($h : \mathbb{R} \to \mathbb{R}, x \mapsto f(x-)$) is right (left) continuous. Moreover, if f has left (right) limit pointwise, then $g(x-)$ ($h(x+)$) also exist. Furthermore, we have $g(x-) = f(x-)$ ($h(x+) = f(x+)$)*

Proof.

1. We just prove the assertion for the right limit function. In fact, by the definition of g, we have: $\forall \varepsilon > 0, \exists \delta > 0$, when $0 < y - x < 2\delta$, then

$$|f(y) - f(x+)| < \varepsilon/2.$$

For any $0 < z - x < \delta$, we can find a sequence $\{y_n\}$ satisfying $0 < y_n - x < 2\delta$ for all $n \in \mathbb{N}_+$ and $y_n \downarrow z$. Replace y with y_n and let $n \to \infty$ in the above inequality to obtain

$$|g(z) - g(x)| = |f(z+) - f(x+)| \leq \varepsilon/2 < \varepsilon.$$

This means g is right continuous.

2. For each $x \in \mathbb{R}$, by the definition of right limit, exists $\Delta_s > 0$, such that

$$|f(x - s + \Delta_s) - f((x - s)+)| < \varepsilon/3.$$

Similarly, exists $\Delta_h > 0$, such that

$$|f(x - h + \Delta_h) - f((x - h)+)| < \varepsilon/3.$$

Since f has left limit pointwise, consider the given $x \in \mathbb{R}$, then exists $\delta > 0$, such that for each $s', h' > 0, |s' - h'| < \delta$, we have

$$|f(x - s') - f(x - h')| < \varepsilon/3.$$

Thus we can let $s, h \in \mathbb{R}_+$ and $|s-h| < \delta$. Moreover, we let Δ_s and Δ_h small enough to satisfy

$$0 < \Delta_s = \Delta_h < \min\{s, h\}.$$

Replace s' with $s - \Delta_s$ and replace h' with $h - \Delta_h$, we will obtain $|s' - h'| = |s - h| < \delta$ and thus

$$\begin{aligned}
&|g(x-s) - g(x-h)| = |f((x-s)+) - f((x-h)+)| \\
\leq &|f((x-s)+) - f(x-s+\Delta_s)| + |f(x-s+\Delta_s) - f(x-h+\Delta_h)| \\
&+ |f(x-h+\Delta_h) - f((x-h)+)| \\
< &\varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon.
\end{aligned}$$

By Cauchy's criterion, we know $\lim_{z \to x-} g(z)$ exists.

3. By the definition of $f(x-)$, for each $\varepsilon > 0$, exists $\delta > 0$, for all $0 < \Delta < \delta$, we have

$$|f(x-\Delta) - f(x-)| < \varepsilon/2.$$

By the definition of g and the definition of $f(x+)$ at $x - \Delta$, for each $\varepsilon > 0, \exists \delta_\Delta > 0$, for all $0 < \Delta' < \delta_\Delta$, we have

$$|g(x-\Delta) - f(x-\Delta+\Delta')| = |f((x-\Delta)+) - f(x-\Delta+\Delta')| < \varepsilon/2.$$

For any $0 < \Delta < \delta$, we pick $0 < \Delta' < \delta_\Delta \wedge \Delta$. Thus $0 < \Delta - \Delta' < \delta$. Combine the above two formulas to obtain

$$\begin{aligned}
&|g(x-\Delta) - f(x-)| = |f((x-\Delta)+) - f(x-)| \\
\leq &|f((x-\Delta)+) - f(x-\Delta+\Delta')| + |f(x-\Delta+\Delta') - f(x-)| \\
< &\varepsilon/2 + \varepsilon/2 = \varepsilon.
\end{aligned}$$

This shows $g(x-)$ exists and $g(x-) = f(x-)$. □

QUESTION 2.14 *Suppose $u : U \to \mathbb{R}$ and $U \subset \mathbb{R}^n$ is an open set. Prove u can be continuously extended to ∂U, namely, $u \in C(\bar{U})$ iff u is uniformly continuous on every bounded subset of U.*

Proof. 1. If $u \in C(\bar{U})$, then for all bounded subset $V \subset U$, we have $u \in C(\bar{V})$. Notice $\bar{V}$ is a compact subset, so u is uniformly continuous on $\bar{V}$, as well as V.

2. For all $x \in \partial U$, consider the γ-neighborhood of x, denoted by $B(x,\gamma) \cap U$. Obviously, $B(x,\gamma) \cap U$ is a bounded subset of U. Hence u is uniformly continuous on $B(x,\gamma) \cap U$. This means: $\forall \varepsilon > 0, \exists \delta > 0$, for each $y, z \in B(x,\gamma) \cap U$ and $|y - z| < \delta$, then $|u(y) - u(z)| < \varepsilon$. So $\forall \varepsilon > 0, \exists \delta > 0$, for each $y, z \in B(x, \delta/2) \cap U$, we have $|y - z| < \delta$, which leads to $|u(y) - u(z)| < \varepsilon$. By Cauchy's criterion, we know $\lim_{U \ni y \to x} u(y)$ exists. Hence u can be continuously extended to ∂U. □

QUESTION 2.15 *Given function $f : \mathbb{R} \to \mathbb{R}, x \mapsto |x|^\alpha$ for each $\alpha \in \mathbb{R}$. Then calculate its derivative.*

Proof. Notice for each $\alpha \in \mathbb{R}$, we have the following equality:

$$\lim_{x \to 0} \frac{(1+x)^\alpha - 1}{x} = \alpha.$$

Then

1. If $x \neq 0$, then for each $\alpha \in \mathbb{R}$, $f(x)$ has definition. Observe when $h > 0$ small enough, we can guarantee $x + h \neq 0$. So $f(x+h)$ has definition as well. Then

$$\begin{aligned} f'(x) &= \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \to 0} \frac{|x+h|^\alpha - |x|^\alpha}{h} \\ &= \lim_{h \to 0} |x|^\alpha \frac{|1 + h/x|^\alpha - 1}{h} \\ &= \lim_{h \to 0} |x|^{\alpha - 1} \mathrm{sgn}(x) \frac{|1 + h/x|^\alpha - 1}{h/x}. \end{aligned}$$

   Observe $h \to 0$ implies $h/x \to 0$ and when h small enough, we have $1 + h/x > 0$. Thus we can remove the absolute value sign and utilize the above equiality to obtain

$$\begin{aligned} f'(x) &= \lim_{h \to 0} |x|^{\alpha - 1} \mathrm{sgn}(x) \frac{(1 + h/x)^\alpha - 1}{h/x} \\ &= \alpha |x|^{\alpha - 1} \mathrm{sgn}(x). \end{aligned}$$

2. If $x = 0$, then $f(x)$ has definition only for $\alpha > 0$. So

$$\begin{aligned} f'(0) &= \lim_{h \to 0} \frac{f(h) - f(0)}{h} = \lim_{h \to 0} \frac{|h|^\alpha - 0}{h} \\ &= \lim_{h \to 0} \left(\mathrm{sgn}(h) \frac{|h|^\alpha - 0}{|h|} \right) \\ &= \lim_{h \to 0} \left(\mathrm{sgn}(h) |h|^{\alpha - 1} \right). \end{aligned}$$

(a) If $\alpha \in (0,1)$, then

$$\operatorname{sgn}(h) = \pm 1, \quad \lim_{h\to 0} |h|^{\alpha-1} = \infty.$$

So the above limit does not exist.

(b) If $\alpha = 1$, then

$$\lim_{h\to 0} \left(\operatorname{sgn}(h)|h|^{\alpha-1}\right) = \lim_{h\to 0} \operatorname{sgn}(h).$$

obviously, this limit does not exist as well.

(c) If $\alpha \in (1,\infty)$, then

$$f'(0) = \lim_{h\to 0} \left(\operatorname{sgn}(h)|h|^{\alpha-1}\right) = 0.$$

□

In conclusion, we the following results:

$\boldsymbol{f(x) = \lvert x\rvert^{\alpha}}$	$\alpha \le 0$	$0 < \alpha \le 1$	$\alpha > 1$
$x \neq 0$	$f'(x) = \alpha\lvert x\rvert^{\alpha-1}\operatorname{sgn}(x)$		
$x = 0$	$f(0)$ is not defined	$f'(0)$ does not exist	$f'(0) = 0$

QUESTION 2.16 *Suppose D is a countable dense subset of $\mathbb{R}$. Then for each $x \in \mathbb{R}$, exists $\{y_n\}$ and $\{z_n\}$ such that $y_n \uparrow x$ and $z_n \downarrow x$.*

Proof. Suppose we do not have such $\{y_n\}$, such that $y_n \uparrow x$. Then exists $\delta > 0$, such that $(x-\delta, x] \cap D = \emptyset$. Select and fix $z \in (x-\delta, x)$. We can find a neighborhood of z, denoted as $U(z)$, such that $U(z) \subset (x-\delta, x]$. So $U(z) \cap D = \emptyset$. This contradicts the density of D in $\mathbb{R}$. □

QUESTION 2.17 *Suppose $D = \{r_n\}$ is a dense subset of $[a,b]$ and $f : [a,b] \to \mathbb{R}$ is right continuous. Write $D_n = [a,b] \cap \{r_1, \cdots, r_n, b\}$. Then*

$$\sup_{x\in U} f(x) = \lim_{n\to\infty} \sup_{s\in D_n} f(s).$$

Proof. Similar to **QUESTION 2.10**, we can easily verify

$$\lim_{n\to\infty} \sup_{s\in D_n} f(s) = \sup_{s\in D\cup\{b\}} f(s).$$

So we only need to prove $\sup_{x\in U} f(x) = \sup_{x\in D\cup\{b\}} f(x)$. By the definition of $\sup_{x\in U} f(x)$, we know exists $\{x_n\} \subset U$ such that $f(x_n) \uparrow \sup_{x\in U} f(x)$. By

the right continuity of f, for each $n \geq 1$, we can find $s_n \in D \cup \{b\}$, such that $x_n \leq s_n$ and $|f(x_n) - f(s_n)| < 1/n$. Thus we have

$$\begin{aligned}|f(s_n) - \sup_{x\in U} f(x)| &\leq |f(s_n) - f(x_n)| + |f(x_n) - \sup_{x\in U} f(x)| \\ &< 1/n + |f(x_n) - \sup_{x\in U} f(x)|.\end{aligned}$$

Let $n \to \infty$, we will obtain $f(s_n) \to \sup_{x\in U} f(x)$. Therefore

$$\sup_{x\in D\cup\{b\}} f(s) \geq \lim_{n\to\infty} f(s_n) = \sup_{x\in U} f(x).$$

It is obvious $\sup_{x\in D\cup\{b\}} f(s) \leq \sup_{x\in U} f(x)$. □

QUESTION 2.18 *Consider the Lebesgue measure space* $(\mathbb{R}, \mathscr{B}(\mathbb{R}), dx)$. *The Dominance convergence theorem implies: if* $f_n \to f$ *uniformly, then for each compact set* $A \in \mathscr{B}(\mathbb{R})$, *we have*

$$\int_A f_n(x)dx \to \int_A f(x)dx.$$

Proof. Since $f_n \to f$ uniformly, then $\forall \varepsilon > 0$, exists $N \geq 1$, when $n > N$, for each $x \in \mathbb{R}$, we have $|f_n(x) - f(x)| < \varepsilon$. Because the limit has nothing to do with the first finite items, we can omit the first N items and $|g_n(x)| := |f_n(x) - f(x)|$ is dominated by the integrable function ε. Thus the Dominance convergence theorem tells us

$$\left|\int_A f_n(x)dx - \int_A f(x)dx\right| \leq \int_A |f_n(x) - f(x)|dx \to 0.$$

□

QUESTION 2.19 *Suppose* $x \in \mathbb{R}$ *and* $|x| \leq 1$. *Let* $P_0(x) = 0$ *and*

$$P_{n+1}(x) = P_n(x) + \frac{1}{2}\left[x^2 - P_n(x)^2\right], \quad n \geq 0.$$

Then $P_n(x) \uparrow |x|$.

Proof. First we verify $P_n(x) \leq |x|$ for each $n \geq 0$. We utilize induction on n. When $n = 0$, the assertion naturally holds. Assume $P_n(x) \leq |x|$, then

$$P_{n+1}(x) = P_n(x) + \frac{1}{2}\left[x^2 - P_n(x)^2\right] = -\frac{1}{2}\left[P_n(x)\right]^2 + P_n(x) + \frac{x^2}{2}$$

is a quadratic function with respect to the variable $P_n(x)$ and interval $P_n(x) \in [-|x|, |x|]$. It is easy to know the quadratic function is increasing in $[-|x|, |x|]$. So

$$P_{n+1}(x) = P_n(x) + \frac{1}{2}\left[x^2 - P_n(x)^2\right] \leq P_{n+1}(x)\Big|_{P_n(x)=|x|} = |x|.$$

Till now, we have proved the assertion. Next, since $P_n(x) \leq |x|$ for each $n \geq 0$, then

$$P_{n+1}(x) - P_n(x) = \frac{1}{2}\left[x^2 - P_n(x)^2\right] \geq 0,$$

which concludes that $\{P_n(x)\}$ is an increasing sequence with upper bound $|x|$. So the limit $z(x) = \lim_{n\to\infty} P_n(x)$ exists. Thus we can let $n \to \infty$ to obtain

$$z(x) = z(x) + \frac{1}{2}\left\{x^2 - [z(x)]^2\right\} \Longrightarrow z(x) = \pm|x|.$$

Notice $\{P_n(x)\}$ is an increasing sequence, thus

$$z(x) = \lim_{n\to\infty} P_n(x) \geq P_1(x) = \frac{x^2}{2} > -|x|.$$

So $z(x) = |x|$. □

QUESTION 2.20 *Suppose $f(x) : \mathbb{R} \to \mathbb{R}$ is a nondecreasing and continuous function along a dense subset Λ of $\mathbb{R}$. Namely, for each $x, y \in \Lambda$ and $x < y$, we have $f(x) \leq f(y)$. Then $f(x)$ is nondecreasing in $\mathbb{R}$.*

Proof. For each $x < y \in \mathbb{R}$, exists $\Lambda \ni x_n \uparrow x$ and $\Lambda \ni y_n \downarrow y$. Therefore, the continuity of $f(x)$ yields

$$f(y) - f(x) = \lim_{n\to\infty}[f(y_n) - f(x_n)] \geq 0.$$

So $f(x)$ is nondecreasing in $\mathbb{R}$. □

QUESTION 2.21 *Suppose for each $n \geq 1, f_n(x) : [0, T] \to \mathbb{R}$ is a nondecreasing function along the finite sequence $(x_i^n : 0 \leq i \leq p_n)$, which satisfies $0 = x_0^n < x_1^n < \cdots < x_{p_n}^n = T$ and $\max_{0\leq i\leq p_n} |x_{i+1}^n - x_i^n| \to 0, n \to \infty$. Besides, we have $(x_i^n : 0 \leq i \leq p_n) \subset (x_i^{n+1} : 0 \leq i \leq p_{n+1})$ for each $n \geq 1$. Assume $f(x) := \lim_{n\to\infty} f_n(x)$ exists pointwise and is continuous, then $f(x) : [0, T] \to \mathbb{R}$ is a nondecreasing function.*

Proof. First we show $f(x)$ is a nondecreasing function along the countable set $\Lambda := \cup_n(x_i^n : 0 \leq i \leq p_n)$. In fact, for each $x_i^n, x_j^m \in \Lambda$ and $x_i^n < x_j^m$, without loss of generality, assume $n \geq m$, then exists $0 \leq j' \leq p_n$, such that $x_j^m = x_{j'}^n$. Thus we have $x_i^n \leq x_{j'}^n = x_j^m$. Utilize the monotonicity of $f_n(x)$ on $(x_i^n : 0 \leq i \leq p_n)$, we have

$$f(x_i^n) = \lim_{k\to\infty} f_k(x_i^n) \leq \lim_{k\to\infty} f_k(x_{j'}^n) = f(x_j^m).$$

This shows $f(x)$ is a nondecreasing function along the countable set Λ.

Then we show Λ is a dense subset of $[0,T]$. In fact, for any $x \in [0,T]$, notice $\max_{0\le i\le p_n} |x_{i+1}^n - x_i^n| \to 0, n \to \infty$, then for any $\varepsilon > 0$, exists $n \ge 1$ such that $\max_{0\le i\le p_n} |x_{i+1}^n - x_i^n| < \varepsilon$. This implies exists $x_i^n \in (x_i^n : 0 \le i \le p_n)$ satisfying $|x - x_i^n| < \varepsilon$. So Λ is a dense subset of $[0,T]$.

Finally, by the previous **QUESTION**, we know $f(x) : [0,T] \to \mathbb{R}$ is a nondecreasing function. □

QUESTION 2.22 *Given two continuous functions $f, g : \mathbb{R} \to \mathbb{R}$. Suppose when $x \to x_0$, we have $f(x) > g(x) + o(1)$. Then $f(x_0) > g(x_0)$.*

Proof. If $f(x_0) = g(x_0)$, consider $f(x) - g(x)$. We have

$$\lim_{x\to x_0} [f(x) - g(x)] = f(x_0) - g(x_0) = 0,$$

which means $f(x) - g(x) = o(1) \Longrightarrow f(x) = g(x) + o(1)$ when $x \to x_0$. This contradicts $f(x) > g(x) + o(1)$. □

QUESTION 2.23 *Given $f(x,y) : \mathbb{R}^2 \to \mathbb{R}$ and define $U = (x' - \alpha, x' + \alpha) \times (y' - \beta, y' + \beta) =: U_x \times U_y$. Assume $\lim_{x\to x'} f(x,y) = g(y)$ and $\lim_{y\to y'} f(x,y) = h(x)$ exist in the region U. And the repeated limits*

$$\lim_{x\to x'} \lim_{y\to y'} f(x,y) \quad \text{and} \quad \lim_{y\to y'} \lim_{x\to x'} f(x,y)$$

also exists in the region U. If one of the above limit is uniform convergence, then we have

$$\lim_{x\to x'} h(x) = \lim_{y\to y'} g(y).$$

Proof. Because of the similarity, we just postulate $\lim_{x\to x'} f(x,y) = g(y)$ is a uniformly convergence. Then $\forall \varepsilon > 0$, exists $\delta \in (0,\alpha)$. Such that for any $y \in U_y$ and any $|x - x'| < \delta$,

$$|f(x,y) - g(y)| < \varepsilon/3.$$

Moreover, by the definition of limit, exists $\delta' \in (0,\delta)$ such that for any $|y - y'| < \delta'$,

$$\left|\lim_{y\to y'} g(y) - g(y)\right| < \varepsilon/3 \quad \text{and} \quad |f(x,y) - h(x)| < \varepsilon/3.$$

Therefore, for any $|x - x'| < \delta'$, we have

$$\begin{aligned}\left|\lim_{y\to y'} g(y) - h(x)\right| &\le \left|\lim_{y\to y'} g(y) - g(y)\right| + |g(y) - f(x,y)| + |f(x,y) - h(x)| \\ &< \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon.\end{aligned}$$

This means $\lim_{x\to x'} h(x) = \lim_{y\to y'} g(y)$. □

QUESTION 2.24 *Given a complex sequence $\{z_n\} \subset \mathbb{C}$. Suppose $\{z_n\}$ is not convergent (in this question, $z_n \to \infty$ is also a convergence sequence), then exists two convergent sub-sequences $\{z_{n'}\}$ and $\{z_{n''}\}$ with disparate limit.*

Proof. For the given sequence $\{z_n\}$, the compactness of $\mathbb{C} \cup \{\infty\}$ implies there is a convergent sub-sequences $\{z_{n'}\}$, say $z_{n'} \to z_1 \in \mathbb{C} \cup \{\infty\}$. We claim exists $\varepsilon > 0$, such that there are infinite items in $B(z_1, \varepsilon)^c$. If not, for any $\varepsilon > 0$, there are finite items in $B(z_1, \varepsilon)^c$, then $z_n \to z_1$. This contradict $\{z_n\}$ is not convergent. So select such $\varepsilon > 0$, consider $\{z_n\} \cap B(z_1, \varepsilon)^c$. The compactness of $\mathbb{C} \cup \{\infty\}$ shows again there is a convergent sub-sequences $\{z_{n''}\}$. obviously, the limits of $\{z_{n'}\}$ and $\{z_{n''}\}$ are different. $\square$

QUESTION 2.25 *Given $n = 3m, m \in \mathbb{N}_+$ and $i, j, k \in \mathbb{N}, i + j + k = n$. Then*

$$\binom{n}{i\ j\ k} \leq \binom{n}{m\ m\ m}.$$

Proof. Notice

$$\binom{n}{i\ j\ k} = \frac{n!}{i!j!k!}.$$

In order to verify the assertion, we just need to verify $(m!)^3 \leq i!j!k!$. In fact, consider the restraint $i, j, k \in \mathbb{N}$ and $i + j + k = n$, without loss of generality, we have three situations:

1. $i = m + n_1, j = m + n_2, k = m - n_1 - n_2$, where $n_1, n_2 \in \mathbb{N}_+$. In this situation, we have

$$\begin{aligned}&\frac{(m!)^3}{i!j!k!}\\ =&\frac{m(m-1)\cdots(m-n_1-n_2+1)}{(m+n_1)(m+n_1-1)\cdots(m+1)(m+n_2)(m+n_2-1)\cdots(m+1)}.\end{aligned}$$

   Observe the numerator and the denominator both has $n_1 + n_2$ items and each item of numerator is smaller than each item of denominator. So $(m!)^3 \leq i!j!k!$ holds.

2. $i = m + n_1 + n_2, j = m - n_1, k = m - n_2$, where $n_1, n_2 \in \mathbb{N}_+$. In this situation, we have

$$\frac{(m!)^3}{i!j!k!} = \frac{m(m-1)\cdots(m-n_1+1)m(m-1)\cdots(m-n_2+1)}{(m+n_1+n_2)(m+n_1+n_2-1)\cdots(m+1)}.$$

Observe the numerator and the denominator both has $n_1 + n_2$ items and each item of numerator is smaller than each item of denominator. So $(m!)^3 \leq i!j!k!$ holds.

3. $i = m, j = m + n_1, k = m - n_1$, where $n_1 \in \mathbb{N}_+$. The same argument will also gives us $(m!)^3 \leq i!j!k!$. □

QUESTION 2.26 *Given $n \in \mathbb{N}_+$ and $i, j, k \in \mathbb{N}, i + j + k = n$. Then*

$$\binom{n}{i\ j\ k} \leq \frac{n!}{\Gamma^3(n/3+1)}.$$

Proof. Observe $\lg \Gamma(x)$ is a convex function, then

$$\begin{aligned}
\lg \binom{n}{i\ j\ k} &= \lg \frac{n!}{i!j!k!} = \lg n! - (\lg i! + \lg j! + \lg k!) \\
&= \lg n! - 3 \times \frac{1}{3}(\lg \Gamma(i+1) + \lg \Gamma(j+1) + \lg \Gamma(k+1)) \\
&\leq \lg n! - 3 \lg \Gamma(\frac{i+j+k+3}{3}) \\
&= \lg n! - 3 \lg \Gamma(\frac{n}{3} + 1) \\
&= \lg \frac{n!}{\Gamma^3(n/3+1)},
\end{aligned}$$

which is equivalent to the assertion we want to prove. When $n = 3m, m \in \mathbb{N}_+$, we will obtain the conclusion of the previous **QUESTION**. □

QUESTION 2.27 *Prove the following equality:*

$$x_0 + \sum_{n=2}^{\infty} x_{n-1} \prod_{i=0}^{n-2} (1 - x_i) = 1 - \prod_{n=0}^{\infty} (1 - x_n),$$

where the left side is convergent.

Proof. Note

$$
\begin{aligned}
&1-\left(x_0+\sum_{n=2}^{\infty} x_{n-1}\prod_{i=0}^{n-2}(1-x_i)\right)\\
=&1-x_0-x_1(1-x_0)-\sum_{n=3}^{\infty} x_{n-1}\prod_{i=0}^{n-2}(1-x_i)\\
=&(1-x_0)\left(1-x_1-\sum_{n=3}^{\infty} x_{n-1}\prod_{i=1}^{n-2}(1-x_i)\right)\\
=&(1-x_0)(1-x_1)\left(1-x_2-\sum_{n=4}^{\infty} x_{n-1}\prod_{i=2}^{n-2}(1-x_i)\right)\\
=&\cdots=\prod_{n=0}^{m}(1-x_n)\left(1-x_{m+1}-\sum_{n=m+3}^{\infty} x_{n-1}\prod_{i=m+1}^{n-2}(1-x_i)\right)\\
=&\prod_{n=0}^{\infty}(1-x_n).
\end{aligned}
$$

The last equality holds since the convergence of

$$
\sum_{n=m+3}^{\infty} x_{n-1}\prod_{i=m+1}^{n-2}(1-x_i) \quad \text{and} \quad \sum_{n=2}^{\infty} x_{n-1}\prod_{i=0}^{n-2}(1-x_i)
$$

are the same. Thus the the desired result follows. □

QUESTION 2.28 *For any real sequence $\{a_n\}_{n\geq 1}$, we have the following ine qualities:*

$$
\varliminf_{n\to\infty}\frac{a_{n+1}}{a_n}\leq \varliminf_{n\to\infty}\sqrt[n]{a_n}\leq \varlimsup_{n\to\infty}\sqrt[n]{a_n}\leq \varlimsup_{n\to\infty}\frac{a_{n+1}}{a_n}.
$$

Proof. The inequality in the middle is trivial, we just prove the first and the final ones.

1. Write $K:=\varliminf_{n\to\infty} a_{n+1}/a_n$, then for any $\varepsilon>0$, exists $N\in\mathbb{N}$ such that when $n>N$,

   $$
   \frac{a_n}{a_{n-1}}>K-\varepsilon \Longrightarrow a_n>(K-\varepsilon)a_{n-1}>\cdots>(K-\varepsilon)^{n-1}a_1.
   $$

   Based on this, we have

   $$
   \begin{aligned}
   &\sqrt[n]{a_n}>(K-\varepsilon)^{1-1/n}\sqrt[n]{a_1}\\
   &\qquad\Longrightarrow \varliminf_{n\to\infty}\sqrt[n]{a_n}\geq \lim_{n\to\infty}(K-\varepsilon)^{1-1/n}\sqrt[n]{a_1}=K-\varepsilon.
   \end{aligned}
   $$

   Let $\varepsilon\to 0$ will complete the proof.

2. Wirte $L := \overline{\lim}_{n\to\infty} a_{n+1}/a_n$, then for any $\varepsilon > 0$, exists $N \in \mathbb{N}$ such that when $n > N$,

$$\frac{a_n}{a_{n-1}} < L + \varepsilon \Longrightarrow a_n < (L+\varepsilon)a_{n-1} < \cdots < (L+\varepsilon)^{n-1}a_1.$$

Based on this, we have

$$\begin{aligned}&\sqrt[n]{a_n} < (L+\varepsilon)^{1-1/n}\sqrt[n]{a_1}\\ &\quad\Longrightarrow \overline{\lim_{n\to\infty}}\sqrt[n]{a_n} \le \lim_{n\to\infty}(L+\varepsilon)^{1-1/n}\sqrt[n]{a_1} = L+\varepsilon.\end{aligned}$$

Let $\varepsilon \to 0$ will complete the proof. □

QUESTION 2.29 *Suppose $f : \mathbb{R} \to \mathbb{R}$ is a uniformly continuous function and the generalized integral*

$$\int_0^\infty f(x)dx$$

exists, then $\lim_{x\to\infty} f(x) = 0$.

Proof. Since the generalized integral exists, then for any $\varepsilon > 0$, exists $M > 0$, when $x > M$ and for any $\Delta > 0$,

$$\left|\int_x^{x+\Delta} f(y)dy\right| < \varepsilon/2.$$

Besides, the uniform continuity of ensure for the $\varepsilon > 0$ mentioned above, exists $\delta > 0$. When $|x-y| < \delta$,

$$|f(x) - f(y)| < \varepsilon/2.$$

Let $\Delta < \delta$, then for any $x > M$, we have

$$\begin{aligned}|f(x)| &\le \frac{1}{\Delta}\int_x^{x+\Delta}|f(x)-f(y)|dy + \left|\int_x^{x+\Delta} f(y)dy\right|\\ &< \frac{1}{\Delta}\Delta\varepsilon/2 + \varepsilon/2 = \varepsilon.\end{aligned}$$

□

QUESTION 2.30 *Suppose $f : \mathbb{R} \to \mathbb{R}$ is an increasing function and the generalized integral*

$$\int_0^\infty f(x)dx$$

exists, then $\lim_{x\to\infty} xf(x) = 0$.

Proof. If $\lim_{x\to\infty} xf(x) \neq 0$, then exists a sequence $\{x_n\}_{n\geq 1}$ such that $\lim_{n\to\infty} x_n f(x_n) = c \neq 0$. Note $f(x)$ is increasing, then

$$\begin{aligned}
&\exists \{x_n\}_{n\geq 1}, \text{ such that } \lim_{n\to\infty} x_n f(x_n) = c \neq 0 \\
&\iff \lim_{x\to\infty} xf(x) = c \neq 0 \\
&\implies \frac{f(x)}{1/x} = c \neq 0.
\end{aligned}$$

Therefore, we know the convergence property of

$$\int_0^\infty f(x)dx \in \mathbb{R} \text{ and } \int_0^\infty \frac{1}{x}dx = \infty$$

are identical. This is a contradiction. □

QUESTION 2.31 *Suppose $\{A_t\}_{t\geq 0} \subset 2^{\mathbb{R}}$ is a decreasing set sequence. Define*

$$\alpha_t := \inf\{x : x \in A_t\} \text{ and } A_{t-} = \bigcap_{0\leq s<t} A_s.$$

Then for any $0 \leq s < t$, we have

$$\lim_{s\uparrow t} \alpha_s \leq \alpha_{t-} := \inf\{x : x \in A_{t-}\}.$$

Proof. Since $\{A_t\}_{t\geq 0}$ is decreasing, then it is trivial to find $t \mapsto \alpha_t$ is increasing (not necessarily strict). First, since $A_{t-} \subset A_s$ for any $0 \leq s < t$, then we can readily obtain

$$\alpha_s \leq \alpha_{t-}, \quad \forall s < t \implies \lim_{s\uparrow t} \alpha_s \leq \alpha_{t-}.$$

Now we give an example to show the strict inequality might happen. Consider

$$A_t = \begin{cases} (1, 6-t) \cup (6, \infty) & \text{, if } 0 \leq t < 5, \\ (6, \infty) & \text{, if } t \geq 5. \end{cases}$$

So $\{A_t\}_{t\geq 0}$ is decreasing and

$$A_{5-} = \bigcap_{0\leq t<5} (1, 6-t) \cup (6, \infty) = (6, \infty).$$

Thus $\alpha_t = 1$ for any $0 \leq t < 5$ and $\alpha_{5-} = \alpha_5 = 6$ implies

$$\lim_{t\uparrow 5} \alpha_t = 1 < 6 = \alpha_{5-}.$$

□

QUESTION 2.32 *If $f(x) \in C_0(\mathbb{R}^n)$, then $f(x)$ is uniformly continuous.*

Proof. For any $\varepsilon > 0$, since $f(x) \to 0, x \to \infty$, then exists $R > 0$ satisfying $|f(x)| < \varepsilon/3, \forall x \in B(0,R)^c$. Since $\overline{B(0,R)}$ is compact, then $f(x)$ is uniformly continuous on $\overline{B(0,R)}$. Thus exists $\delta > 0$, for any $|x-y| < \delta$ and $x, y \in \overline{B(0,R)}$, we have $|f(x) - f(y)| < \varepsilon/3$. Then we claim for any $|x-y| < \delta$ and $x, y \in \mathbb{R}^n$, we have $|f(x) - f(y)| < \varepsilon$. In fact,

1. If $x, y \in \overline{B(0,R)}$, then the above argument shows $|f(x)-f(y)| < \varepsilon/3 < \varepsilon$.
2. If $x \in \overline{B(0,R)}$ and $y \in \overline{B(0,R)}^c \subset B(0,R)^c$. Then we can find $z \in \partial B(0,R) = \overline{B(0,R)} \cap B(0,R)^c$ and $|x-z| + |z-y| = |x-y|$. Thus $|x-z| < \delta$ and
$$\begin{aligned}|f(x) - f(y)| &\le |f(x) - f(z)| + |f(z) - f(y)| \\ &< \varepsilon/3 + |f(z)| + |f(y)| < \varepsilon.\end{aligned}$$
3. If $x, y \in B(0,R)^c$, then $|f(x) - f(y)| \le |f(x)| + |f(y)| < 2\varepsilon/3 < \varepsilon$. □

QUESTION 2.33 *Given $f(x,y) : \mathbb{R}^m \times \mathbb{R}^n \to \mathbb{R}$. If $f(\cdot, y)$ is uniformly continuous with respect to y and $f(x, \cdot)$ is continuous, then $f(x,y)$ is continuous with respect to (x,y).*

Proof. For any $\varepsilon > 0$, since $f(\cdot, y)$ is uniformly continuous with respect to y, then for the given $x \in \mathbb{R}^n$, exists $\delta_x > 0$, such that for any $y \in \mathbb{R}^n$, we have

$$|f(x', y) - f(x,y)| < \varepsilon/2, \quad \forall |x' - x| < \delta_x.$$

And note $f(x, \cdot)$ is continuous, then for the given $x \in \mathbb{R}^n$, exists $\delta_{x,y} > 0$ such that

$$|f(x, y') - f(x,y)| < \varepsilon/2, \quad \forall |y' - y| < \delta_{x,y}.$$

So when $|x - x'| < \delta_x$ and $|y' - y| < \delta_{x,y}$ we have

$$|f(x', y') - f(x,y)| \le |f(x', y') - f(x, y')| + |f(x, y') - f(x,y)| < \varepsilon.$$

So the claim is complete. □

QUESTION 2.34 *Given a complex measurable function $h = f + ig$. Then exists a real measurable function H such that $|h| \le H$ iff exists two measurable function F and G such that $|f| \le F$ and $|g| \le G$.*

Proof. Observe

$$|f| \vee |g| \leq |h| = \sqrt{f^2 + g^2} \leq |f| + |g|.$$

If $|h| \leq H$, then $|f| \vee |g| \leq |h| \leq H$. Let $F = G = H$ will be fine. On the contrary, if $|f| \leq F$ and $|g| \leq G$, then $|h| \leq |f| + |g| \leq F + G$. Let $H = F + G$ will be fine. □

QUESTION 2.35 *Define function* $f : \mathbb{R}^d \to \mathbb{R}$,

$$f(x) = \prod_{k=1}^{d} \frac{\sin x_k}{x_k}.$$

Then we have $f(x) = 1 - |x|^2/6 + O(|x|^4)$ *as* $|x| \to 0$.

Proof. As $|x_k| \to 0$, we have

$$\frac{\sin x_k}{x_k} = \frac{1}{x_k}\left(x_k - \frac{1}{6}x_k^3 + O(|x_k|^5)\right) = 1 - \frac{1}{6}x_k^2 + O(|x_k|^4).$$

Then as $|x| \to 0$, which implies $|x_k| \to 0$ for any $1 \leq k \leq d$, we have

$$\begin{aligned}\prod_{k=1}^{d} \frac{\sin x_k}{x_k} &= \prod_{k=1}^{d}\left(1 - \frac{1}{6}x_k^2 + O(|x_k|^4)\right) \\ &= 1 - \frac{1}{6}\sum_{k=1}^{d} x_k^2 + \frac{1}{36}\sum_{i,j=1}^{d}(x_i x_j)^2 - \frac{1}{6}\sum_{i,j=1}^{d} x_i^2 O(|x_j|^4) \cdots\end{aligned}$$

By the mean value inequality, note

$$\frac{(x_i x_j)^2}{|x|^4} \leq \frac{1}{|x|^4}\left(\sqrt{\frac{x_i^2 + x_j^2}{2}}\right)^4 \leq \frac{1}{|x|^4}\left(\sqrt{\frac{1}{2}\sum_{k=1}^{d} x_k^2}\right)^4 = \frac{1}{4}$$

and

$$\frac{x_i^2 O(|x_j|^4)}{|x|^4} \leq \frac{|x|^2 O(|x_j|^4)}{|x_j|^4} \leq M|x|^2.$$

So $(x_i x_j)^2 = O(|x|^4)$ and $x_i^2 O(|x_j|^4) = O(|x|^4)$. And the result follows. □

QUESTION 2.36 *Given two non negative series satisfying* $\sum_n a_n = \infty$ *and* $\sum_n b_n = \infty$. *If* $\lim_n a_n = \ell > 0$ *(or equivalently* $\lim_n b_n = \ell > 0$*), then* $\sum_n a_n \wedge b_n = \infty$. *Exemplify* $\sum_n a_n \wedge b_n = \infty$ *might not happen when* $\lim_n a_n = \ell > 0$ *is not correct.*

Proof. Write $c_n := a_n \wedge b_n$. If there are infinite items of $\{c_n\}_{n\geq 1}$ satisfying $c_{n_k} = a_{n_k}, k \geq 1$, note $a_{n_k} \to \ell > 0, k \to \infty$, then

$$\sum_n a_n \wedge b_n = \sum_n c_n \geq \sum_k c_{n_k} = \sum_k a_{n_k} = \infty.$$

If there are just finite items of $\{c_n\}_{n\geq 1}$ satisfying $c_n = a_n$, then there is a $N \geq 1$ such that for any $n \geq N$, we have $c_n = b_n$. Therefore

$$\sum_n a_n \wedge b_n = \sum_n c_n \geq \sum_{n\geq N} c_n = \sum_{n\geq N} b_n = \infty.$$

1. When $\lim_n a_n$ does not exists. Consider

$$a_n = \begin{cases} 1/2^n, & \text{if } n \text{ is even,} \\ 1/n, & \text{if } n \text{ is odd} \end{cases} \text{ and } b_n = \begin{cases} 1/n, & \text{if } n \text{ is even,} \\ 1/2^n, & \text{if } n \text{ is odd.} \end{cases}$$

   Thus $\sum_n a_n = \infty$ and $\sum_n b_n = \infty$. However $a_n \wedge b_n = 1/2^n$ yields $\sum_n a_n \wedge b_n = \sum_n 1/2^n < \infty$.

2. When $\lim_n a_n = 0$. Consider

$$a_n = 1/n \text{ and } b_n = \begin{cases} n, & \text{if } n = 2^k, k \geq 1, \\ 0, & \text{otherwise.} \end{cases}$$

   Thus $\sum_n a_n = \infty$ and $\sum_n b_n = \infty$. However

$$a_n \wedge b_n = \begin{cases} 1/n, & \text{if } n = 2^k, k \geq 1, \\ 0, & \text{otherwise.} \end{cases}$$

   Therefore $\sum_n a_n \wedge b_n = \sum_n 1/2^n < \infty$. □

QUESTION 2.37 *Given an increasing function $g : \mathbb{R}_+ \to \mathbb{R}$ satisfying $g(x) \to \infty, x \to \infty$ and define*

$$f(x) := \int_0^x g(u)du, \quad x \in \mathbb{R}_+.$$

Then f is increasing, convex and satisfies $f(x)/x \to \infty, x \to \infty$.

Proof. The increasing property of f is obvious. For any two nonnegative real numbers $x < y$, we have

$$g(x)(y-x) \leq f(y) - f(x) = \int_x^y g(u)du \leq g(y)(y-x).$$

Thus for any three nonnegative real numbers $x < y < z$, we have

$$\frac{f(y) - f(x)}{y - x} \leq g(y) \leq \frac{f(z) - f(y)}{z - y},$$

which implies f is convex. Note that for any $0 \leq y < x$,

$$\frac{f(x)}{x} = \frac{f(x) - f(y)}{x} + \frac{f(y)}{x} \geq g(y)\frac{x - y}{x} + \frac{f(y)}{x}.$$

Let $x \to \infty$, we will obtain

$$\lim_{x\to\infty} \frac{f(x)}{x} \geq \lim_{x\to\infty} \left[g(y)\frac{x - y}{x} + \frac{f(y)}{x}\right] = g(y).$$

Let $y \to \infty$ and apply $g(y) \to \infty, y \to \infty$, we will obtain the desired results. □

QUESTION 2.38 *Suppose $f : \mathbb{R} \to \mathbb{R}$ is an increasing function. Define*

$$g(y) := \inf\{x \in \mathbb{R} : f(x) \geq y\}, \quad h(y) := \sup\{x \in \mathbb{R} : f(x) < y\} \quad y \in \mathbb{R}$$

and

$$\eta(y) := \inf\{x \in \mathbb{R} : f(x) > y\}, \quad \theta(y) := \sup\{x \in \mathbb{R} : f(x) \leq y\} \quad y \in \mathbb{R}.$$

Then we have

1. *$g = h$ and $\eta = \theta$.*
2. *$g \leq \eta$ (or $h \leq \theta$).*
3. *$g(y) > x$ implies $y > f(x)$ (or $y \leq f(x)$ implies $g(y) \leq x$). If furthermore f is right-continuous, then $g(y) > x$ iff $y > f(x)$ (or $g(y) \leq x$ iff $y \leq f(x)$).*
4. *$\eta(y) < x$ implies $y < f(x)$ (or $y \geq f(x)$ implies $\eta(y) \geq x$). If furthermore f is left-continuous, then $\eta(y) < x$ iff $y < f(x)$ (or $\eta(y) \geq x$ iff $y \geq f(x)$).*

Proof.

1. Note for any $z_1 \in \{x \in \mathbb{R} : f(x) \geq y\}$ and $z_2 \in \sup\{x \in \mathbb{R} : f(x) < y\}$, we have $z_1 \geq z_2$ by the increasing of f. Thus $g(y) \geq h(y)$. If $g(y) > h(y)$, then there exists $h(y) < z < g(y)$. Using the definition of g and h, we know $h(y) < z \implies f(z) \geq y$ and $z < g(y) \implies f(x) < y$. And $y \leq f(z) < y$ is a contradiction. In conclusion, $g = h$. A similar argument yields $\eta = \theta$.
2. The proof of $g \leq \eta$ (or $h \leq \theta$) is trivial.

3. Since $g(y) > x$, by the definition of infimum, for every z such that $f(z) \geq y$, we have $f(x) < y$. On the contrary, if $y > f(x)$, the increasing property of f shows $g(y) \geq x$. However, we claim $g(y) = x$ is impossible. In fact, if $g(y) = x$, then there exists $(z_n^+, n \geq 1)$ such that $f(z_n^+) \geq y, z_n^+ > x$ for any $n \geq 1$ and $z_n^+ \downarrow x, n \to \infty$. The right-continuity of f yields $f(x) \geq y$, which contradicts $y > f(x)$.

4. The definition of η and the increasing property of f shows $f(x) > y$. On the contrary, if $y < f(x)$, then $\eta(y) \leq x$. We claim $\eta(y) = x$ is impossible. In fact, if $\eta(y) = x$, then there exists $(z_n^-, n \geq 1)$ such that $f(z_n^-) \leq y, z_n^- < x$ for any $n \geq 1$ and $z_n^- \uparrow x, n \to \infty$. The left-continuity of f yields $f(x) \leq y$, which contradicts $y < f(x)$. □

QUESTION 2.39 *Suppose $f : \mathbb{R} \to \mathbb{R}$ is a decreasing function. Define*

$$g(y) := \inf\{x \in \mathbb{R} : f(x) \leq y\}, \quad h(y) := \sup\{x \in \mathbb{R} : f(x) > y\} \quad y \in \mathbb{R}$$

and

$$\eta(y) := \inf\{x \in \mathbb{R} : f(x) < y\}, \quad \theta(y) := \sup\{x \in \mathbb{R} : f(x) \geq y\} \quad y \in \mathbb{R}.$$

Then we have

1. *$g = h$ and $\eta = \theta$.*

2. *$g \leq \eta$ (or $h \leq \theta$).*

3. *$g(y) > x$ implies $y < f(x)$ (or $y \geq f(x)$ implies $g(y) \leq x$). If furthermore f is right-continuous, then $g(y) > x$ iff $y < f(x)$ (or $g(y) \leq x$ iff $y \geq f(x)$).*

4. *$\eta(y) < x$ implies $y > f(x)$ (or $y \leq f(x)$ implies $\eta(y) \geq x$). If furthermore f is left-continuous, then $\eta(y) < x$ iff $y > f(x)$ (or $\eta(y) \geq x$ iff $y \leq f(x)$).*

Proof. Apply a similar argument of the previous QUESTION. □

QUESTION 2.40 *Let c be an increasing right-continuous function from $\mathbb{R}_+$ into $\bar{\mathbb{R}}_+$. Define*

$$a(u) := \inf\{t \in \mathbb{R}_+ : c(t) > u\}, \quad u \in \mathbb{R}_+,$$

with the usual convention that $\inf \emptyset = \infty$. Then

1. *the function $a : \mathbb{R}_+ \to \bar{\mathbb{R}}_+$ is also increasing and right-continuous, and that*

$$c(t) := \inf\{u \in \mathbb{R}_+ : a(u) > t\}, \quad t \in \mathbb{R}_+.$$

Thus a and c are right-continuous 'functional inverse' of each other.

2. *d*

Proof. 1. The increasing property of a and c are both obvious.

2. Given $u \in \mathbb{R}_+$. By the definition of $a(u)$, there exist $(t_n)_{n\geq 1}$ such that $c(t_n) > u$ and $t_n \downarrow a(u)$ as $n \to \infty$. The right-continuity of c yields $c(a(u)) = \lim_{n\to\infty} c(t_n) \geq u$.

   (a) If $c(a(u)) > u$, then we can find $v \in \mathbb{R}_+$ such that $c(a(u)) > v > u$. On one side, the increasing property of a shows $a(v) \geq a(u)$. On another side, $c(a(u)) > v$ shows $a(v) \leq a(u)$. Thus $a(u) = a(v)$. In short, we have shown that, for any $\varepsilon > 0$, there exists $\delta = c(a(u)) - u > 0$. When u satisfies $0 < v - u < \delta$, then $a(v) - a(u) = 0 < \varepsilon$.

   (b) If $c(a(u)) = u$, there exist $(t_n)_{n\geq 1}$ such that $c(t_n) > u$ and $t_n \downarrow a(u)$ as $n \to \infty$. Thus for any $\varepsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $0 < t_n - a(u) < \varepsilon$ for any $n \geq N$. For any $u < v < c(t_n)$, the definition of a yields $a(v) \leq t_n$. Thus $0 < a(v) - a(u) \leq t_n - a(u) < \varepsilon$. In short, we have shown that, for any $\varepsilon > 0$, there exists $\delta = c(t_n) - u > 0$. When v satisfies $0 < v - u < \delta$, then $0 < a(v) - a(u) < \varepsilon$.

   The above arguments together show a is right-continuous.

3. Write $c'(t) := \inf\{u \in \mathbb{R}_+ : a(u) > t\}$. Then what we need to show is $c = c'$. For any $u \in \mathbb{R}_+$ such that $a(u) > t$, the definition of a and the increasing property of c yield $c(t) \leq u$. Thus $c(t) \leq c'(t)$. On the contrary, applying the definition of c', we have $a(c'(t) - \varepsilon) \leq t$ for any $\varepsilon > 0$. The definition of a again shows $c(t) \geq c'(t) - \varepsilon$. Let $\varepsilon \to 0$ to obtain $c(t) \geq c'(t)$. Therefore $c = c'$. □

QUESTION 2.41 *We say a function $f : \mathbb{R} \to \mathbb{R}$ is convex if for any $x, y \in \mathbb{R}$ and $\theta \in [0, 1]$, we have*

$$(*) \qquad f(\theta x + (1-\theta)y) \leq \theta f(x) + (1-\theta) f(y).$$

Prove the following statements are all equivalent to the convexity:

1. *For any* $x, y \in \mathbb{R}$*, we have*

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x)+f(y)}{2}.$$

2. *For any* $n \in \mathbb{Z}_+$ *and* $x_1, \cdots, x_n \in \mathbb{R}$*, we have*

$$(*) \qquad f\left(\frac{1}{n}\sum_{k=1}^{n} x_k\right) \leq \frac{1}{n}\sum_{k=1}^{n} f(x_k).$$

3. *For any* $x_1 < x_2 < x_3$*, we have*

$$\frac{f(x_2)-f(x_1)}{x_2-x_1} \leq \frac{f(x_3)-f(x_2)}{x_3-x_2}.$$

Furthermore, if $f \in C^1(\mathbb{R})$*, the convexity is also equivalent to*

4. *For any* $x, y \in \mathbb{R}$*, we have*

$$f(y) \geq f(x) + f'(x)(y-x).$$

5. $f'(x)$ *is increasing on* $\mathbb{R}$*.*

If $f \in C^2(\mathbb{R})$*, the convexity is also equivalent to*

6. $f''(x) > 0$ *for any* $x \in \mathbb{R}$*.*

Proof. 1. $1 \Leftrightarrow 2$. We just need to show $1 \Rightarrow 2$. We use induction on n to show

$$f\left(\frac{1}{2^n}\sum_{k=1}^{2^n} x_k\right) \leq \frac{1}{2^n}\sum_{k=1}^{2^n} f(x_k), \quad \forall n \in \mathbb{Z}_+.$$

When $n = 1$, it is obviously true. Suppose the inequality holds for $n = \ell$. When $n = \ell + 1$, using the hypothesis assumption, we have

$$\begin{aligned} f\left(\frac{1}{2^{\ell+1}}\sum_{k=1}^{2^{\ell+1}} x_k\right) &= f\left(\frac{\frac{1}{2^\ell}\sum_{k=1}^{2^\ell} x_k + \frac{1}{2^\ell}\sum_{k=2^\ell+1}^{2^{\ell+1}} x_k}{2}\right) \\ &\leq \frac{1}{2}\left[f\left(\frac{1}{2^\ell}\sum_{k=1}^{2^\ell} x_k\right) + f\left(\frac{1}{2^\ell}\sum_{k=2^\ell+1}^{2^{\ell+1}} x_k\right)\right] \\ &\leq \frac{1}{2^{\ell+1}}\sum_{k=1}^{2^{\ell+1}} f(x_k). \end{aligned}$$

Then we show that, if $(*)$ holds for $n+1$, then $(*)$ also holds for n. Let $t = \frac{1}{n}\sum_{k=1}^{n} x_k$, then $nt = \sum_{k=1}^{n} x_k \Longrightarrow t = \frac{1}{n+1}(\sum_{k=1}^{n} x_k + t)$. Thus

$$f(t) = f\left(\frac{1}{n+1}\left(\sum_{k=1}^{n} x_k + t\right)\right) \leq \frac{1}{n+1}\left[\sum_{k=1}^{n} f(x_k) + f(t)\right].$$

This is equivalent to

$$f\left(\frac{1}{n}\sum_{k=1}^{n} x_k\right) = f(t) \leq \frac{1}{n}\sum_{k=1}^{n} f(x_k).$$

Finally, the above arguments show $(*)$ holds for any $n \in \mathbb{Z}_+$.

2. $(*) \Rightarrow 1$. We just need to let $\theta = 1/2$.

3. $2 \Rightarrow (*)$. First we assume $\lambda = n/m \in \mathbb{Q}$, where $n, m \in \mathbb{Z}_+, n \leq m$ and $(n, m) = 1$. Then

$$\begin{aligned} f(\theta x + (1-\theta)y) &= f\left(\frac{n}{m}x + \frac{m-n}{m}y\right) \\ &= f\left(\frac{nx + (m-n)y}{m}\right) \\ &\overset{2}{\leq} \frac{1}{m}\left[nf(x) + (m-n)y\right] \\ &= \theta f(x) + (1-\theta)f(y). \end{aligned}$$

Since the convex has a left derivative and a right derivative, it must be left continuous and right continuous, i.e., continuous. Next, for any $\theta \in [0,1]$, we can find a rational sequence $(\theta_n)_{n\geq 1} \subset [0,1] \cap \mathbb{Q}$ satisfying $\theta_n \to \theta$. So

$$\begin{aligned} f(\theta x + (1-\theta)y) &= \lim_{n\to\infty} f(\theta_n x + (1-\theta_n)y) \\ &\leq \lim_{n\to\infty} \theta_n f(x) + (1-\theta_n)f(y) \\ &= \theta f(x) + (1-\theta)f(y). \end{aligned}$$

4. $(*) \Leftrightarrow 3$. For $\Rightarrow$, let $\theta = \frac{x_3 - x_2}{x_3 - x_1}$, then $x_2 = \theta x_1 + (1-\theta)x_3$, then

$$\begin{aligned} f(x_2) &= f(\theta x_1 + (1-\theta)x_3) \leq \theta f(x_1) + (1-\theta)f(x_3) \\ &= \frac{x_3 - x_2}{x_3 - x_1} f(x_1) + \frac{x_2 - x_1}{x_3 - x_1} f(x_3). \end{aligned}$$

This is equivalent to either of the following inequality:

$$\begin{aligned} [(x_3 - x_2) + (x_2 - x_1)]f(x_2) &\leq (x_3 - x_2)f(x_1) + (x_2 - x_1)f(x_3) \\ \Longleftrightarrow \frac{f(x_2) - f(x_1)}{x_2 - x_1} &\leq \frac{f(x_3) - f(x_2)}{x_3 - x_2} \end{aligned}$$

For $\Leftarrow$, define $x_1 := x, x_2 := \theta x + (1-\theta)y, x_3 := y$. Then $\theta = \frac{x_3 - x_2}{x_3 - x_1}$. We can reverse the whole process above to obtain

$$f(x_2) \leq \frac{x_3 - x_2}{x_3 - x_1} f(x_1) + \frac{x_2 - x_1}{x_3 - x_1} f(x_3).$$

However, this is equivalent to

$$f(\theta x + (1-\theta)y) = \theta f(x_1) + (1-\theta) f(x_3) = \theta f(x) + (1-\theta) f(y).$$

5. $3 \Rightarrow 4$. Without loss of generality, we assume $y < x$. Arbitrarily pick $y < z < x$, then we have

$$\frac{f(x) - f(y)}{x - y} \leq \frac{f(x) - f(z)}{x - z}.$$

Let $z \to x-$, we have

$$\begin{aligned} \frac{f(x) - f(y)}{x - y} \leq f'(x) &\iff f(x) \leq f(y) + f'(x)(x - y) \\ &\iff f(y) \geq f(x) + f'(x)(y - x). \end{aligned}$$

6. $4 \Rightarrow (*)$. Let $z = \theta x + (1-\theta)y$, then we have

$$\begin{cases} f(x) \geq f(z) + f'(z)(x - z) \\ f(y) \geq f(z) + f'(z)(y - z). \end{cases}$$

Plug $z = \theta x + (1-\theta)y$ to the above inequalities, then this is equivalent to

$$\begin{cases} f(x) \geq f(z) + (1-\theta) f'(z)(x - y) \\ f(y) \geq f(z) + \theta f'(z)(y - x). \end{cases}$$

This is again equivalent to

$$\begin{cases} \theta f(x) \geq \theta f(z) + \theta(1-\theta) f'(z)(x - y) \\ (1-\theta) f(y) \geq (1-\theta) f(z) + \theta(1-\theta) f'(z)(y - x). \end{cases}$$

Add these two formulas together to obtain

$$\theta f(x) + (1-\theta) f(y) \geq f(z) = f(\theta x + (1-\theta)y).$$

7. $3 \Leftrightarrow 5$. For $\Rightarrow$, for any $x, y \in \mathbb{R}$ and $x < y$, we have

$$\frac{f(x) - f(x-h)}{h} \leq \frac{f(x) - f(y)}{x - y} \leq \frac{f(y+h) - f(y)}{h}.$$

Let $h \to 0$, we have

$$f'(x) \leq \frac{f(x) - f(y)}{x - y} \leq f'(y).$$

For $\Leftarrow$, apply the mean value theorem of differential, we have

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(\xi), \quad \frac{f(x_3) - f(x_2)}{x_3 - x_2} = f'(\eta),$$

where $x_1 \leq \xi \leq x_2 \leq \eta \leq x_3$. The increasing property of f' yields the desired result.

8. $5 \Leftrightarrow 6$. We just need to note that, if $f \in C^2(\mathbb{R})$, then f' is increasing iff $(f')' = f'' \geq 0$. □

QUESTION 2.42 *A function $f : \mathbb{R}^n \to \mathbb{R}$ is called semi-upper continuous at $x = x_0$ if $\overline{\lim}_{x \to x_0} f(x) \leq f(x_0)$. Prove the following three definitions are equivalent:*

1. *f is semi-upper continuous at $x = x_0$.*
2. *For any $\varepsilon > 0$, there exists $\delta > 0$, such that $f(x) < f(x_0) + \varepsilon$ for any $x \in B(x_0, \delta)$.*
3. *The upper contour set $U_w(x_0) := \{x \in \mathbb{R}^n : f(x) \geq f(x_0)\}$ is closed.*
4. *The strict lower contour set $L_s(x_0) := \{x \in \mathbb{R}^n : f(x) < f(x_0)\}$ is open.*

Proof. 1. $1 \Leftrightarrow 2$. First, we show $1 \Rightarrow 2$. Recall the definition of the upper limit:

$$\overline{\lim_{x \to x_0}} f(x) = \lim_{r \to 0} \sup_{x \in B(x_0, r)} f(x) = \inf_{r > 0} \sup_{x \in B(x_0, r)} f(x).$$

Thus for any $\varepsilon > 0$, there exists $\delta > 0$, for any $0 < r < \delta$, we have

$$\sup_{x \in B(x_0, r)} f(x) < \overline{\lim_{x \to x_0}} f(x) + \varepsilon \leq f(x_0) + \varepsilon.$$

Thus

$$f(x) \leq f(x_0) + \varepsilon, \quad \forall x \in B(x_0, r)$$

As for $2 \Rightarrow 1$, since for any $\varepsilon > 0$, there exists $\delta > 0$, such that $f(x) < f(x_0) + \varepsilon, \forall x \in B(x_0, \delta)$, we have

$$\sup_{x \in B(x_0, r)} f(x) \leq f(x_0) + \varepsilon.$$

Furthermore,

$$\overline{\lim_{x\to x_0}} f(x) = \inf_{r>0} \sup_{x\in B(x_0,r)} f(x) \le f(x_0) + \varepsilon.$$

Letting $\varepsilon \to 0+$ will yield the result.

2. $3 \Leftrightarrow 4$. This is obvious.

3. $2 \Leftrightarrow 4$. First, we show $2 \Rightarrow 4$. For any $x \in L_s(x_0)$, we have $f(x) < f(x_0)$. Let $\varepsilon = [f(x_0) - f(x)]/2$. There exists $\delta > 0$ such that

$$f(y) < f(x) + \varepsilon < f(x_0), \quad \forall x \in B(x, \delta).$$

This means $B(x, \delta) \subset L_s(x_0)$ and thus $L_s(x_0)$ is open. As for $4 \Rightarrow 2$, consider □

QUESTION 2.43 *Given a random variable X on the probability space $(\Omega, \mathscr{G}, \mathbb{P})$ and valued on the measurable space $(E, \mathscr{E})$. Suppose $(F, \mathscr{F})$ is another measurable space and $f : E \times F \to \mathbb{R}$ is measurable, then*

$$\sup_{y\in F} \mathbb{P}[f(X, y)] \le \mathbb{P}\left[\sup_{y\in F} f(X, y)\right].$$

Proof. On the one hand, for any $y \in F$,

$$\mathbb{P}[f(X, y)] \le \mathbb{P}\left[\sup_{y\in F} f(X, y)\right].$$

Taking $\sup_{y\in F} \cdot$ on two sides of the previous formula yields the inequality. On the other hand, consider $\Phi(x) =$. We have

$$\mathbb{P}[f(X, y)] = 1/y - 1/y = 0, \quad \forall y \neq 0.$$

Therefore $\sup_{y\in F} \mathbb{P}[f(X, y)] = 0$. However,

$$\sup_{y\in F} f(X, y) = \sup_{y\in F} X/y = +\infty \Longrightarrow \mathbb{P}\left[\sup_{y\in F} f(X, y)\right] = +\infty.$$

The proof is completed. □

QUESTION 2.44 *Given an increasing function $\Phi : \mathbb{R} \to \mathbb{R}$ and $A \subset \mathbb{R}$. Then*

$$\sup_{x\in A} \Phi(x) \le \Phi\left(\sup_{x\in A} x\right).$$

Proof. On the one hand, for any $x \in A$,

$$\Phi(x) \leq \Phi\left(\sup_{x\in A} x\right).$$

Taking $\sup_{x\in A} \cdot$ on two sides of the previous formula yields the inequality. On the other hand, consider an increasing function $\Phi(x) = \mathbb{1}_{[1,\infty)}(x)$ and $A = (-\infty, 1)$. Then

$$\sup_{x<1} \mathbb{1}_{[1,\infty)}(x) = 0 < 1 = \mathbb{1}_{[1,\infty)}(1) = \mathbb{1}_{[1,\infty)}\left(\sup_{x<1} x\right).$$

The proof is completed. □

QUESTION 2.45 *Given a differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ satisfying $|f(x) - f(y)| \leq L||x - y||$, namely, L-Lipschitz with respect to Euclidian norm. Then $||\nabla f|| \leq L$.*

Proof. Since f is differentiable, we have for any $y \in \mathbb{R}^n$,

$$|f(x) - f(y)| = |\nabla f(y) \cdot (x - y) + o(||x - y||)| \leq L||x - y||.$$

Divide both sides by $||x - y||$ to obtain

$$\frac{|\nabla f(y) \cdot (x - y) + o(||x - y||)|}{||x - y||} \leq L.$$

Replacing x with $x_n = \nabla f(y)/n + y$ and letting $n \to \infty$, we will have $x_n \to y$ and

$$\left|\nabla f(y) \cdot \frac{\nabla f(y)}{||\nabla f(y)||} + o(1)\right| \to ||\nabla f(y)|| \leq L.$$

□

QUESTION 2.46 *For any cádlág function $f \in D[0, 1]$ and any $\varepsilon > 0$, there must exist finite points $0 = x_0 < x_1 < \cdots < x_{n-1} < x_n = 1$ such that $\sup_{x,y\in[x_i,x_{i+1})} |f(x) - f(y)| < \varepsilon$ for any $0 \leq i \leq n - 1$ (we say f is finitely continuous in $[0, 1]$).*

Proof. If not, we divide $[0, 1]$ into $[0, 1/2) \cup [1/2, 1]$. Then f is not finitely continuous in either $[0, 1/2)$ or $[1/2, 1]$, say $A_1 := [0, 1/2)$. We can divide $[0, 1/2)$ into $[0, 1/4) \cup (1/4, 1/2)$ again and f is not finitely continuous in either $[0, 1/4)$ or $[1/4, 1/2)$, the corresponding interval is denoted by A_2. Applying induction, we can get a decreasing sequence $\{A_n\}_{n\in\mathbb{Z}}$ and $\operatorname{diam}(A_n) \to 0$. The *Nested Interval Theorem* shows there exists a unique $x^* \in \cap_n A_n$. Without loss of generality, we assume $x^* \in (0, 1)$. Since f is cádlág at x^*, for the given

$\varepsilon > 0$, there exists $\delta > 0$, such that $|f(x) - f(x^*)| < \varepsilon/2, \forall x - x^* < 2\delta$ and $|f(x) - f(x^*-)| < \varepsilon/2, \forall x^* - x < 2\delta$. Therefore, letting $x^* - \delta = x_0 < x_1 = x^* < x + \delta$, we have

$$\sup_{x,y\in[x_i,x_{i+1})} |f(x) - f(y)| < \varepsilon, \quad i = 0, 1.$$

So f is finitely continuous in $[x^* - \delta, x^* + \delta]$. However, when n big enough, we can select A_n such that $A_n \subset [x^* - \delta, x^* + \delta]$. The definition of A_n shows f is not finitely continuous in A_n, which is a contradiction. □

QUESTION 2.47 *Assume $f \in C[0,1]$ and $f \geq 0$. What are the limits of*

$$\left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \tag{1}$$

when $\varepsilon \to 0, \varepsilon \to 1$ and $\varepsilon \to \infty$?

Proof. (1) Assume $\varepsilon \to 0$. Now

$$\int_0^1 f(x)^\varepsilon dx \to 1, \quad 1/\varepsilon \to \infty.$$

This is a 1^∞ type indeterminate form. Take $\log(\cdot)$ and apply the L'Hôspital rule to obtain

$$\lim_{\varepsilon\to 0} \frac{1}{\varepsilon} \log\left(\int_0^1 f(x)^\varepsilon dx\right) = \lim_{\varepsilon\to 0} \frac{\int_0^1 f(x)^\varepsilon \log f(x) dx}{\int_0^1 f(x)^\varepsilon dx} = \int_0^1 \log f(x) dx.$$

So,

$$\left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \to \exp\left\{\int_0^1 \log f(x) dx\right\}, \quad \varepsilon \to 0.$$

This is the Cobb-Douglas case.

(2) Assume $\varepsilon \to 1$. Obviously,

$$\left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \to \int_0^1 f(x) dx, \quad \varepsilon \to 1.$$

This is a perfectly substitutable case.

(3) Assume $\varepsilon \to \infty$. Since $f \in C[0,1]$, we can define

$$\sup_{y\in[0,1]} f(y) = \max_{y\in[0,1]} f(y) =: \bar{f}$$

Moreover, write $z = \arg\max_{y\in[0,1]} f(y)$. For any small enough $\gamma > 0$, we can find $\delta(\gamma) > 0$ such that

$$f(x) > \bar{f} - \gamma > 0, \quad \forall x \in [z - \delta(\gamma), z + \delta(\gamma)], \tag{2}$$

On one side, obviously,

$$\left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \le \left[\int_0^1 \bar{f}^\varepsilon dx\right]^{1/\varepsilon} = \bar{f}.$$

Letting $\varepsilon \to \infty$ yields

$$\overline{\lim_{\varepsilon\to\infty}} \left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \le \bar{f}. \tag{3}$$

On the other side, using (2),

$$\left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \ge \left[\int_{z-\delta(\gamma)}^{z+\delta(\gamma)} (\bar{f} - \gamma)^\varepsilon dx\right]^{1/\varepsilon} = (\bar{f} - \gamma)[2\delta(\gamma)]^{1/\varepsilon}.$$

Let $\varepsilon \to \infty$ on both sides, we will get

$$\underline{\lim_{\varepsilon\to\infty}} \left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \ge \bar{f} - \gamma.$$

Let $\gamma \to 0$, we can get

$$\underline{\lim_{\varepsilon\to\infty}} \left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \ge \bar{f}. \tag{4}$$

Combine (3) and (4) to obtain

$$\bar{f} \le \underline{\lim_{\varepsilon\to\infty}} \left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \le \overline{\lim_{\varepsilon\to\infty}} \left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \le \bar{f}.$$

This implies

$$\left[\int_0^1 f(x)^\varepsilon dx\right]^{1/\varepsilon} \to \bar{f}, \quad \varepsilon \to \infty.$$

This is the Leontief case. □

QUESTION 2.48 *If $f : E \to \mathbb{R}^d$ is a concave function, where E is a convex linear space, then the local maximum is the global maximum.*

Proof. If $x \in E$ is the local maximum, then there exists a neighborhood of x, say U_x, such that $f(x) \geq f(y), \forall y \in U_x$. If x is not a global maximum, there exists $x' \in E$ such that $f(x') > f(x)$. For any $\theta \in (0,1)$ small enough, we have $\theta x' + (1-\theta)x \in U_x$. And, the definition of concavity yields

$$f(\theta x' + (1-\theta)x) \geq \theta f(x') + (1-\theta) f(x) > f(x).$$

This contradicts the fact that x is a local maximum. □

QUESTION 2.49 *Let $f : \mathbb{R}^n \to (-\infty, +\infty]$ be an extended real-valued function. Recall the* epigraph *and* sublevel sets*:*

$$\operatorname{epi} f := \{(x,t) \in \mathbb{R}^{n+1} : f(x) \leq t\}, \qquad L_\alpha := \{x \in \mathbb{R}^n : f(x) \leq \alpha\}\,(\alpha \in \mathbb{R}).$$

Prove the following equivalences.

(A) f is lower semicontinuous (l.s.c.), i.e., $f(x) \leq \liminf_{k\to\infty} f(x_k)$ for every sequence $x_k \to x$, if and only if $\operatorname{epi} f$ *is closed in $\mathbb{R}^{n+1}$.*

(B) f is lower semicontinuous if and only if every sublevel set L_α is closed in $\mathbb{R}^n$.

Proof. We work in the Euclidean topology, so sequential arguments suffice.

(A) "⇒": Assume f is l.s.c. Let $(x_k, t_k) \in \operatorname{epi} f$ with $(x_k, t_k) \to (x,t)$. Since $f(x_k) \leq t_k$ for all k,

$$f(x) \;\leq\; \liminf_{k\to\infty} f(x_k) \;\leq\; \liminf_{k\to\infty} t_k \;=\; t,$$

hence $(x,t) \in \operatorname{epi} f$. Therefore $\operatorname{epi} f$ is closed.

(A) "⇐": Assume $\operatorname{epi} f$ is closed. Let $x_k \to x$ and set $s := \liminf_{k\to\infty} f(x_k)$. Choose a subsequence k_j with $f(x_{k_j}) \to s'$ where $s' = s$ (or $s' = +\infty$). Then

$$(x_{k_j},\, f(x_{k_j})) \in \operatorname{epi} f, \qquad (x_{k_j},\, f(x_{k_j})) \to (x,\, s').$$

By closedness, $(x, s') \in \operatorname{epi} f$, hence $f(x) \leq s' = \liminf_{k\to\infty} f(x_k)$. Thus f is l.s.c.

(B) "⇒": Assume f is l.s.c. Fix $\alpha \in \mathbb{R}$ and take $x_k \in L_\alpha$ with $x_k \to x$. Then

$$f(x) \;\leq\; \liminf_{k\to\infty} f(x_k) \;\leq\; \alpha,$$

so $x \in L_\alpha$. Hence each L_α is closed.

(B) "⇐": Assume every L_α is closed. Let $x_k \to x$ and set $s := \liminf_{k\to\infty} f(x_k)$. If $f(x) > s$, choose α with $s < \alpha < f(x)$. Then $f(x_k) \leq \alpha$ holds for infinitely

many k, so a subsequence $x_{k_j} \in L_\alpha$ converges to x. Closedness of L_α implies $x \in L_\alpha$, i.e., $f(x) \le \alpha < f(x)$, a contradiction. Thus $f(x) \le \liminf_{k\to\infty} f(x_k)$, and f is l.s.c. □

QUESTION 2.50 *Prove the following inequality:*

$$\sin(\sin x) < \cos(\cos x), \quad \forall x \in \mathbb{R}.$$

Proof. Since $\sin x + \cos x = \sqrt{2}\left(\frac{\sqrt{2}}{2}\sin x + \frac{\sqrt{2}}{2}\cos x\right) = \sqrt{2}\sin\left(x + \frac{\pi}{4}\right) \le \sqrt{2} < \frac{\pi}{2}$, and $x \mapsto \sin x$ is strictly increasing in $[-1, 1]$, we have

$$\sin(\sin x) < \sin\left(\frac{\pi}{2} - \cos x\right) = \cos(\cos x).$$

□

QUESTION 2.51 *Prove the limit* $\lim_{n\to\infty} \sin n$ *does not exist.*

Proof. We prove this by contradiction. If the limit exists, denoted by $\lim_{n\to\infty} \sin n = a$, we can let $n \to \infty$ in the following equality

$$\sin(n+1) + \sin(n-1) = 2\sin n \cos 1$$

to obtain $2a = 2a\cos 1$. If $a \neq 0$, we have $\cos 1 = 1$, yielding a contradiciton. If $a = 0$, we let $n \to \infty$ in the following equality

$$\sin(n+1) - \sin(n-1) = 2\cos n \sin 1$$

to obtain $0 = 2\lim_{n\to\infty} \cos n \sin 1 \implies \cos n \to 0, n \to \infty$. Letting $n \to \infty$ in the following equality

$$\sin^2 n + \cos^2 n = 1,$$

we will get $0 = 1$, which is also a contradiction. □

QUESTION 2.52 (Deterministic $o(n^{-1/2})$ equivalence) *Let* $(E, \|\cdot\|)$ *be a normed linear space and let* $(b_n)_{n\ge1}$ *be a* deterministic *sequence in* E. *The following are equivalent:*

1. $\|b_n\| = o(n^{-1/2})$ *as* $n \to \infty$*;*
2. *There exists a nonnegative real sequence* $(a_n)_{n\ge1}$ *with* $a_n = o(1)$ *such that for all* n*,*

$$\|b_n\| \le a_n n^{-1/2}.$$

Proof. *(1 ⇒ 2)* Assume $\|b_n\| = o(n^{-1/2})$. Define

$$a_n := n^{1/2}\,\|b_n\| \;\geq\; 0.$$

Then $a_n \to 0$ and $\|b_n\| = a_n\, n^{-1/2}$, hence in particular $\|b_n\| \leq a_n\, n^{-1/2}$ for all n.

(2 ⇒ 1) If there exists $a_n \geq 0$ with $a_n = o(1)$ and $\|b_n\| \leq a_n\, n^{-1/2}$ for all n, then multiplying both sides by $n^{1/2}$ gives

$$n^{1/2}\,\|b_n\| \;\leq\; a_n \;\longrightarrow\; 0,$$

so $\|b_n\| = o(n^{-1/2})$. □

Remark 2.1 The statement is specific to *deterministic* sequences. For random sequences (X_n), the statement $\|X_n\| = o_P(n^{-1/2})$ is *not* equivalent to the existence of a deterministic (a_n) with $\|X_n\| \leq a_n n^{-1/2}$ almost surely for all n; a high-probability formulation is the correct analogue: $\exists\, a_n \downarrow 0$ such that $\mathbb{P}\big(\|X_n\| \leq a_n n^{-1/2}\big) \to 1$.

QUESTION 2.53 ((log-concavity of F and $1 - F$)) *Let $f : \mathbb{R} \to [0, \infty)$ be a one-dimensional probability density function. Assume that $f > 0$ on the interior of its support, that f is continuously differentiable there, and that f is log-concave, i.e. $\log f$ is a concave function on the interior of the support. Define*

$$F(x) := \int_{(-\infty,x]} f(t)\,dt, \qquad 1 - F(x) := \int_{[x,\infty)} f(t)\,dt.$$

Then F and $1 - F$ are log-concave on the interior of the support.

Proof. Because f is log-concave, there exists a concave function $\varphi : \mathbb{R} \to [-\infty, \infty)$ such that $f(x) = e^{\varphi(x)}$. For any $c > 0$ the super-level set

$$A_c := \{x \in \mathbb{R} : f(x) \geq c\} = \{x : \varphi(x) \geq \log c\}$$

is convex, since φ is concave and hence its super-level sets are convex. In one dimension this means that A_c is an interval (possibly empty, a bounded interval, a half-line or all of $\mathbb{R}$).

Suppose that $\overline{\lim}_{x\to+\infty} f(x) > 0$. Then there exist $c > 0$ and a sequence $x_n \to +\infty$ such that $f(x_n) \geq c$ for all n. Hence A_c is unbounded to the right. Because A_c is a convex subset of $\mathbb{R}$ that is unbounded to the right, there exists $R \in \mathbb{R}$ such that

$$[R, \infty) \subset A_c.$$

Consequently

$$\int_{\mathbb{R}} f(x)\,dx \;\geq\; \int_R^{\infty} f(x)\,dx \;\geq\; \int_R^{\infty} c\,dx = \infty,$$

which contradicts that f is a probability density. Therefore $\overline{\lim}_{x\to+\infty} f(x) \leq 0$. Since $f(x) \geq 0$, we obtain $\lim_{x\to+\infty} f(x) = 0$. The same argument applied to $x \to -\infty$ yields $\lim_{x\to-\infty} f(x) = 0$.

On the interior of the support we have $F'(x) = f(x)$ and $F''(x) = f'(x)$. Moreover $\log f$ is concave and twice differentiable there, so

$$(\log f)'(x) = \frac{f'(x)}{f(x)}$$

is nonincreasing as a function of x.

For any $t \leq x$ the monotonicity of f'/f gives

$$\frac{f'(t)}{f(t)} \geq \frac{f'(x)}{f(x)}.$$

Multiplying by $f(t)f(x) > 0$ yields

$$f'(t)f(x) \geq f'(x)f(t).$$

Integrating both sides from $t = -\infty$ to $t = x$ and using the tail limit from Step 1, we obtain

$$\int_{-\infty}^{x} f'(t)f(x)\,dt \;\geq\; \int_{-\infty}^{x} f'(x)f(t)\,dt \quad\Longrightarrow\quad f(x)^2 \geq f'(x)F(x).$$

Now compute the second derivative of $\log F$:

$$(\log F)'(x) = \frac{F'(x)}{F(x)} = \frac{f(x)}{F(x)},$$

$$(\log F)''(x) = \frac{F''(x)F(x) - F'(x)^2}{F(x)^2} = \frac{f'(x)F(x) - f(x)^2}{F(x)^2} \leq 0,$$

where the inequality follows from $f'(x)F(x) - f(x)^2 \leq 0$ proved above. Thus $\log F$ is concave on the interior of the support, i.e. F is log-concave there.

Define the survival function

$$G(x) := 1 - F(x) = \int_x^{\infty} f(t)\,dt.$$

Then $G'(x) = -f(x)$ and $G''(x) = -f'(x)$, so

$$(\log G)'(x) = \frac{G'(x)}{G(x)} = -\frac{f(x)}{G(x)},$$

$$(\log G)''(x) = \frac{G''(x)G(x) - G'(x)^2}{G(x)^2} = \frac{-f'(x)G(x) - f(x)^2}{G(x)^2}.$$

We again use the monotonicity of f'/f, now for $t \geq x$:

$$\frac{f'(t)}{f(t)} \leq \frac{f'(x)}{f(x)} \quad \Rightarrow \quad f'(t)f(x) \leq f'(x)f(t).$$

Integrating from $t = x$ to $t = \infty$ and using the tail limit, we obtain

$$\int_x^\infty f'(t)f(x)\,dt \;\leq\; \int_x^\infty f'(x)f(t)\,dt \quad \Longrightarrow \quad -f(x)^2 \leq f'(x)G(x).$$

Rewriting this inequality as

$$f'(x)G(x) + f(x)^2 \geq 0$$

and substituting into the formula for $(\log G)''(x)$ gives

$$(\log G)''(x) = \frac{-f'(x)G(x) - f(x)^2}{G(x)^2} \leq 0.$$

Hence $\log G$ is concave on the interior of the support, so $G(x) = 1 - F(x)$ is log-concave there. □

§ 3 Algebra

QUESTION 3.1 *Given two positive definite matrices A and B. If $A - B$ is nonnegative definite, we write $A \geq B$. Prove:*

(a) The order $\geq$ is steady under congruence transformation, i.e., for any invertible matrix U,

$$A \geq B \Longleftrightarrow U'AU \geq U'BU.$$

(b) If $B = I$, then the order $\geq$ is steady under inverse operation, i.e.,

$$A \geq I \Longleftrightarrow I \geq A^{-1}.$$

(c) The order $\geq$ is steady under inverse operation, i.e.,

$$A \geq B \Longleftrightarrow B^{-1} \geq A^{-1}.$$

Proof. (a) Since $A \geq B$ iff $v'(A-B)v \geq 0$ for any $v \in \mathbb{R}^n$. Since each $v \in \mathbb{R}^n$, there exists $u \in \mathbb{R}^n$ such that $v = Uu$. The latter one is equivalent to $u'U'(A-B)Uu = u'(U'AU - U'BU)u \geq 0$ for any $u \in \mathbb{R}^n$. So, $U'AU \geq U'BU$.

(b) Noting A is positive definite, there exists positive definite matrix $A^{1/2}$ such that $A = A^{1/2}A^{1/2} \Longleftrightarrow A^{-1} = A^{-1/2}A^{-1/2}$. Moreover, we can find

$$I - A^{-1} = A^{-1/2}(A-I)A^{-1/2} = A^{-1/2}AA^{-1/2} - A^{-1/2}IA^{-1/2}.$$

Letting $U = A^{-1/2}$ and applying (a), we know $I \geq A^{-1}$.

(c) Applying (a) and (b), we have

$$\begin{aligned} A \geq B &\Longleftrightarrow B^{-1/2}AB^{-1/2} \geq I \\ &\Longleftrightarrow I \geq B^{1/2}A^{-1}B^{1/2} \Longleftrightarrow B^{-1} = B^{-1/2}IB^{-1/2} \geq A^{-1}. \end{aligned}$$

□

QUESTION 3.2 *Suppose $A \in \mathbb{R}^{m\times m}$ is positive definite and $B \in \mathbb{R}^{m\times n}$ has rank n, then*

$$(B'B)^{-1}B'AB(B'B)^{-1} - (B'A^{-1}B)^{-1}$$

is a positive definite matrix.

Proof. Based on the last QUESTION, we know what we need to do is equivalent to prove

$$B'B(B'AB)^{-1}B'B - B'A^{-1}B$$

is positive semi-definite. In fact, we can find

$$\begin{aligned} &B'B(B'AB)^{-1}B'B - B'A^{-1}B \\ &\quad = B'A^{-1/2}[A^{1/2}B(B'AB)^{-1}B'A^{1/2} - I]A^{-1/2}B \\ &\quad = B'A^{-1/2}\Omega A^{-1/2}B. \end{aligned}$$

We can easily verify that $\Omega = \Omega'$ and $\Omega = \Omega^2$. So,

$$\begin{aligned} &B'B(B'AB)^{-1}B'B - B'A^{-1}B \\ &\quad = B'A^{-1/2}\Omega'\Omega A^{-1/2}B = (\Omega A^{-1/2}B)'(\Omega A^{-1/2}B), \end{aligned}$$

which shows the above matrix is positive semi-definite. □

QUESTION 3.3 *For any $x \in \mathbb{R}^n$, the rank-one matrix $A := xx^\top$ satisfies*

$$\|A\|_F = \|x\|_2^2.$$

Proof. By definition of the Frobenius norm,

$$\|xx^\top\|_F^2 = \operatorname{tr}\big((xx^\top)^\top(xx^\top)\big) = \operatorname{tr}(xx^\top xx^\top).$$

Since $x^\top x$ is a scalar, we may regroup and commute it with matrices:

$$\operatorname{tr}(xx^\top xx^\top) = \operatorname{tr}\big(x(x^\top x)x^\top\big) = \operatorname{tr}\big((x^\top x)\,xx^\top\big) = (x^\top x)\ \operatorname{tr}(xx^\top).$$

Using the cyclic property of the trace, $\operatorname{tr}(xx^\top) = \operatorname{tr}(x^\top x)$, and $\operatorname{tr}(x^\top x) = x^\top x$ because $x^\top x$ is a 1×1 matrix. Hence

$$\|xx^\top\|_F^2 = (x^\top x)\ \operatorname{tr}(x^\top x) = (x^\top x)\,(x^\top x) = (x^\top x)^2 = \|x\|_2^4.$$

Taking square roots on both sides gives $\|xx^\top\|_F = \|x\|_2^2$. □

QUESTION 3.4 *For any normed linear space $(V, ||\cdot||)$, the norm function $f \mapsto ||f||$ is convex.*

Proof. For any $t \in [0,1]$ and $f, g \in V$, we have

$$||tf + (1-t)g|| \le ||tf|| + ||(1-t)g|| = t||f|| + (1-t)||g||.$$

□

QUESTION 3.5 *For any $A \in M_n(\mathbb{R})$, define*

$$||A||_{2\to 2} := \sup_{||x||_2 \neq 0} \frac{||Ax||_2}{||x||_2}.$$

Then

- $||A||_{2\to 2} = \sqrt{\sigma_{\max}(A)}$, *where $\sigma_{\max}(A)$ is the biggest singular value of A.*
- *Moreover, when A is symmetric, $||A||_{2\to 2} = |\lambda_{\max}(A)|$, where $\lambda_{\max}(A)$ is the biggest eigenvalue of A.*
- *When $A = BB'$, where $B \in M_{n\times k}(\mathbb{R})$, we have $||A||_{2\to 2} = ||B||_{2\to 2}^2$.*
- *When $A = xx'$, where $x \in \mathbb{R}^n$, we have $||A||_{2\to 2} = ||x||_2^2$.*

Proof. Throughout the proof, we adopt the convention

$$\sigma_{\max}(A) := \lambda_{\max}(AA^\top).$$

i.e. the (squared) largest singular value is the largest eigenvalue of $AA^\top$.

- By the definition of the operator norm and the Euclidean inner product,

$$\|A\|_{2\to 2}^2 = \sup_{\|x\|_2=1} \|Ax\|_2^2 = \sup_{\|x\|_2=1} x^\top A^\top Ax.$$

The variational characterization of the largest eigenvalue yields

$$\sup_{\|x\|_2=1} x^\top A^\top Ax = \lambda_{\max}(A^\top A).$$

It is a standard fact that $A^\top A$ and $AA^\top$ have the same nonzero eigenvalues (with the same multiplicities). Hence

$$\|A\|_{2\to 2}^2 = \lambda_{\max}(A^\top A) = \lambda_{\max}(AA^\top) = \sigma_{\max}(A),$$

and therefore $\|A\|_{2\to 2} = \sqrt{\sigma_{\max}(A)}$.

- If A is symmetric, then $A^\top A = A^2$, so

$$\|A\|_{2\to 2}^2 = \lambda_{\max}(A^2) = \max_{1\le i\le n} \lambda_i(A)^2.$$

Taking square roots gives

$$\|A\|_{2\to 2} = \max_{1\le i\le n} |\lambda_i(A)|.$$

In particular, if $A \succeq 0$, then $\lambda_i(A) \ge 0$ and $\|A\|_{2\to 2} = \lambda_{\max}(A)$.

- If $A = BB^\top$ with $B \in \mathbb{R}^{n\times k}$, then $A \succeq 0$, so by Item 2,

$$\|A\|_{2\to 2} = \lambda_{\max}(A) = \lambda_{\max}(BB^\top).$$

On the other hand,

$$\|B\|_{2\to 2}^2 = \lambda_{\max}(B^\top B) = \lambda_{\max}(BB^\top),$$

since $B^\top B$ and $BB^\top$ have the same nonzero eigenvalues. Hence $\|A\|_{2\to 2} = \|B\|_{2\to 2}^2$.

- If $A = xx^\top$ with $x \in \mathbb{R}^n$, then $A \succeq 0$ and $\operatorname{rank}(A) = 1$. We have

$$AA^\top = (xx^\top)(xx^\top) = x(x^\top x)x^\top,$$

so the only nonzero eigenvalue of $AA^\top$ (hence of A) is $x^\top x = \|x\|_2^2$. Therefore,

$$\|xx^\top\|_{2\to 2} = \lambda_{\max}(A) = \|x\|_2^2.$$

This also follows by applying Item 3 with $B = x$ viewed as an $n \times 1$ matrix. Or, since $xx^\top$ is symmetric,

$$\begin{aligned} ||A||_{2\to 2} = \lambda_{\max}(A) &= \sup_{||y||_2=1} y^\top (xx^\top) y \\ &= \sup_{||y||_2=1} (y^\top x)(x^\top y) = \sup_{||y||_2=1} (x^\top y)^2 \\ &= \sup_{||y||_2=1} ||x||_2^2 \, ||y||_2^2 = ||x||_2^2, \end{aligned}$$

where $y^\top x = x^\top y \in \mathbb{R}$ and $x^\top y \leq ||x||_2 \, ||y||_2$ by Cauchy-Schewarz inequality. □

QUESTION 3.6 *Let $A \in \mathbb{R}^{m\times n}$. Then their nonzero eigenvalues (counting geometric multiplicities) coincide. Namely, let $\lambda > 0$. Denote the λ-eigenspace of a square matrix M by*

$$\mathcal{E}_\lambda(M) := \{v \neq 0 : Mv = \lambda v\} \cup \{0\}.$$

Then

$$\dim \mathcal{E}_\lambda(A^\top A) = \dim \mathcal{E}_\lambda(AA^\top).$$

Proof. Let $x \in \mathbb{R}^n$ satisfy $A^\top Ax = \lambda x$ with $\lambda > 0$. Then

$$\|Ax\|_2^2 = x^\top A^\top Ax = \lambda \, \|x\|_2^2 > 0,$$

so $y := Ax \neq 0$. Moreover,

$$AA^\top y = AA^\top Ax = A(A^\top Ax) = A(\lambda x) = \lambda(Ax) = \lambda y,$$

showing y is an eigenvector of $AA^\top$ with eigenvalue λ. Conversely, if $AA^\top y = \lambda y$ with $\lambda > 0$, then

$$\|A^\top y\|_2^2 = y^\top AA^\top y = \lambda \, \|y\|_2^2 > 0,$$

so $x := A^\top y \neq 0$ and

$$A^\top Ax = A^\top AA^\top y = A^\top (AA^\top y) = A^\top(\lambda y) = \lambda x.$$

Thus positive eigenvalues and their algebraic multiplicities match between $A^\top A$ and $AA^\top$.

This defines two linear maps

$$T : \mathcal{E}_\lambda(A^\top A) \to \mathcal{E}_\lambda(AA^\top), \qquad T(x) = Ax$$

and

$$S:\ \mathcal{E}_\lambda(AA^\top) \to \mathcal{E}_\lambda(A^\top A), \qquad S(y) = \tfrac{1}{\lambda}A^\top y.$$

Finally,

$$\begin{aligned} S(T(x)) &= \tfrac{1}{\lambda}A^\top(Ax) = \tfrac{1}{\lambda}(A^\top Ax) = x, \\ T(S(y)) &= A\big(\tfrac{1}{\lambda}A^\top y\big) = \tfrac{1}{\lambda}(AA^\top y) = y, \end{aligned}$$

so T and S are mutual inverses. Hence T is a linear isomorphism between the two eigenspaces, and therefore

$$\dim \mathcal{E}_\lambda(A^\top A) = \dim \mathcal{E}_\lambda(AA^\top).$$

□

§ 4 Topology

QUESTION 4.1 *Suppose X and Y are two metric spaces. Find a linear operator $A : X \to Y$ which is a close mapping but not an open mapping.*

Proof. Consider $X = \mathbb{R}$ and $Y = \mathbb{R}^2$ with Euclidean topology, then we claim the following linear operator A is a close mapping but not an open mapping:

$$A : \mathbb{R} \to \mathbb{R}^2,\ x \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}.$$

The linearity of A is obvious. For all set $T \subset \mathbb{R}$, we have

$$A(T) = T \times \{0\}.$$

When T is a close set, then $T \times \{0\}$ is a closed set in $\mathbb{R}^2$. When T is an open set, then $T \times \{0\}$ is a non-open and non-closed set in $\mathbb{R}^2$. These facts indicate that A is a close mapping but not an open mapping. □

QUESTION 4.2 *Suppose (E, d) and (F, ρ) are two metric spaces and $f : E \to F$ is continuous. Given $A \subset F$, then $\partial f^{-1}(A) \subset f^{-1}(\partial A)$. Moreover, show $\partial f^{-1}(A) \supset f^{-1}(\partial A)$ is not correct.*

Proof.

1. Since $x \in \partial f^{-1}(A)$ iff exist $\{x_n\} \subset f^{-1}(A)$ and $\{x'_n\} \subset [f^{-1}(A)]^c = f^{-1}(A^c)$, such that $x_n \to x$ and $x'_n \to x$. Note the continuity of f, so exist $\{y_n\} \subset A$ and $\{y'_n\} \subset A^c$ such that $y_n = f(x_n) \to f(x)$ and $y'_n = f(x'_n) \to f(x)$. Therefore $f(x) \in \partial A \iff x \in f^{-1}(\partial A)$.

2. Consider $E = F = \mathbb{R}, f(x) = x^2$ and $A = [0, 1]$. Then $\partial f^{-1}[0, 1] = \{-1, 1\} \subset \{-1, 0, 1\} = f^{-1}(\partial[0, 1])$. So $\partial f^{-1}(A) \supset f^{-1}(\partial A)$ is not correct. □

QUESTION 4.3 *Given two metric spaces (E, d). Then for any $A, B \subset E$, we have $\partial(A\cap B)\cup\partial(A\cup B) \subset \partial A\cup\partial B$. Moreover, generally $\partial(A\cap B)\cap\partial(A\cup B) \neq \emptyset$.*

Proof. For any $x \in \partial(A\cap B)$, exist $\{y_n\}_{n\geq 1} \subset A\cap B$ and $\{z_n\}_{n\geq 1} \subset (A\cap B)^c = A^c \cup B^c$ such that $y_n \to x$ and $z_n \to x$. There are infinite items of $\{z_n\}_{n\geq 1}$ in A^c or B^c. Without loss of generality, we can pick a subsequence $\{z_{n_k}\}_{k\geq 1} \subset A^c$. Thus

$$A \ni y_{n_k} \to x, \quad A^c \ni z_{n_k} \to x, \quad k \to \infty.$$

So $x \in \partial A \subset \partial A\cup\partial B$ and therefore $\partial(A\cap B) \subset \partial A\cup\partial B$. The similar argument also shows $\partial(A \cup B) \subset \partial A \cup \partial B$. So $\partial(A \cap B) \cup \partial(A \cup B) \subset \partial A \cup \partial B$.

Consider $A = B(-2, 3) \subset \mathbb{R}^2$ and $B = B(2, 3) \subset \mathbb{R}^2$. Then the intersection point (x, y) are determined by

$$\begin{cases} (x+2)^2 + y^2 = 9, \\ (x-2)^2 + y^2 = 9. \end{cases} \Longrightarrow (x, y) = (0, \pm\sqrt{5}).$$

So $\partial(A \cap B) \cap \partial(A \cup B) = \{(0, \sqrt{5}), (0, -\sqrt{5})\} \neq \emptyset$. □

QUESTION 4.4 *Suppose (E, d) is a metric space and $\{K_n\}_{n\geq 1} \subset E$ is a sequence of compact sets satisfying $K_n \uparrow E$. If $K \subset E$ is a compact, use example to show the fact: there exists $m \geq 1$, such that $K \subset K_m$ might not hold.*

Proof. Way 1. Consider the subspace $E := [0, 1]$ of $\mathbb{R}$ with subspace topology. Then E itself is a compact space. Consider $K_n = [0, 1 - 1/n] \cup \{1\}$ for each $n \geq 1$. Then $\{K_n\}_{n\geq 1}$ is an increasing compact set sequence satisfying

$$\bigcup_{n\geq 1} K_n = [0, 1) \cup \{1\} = E.$$

Consider the compact set $K := E$. Obviously there are no E_n satisfying $K \subset E_n$.

Way 2. Consider $E := \mathbb{S}^2$. Denote $\mathbb{R}^2\cup\{\infty\}$ the one-point compactification of $\mathbb{R}^2$. So $E = \mathbb{S}^2 \cong \mathbb{R}^2 \cup \{\infty\}$. Write the homeomorphic mapping as $\varphi : \mathbb{S}^2 \to \mathbb{R}^2 \cup \{\infty\}$. By the definition of one-point compactification, we know $K_n := \varphi(\{x \in \mathbb{R}^2 : |x| \leq n\}) \cup \{N\}$ is an increasing compact set sequence of $\mathbb{S}^2$, where N is the north pole satisfying

$$\bigcup_{n\geq 1} K_n = \varphi\left(\bigcup_{n\geq 1}\{x \in \mathbb{R}^2 : |x| \leq n\}\right) \cup \{N\} = \mathbb{S}^2.$$

Apply the definition of one-point compactification again, consider the compact set $K := \varphi(\{x \in \mathbb{R}^2 : |x| \geq 10\}) \cup \{N\}$. Obviously there are no E_n satisfying $K \subset E_n$. □

QUESTION 4.5 *Let x be a point in the normed linear space $(E, ||\cdot||)$. For any $\delta > 0$, we have $B(x+y,\delta) = B(x,\delta) + y$.*

Proof. For any $z \in B(x+y,\delta)$, write $z = (z-y)+y =: v+y$. Note

$$||v-x|| = ||(v+y)-(x+y)|| = ||z-(x+y)|| < \delta.$$

So $v \in B(x,\delta)$. The decomposition $z = v+y$ yields that $z \in B(x,\delta)+y$. On the contrary, for any $z \in B(x,\delta)+y$, there exists $u \in B(x,\delta)$ such that $z = u+y$. Note

$$||z-(x+y)|| = ||(u+y)-(x+y)|| = ||u-x|| < \delta.$$

So $z \in B(x+y,\delta)$. □

QUESTION 4.6 *Let A, B be two subsets of the normed linear space $(E, ||\cdot||)$. Let $C := A+B := \{x+y : x \in A, y \in B\}$. Then $(\mathring{A}+B) \cup (A+\mathring{B}) \subset \mathring{C}$.*

Proof. Let $x \in \mathring{A}$, there exists $\delta > 0$ such that $B(x,\delta) \subset A$. Fix $y \in B$. We claim $x+y \in \mathring{C}$. In fact, the previous **QUESTION** shows $B(x+y,\delta) = B(x,\delta)+y \subset A+B = C$. Similarly, we can show $A+\mathring{B} \subset \mathring{C}$. Hence $(\mathring{A}+B) \cup (A+\mathring{B}) \subset \mathring{C}$. □

QUESTION 4.7 *Assume $A \subset \mathbb{R}^n$ is convex. If $x \in \mathrm{Int}(A)$ and $y \in A$, then $\theta x + (1-\theta)y \in \mathrm{Int}(A)$ for any $\theta \in (0,1)$.*

Proof. Since $x \in \mathrm{Int}(A)$, there exists $r > 0$ such that $B(x,r) \subset A$. For any $x' \in B(x,r)$, there exists $\varepsilon \in [0,r]$ and a direction vector β satisfying $||\beta||_{2,n} = 1$, such that $x' = x + \varepsilon\beta$. The convexity of A yields

$$(*) \quad \begin{aligned} \gamma x' + (1-\gamma)y &= \gamma(x+\varepsilon\beta) + (1-\gamma)y \\ &= \gamma x + (1-\gamma)y + \gamma\varepsilon\beta \in A, \quad \forall \gamma \in [0,1], \forall \varepsilon \in [0,r]. \end{aligned}$$

For any $\theta \in (0,1)$, consider $z = \theta x + (1-\theta)y$. We claim $B(z,\theta r) \subset A$. In fact, for any $z' \in B(z,\theta r)$, there exists $\delta \in [0,\theta r]$ and a direction vector β satisfying $||\beta||_{2,n} = 1$, such that

$$\begin{aligned} z' &= z + \delta\beta = \theta x + (1-\theta)y + \delta\beta \\ &= \theta x + (1-\theta)y + \theta\left(\frac{\delta}{\theta}\right)\beta, \quad \frac{\delta}{\theta} \in [0,r]. \end{aligned}$$

Noticing z' has the form just as $(*)$, thus $z' \in A$. Namely, $B(z,\theta r) \subset A$. Finally, we have shown $\theta x + (1-\theta)y \in \mathrm{Int}(A)$ for any $\theta \in (0,1)$. □

§ 5 Complex Analysis

QUESTION 5.1 *For a complex function $f(z) = u(x, y) + iv(x, y)$, we have*

$$\overline{\partial_{\overline{z}} f} = \partial_z \overline{f}.$$

Thus we have

$$\partial^2_{z\overline{z}} f = \overline{\partial^2_{z\overline{z}} \overline{f}}.$$

Proof. Note

$$\partial_z = (\partial_x - i\partial_y)/2, \quad \partial_{\overline{z}} = (\partial_x + i\partial_y)/2.$$

Thus we have

$$\overline{\partial_{\overline{z}} f} = \overline{(\partial_x + i\partial_y)(u + iv)/2} = (\partial_x - i\partial_y)(u - iv)/2 = \partial_z \overline{f}.$$

Therefore $\overline{\partial_{\overline{z}} f} = \partial_z \overline{f}$. Finally,

$$\partial^2_{z\overline{z}} f = \partial_z(\partial_{\overline{z}} f) = \partial_z \left(\overline{\partial_z \overline{f}}\right) = \overline{\partial_{\overline{z}}(\partial_z \overline{f})} = \overline{\partial^2_{z\overline{z}} \overline{f}}.$$

□

QUESTION 5.2 *Let $z = re^{i\theta}$ and $f(z) = u(r, \theta) + iv(r, \theta)$. Prove the Cauchy-Riemann equation is equivalent to*

$$(*) \qquad \partial_r u = \partial_\theta v/r, \quad \partial_r v = -\partial_\theta u/r.$$

Proof. Note $x = r\cos\theta$ and $y = r\sin\theta$. Then

$$\begin{cases} \partial_r u = \cos\theta \partial_x u + \sin\theta \partial_y u, \\ \partial_\theta u = -r\sin\theta \partial_x u + r\cos\theta \partial_y u. \end{cases}$$

Similarly, we can replace all the u in above formula into v to obtain the formula for v. Apply the Cauchy-Riemann equation

$$\partial_x u = \partial_y v, \quad \partial_y u = -\partial_x v,$$

we will get

$$\begin{cases} \partial_r u = \cos\theta \partial_y v - \sin\theta \partial_x v = \partial_\theta v/r, \\ \partial_r v = -\cos\theta \partial_y u + \sin\theta \partial_x u = -\partial_\theta u/r. \end{cases}$$

On the contrary, if $(*)$ holds, we will obtain

$$\begin{cases} \partial_r u = \cos\theta \partial_x u + \sin\theta \partial_y u = -\sin\theta \partial_x v + \cos\theta \partial_y v = \partial_\theta v/r, \\ \partial_r v = \cos\theta \partial_x v + \sin\theta \partial_y v = \sin\theta \partial_x u - \cos\theta \partial_y u = -\partial_\theta u/r. \end{cases}$$

It is equivalent to

$$\begin{cases} (\partial_x u - \partial_y v)\cos\theta + (\partial_y u + \partial_x v)\sin\theta = 0, \\ (\partial_x u - \partial_y v)(-\sin\theta) + (\partial_y u + \partial_x v)\cos\theta = 0. \end{cases}$$

Observe the homogeneous equations

$$\begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

has only zero solution. So we must have

$$\partial_x u - \partial_y v = 0, \quad \partial_y u + \partial_x v = 0.$$

Namely, the Cauchy-Riemann equation holds. □

QUESTION 5.3 *Let $r = re^{i\theta}$. Prove*

$$\partial_{\overline{z}} = e^{i\theta}(\partial_r + i/r\partial_\theta)/2, \quad \partial_z = e^{-i\theta}(\partial_r - i/r\partial_\theta)/2.$$

Hence for any $f \in H(D)$, we have

$$\begin{aligned} f' = \partial_z f &= r/z(\partial_r u + i\partial_r u) = r/z(\partial_\theta v/r + i\partial_r u) \\ &= r/z(\partial_r u - i\partial_\theta u/r) = r/z(\partial_\theta v/r - i\partial_\theta u/r). \end{aligned}$$

Proof. In the previous question we have obtained

$$\partial_r = \cos\theta\partial_x + \sin\theta\partial_y, \quad \partial_\theta = -r\sin\theta\partial_x + r\cos\theta\partial_y.$$

Thus

$$\begin{aligned} e^{i\theta}(\partial_r + i/r\partial_\theta)/2 &= (\cos\theta + i\sin\theta)(\partial_r + i/r\partial_\theta)/2 \\ &= [(\cos\theta\partial_r - \sin\theta/r\partial_\theta) + i(\cos\theta/r\partial_\theta + \sin\theta\partial_r)]/2 \\ &= (\partial_x + i\partial_y)/2 = \partial_{\overline{z}} \end{aligned}$$

and

$$\begin{aligned} e^{-i\theta}(\partial_r - i/r\partial_\theta)/2 &= (\cos\theta - i\sin\theta)(\partial_r - i/r\partial_\theta)/2 \\ &= [(\cos\theta\partial_r - \sin\theta/r\partial_\theta) - i(\cos\theta/r\partial_\theta + \sin\theta\partial_r)]/2 \\ &= (\partial_x - i\partial_y)/2 = \partial_z. \end{aligned}$$

Therefore, we have

$$\begin{aligned} f' = \partial_z f &= e^{-i\theta}(\partial_r - i/r\partial_\theta)(u + iv)/2 \\ &= e^{-i\theta}[(\partial_r u + \partial_\theta v/r) + i(\partial_r v - \partial_\theta u/r)]/2. \end{aligned}$$

Apply the polar coordinate form Cauchy-Riemann equation

$$\partial_r u = \partial_\theta v/r, \quad \partial_r v = -\partial_\theta u/r,$$

we will get

$$\begin{aligned} f' = \partial_z f &= r/z(\partial_r u + i\partial_r u) = r/z(\partial_\theta v/r + i\partial_r u) \\ &= r/z(\partial_r u - i\partial_\theta u/r) = r/z(\partial_\theta v/r - i\partial_\theta u/r). \end{aligned}$$

□

QUESTION 5.4 *Suppose $\gamma : I \to \mathbb{C}$ is a rectifiable curve and φ is holomorphic on γ. Write $\Gamma = \varphi(\gamma)$. Prove*

1. *Γ is also a rectifiable curve.*
2. *If f is continuous on Γ, then*

$$\int_\Gamma f(w)dw = \int_\gamma f(\varphi(z))\varphi'(z)dz.$$

3. *If γ is a smooth curve (i.e. $\gamma \in C^1(I)$ and $\gamma' \not\equiv 0$), then Γ is also a smooth curve and the above formula also holds.*

Proof. 1. Since φ is holomorphic on γ and γ is compact, then exists $M > 0$ such that $|\varphi'(w)| \le M, \forall w \in \gamma(I)$. So for any division $0 = t_0 < t_1 < \cdots < t_n < t_{n+1} = 1$, write γ_k^{k+1} for the sub-curve $\gamma : [t_k, t_{k+1}] \to \mathbb{C}$ and write $|\gamma|$ for the length of curve γ, then we have

$$\begin{aligned} \sum_{k=0}^n |\Gamma(t_{k+1}) - \Gamma(t_k)| &= \sum_{k=0}^n |\varphi(\gamma(t_{k+1})) - \varphi(\gamma(t_k))| \\ &= \sum_{k=0}^n \left| \int_{\gamma(t_k)}^{\gamma(t_{k+1})} \varphi'(z)dz \right| \le \sum_{k=0}^n \int_{\gamma_k^{k+1}} |\varphi'(z)|ds \\ &\le \sum_{k=0}^n M|\gamma_k^{k+1}| = M|\gamma|. \end{aligned}$$

Therefore Γ is also rectifiable.

2. For any $\varepsilon > 0$, there exists $\delta > 0$ and a division of I as $0 = t_0 < t_1 < \cdots < t_{n-1} < t_n = 1$ satisfying $\lambda := \max_{1\le k\le n} |t_k - t_{k-1}| < \delta$ such that

$$\left| \int_\Gamma f(w)dw - \sum_{k=1}^n f(\varphi(\gamma(t_{k-1})))[\varphi(\gamma(t_k)) - \varphi(\gamma(t_{k-1}))] \right| < \varepsilon$$

and

$$\left| \int_\gamma f(\varphi(z))\varphi'(z)dz \right.$$
$$\left. - \sum_{k=1}^{n} f(\varphi(\gamma(t_{k-1})))\varphi'(\gamma(t_{k-1}))[\gamma(t_k) - \gamma(t_{k-1})] \right| < \varepsilon.$$

Moreover, we can let $\delta > 0$ small enough, apply the uniform continuity of φ' on $\gamma(I)$, such that for any $1 \le k \le n$,

$$|\varphi'(w_1) - \varphi'(w_2)| < \varepsilon, \quad \forall w_1, w_2 \in \gamma_{k-1}^k.$$

Therefore

$$\begin{aligned}
&\left| \sum_{k=1}^{n} f(\varphi(\gamma(t_{k-1})))[\varphi(\gamma(t_k)) - \varphi(\gamma(t_{k-1}))] \right. \\
&\qquad \left. - \sum_{k=1}^{n} f(\varphi(\gamma(t_{k-1})))\varphi'(\gamma(t_{k-1}))[\gamma(t_k) - \gamma(t_{k-1})] \right| \\
=&\left| \sum_{k=1}^{n} f(\varphi(\gamma(t_{k-1}))) \int_{\gamma_{k-1}^k} \varphi'(z)dz \right. \\
&\qquad \left. - \sum_{k=1}^{n} f(\varphi(\gamma(t_{k-1}))) \int_{\gamma_{k-1}^k} \varphi'(\gamma(t_{k-1}))dz \right| \\
\le& \sum_{k=1}^{n} f(\varphi(\gamma(t_{k-1}))) \int_{\gamma_{k-1}^k} |\varphi'(z) - \varphi'(\gamma(t_{k-1}))|ds \\
<& \varepsilon \sum_{k=1}^{n} |f(\varphi(\gamma(t_{k-1})))||\gamma_{k-1}^k| \\
\le& \left(\max_{z\in\Gamma} |f(z)||\gamma| \right) \varepsilon.
\end{aligned}$$

So we have obtained the desired result.

3. $\Gamma \in C^1(I)$ is Obvious. Moreover, since φ is holomorphic on γ, by the uniqueness of holomorphic function, $\varphi'(w) \not\equiv 0, \forall w \in \gamma(I)$. Therefore $\Gamma'(t) = \varphi'(\gamma(t))\gamma'(t) \not\equiv 0, \forall t \in I$. So Γ is also a smooth curve. Then

$$\begin{aligned}
\int_\Gamma f(w)dw &= \int_I f(\Gamma(t))\Gamma'(t)dt = \int_I f(\varphi(\gamma(t)))\varphi'(\gamma(t))\gamma'(t)dt \\
&= \int_\gamma f(\varphi(z))\varphi'(z)dz.
\end{aligned}$$

□

§ 6 Probability

QUESTION 6.1 *Suppose ξ, η are two identically distributed random variables on $(\Omega, \mathscr{F}, \mathbb{P})$ and sub σ-algebra $\mathscr{A} \subset \mathscr{F}$. Use an example to show*

$$\mathbb{P}(\xi|\mathscr{A}) = \mathbb{P}(\eta|\mathscr{A})$$

is not true. However, if $\xi = \eta, \mathbb{P}$-a.s., then $\mathbb{P}(\xi|\mathscr{A}) = \mathbb{P}(\eta|\mathscr{A})$.

Proof. Consider $\xi \sim N(0,1)$ and let $\eta = -\xi$, then $\eta \sim N(0,1)$. Moreover, let $\mathscr{A} = \sigma(\xi)$, we will have

$$\mathbb{P}(\xi|\xi) = \xi \neq \eta = -\xi = -\mathbb{P}(\xi|\xi) = \mathbb{P}(-\xi|\xi) = \mathbb{P}(\eta|\xi).$$

This is a smart example. If $\xi = \eta, \mathbb{P}$-a.s., then for any $A \in \mathscr{A}$, we have $\mathbb{P}(\xi 1_A) = \mathbb{P}(\eta 1_A)$. This shows $\mathbb{P}(\xi|\mathscr{A}) = \mathbb{P}(\eta|\mathscr{A})$. □

QUESTION 6.2 *Consider the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and the sub σ-algebra $\mathscr{A}_1, \mathscr{A}_2$. Suppose ξ is a random variable. Use an example to show*

$$\mathbb{P}\left[\mathbb{P}(\xi|\mathscr{A}_1)|\mathscr{A}_2\right] = \mathbb{P}(\xi|\mathscr{A}_1 \cap \mathscr{A}_2)$$

is not true.

Proof. Consider $\Omega = \{0,1,2\}, \mathscr{F} = 2^\Omega, \xi = \mathbf{1}_{\{0\}}$,

$$\mathscr{A}_1 = \{\{1\}, \{0,2\}, \Omega, \emptyset\}, \quad \mathscr{A}_2 = \{\{2\}, \{0,1\}, \Omega, \emptyset\},$$

and $\mathbb{P}(\{0\}) = 1/2, \mathbb{P}(\{1\}) = \mathbb{P}(\{2\}) = 1/4$. Observe $\mathscr{A}_1 \cap \mathscr{A}_2 = \{\Omega, \emptyset\}$, then ξ is independent of $\mathscr{A}_1 \cap \mathscr{A}_2$ and hence

$$\mathbb{P}(\xi|\mathscr{A}_1 \cap \mathscr{A}_2) = \mathbb{P}(\xi) = 1/2.$$

Moreover, notice $\{1\}$ and $\{0,2\}$ are two atoms in $\mathscr{A}_1$, so

$$\begin{aligned}
\mathbb{P}(\xi|\mathscr{A}_1) &= \mathbb{P}(\xi|\{1\})\mathbb{1}_{\{1\}} + \mathbb{P}(\xi|\{0,2\})\mathbb{1}_{\{0,2\}} \\
&= \mathbb{P}(\{0\}|\{1\})\mathbb{1}_{\{1\}} + \mathbb{P}(\{0\}|\{0,2\})\mathbb{1}_{\{0,2\}} \\
&= 0\mathbb{1}_{\{1\}} + \frac{\mathbb{P}(\{0\})}{\mathbb{P}(\{0,2\})}\mathbb{1}_{\{0,2\}} \\
&= \frac{1/2}{1/2 + 1/4}\mathbb{1}_{\{0,2\}} = 2/3\mathbb{1}_{\{0,2\}}.
\end{aligned}$$

Note $\{2\}$ and $\{0,1\}$ are two atoms in $\mathscr{A}_2$, so

$$\begin{aligned}
\mathbb{P}\left[\mathbb{P}(\xi|\mathscr{A}_1)|\mathscr{A}_2\right] &= \mathbb{P}\left(2/3\mathbb{1}_{\{0,2\}}\middle|\mathscr{A}_2\right) \\
&= 2/3\left[\mathbb{P}\left(\{0,2\}|\{2\}\right)\mathbb{1}_{\{2\}} + \mathbb{P}\left(\{0,2\}|\{0,1\}\right)\mathbb{1}_{\{0,1\}}\right] \\
&= \frac{2}{3}\left(\mathbb{1}_{\{2\}} + \frac{\mathbb{P}(\{0\})}{\mathbb{P}(\{0,1\})}\mathbb{1}_{\{0,1\}}\right) \\
&= \frac{2}{3}\left(\mathbb{1}_{\{2\}} + \frac{1/2}{1/2+1/4}\mathbb{1}_{\{0,1\}}\right) \\
&= \frac{2}{3}\left(\mathbb{1}_{\{2\}} + \frac{2}{3}\mathbb{1}_{\{0,1\}}\right) \\
&= \frac{2}{3}\mathbb{1}_{\{2\}} + \frac{4}{9}\mathbb{1}_{\{0,1\}}.
\end{aligned}$$

Hence

$$\mathbb{P}(\xi|\mathscr{A}_1 \cap \mathscr{A}_2) = \frac{1}{2} \neq \frac{2}{3}\mathbb{1}_{\{2\}} + \frac{4}{9}\mathbb{1}_{\{0,1\}} = \mathbb{P}\left[\mathbb{P}(\xi|\mathscr{A}_1)|\mathscr{A}_2\right].$$

□

QUESTION 6.3 *1. Consider the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Given A, B and $C \in \mathscr{F}$, if A is independent to B, C or C is independent to A, B, then A and C are independent conditional on B.*

2. Suppose conditional on B, A and C are conditionally independent, namely, $\mathbb{P}(A \cap C|B) = \mathbb{P}(A|B)\mathbb{P}(C|B)$, then $\mathbb{P}(A|B \cap C) = \mathbb{P}(A|B)$.

Proof. 1. Without loss of generality, we assume A is independent to B, C. Then we have

$$\mathbb{P}(A \cap B|C) = \frac{\mathbb{P}(A \cap B \cap C)}{\mathbb{P}(C)} = \frac{\mathbb{P}(A)\mathbb{P}(B \cap C)}{\mathbb{P}(C)}$$

and

$$\begin{aligned}
\mathbb{P}(A|C)\mathbb{P}(B|C) &= \frac{\mathbb{P}(A \cap C)\mathbb{P}(B \cap C)}{[\mathbb{P}(C)]^2} = \frac{\mathbb{P}(A)\mathbb{P}(C)\mathbb{P}(B \cap C)}{[\mathbb{P}(C)]^2} \\
&= \frac{\mathbb{P}(A)\mathbb{P}(B \cap C)}{\mathbb{P}(C)}.
\end{aligned}$$

2. We have

$$\begin{aligned}
\mathbb{P}(A|B \cap C) &= \frac{\mathbb{P}(A \cap B \cap C)}{\mathbb{P}(B \cap C)} = \frac{\mathbb{P}(A \cap C|B)\mathbb{P}(B)}{\mathbb{P}(B \cap C)} \\
&= \frac{\mathbb{P}(A|B)\mathbb{P}(C|B)\mathbb{P}(B)}{\mathbb{P}(B \cap C)} = \mathbb{P}(A|B).
\end{aligned}$$

□

QUESTION 6.4 *Suppose A is a subset on $\mathbb{R}^n$. If $D \subset A$ and $dx(D) = dx(A)$, where dx is the Lebesgue measure, then D is dense in A.*

Proof. For all $x \in A$ and for all $\varepsilon > 0$, consider the neighborhood $B(x, \varepsilon) \subset A$. We claim there must exist $a \in A$, such that $a \in B(x, \varepsilon)$. If not, then $D \subset B(x, \varepsilon)^c$. Hence $dx(D) \leq dx(A) - n\alpha(n)\varepsilon^n < dx(A)$, which is a contradiction. □

QUESTION 6.5 *Let*

$$\begin{aligned}
G_1 &= \left(\frac{1}{3}, \frac{2}{3}\right), \\
G_2 &= \left(\frac{1}{3^2}, \frac{2}{3^2}\right) \bigcup \left(\frac{2}{3} + \frac{1}{3^2}, \frac{2}{3} + \frac{2}{3^2}\right), \\
&\vdots \\
G_n &= \bigcup \left\{ \left(\sum_{k=1}^{n-1} \frac{a_k}{3^k} + \frac{1}{3^n}, \sum_{k=1}^{n-1} \frac{a_k}{3^k} + \frac{2}{3^n} \right) : a_k = 0 \text{ or } 2 \right\}.
\end{aligned}$$

Then $G := \cup_n G_n$ is an open subset of $[0, 1]$. The closed set $C = [0, 1] - G$ is called the Cantor set. Let

$$F_0(u) = \sum_{k=1}^{n-1} \frac{a_k}{2^k} + \frac{1}{2^n}, \quad \sum_{k=1}^{n-1} \frac{a_k}{3^k} + \frac{1}{3^n} < u < \sum_{k=1}^{n-1} \frac{a_k}{3^k} + \frac{2}{3^n}.$$

We can extend F to a probability distribution function on $\mathbb{R}$ by setting

$$F(x) := \begin{cases} 1 & \text{, if } x \geq 1 \\ \inf\{F_0(y) : y \in G \text{ and } y > x\} & \text{, if } x < 1. \end{cases}$$

Show F is uniformly continuous on $\mathbb{R}$.

Proof. We first prove F_0 is uniformly continuous on G. Observe F_0 is increasing on G and for all $n \in \mathbb{N}$, the distances between each pair of adjacent open intervals in $\cup_{k=1}^n G_k$ equal to $1/3^n$. By the definition of F_0, the variation in each closed set on $[0, 1] - \cup_{k=1}^n G_k$ is $1/2^n$. Thus for every $x, y \in [0, 1]$ and $|x - y| < 1/3^n$, there are three situations might happens:

1. x and y all lie in one of open intervals in $\cup_{k=1}^n G_k$. Since F_0 stays constant in the open interval, we have $|F_0(x) - F_0(y)| = 0$.

2. x lies in one of open intervals in $\cup_{k=1}^n G_k$ and y lies in one of closed intervals in $[0, 1] - \cup_{k=1}^n G_k$, we have $|F_0(x) - F_0(y)| < 1/2^n$.

3. x and y are all lie in one of closed intervals in $[0,1]-\cup_{k=1}^{n}G_k$, we also have $|F_0(x)-F_0(y)|<1/2^n$.

In short, we know if $x,y\in G$ and $|x-y|<1/3^n$, then $|F_0(x)-F_0(y)|<1/2^n$, which means F is uniformly continuous on G. Since $dx(G)=1$, by **QUESTION 1.4** and **QUESTION 3.3**, we know G is dense in $[0,1]$ and F is uniformly continuous on $\mathbb{R}$. □

QUESTION 6.6 *Suppose $\mathscr{B}$ is the Borel σ-algebra on $\mathbb{R}$. Then $\mathscr{B}[0,1]=\mathscr{B}\cap[0,1]$.*

Proof. Write $\mathscr{O}$ for all the open set on $\mathbb{R}$, then $\mathscr{B}=\sigma(\mathscr{O})$. We can rewrite $\mathscr{B}[0,1]=\mathscr{B}\cap[0,1]$ as $\sigma(\mathscr{O}\cap[0,1])=\sigma(\mathscr{O})\cap[0,1]$. Observe $\mathscr{O}\cap[0,1]\subset\sigma(\mathscr{O})\cap[0,1]$ and $\sigma(\mathscr{O})\cap[0,1]$ is a σ-algebra on $[0,1]$, therefore $\sigma(\mathscr{O}\cap[0,1])\subset\sigma(\mathscr{O})\cap[0,1]$. On the contrary, consider

$$\mathscr{A}=\{A\in\sigma(\mathscr{O}):A\cap[0,1]\in\sigma(\mathscr{O}\cap[0,1])\}.$$

Obviously $\mathscr{O}\subset\mathscr{A}$ and it is easy to verify $\mathscr{A}$ is a σ-algebra. Thus $\forall A\in\sigma(\mathscr{O}), A\cap[0,1]\in\sigma(\mathscr{O}\cap[0,1])$, i.e., $\sigma(\mathscr{O}\cap[0,1])\supset\sigma(\mathscr{O})\cap[0,1]$. □

QUESTION 6.7 *Suppose $(\Omega,\mathscr{F})$ and $(E,\mathscr{E})$ are two measurable spaces and $\{X_i\}_{i=1}^n$ are measurable mappings with respect to $\mathscr{F}$ and $\mathscr{E}$. Consider the measurable vector $(X_1,\cdots,X_n):(\Omega,\mathscr{F})\to(E^n,\mathscr{E}^n)$, then we have*

$$\sigma(X_1,\cdots,X_n)=\sigma\left(\bigcup_{i=1}^{n}\sigma(X_i)\right).$$

Proof. For each $1\le i\le n$ and $B_i\in\mathscr{E}$, we have

$$\{X_i\in B_i\}=\{(X_1,\cdots,X_n)\in E\times\cdots\times B_i\times\cdots\times E\}.$$

Then $\sigma(X_i)\subset\sigma(X_1,\cdots,X_n)\implies\cup_{i=1}^n\sigma(X_i)\subset\sigma(X_1,\cdots,X_n)$. Thus we have $\sigma(X_i,1\le i\le n)\subset\sigma(X_1,\cdots,X_n)$. On the contrary, the above equality means

$$\{(X_1,\cdots,X_n)\in B_1\times\cdots\times B_n\}=\bigcap_{i=1}^{n}\{X_i\in B_i\}\in\sigma\left(\bigcup_{i=1}^{n}\sigma(X_i)\right).$$

Consider the following set:

$$\mathscr{A}:=\{B\in\mathscr{E}^n:\{(X_1,\cdots,X_n)\in B\}\in\sigma(X_i,1\le i\le n)\}.$$

It is easy to prove $\mathscr{A}$ is a σ-algebra and the previous formula shows all the measurable rectangles are contained in $\mathscr{A}$. Then $\mathscr{A}=\mathscr{E}^n$, namely, $\sigma(X_1,\cdots,X_n)\subset\sigma(X_i,1\le i\le n)$. □

QUESTION 6.8 *Suppose $(X_t)_{t\in I}$ is a sequence of random variables from the measurable space $(\Omega, \mathscr{F})$ to $(E, \mathscr{E})$, where I is a index set. Then*

$$\begin{aligned}&\sigma(X_t : t \in I)\\ =&\sigma\left(\{\{X_{t_1} \in A_1, \cdots, X_{t_n} \in A_n\} : A_1, \cdots, A_n \in \mathscr{E}, (t_1, \cdots, t_n) \in S(I)\}\right)\\ =&: \mathscr{A}.\end{aligned}$$

where $S(I)$ denotes all the ordered finite subset of I.

Proof. Define $\mathscr{C} = \{A_1 \times \cdots \times A_n : A_1, \cdots, A_n \in \mathscr{E}\}$, then $\sigma(\mathscr{C}) = \mathscr{E}^n$. Obviously, for each $A_1, \cdots, A_n \in \mathscr{E}, (t_1, \cdots, t_n) \in S(I)$, we have

$$\{X_{t_1} \in A_1, \cdots, X_{t_n} \in A_n\} = \bigcap_{k=1}^{n} \{X_{t_k} \in A_k\} \in \sigma(X_t : t \in I).$$

Thus $\mathscr{A} \subset \sigma(X_t : t \in I)$. On the contrary, for each $t \in I$, obviously $\sigma(X_t) \subset \mathscr{A} \Longrightarrow \cup_{t\in I}\sigma(X_t) \subset \mathscr{A}$ and therefore $\sigma(X_t : t \in I) \subset \mathscr{A}$. □

QUESTION 6.9 *Suppose $(X_t^i)_{t\in I}, i = 1, \cdots, d$ are d sequences of random variables from the measurable space $(\Omega, \mathscr{F})$ to $(E, \mathscr{E})$. Then*

$$\begin{aligned}&\sigma(X_t^i : t \in I, i = 1, \cdots, d)\\ =&\sigma\left(\left\{\bigcup_{i=1}^{d}\{X_{t_1}^i \in A_1^i, \cdots, X_{t_n}^i \in A_n^i\} : A_1^i, \cdots, A_n^i \in \mathscr{E}, (t_1, \cdots, t_n) \in S(I)\right\}\right).\end{aligned}$$

Proof. The proof is similar to the previous question. □

QUESTION 6.10 *Suppose $(\Omega, \mathscr{F})$ and $(E, \mathscr{E})$ are two measurable spaces. Then $X : (\Omega, \mathscr{F}) \to (E, \mathscr{E})$ is measurable iff $f(X) : (\Omega, \mathscr{F}) \to (\mathbb{R}, \mathscr{B}(\mathbb{R}))$ is measurable for each $f \in \mathrm{b}\mathscr{E}$.*

Proof. If $f(X) \in \mathscr{F}$ for each $f \in \mathrm{b}\mathscr{E}$, then for all $A \in \mathscr{E}$, choose $f = \mathbf{1}_A$, we have $f(X) = \mathbf{1}_{\{X\in A\}} \in \mathscr{F}$. Since $\{1\} \in \mathscr{B}(\mathbb{R})$, then

$$\{X \in A\} = \{\mathbf{1}_{\{X\in A\}} = 1\} \in \mathscr{F},$$

which means $X \in \mathscr{F}$. On the contrary, if $X \in \mathscr{F}$, obviously $f(X) \in \mathscr{F}$ for each $f \in \mathrm{b}\mathscr{E}$. □

QUESTION 6.11 *Suppose $\mathscr{F}$ and $\mathscr{G}$ are two σ algebras on Ω. Then $\mathscr{F} \subset \mathscr{G}$ iff for each $F \in \mathrm{b}\mathscr{F}$, we have $F \in \mathrm{b}\mathscr{G}$.*

Proof. If $\mathscr{F} \subset \mathscr{G}$, then obviously $F \in \mathrm{b}\mathscr{F}$ implies $F \in \mathrm{b}\mathscr{G}$. On the contrary, if for each $F \in \mathrm{b}\mathscr{F}$, we have $F \in \mathrm{b}\mathscr{G}$, we can choose $F = 1_A$ for each $A \in \mathscr{F}$. Then $1_A \in \mathrm{b}\mathscr{G}$ and thus $A \in \mathscr{G}$. So $\mathscr{F} \subset \mathscr{G}$. □

QUESTION 6.12 *Suppose* $(\Omega, \mathscr{F}, \mathbb{P})$ *is a probability space and* X *is an integral random variable on* $(\Omega, \mathscr{F}, \mathbb{P})$*. Assume* $\mathscr{G} \subset \mathscr{F}$ *is also a* σ*-algebra, then* $Z = \mathbb{P}(X|\mathscr{G})$ *iff* $Z \in \mathscr{G}$ *and for each measurable function* $F \in \mathrm{b}\mathscr{G}$*,*

$$\mathbb{P}(XF) = \mathbb{P}(ZF).$$

Proof. If the above equality holds, we can choose $F = 1_A$ for each $A \in \mathscr{G}$, the $Z = \mathbb{P}(X|\mathscr{G})$ follows. On the contrary, consider the following set:

$$\mathscr{L} := \{F \in \mathrm{b}\mathscr{G} : \mathbb{P}(XF) = \mathbb{P}(ZF)\}.$$

By the definition of $\mathbb{P}(X|\mathscr{G})$, we know $\{1_A : A \in \mathscr{G}\} \subset \mathscr{L}$. If $F, G \in \mathscr{L}$ and $\alpha, \beta \in \mathbb{R}$, we have

$$\begin{aligned}\mathbb{P}[\mathbb{P}(X|\mathscr{G})(\alpha F + \beta G)] &= \alpha\mathbb{P}[\mathbb{P}(X|\mathscr{G})F] + \beta\mathbb{P}[\mathbb{P}(X|\mathscr{G})G] \\ &= \alpha\mathbb{P}(XF) + \beta\mathbb{P}(XG) = \mathbb{P}[X(\alpha F + \beta G)].\end{aligned}$$

Moreover, if $\{F_n\} \subset \mathscr{L}, F_n \geq 0, F \in \mathrm{b}\mathscr{G}$ and $F_n \uparrow F$, then exists $M > 0$ such that $|F| \leq M$. Observe $|\mathbb{P}(X|\mathscr{G})F| \leq M\mathbb{P}(|X||\mathscr{G})$ and

$$\mathbb{P}[M\mathbb{P}(|X||\mathscr{G})] = M\mathbb{P}(|X|) < \infty.$$

Therefore by Dominance convergence theorem, we have

$$\mathbb{P}\left[\mathbb{P}(X|\mathscr{G})F\right] = \lim_{n\to\infty} \mathbb{P}[\mathbb{P}(X|\mathscr{G})F_n] = \lim_{n\to\infty} \mathbb{P}(XF_n) = \mathbb{P}(XF).$$

In conclusion, the Monotone class theorem of the function form means $\mathscr{L}$ contains all the $\mathrm{b}\mathscr{G}$ measurable functions. □

QUESTION 6.13 *Suppose* $(\Omega, \mathscr{F}, \mathbb{P})$ *is a probability space and* X, Y *are two random variables on* $(E, \mathscr{E})$ *and* $(G, \mathscr{G})$ *respectively. Then* X *and* Y *are independent iff for each* $f \in \mathrm{b}\mathscr{E}, g \in \mathrm{b}\mathscr{G}$*, we have*

$$\mathbb{P}[f(X)g(Y)] = \mathbb{P}[f(X)]\mathbb{P}[g(Y)].$$

Proof. We can let $f = 1_A$ and $g = 1_B$, where $A \in \mathscr{E}, B \in \mathscr{G}$. So the definition of independence between X and Y guarantees the above equality holds. On the contrary, if X and Y are independent, then $f(X)$ and $g(Y)$ are independent, the above equality naturally holds. Thus the theorem is completed. □

QUESTION 6.14 *Suppose* $(\Omega, \mathscr{F}, \mathbb{P})$ *is a complete measure space. If* $A, B \in \mathscr{F}$ *and* $\mathbb{P}(A\Delta B) = 0$*, then* $\mathbb{P}(X; A) = \mathbb{P}(X; B)$*.*

Proof. We first prove: if $\mathbb{P}(N) = 0$, then for each $U \in \mathscr{F}, \mathbb{P}(X; U \cup N) = \mathbb{P}(X; U)$. In fact,

$$\int_U X^{\pm} d\mathbb{P} \leq \int_{U \cup N} X^{\pm} d\mathbb{P} \leq \int_U X^{\pm} d\mathbb{P} + \int_N X^{\pm} d\mathbb{P} = \int_U X^{\pm} d\mathbb{P}.$$

Thus $\mathbb{P}(X^{\pm}; U) = \mathbb{P}(X^{\pm}, U \cup N)$ and furthermore we have proved the assertion. Finally, observe $\mathbb{P}(A \Delta B) = 0$, then we have

$$\int_A X d\mathbb{P} = \int_{(A\Delta B)\Delta B} X d\mathbb{P} = \int_{(A\Delta B)\cup B} X d\mathbb{P} - \int_{(A\Delta B)\cap B} X d\mathbb{P} = \int_B X d\mathbb{P}.$$

The last equality holds because $\mathbb{P}\{(A\Delta B) \cap B\} = 0$. □

QUESTION 6.15 *Suppose $\{\mu_n\}$ is a measure sequence on the measure space $(\Omega, \mathscr{F})$. Define $\mu := \sum_n \mu_n$ as follows:*

$$\mu(A) := \sum_{n=1}^{\infty} \mu_n(A), \quad \forall A \in \mathscr{F}.$$

Show μ is a well-defined measure. Besides, for each $f \in \mathscr{F}$, if $\mu(f)$ makes sense, then for each $n \geq 1, \mu_n(f)$ makes sense and we have

$$\int f d\mu = \sum_{n=1}^{\infty} \int f d\mu_n.$$

Proof. Given $A \in \mathscr{F}$, for each $n \geq 1$, we know $\mu_n(A) \geq 0$. So $\mu(A)$ either converge to a finite positive number or equals to ∞. Thus μ is well-defined. Moreover, for each disjoint sets $\{A_i\} \subset \mathscr{F}$, by the definition of μ, notice μ_n is a measure and the Fubini's theorem, we have

$$\mu\left(\sum_{i=1}^{\infty} A_i\right) = \sum_{n=1}^{\infty} \mu_n\left(\sum_{i=1}^{\infty} A_i\right) = \sum_{n=1}^{\infty}\sum_{i=1}^{\infty} \mu_n(A_i) = \sum_{i=1}^{\infty}\sum_{n=1}^{\infty} \mu_n(A_i) = \sum_{i=1}^{\infty} \mu(A_i).$$

Thus μ is a measure. Next we prove the second assertion. When $f = \mathbf{1}_A$, where $A \in \mathscr{F}$, we have

$$\int \mathbf{1}_A d\mu = \mu(A) = \sum_{n=1}^{\infty} \mu_n(A) = \sum_{n=1}^{\infty} \int \mathbf{1}_A d\mu_n.$$

Actually, this is the definition of μ. When $f = \sum_{i=1}^{j} \alpha_i \mathbf{1}_{A_i}$, where $\alpha_i \geq 0$ and $A_i \in \mathscr{F}$ for each $i = 1, \cdots, j$, we can use the previous equality to obtain

$$\begin{aligned}\int \sum_{i=1}^{j} \alpha_i \mathbf{1}_{A_i} d\mu &= \sum_{i=1}^{j} \alpha_j \int \mathbf{1}_{A_i} d\mu = \sum_{i=1}^{j} \alpha_j \sum_{n=1}^{\infty} \int \mathbf{1}_{A_i} d\mu_n \\ &= \sum_{n=1}^{\infty}\sum_{i=1}^{j} \alpha_j \int \mathbf{1}_{A_i} d\mu_n = \sum_{n=1}^{\infty} \int \sum_{i=1}^{j} \alpha_j \mathbf{1}_{A_i} d\mu_n.\end{aligned}$$

When $f \in \mathrm{p}\mathscr{F}$, we can find positive simple function $\{f_i\}$ satisfying $f_i \uparrow f$. Hence use Monotonicity convergence theorem twice, we will acquire

$$\int f d\mu = \lim_{i\to\infty} \int f_i d\mu = \lim_{i\to\infty} \sum_{n=1}^{\infty} \int f_i d\mu_n = \sum_{n=1}^{\infty} \lim_{i\to\infty} \int f_i d\mu_n = \sum_{n=1}^{\infty} \int f d\mu_n.$$

When $f \in \mathscr{F}$, since $\mu(f)$ makes sense, then either $\mu(f^+) < \infty$ or $\mu(f^-) < \infty$. Without loss of generality, we assume $\mu(f^-) < \infty$, then

$$\mu(f^-) = \int f^- d\mu = \sum_{n=1}^{\infty} \int f^- d\mu_n < \infty \Longrightarrow \int f^- d\mu_n < \infty, \quad \forall n \geq 1.$$

Thus

$$\begin{aligned}\int f d\mu &= \int f^+ d\mu - \int f^- d\mu = \sum_{n=1}^{\infty} \int f^+ d\mu_n - \sum_{n=1}^{\infty} \int f^- d\mu_n \\ &= \sum_{n=1}^{\infty} \left(\int f^+ d\mu_n - \int f^- d\mu_n \right) = \sum_{n=1}^{\infty} \int f d\mu_n.\end{aligned}$$

Till now, we have completed the proof. □

QUESTION 6.16 *Suppose $\mathscr{F}_1 = \sigma(\mathscr{G}_1)$ and $\mathscr{F}_2 = \sigma(\mathscr{G}_2)$. Then $\mathscr{F}_1 \times \mathscr{F}_2 = \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\})$.*

Proof. Notice $\mathscr{F}_1 \times \mathscr{F}_2 = \sigma(\{F_1 \times F_2 : F_1 \in \mathscr{F}_1, F_2 \in \mathscr{F}_2\})$ and $\mathscr{G}_1 \subset \mathscr{F}_1, \mathscr{G}_2 \subset \mathscr{F}_2$, then $\mathscr{F}_1 \times \mathscr{F}_2 \supset \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\})$. On the contrary, for all $F_1 \in \mathscr{F}_1$, we have

$$F_1 \times \Omega_2 \subset \sigma(\mathscr{G}_1) \times \Omega_2 \subset \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\}),$$

where $\sigma(\mathscr{G}_1) \times \Omega_2 = \sigma(\{G_1 \times \Omega_2 : G_1 \in \mathscr{G}_1\})$. Similarly, for each $F_2 \in \mathscr{F}_2$,

$$\Omega_1 \times F_2 \subset \Omega_1 \times \sigma(\mathscr{G}_2) \subset \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\}).$$

Thus $F_1 \times F_2 = (F_1 \times \Omega_2) \cap (\Omega_1 \times F_2) \subset \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\})$. This means

$$\{F_1 \times F_2 : F_1 \in \mathscr{F}_1, F_2 \in \mathscr{F}_2\} \subset \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\})$$

and furthermore

$$\begin{aligned}\mathscr{F}_1 \times \mathscr{F}_2 &= \sigma(\{F_1 \times F_2 : F_1 \in \mathscr{F}_1, F_2 \in \mathscr{F}_2\}) \\ &\subset \sigma(\{G_1 \times G_2 : G_1 \in \mathscr{G}_1, G_2 \in \mathscr{G}_2\}).\end{aligned}$$

We finish the proof. □

QUESTION 6.17 *Suppose $\xi : (\Omega, \mathscr{F}) \to (E, \mathscr{E})$ is a random variable. Then for each $U \in \mathscr{E}$, the restriction mapping $\xi : (\xi^{-1}(U), \mathscr{F} \cap \xi^{-1}(U)) \to (U, \mathscr{E} \cap U)$ is also a random variable.*

Proof. For each $A \in \mathscr{E}$, we have

$$\xi^{-1}(A \cap U) = \xi^{-1}(A) \cap \xi^{-1}(U) \in \mathscr{F} \cap \xi^{-1}(U).$$

So the restriction mapping $\xi : (\xi^{-1}(U), \mathscr{F} \cap \xi^{-1}(U)) \to (U, \mathscr{E} \cap U)$ is also a random variable. □

QUESTION 6.18 *Suppose $(\Omega, \mathscr{F}, \mathbb{P})$ is a probability space and $\{X_n\}$ is a random variable sequence. Then $X_n \Rightarrow X$ does not implies $\mathbb{P}(X_n) \to \mathbb{P}(X)$.*

Proof. Consider the following random variable sequence:

$$X_n = \begin{cases} n^2 & \text{, with probability } 1/n \\ 0 & \text{, with probability } 1 - 1/n. \end{cases}$$

Since for each $\varepsilon \in (0, 1)$, we have

$$\mathbb{P}(|X_n| > \varepsilon) = 1/n \to 0, \quad n \to \infty.$$

Thus $X_n \xrightarrow{\mathbb{P}} 0$. Thus we can find a subsequence $\{X_{n_k}\}$ such that $X_{n_k} \xrightarrow{\text{a.s.}} 0$. However, we know for each $k \geq 1$,

$$\mathbb{P}(X_{n_k}) = \int X_{n_k} d\mathbb{P} = \frac{n_k^2}{n_k} = n_k.$$

Thus $\mathbb{P}(X_{n_k}) \to \infty \neq \mathbb{P}(0) = 0$. □

QUESTION 6.19 *Suppose $\{X_n\}$ and $\{Y_n\}$ are two sequences of random variable. If X_n and Y_n are independent for each $n \geq 1$ and $X_n \xrightarrow{\text{a.s.}} X, Y_n \xrightarrow{\text{a.s.}} Y$. Then X and Y are independent.*

Proof. We can utilize the characteristic function and Dominance convergence theorem to obtain

$$\begin{aligned} \mathbb{P}\left[e^{i(t_1 X + t_2 Y)}\right] &= \lim_{n\to\infty} \mathbb{P}\left[e^{i(t_1 X_n + t_2 Y_n)}\right] = \lim_{n\to\infty} \mathbb{P}\left(e^{it_1 X_n}\right) \mathbb{P}\left(e^{it_2 Y_n}\right) \\ &= \mathbb{P}\left(e^{it_1 X}\right) \mathbb{P}\left(e^{it_2 Y}\right). \end{aligned}$$

Thus X and Y are independent. □

QUESTION 6.20 *Suppose μ is a measure on $(\mathbb{R}, \mathscr{B}(\mathbb{R}))$ and dx is the Lebesgue measure. Define $I_n := n\mathbb{1}_{[0,1/n]}$ and*

$$I_n * \mu(x) = \int_{\mathbb{R}} I_n(t)\mu(x - dt).$$

Then discuss: under what conditions the weak convergence $(I * \mu)dx \Rightarrow \mu(dx)$ holds?

Proof. Since

$$\begin{aligned} I_n * \mu(x) &= \int_{\mathbb{R}} I_n(t)\mu(x - dt) = \int_{\mathbb{R}} n\mathbb{1}_{[0,1/n]}(t)\mu(x - dt) \\ &= n\mu(x - [0, 1/n]) = n\mu[x - 1/n, x]. \end{aligned}$$

First we claim $I_n * \mu \in \mathscr{B}(\mathbb{R})$. In fact,

$$I_n * \mu(x) = n\mu[x - 1/n, x] = n\left[F_\mu(x) - F_\mu((x - 1/n)-)\right],$$

where F_μ is the distribution function of μ, namely $F_\mu(x) = \mu(x_0, x]$ for a $x_0 \in \mathbb{R}$ such that $F_\mu(x) < \infty$ for each $x \in \mathbb{R}$. Since F_μ is a non-decreasing function, so $F_\mu \in \mathscr{B}(\mathbb{R})$. Moreover, for each $x \in \mathbb{R}$,

$$F_\mu(x-) = \lim_{y \uparrow x} F_\mu(y) = \lim_{\mathbb{Q} \ni q \uparrow x} F_\mu(q) \in \mathscr{B}(\mathbb{R}).$$

Thus $F_\mu((x - 1/n)-) \in \mathscr{B}(\mathbb{R})$ and therefore $I_n * \mu \in \mathscr{B}(\mathbb{R})$. Suppose μ satisfying for each $f \in \mathrm{b}C(\mathbb{R})$,

$$\int_{\mathbb{R}} \sup_{x \in [t,t+1]} |f(x)|\mu(dt) < \infty.$$

Then we have

$$\begin{aligned} \int_{\mathbb{R}} f(x) I_n * \mu(x)dx &= \int_{\mathbb{R}} nf(x) \int_{[x-1/n,x]} \mu(dt)dx \\ &= \int_{\mathbb{R}} \left(\int_{[t,t+1/n]} nf(x)dx \right) \mu(dt) \\ &= \int_{\mathbb{R}} \left(\fint_{[t,t+1/n]} f(x)dx \right) \mu(dt) \\ &\to \int_{\mathbb{R}} f(t)\mu(dt), \quad n \to \infty. \end{aligned}$$

1. The second equality holds since

$$\begin{aligned} \int_{\mathbb{R}} n|f(x)| \int_{[x-1/n,x]} \mu(dt)dx &= \int_{\mathbb{R}} \left(\int_{[t,t+1/n]} n|f(x)|dx \right) \mu(dt) \\ &\le \int_{\mathbb{R}} \sup_{x \in [t,t+1]} |f(x)|\mu(dt) < \infty. \end{aligned}$$

   Then we can use Fubini's theorem.

2. The last limit holds since

$$\fint_{[t,t+1/n]} f(x)dx \leq \sup_{x\in[t,t+1]} |f(x)|$$

and $\sup_{x\in[t,t+1]} |f(x)|$ is integrable with respect to μ. So we can utilize the Dominance convergence theorem and the integral mean value theorem.

In conclusion, if we suppose μ satisfying for each $f \in \mathrm{b}C(\mathbb{R})$,

$$\int_{\mathbb{R}} \sup_{x\in[t,t+1]} |f(x)|\mu(dt) < \infty,$$

then the weak convergence $(I * \mu)dx \Rightarrow \mu(dx)$ holds. Actually, we claim the above requirement is equivalent to $\mu(\mathbb{R}) < \infty$. In fact, we can choose $1 \in$ b$C(\mathbb{R})$, then

$$\int_{\mathbb{R}} \sup_{x\in[t,t+1]} 1\mu(dt) = \mu(\mathbb{R}) < \infty.$$

The inverse direction is obvious. □

QUESTION 6.21 *Suppose $(\Omega, \mathscr{F}, \mathbb{P})$ and $(E, \mathscr{E})$ are two probability spaces and X, Y are two random variables with respect to $\mathscr{F}$ and $\mathscr{E}$. Assume $\mathscr{G} \subset \mathscr{F}$ is a sub σ-algebra and $X \in \mathscr{G}$, then for each $f \in \mathrm{b}\mathscr{E}^2$, we have*

$$\mathbb{P}[f(X,Y)|\mathscr{G}] = \mathbb{P}[f(x,Y)|\mathscr{G}]|_{x=X}.$$

Moreover, if Y is independent to $\mathscr{G}$, then we have

$$\mathbb{P}[f(X,Y)|\mathscr{G}] = \mathbb{P}[f(x,Y)]|_{x=X}.$$

Especially, if we let $F(x) = \mathbb{P}(x \leq Y)$, then we have

$$\mathbb{P}[F(X)] = \mathbb{P}(X \leq Y).$$

Proof. When $f(x,y) = \mathbb{1}_{A\times B}(x,y)$, where $A, B \in \mathscr{E}$, we have

$$\begin{aligned}\mathbb{P}[f(X,Y)|\mathscr{G}] &= \mathbb{P}[\mathbb{1}_{\{X\in A\}}\mathbb{1}_{\{Y\in B\}}|\mathscr{G}] = \mathbb{1}_{\{X\in A\}}\mathbb{P}[\mathbb{1}_{\{Y\in B\}}|\mathscr{G}] \\ &= \mathbb{1}_A(x)\mathbb{P}[\mathbb{1}_{\{Y\in B\}}|\mathscr{G}]|_{x=X} = \mathbb{P}[\mathbb{1}_A(x)\mathbb{1}_{\{Y\in B\}}|\mathscr{G}]|_{x=X} \\ &= \mathbb{P}[f(x,Y)|\mathscr{G}]|_{x=X}.\end{aligned}$$

Consider the following set

$$\mathscr{L} := \{f(x,y) \in \mathrm{b}\mathscr{E}^2 : \mathbb{P}[f(X,Y)|\mathscr{G}] = \mathbb{P}[f(x,Y)|\mathscr{G}]|_{x=X}\}$$

and write $\mathscr{A} := \{\mathbb{1}_{A\times B}(x,y) : A, B \in \mathscr{E}\}$. Then we know $\mathscr{A} \subset \mathscr{L}$. Obviously, $1 \in \mathscr{L}$, for each $f, g \in \mathscr{L}$ and $\alpha, \beta \in \mathbb{R}$, we have

$$\begin{aligned}\mathbb{P}[\alpha f(X,Y) + \beta g(X,Y)|\mathscr{G}] &= \alpha\mathbb{P}[f(X,Y)|\mathscr{G}] + \beta\mathbb{P}[g(X,Y)|\mathscr{G}] \\ &= \alpha\mathbb{P}[f(x,Y)|\mathscr{G}]|_{x=X} + \beta\mathbb{P}[g(x,Y)|\mathscr{G}]|_{x=X} \\ &= \mathbb{P}[\alpha f(x,Y) + \beta g(x,Y)|\mathscr{G}]|_{x=X}.\end{aligned}$$

So $\mathscr{L}$ is a linear space. Moreover, for $\{f_n\} \subset \mathscr{L}, f \in \mathrm{b}\mathscr{E}$ and $0 \le f_n \uparrow f$, by Monotone convergence theorem, we have

$$\begin{aligned}\mathbb{P}[f(X,Y)|\mathscr{G}] &= \lim_{n\to\infty}\mathbb{P}[f_n(X,Y)|\mathscr{G}] = \lim_{n\to\infty}\mathbb{P}[f_n(x,Y)|\mathscr{G}]|_{x=X} \\ &= \mathbb{P}[f(x,Y)|\mathscr{G}]|_{x=X}.\end{aligned}$$

Define $\mathscr{C} := \{A \times B : A, B \in \mathscr{E}\}$. The Monotone class theorem of function form gives $\mathscr{L}$ contains all the $\sigma(\mathscr{C}) = \mathscr{E}^2$ measurable mappings. The second assertion is trivial.

Observe X and Y are independent, then

$$F(X) = \mathbb{P}(x \le Y)|_{x=X} = \mathbb{P}(\mathbb{1}_{\{X\le Y\}}|X).$$

Therefore,

$$\mathbb{P}[F(X)] = \mathbb{P}[\mathbb{P}(\mathbb{1}_{\{X\le Y\}}|X)] = \mathbb{P}(X \le Y).$$

□

QUESTION 6.22 *Suppose X is a random variable on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and $\mathscr{A}, \mathscr{G}$ are two sub σ-algebras of $\mathscr{F}$. Assume $\mathscr{G}$ and $\sigma(\mathscr{A} \cup \sigma(X))$ are independent, then we have*

$$\mathbb{P}[X|\sigma(\mathscr{A} \cup \mathscr{G})] = \mathbb{P}(X|\mathscr{A}).$$

Proof. For each $A \in \mathscr{A}, G \in \mathscr{G}$, we have

$$\mathbb{P}(X\mathbb{1}_{A\cap G}) = \mathbb{P}(X1_A)\mathbb{P}(G) = \mathbb{P}[\mathbb{P}(X|\mathscr{A})1_A]\mathbb{P}(G) = \mathbb{P}[\mathbb{P}(X|\mathscr{A})\mathbb{1}_{A\cap G}].$$

Define the π-system $\mathscr{C} := \{A \cap G : A \in \mathscr{A}, G \in \mathscr{G}\}$, then we have $\sigma(\mathscr{C}) = \sigma(\mathscr{A} \cup \mathscr{G})$. So the Monotone class theorem yields that the above formula holds for each $B \in \sigma(\mathscr{A} \cup \mathscr{G})$. Obviously $\mathbb{P}(X|\mathscr{A}) \in \sigma(\mathscr{A} \cup \mathscr{G})$, so $\mathbb{P}[X|\sigma(\mathscr{A} \cup \mathscr{G})] = \mathbb{P}(X|\mathscr{A})$. □

QUESTION 6.23 *Suppose $(\Omega, \mathscr{F}, \mathbb{P})$ is a probability space and $\mathscr{A}, \mathscr{G}$ are two independent sub σ-algebras of $\mathscr{F}$. Given two random variables $X \in \mathscr{A}, Y \in \mathscr{G}$ and two σ-algebra $\mathscr{A}' \subset \mathscr{A}, \mathscr{G}' \subset \mathscr{G}$, then we have*

$$\mathbb{P}[XY|\sigma(\mathscr{A}' \cup \mathscr{G}')] = \mathbb{P}(X|\mathscr{A}')\mathbb{P}(Y|\mathscr{G}').$$

Proof. For each $A \in \mathscr{A}', G \in \mathscr{G}'$, observe $X1_A \in \mathscr{A}, Y1_G \in \mathscr{G}$ and the independence between $\mathscr{A}, \mathscr{G}$, we have

$$\begin{aligned}\mathbb{P}(XY1_{A\cap G}) &= \mathbb{P}(X1_A Y1_G) = \mathbb{P}(X1_A)\mathbb{P}(Y1_G)\\ &=\mathbb{P}[\mathbb{P}(X|\mathscr{A}')1_A]\mathbb{P}[\mathbb{P}(Y|\mathscr{G}')1_G] = \mathbb{P}[\mathbb{P}(X|\mathscr{A}')\mathbb{P}(Y|\mathscr{G}')\mathbb{1}_{A\cap G}].\end{aligned}$$

Define the π-system $\mathscr{C} := \{A \cap G : A \in \mathscr{A}', G \in \mathscr{G}'\}$, then we have $\sigma(\mathscr{C}) = \sigma(\mathscr{A}' \cup \mathscr{G}')$. So the Monotone class theorem yields that the above formula holds for each $B \in \sigma(\mathscr{A}' \cup \mathscr{G}')$. Obviously $\mathbb{P}(X|\mathscr{A}')\mathbb{P}(Y|\mathscr{G}') \in \sigma(\mathscr{A}' \cup \mathscr{G}')$, so $\mathbb{P}[XY|\sigma(\mathscr{A}' \cup \mathscr{G}')] = \mathbb{P}(X|\mathscr{A}')\mathbb{P}(Y|\mathscr{G}')$. □

QUESTION 6.24 *Suppose X, Y are two random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and $\mathscr{G}$ is a sub σ-sigma algebra of $\mathscr{F}$. Write $\mathscr{N}$ for all the $\mathbb{P}$-zero set and define the enlargement σ-algebra $\mathscr{G}^* = \sigma(\mathscr{G} \cup \mathscr{N})$. Assume $X = Y, \mathbb{P}$-a.s. and $\mathbb{E}X$ exists, then we have*

$$\mathbb{P}(X|\mathscr{G}) = \mathbb{P}(Y|\mathscr{G}^*), \quad \mathbb{P}\text{-}a.s.$$

Proof. First we have $\mathbb{P}(X|\mathscr{G}) \in \mathscr{G} \subset \mathscr{G}^*$ and $\mathbb{P}(Y|\mathscr{G}^*) \in \mathscr{G}^*$. Moreover, for each $G^* \in \mathscr{G}^*$, exist $G \in \mathscr{G}$ and $N \in \mathscr{N}$ such that $G^* = G\Delta N$. So

$$\mathbb{P}[\mathbb{P}(X|\mathscr{G})\mathbb{1}_{G^*}] = \mathbb{P}[\mathbb{P}(X|\mathscr{G})\mathbb{1}_G] = \mathbb{P}(X1_G) = \mathbb{P}(Y1_G) = \mathbb{P}(Y\mathbb{1}_{G^*}).$$

This yields $\mathbb{P}(X|\mathscr{G}) = \mathbb{P}(Y|\mathscr{G}^*), \mathbb{P}$-a.s.. □

QUESTION 6.25 *Prove $\mathscr{B}(\mathbb{R}_+) = \sigma(\cup_{t>0}\mathscr{B}[0,t))$.*

Proof. Obviously $\sigma(\cup_{t>0}\mathscr{B}[0,t)) \subset \mathscr{B}(\mathbb{R}_+)$. On the contrary, for each $0 \le a < b < \infty$, then exists $t \ge 0$ such that $(a,b] \subset [0,t)$. So $(a,b] \in \mathscr{B}[0,t) \subset \sigma(\cup_{t>0}\mathscr{B}[0,t))$. Therefore

$$\mathscr{B}(\mathbb{R}_+) = \sigma(\{(a,b] : 0 \le a < b < \infty\}) \subset \sigma(\cup_{t>0}\mathscr{B}[0,t)).$$

Thus $\mathscr{B}(\mathbb{R}_+) = \sigma(\cup_{t>0}\mathscr{B}[0,t))$ holds. □

QUESTION 6.26 *Give an example to show X and Y are uncorrelated but not independent.*

Proof. Consider ξ is a random variable satisfying $\mathbb{P}(\xi = 1) = 1/2$ and $\mathbb{P}(\xi = -1) = 1/2$. Moreover, let X is a random variable independent to ξ. Define $Y = \xi X$, the we show X and Y are uncorrelated but not independent. In fact,

$$\mathbb{P}(XY) = \mathbb{P}(\xi X^2) = \mathbb{P}(\xi)\mathbb{P}(X^2) = 0 \Longleftrightarrow \mathrm{Cov}(XY) = 0.$$

However, obviously X and $Y = \xi X$ are not independent in general. □

QUESTION 6.27 *Consider the measurable space $(E, \mathscr{B}(E))$, where E is a metric space and $\mathscr{B}(E)$ is the Borel σ algebra of E. When $A \in \mathscr{B}(E)$ is open or closed, give some examples to show 1_A can be a pointwise limit of Lipschitz continuous function sequence.*

Proof. 1. When A is an open set.

(a) Consider the following continuous function sequence:

$$F_n(x) = [nd(x, A^c)] \wedge 1, \quad x \in E, n \geq 1.$$

Since A^c is a closed subset, then $d(x, A^c) > 0$ iff $x \in A$. Then it is easy to find $F_n \uparrow 1_A$. Besides, F_n is Lipschitz continuous. We just need to notice **QUESTION 2.1**, we have

$$\begin{aligned} |F_n(x) - F_n(y)| &= |[nd(x, A^c)] \wedge 1 - [nd(y, A^c)] \wedge 1| \\ &\leq |nd(x, A^c) - nd(y, A^c)| = nd(x, y). \end{aligned}$$

(b) Consider the following continuous function sequence:

$$F_n(x) = [nd(x, A^c) - 1]^+ \wedge 1, \quad x \in E, n \geq 1.$$

Since A^c is a closed subset, then $d(x, A^c) > 0$ iff $x \in A$. Then it is easy to find $F_n \uparrow 1_A$. Besides, F_n is Lipschitz continuous. We just need to notice **QUESTION 2.1**, we have

$$\begin{aligned} |F_n(x) - F_n(y)| &= |[nd(x, A^c) - 1]^+ \wedge 1 - [nd(x, A^c) - 1]^+ \wedge 1| \\ &\leq |[nd(x, A^c) - 1]^+ - [nd(t, A^c) - 1]^+| \\ &\leq |nd(x, A^c) - nd(y, A^c)| = nd(x, y). \end{aligned}$$

2. When A is a closed set.

(a) Consider the following continuous function sequence:

$$F_n(x) = [1 - nd(x, A)]^+, \quad x \in E, n \geq 1.$$

Since A is a closed subset, then $d(x, A) = 0$ iff $x \in A$. Then it is easy to find $F_n \uparrow 1_A$. Besides, F_n is Lipschitz continuous. We just need to notice **QUESTION 2.1**, we have

$$\begin{aligned} |F_n(x) - F_n(y)| &= |[1 - nd(x, A)]^+ - [1 - nd(y, A)]^+| \\ &\leq |nd(x, A) - nd(y, A)| = nd(x, y). \end{aligned}$$

(b) Consider the following continuous function sequence:

$$F_n(x) = \frac{1}{1 + nd(x, A)}, \quad x \in E, n \geq 1.$$

Since A is a closed subset, then $d(x, A) = 0$ iff $x \in A$. Then it is easy to find $F_n \uparrow 1_A$. Besides, F_n is Lipschitz continuous. In fact, we have

$$\begin{aligned} |F_n(x) - F_n(y)| &= \left| \frac{1}{1 + nd(x, A)} - \frac{1}{1 + nd(y, A)} \right| \\ &= \left| \frac{n[d(y, A) - d(x, A)]}{[1 + nd(x, A)][1 + nd(y, A)]} \right| \\ &\leq |nd(x, A) - nd(y, A)| = nd(x, y). \end{aligned}$$

□

QUESTION 6.28 *Prove a σ-algebra $\mathscr{F}$ is either finite or uncountable.*

Proof. We prove this by contradiction. Assume $|\mathscr{F}| = \aleph_0$. Define

$$\mathscr{G} := \{A \in \mathscr{F} : \text{there is no } B_1, B_2 \subsetneq A \text{ such that } A = B_1 \cup B_2 \backslash \{\emptyset\}.$$

Obviously $|\mathscr{G}| \leq \aleph_0$, say $\mathscr{G} = \{A_k\}_{k \geq 1}$. We claim $A_i \cap A_j = \emptyset$ for any $i \neq j$. We prove this by contradiction. If this is not the case, we have $A_i \cap A_j \neq \emptyset$.

1. Without loss of generality, if $A_i \cap A_j = A_i$, then

$$A_j = A_i \cup (A_j \backslash A_i).$$

Noting $A_i \neq A_j$ and $A_i, A_j \backslash A_j \in \mathscr{F}$, this contradict the fact $A_j \in \mathscr{G}$.

2. If $A_i \cap A_j \neq A_i$ and $A_i \cap A_j \neq A_j$, we have

$$A_i = (A_i \cap A_j) \cup (A_i \backslash A_j).$$

Noting $A_i \cap A_j \in \mathscr{F}$ and $A_i \backslash A_j \in \mathscr{F}$, this contradict the fact $A_i \in \mathscr{G}$.

In conclusion, we have shown the elements in $\mathscr{G}$ are mutually disjoint.

Now, we consider several situations:

1. If $\sigma(\mathscr{G}) = \mathscr{F}$, noting $|\mathscr{F}| = \aleph_0$, we must have $|\mathscr{G}| = \aleph_0$. since the elements in $\mathscr{G}$ are mutually disjoint, the well-known result shows

$$\mathscr{F} = \sigma(\mathscr{G}) = \left\{ \sum_{k \in I} A_k : I \subset \mathbb{Z}_+ \right\}.$$

So we can construct a bijection:

$$\varphi : 2^{\mathbb{Z}_+} \to \mathscr{F}, I \mapsto \sum_{k\in I} A_k,$$

where $2^{\mathbb{Z}_+}$ is the set of all the subset of $\mathbb{Z}_+$. This implies $|\mathscr{F}| = |\sigma(\mathscr{G})| = \aleph$, which contradicts $|\mathscr{F}| = \aleph_0$.

2. If $\sigma(\mathscr{G}) \subsetneq \mathscr{F}$, there must exists $A \in \mathscr{F}$ such that $A \notin \sigma(\mathscr{G})$. Since $A \notin \mathscr{G}$, there exist $A_i^{(1)} \in \mathscr{F}$ and $A_i^{(1)} \subsetneq A$ $(i = 1, 2)$ such that $A = A_1^{(1)} \cup A_2^{(1)}$. Noting $A \notin \sigma(\mathscr{G})$, so there exists $n_1 = 1$ or 2 such that $A_{n_1}^{(1)} \notin \sigma(\mathscr{G})$ and thus $A_{n_1}^{(1)} \notin \mathscr{G}$. So there exist $A_i^{(2)} \in \mathscr{F}$ and $A_i^{(2)} \subsetneq A_{n_1}^{(1)}$ $(i = 1, 2)$ such that $A_{n_1}^{(1)} = A_1^{(2)} \cup A_2^{(2)}$. Again, $A_{n_1}^{(1)} \notin \sigma(\mathscr{G})$ yields that there exists $n_2 = 1$ or 2 such that $A_{n_2}^{(2)} \notin \sigma(\mathscr{G})$. Using induction, we can obtain a sequence $\{A_{n_k}^{(k)}\}_{k\geq 1} \subset \mathscr{F}$ satisfying

$$A_{n_0}^{(0)} := A \supsetneq A_{n_1}^{(1)} \supsetneq A_{n_2}^{(2)} \supsetneq \cdots \supsetneq A_{n_k}^{(k)} \cdots .$$

Define for any $k \geq 1$,

$$C_k := A_{n_{k-1}}^{(k-1)} - A_{n_k}^{(k)}.$$

Then $\{C_k\}_{k\geq 1}$ is a disjoint set sequence. Since $\{C_k\}_{k\geq 1} \subset \mathscr{F}$, then

$$\sigma\left(\{C_k\}_{k\geq 1}\right) = \left\{\sum_{k\in I} C_k : I \subset \mathbb{Z}_+\right\} \subset \mathscr{F}$$

A similar argument will yield contradiction again.

In conclusion, $|\mathscr{F}| \neq \aleph_0$ for any σ-algebra $\mathscr{F}$.

A fake proof. If $\mathscr{G}$ is countable, denoted as $\mathscr{G} = \{A_n\}_{n\geq 1}$. Since $\cup_n A_n \in \mathscr{G}$, then exists $n_0 \geq 1$ such that $\cup_n A_n = A_{n_0}$. Note $A_{n_0}^c \in \mathscr{G}$, then exists $n_1 \geq 1$ such that $A_{n_0}^c = A_{n_1}$. So

$$A_{n_0}^c = A_{n_1} \subset \bigcup_{n\geq 1} A_n = A_{n_0}.$$

This is a contradiction. This proof is not grounded because $\cup_n A_n = A_{n_0} = \Omega$ and $A_{n_1} = A_{n_0}^c = \emptyset$. So $A_{n_0}^c = \emptyset \subset A_{n_0} = \Omega$ naturally true and there is no contradiction. □

QUESTION 6.29 *Suppose D is a dense subset of $\mathbb{R}$. If the distribution function $F_n(x) \to F(x)$ for each $x \in D$ and F is continuous, then*

$$\sup_{x\in\mathbb{R}} |F_n(x) - F(x)| \to 0, \qquad n \to \infty.$$

Thus $F_n(x) \xrightarrow{\text{w}} F(x)$ naturally.

Proof. At first, we show if $\exists M \in D \cap [0,\infty)$, s.t. $F_n(x) \to F(x), \forall x \in D \cap [-M, M]$, then F_n converges to F uniformly on $D \cap [-M, M]$. In fact, since F is uniformly continuous on $[-M, M]$, then $\forall \varepsilon > 0, \exists \delta > 0, \forall x, y \in [-M, M]$ and $|x - y| < \delta$, then $|F(x) - F(y)| < \varepsilon$. Hence we can separate $[-M, M]$ into n subintervals: $-M = x_0 < x_1 < \cdots < x_n = M$ satisfying $\{x_j\}_{j=0}^n \subset D \cap [-M, M]$ and $\max_{0\le j\le n-1} |x_{j+1} - x_j| < \delta$. Moreover, since $F_n(x_j) \to F(x_j), \forall j \in \{0, \cdots, n\}$, then $\exists N_1 \in \mathbb{N}, \forall n \ge N_1$, we have $|F_n(x_j) - F(x_j)| < \varepsilon, \forall j \in \{0, \cdots, n\}$. When $n \ge N_1$, for each $x \in [-M, M]$, observe $\exists j \in \{0, \cdots, n\}$, s.t. $x \in [x_n, x_{n+1}]$, then

$$\begin{aligned}
&|F_n(x) - F(x)| \\
&\quad \le |F_n(x) - F_n(x_n)| + |F_n(x_n) - F(x_n)| + |F(x_n) - F(x)| \\
&\quad < |F_n(x_{n+1}) - F_n(x_n)| + \varepsilon + \varepsilon \\
&\quad \le |F_n(x_{n+1}) - F(x_{n+1})| + |F(x_{n+1}) - F(x_n)| + |F(x_n) - F_n(x_n)| + 2\varepsilon \\
&\quad < \varepsilon + \varepsilon + \varepsilon + 2\varepsilon = 5\varepsilon.
\end{aligned}$$

This implies F_n converges to F uniformly on $[-M, M]$.

Next, notice $\lim_{x\to-\infty} F(x) = 0, \lim_{x\to\infty} F(x) = 1$. Then $\exists M \in D \cap [0,\infty)$, s.t. $F(-M) < \varepsilon$ and $1 - F(M) < \varepsilon$. Note $F_n(\pm M) \to F(\pm M)$, then $\exists N_2 \in \mathbb{N}, \forall n \ge N_2$, then $|F_n(\pm M) - F(\pm M)| < \varepsilon$. So when $n \ge N_2$, we have

$$F_n(-M) \le |F_n(-M) - F(-M)| + F(-M) < \varepsilon + \varepsilon = 2\varepsilon$$

and

$$1 - F_n(M) \le 1 - F(M) + |F(M) - F_n(M)| < \varepsilon + \varepsilon = 2\varepsilon.$$

Whence $\forall x \le -M$, we have

$$|F_n(x) - F(x)| \le F_n(x) + F(x) \le F_n(-M) + F(-M) < 2\varepsilon + \varepsilon = 3\varepsilon.$$

And $\forall x \ge M$, we have

$$\begin{aligned}
|F_n(x) - F(x)| &\le |F_n(x) - 1| + |1 - F(x)| \\
&\le [1 - F_n(M)] + [1 - F(M)] < 2\varepsilon + \varepsilon = 3\varepsilon.
\end{aligned}$$

This implies F_n converges to F uniformly on $[-M, M]^c$.

In conclusion F_n converges to F uniformly on $\mathbb{R}$, namely, $\sup_{x\in\mathbb{R}} |F_n(x) - F(x)| \to 0$. As for the last assertion, since $F(x)$ is continuous on $\mathbb{R}$, then for each $x \in \mathbb{R}$, we have

$$|F_n(x) - F(x)| \le \sup_{x\in\mathbb{R}} |F_n(x) - F(x)| \to 0,$$

which implies $F_n(x) \xrightarrow{\text{w}} F(x)$. $\square$

QUESTION 6.30 *Suppose $\{\nu_n\}$ is a sequence of normal probability distributions $N(\mu_n, \sigma_n^2)$ and $\nu_n \xrightarrow{\text{w}} \nu$. Then ν is also a normal probability distribution.*

Proof. Write $f_n(x) = \int e^{itx}\nu_n(dx)$ and $f(x) = \int e^{itx}\nu(dx)$. Then the weak convergence implies

$$f_n(x) = e^{i\mu_n x - \sigma_n^2 x^2/2} \to f(x).$$

Therefore,

$$|f_n(x)| = e^{-\sigma_n^2 x^2/2} \to |f(x)|.$$

By the continuity of $f(x)$ at $x = 0$ and note $f(0) = 1$, we know exists $x_0 \neq 0$ small enough, such that $f(x_0) \neq 0$. Thus

$$-\sigma_n^2 x_0^2/2 \to \log|f(x_0)| \neq -\infty,$$

which means $\lim_{n\to\infty} \sigma_n^2$ exists and denoted as σ^2. Till now, we have

$$e^{i\mu_n x} = f_n(x)e^{\sigma_n^2 x^2/2} \to f(x)e^{\sigma^2 x^2/2}.$$

We still use the above $x_0 \neq 0$, then $f(x_0) \neq 0$. Therefore

$$\mu_n \to \frac{1}{ix_0}\text{Log} f(x_0)e^{\sigma^2 x_0^2/2} \in \mathbb{R},$$

where $\text{Log}(\cdot)$ is the main value of the complex logarithmic function. This implies $\mu := \lim_{n\to\infty} \mu_n$ exists. Moreover, by the uniqueness of limit, we know $\lim_{n\to\infty} \mu_n$ does not depends on the selection of x_0. In conclusion, we have

$$f_n(x) = e^{i\mu_n x - \sigma_n^2 x^2/2} \to e^{i\mu x - \sigma^2 x^2/2} = f(x),$$

which means $\nu(dx) \sim N(\mu, \sigma^2)$. □

QUESTION 6.31 *Suppose X, Y are two $(E, \mathscr{E})$ random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Assume $\mathscr{G} \subset \mathscr{F}$ is a sub σ algebra and for each $f, g \in \text{b}\mathscr{E}$,*

$$\mathbb{P}[f(X)g(Y)|\mathscr{G}] = \mathbb{P}[f(X)]\mathbb{P}[g(Y)].$$

Then X and Y are independent. Especially, assume $\{C_k\}_{k\geq 1}$ is a partition of Ω and $\mathbb{P}(C_k) > 0, \forall k \geq 1$. If for each $k \geq 1$ and $A, B \in \mathscr{E}$,

$$\mathbb{P}(X \in A, Y \in B|C_k) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B),$$

then X and Y are independent.

Proof. Take expectation on both sides of $\mathbb{P}[f(X)g(Y)|\mathscr{G}] = \mathbb{P}[f(X)]\mathbb{P}[g(Y)]$ to obtain

$$\mathbb{P}[f(X)g(Y)] = \mathbb{P}\{\mathbb{P}[f(X)g(Y)|\mathscr{G}]\} = \mathbb{P}[f(X)]\mathbb{P}[g(Y)].$$

This implies X and Y are independent. Especially, we let $f = 1_A, g = 1_B$ and $\mathscr{G} = \sigma(C_k : k \geq 1)$. Then $\mathbb{P}(X \in A, Y \in B|C_k) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$ is equivalent to $\mathbb{P}[f(X)g(Y)|\mathscr{G}] = \mathbb{P}[f(X)]\mathbb{P}[g(Y)]$. So the result follows. □

QUESTION 6.32 *Suppose X is an exponential distribution random variable with parameter μ and Y is an independent variable on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Then for each $t > 0$, we have*

$$\mathbb{P}(X > t + Y|X > Y, Y \geq 0) = \mathbb{P}(X > t).$$

Proof. When Y is a nonnegative constant, this is the well-known memoryless property. When Y is a random variable, write its distribution as $\nu(dx)$. Observe for each $t \geq 0$, the independence between X and Y impies

$$\begin{aligned}
\mathbb{P}(X > t + Y, Y \geq 0) &= \mathbb{P}(\mathbb{1}_{\{X>t+Y\}}\mathbb{1}_{\{Y\geq 0\}}) \\
&= \mathbb{P}[\mathbb{P}(\mathbb{1}_{\{X>t+Y\}}\mathbb{1}_{\{Y\geq 0\}}|Y)] \\
&= \mathbb{P}[\mathbb{1}_{\{Y\geq 0\}}\mathbb{P}(\mathbb{1}_{\{X>t+Y\}}|Y)] \\
&= \mathbb{P}[\mathbb{1}_{\{Y\geq 0\}}\mathbb{P}(\mathbb{1}_{\{X>t+y\}}|Y)|_{y=Y}] \\
&= \mathbb{P}[\mathbb{1}_{\{Y\geq 0\}}\mathbb{P}(\mathbb{1}_{\{X>t+y\}})|_{y=Y}] \\
&= \mathbb{P}[\mathbb{1}_{\{Y\geq 0\}}\mathbb{P}(X > t + y)|_{y=Y}] \\
&= \int_{-\infty}^{\infty} \mathbb{1}_{\{y\geq 0\}}\mathbb{P}(X > t + y)\mu(dy) \\
&= \int_{0}^{\infty} \mathbb{P}(X > t + y)\mu(dy).
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
\mathbb{P}(X >t + Y|X > Y, Y \geq 0) &= \frac{\mathbb{P}(X > t + Y, Y \geq 0)}{\mathbb{P}(X > Y, Y \geq 0)} \\
&= \int_0^\infty \mathbb{P}(X > t + y)\nu(dy) \Big/ \int_0^\infty \mathbb{P}(X > y)\nu(dy) \\
&= \int_0^\infty e^{-\mu(t+y)}\nu(dy) \Big/ \int_0^\infty e^{-\mu y}\nu(dy) \\
&= e^{-\mu t}\int_0^\infty e^{-\mu y}\nu(dy) \Big/ \int_0^\infty e^{-\mu y}\nu(dy) \\
&= e^{-\mu t} = \mathbb{P}(X > t).
\end{aligned}$$

This shows a strong version of memoryless property. □

QUESTION 6.33 *Suppose X and Y are two $(E, \mathscr{E})$ random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. If exists a distribution $\nu(dx)$ on $(E, \mathscr{E})$ such that for each $A \in \mathscr{E}$, we have $\mathbb{P}(X \in A|Y) = \nu(A)$, then X and Y are independent. Moreover, the distribution of X is $\nu(dx)$.*

Proof. Take the expectation on both sides of $\mathbb{P}(X \in A|Y) = \nu(A)$, we have $\mathbb{P}(X \in A) = \nu(A)$. Therefore, the distribution of X is $\nu(dx)$. Since for each $B \in \mathscr{E}$, we have

$$\begin{aligned}\mathbb{P}(X \in A, Y \in B) &= \mathbb{P}[\mathbb{P}(X \in A, Y \in B|Y)] \\ &= \mathbb{P}[\mathbb{1}_{\{Y \in B\}}\mathbb{P}(X \in A|Y)] \\ &= \mathbb{P}[\mathbb{1}_{\{Y \in B\}}\nu(A)] \\ &= \nu(A)\mathbb{P}(Y \in B).\end{aligned}$$

The desired result follows. □

QUESTION 6.34 *Given a sequence of integrable i.i.d. random variables $\{X_n\}_{n\geq 1} : (\Omega, \mathscr{F}, \mathbb{P}) \to (\mathbb{R}, \mathscr{B}(\mathbb{R}))$. Then for each sequence of random times $\{T_m\}_{m\geq 1} : \Omega \to \mathbb{Z}_+$, where $T_1 < T_2 < \cdots < T_m < \cdots$, we have*

$$\mathbb{P}\left(\sum_{m=1}^{\infty} X_{T_m} = \infty\right) = 1.$$

Proof. For any subsequence $\{n_m\}_{m\geq 1}$ of $\mathbb{Z}_+$, we know $\{X_{n_m}\}_{m\geq 1}$ is also an integrable i.i.d. random variable sequence. Thus, by the strong law of large numbers, we have

$$\mathbb{P}\left(\sum_{m=1}^{\infty} X_{n_m} = \infty\right) = 1.$$

Write N for the exceptional set with $\mathbb{P}(N) = 0$. Then

$$\sum_{m=1}^{\infty} X_{n_m}(\omega) = \infty, \quad \forall \omega \in N^c.$$

Given any $\omega \in N^c$, notice $\{T_m(\omega)\}_{m\geq 1}$ is a subsequence of $\mathbb{Z}_+$, then

$$\sum_{m=1}^{\infty} X_{T_m(\omega)}(\omega) = \infty,$$

which implies

$$\mathbb{P}\left(\sum_{m=1}^{\infty} X_{T_m} = \infty\right) = 1.$$

□

QUESTION 6.35 *Suppose F is a distribution function of the random variable X, i.e., $F(x) = \mathbb{P}(-\infty, x]$ and define:*

$$F^{-1}(u) = \inf\{x \in \mathbb{R} : F(x) \geq u\}.$$

Then $F(x) \geq u$ iff $x \geq F^{-1}(u)$.

Proof. By the definition of infimum, we know $F(x) \geq u$ implies $x \geq F^{-1}(u)$. On the contrary, by the right continuity of F and the definition of infimum, exists $\{x_n\} \subset \mathbb{R}$ such that $F(x_n) \geq u$ and $x_n \downarrow F^{-1}(u)$. Moreover, observe F is increasing, then $F(x) \geq F(F^{-1}(u)) = \lim_{n\to\infty} F(x_n) \geq u$.. □

QUESTION 6.36 *Suppose $T > 0$. Assume $F : [0, T] \to \mathbb{R}$ is left continuous and $F(0) = 0$. Then the signed measure μ induced by F has no atoms, where $\mu[0, t] = F(t), \forall t \in [0, T]$.*

Proof. Actually, $F : [0, T] \to \mathbb{R}$ must be continuous. In fact, the left continuity is given. The right continuity is obtained by noticing $F(t) = \mu[0, t]$.

For each $t \in (0, T]$, consider $[0, t - 1/n]$ for $n > 1/t$. Then we have

$$\begin{aligned}\mu(\{t\}) &= \lim_{n\to\infty} \mu(t - 1/n, t] = \mu[0, t] - \lim_{n\to\infty} \mu[0, t - 1/n] \\ &= F(t) - \lim_{n\to\infty} F(t - 1/n) = F(t) - F(t-) = 0.\end{aligned}$$

For $t = 0$, we have $\mu(\{0\}) = \mu[0, 0] = F(0) = 0$. Therefore $\mu(\{t\}) = 0$ for each $t \in [0, T]$. Namely, the signed measure μ induced by F has no atoms. □

QUESTION 6.37 *Suppose $F : \mathbb{R}_+ \to \mathbb{R}$ is a right continuous non-decreasing function. Then exists a unique σ-finite measure μ on $(\mathbb{R}_+, \mathscr{B}(\mathbb{R}_+))$ such that for each $T > 0, \mu|_{[0,T]}$ is the measure induced by $F|_{[0,T]}$, denoted as $\mu_F|_{[0,T]}$ (since all the definitions on finite interval $[0, T]$ is already done).*

Proof. Define

$$\mathscr{C} = \{(a, b] : 0 < a \leq b \leq \infty\} \cup \{[0, c] : c \geq 0\}.$$

Then $\mathscr{C}$ is a semi-set algebra on $\mathbb{R}_+$ and $\sigma(\mathscr{C}) = \mathscr{B}(\mathbb{R}_+)$. Define

$$\begin{aligned}&\mu(\mathbb{R}_+) := \lim_{t\to\infty} F(t) - F(0), \quad \text{the limit exists since } F \text{ is non-decreasing,} \\ &\mu(a, b] := F(b) - F(a), \quad \mu[0, c] := F(c) - F(0).\end{aligned}$$

Then μ is indeed a finite measure on $\mathscr{C}$. So the Measure extension theorem shows exists a unique measure μ on $(\mathbb{R}_+, \mathscr{B}(\mathbb{R}_+))$. Moreover μ is σ-finite obviously. Since for each $T > 0$, consider

$$\mathscr{C}_T = \{(a, b] : 0 < a \leq b \leq T_1\} \cup \{[0, c] : 0 \leq c \leq T_1\}.$$

Then $\forall A \in \mathscr{C}_T$, we have $\mu(A) = \mu_F|_{[0,T]}(A)$. The uniqueness of Measure extension theorem still yields $\mu|_{[0,T]} = \mu_F|_{[0,T]}$. □

QUESTION 6.38 *Suppose Ω is the sample space and μ is a measure on semi-set algebra $\mathscr{A}$. Denote also by μ the extension of μ to $\sigma(\mathscr{A})$. Define the inner measure as follows:*

$$\mu_*(A) = \sup\left\{\sum_n \mu(A_n) : \{A_n\} \subset \mathscr{A} \text{ and } \bigcup_n A_n \subset A\right\}.$$

Then

$$\mu^*(A) = \inf\{\mu(B) : A \subset B, B \in \sigma(\mathscr{A})\} =: \lambda^*(A)$$

and

$$\mu_*(A) = \sup\{\mu(C) : C \subset A, C \in \sigma(\mathscr{A})\} =: \lambda_*(A).$$

Moreover, the infimum and supremum are both achieved. Finally, we have A is μ^*-measurable iff $\mu^*(A) = \mu_*(A)$.

Proof. For each $A \subset \Omega$, by the definition of the outer measure, for each $\varepsilon > 0$, exists $\{A_n\} \subset \mathscr{A}$ and $A \subset \cup_n A_n$ such that

$$\sum_n \mu(A) < \mu^*(A) + \varepsilon.$$

Note $\cup_n A_n \in \sigma(\mathscr{A})$, then

$$\lambda^*(A) \leq \mu\left(\bigcup_n A_n\right) \leq \sum_n \mu(A) < \mu^*(A) + \varepsilon.$$

The arbitrariness of ε shows $\lambda^*(A) \leq \mu^*(A)$. On the contrary, for each $B \in \sigma(\mathscr{A})$, $A \subset B$, we have

$$\mu^*(A) \leq \mu^*(B) = \mu(B).$$

Take $\inf_{A\subset B, B\in\sigma(\mathscr{A})}$ on both sides of the above inequality yields $\mu^*(A) \leq \lambda^*(A)$. In conclusion, we have $\mu^*(A) = \lambda^*(A)$. Let $\varepsilon = 1/k$, then exists $\{A_n^k\} \subset \mathscr{A}$ cover A and

$$\mu^*(A) \leq \mu\left(\bigcup_n A_n^k\right) < \mu^*(A) + 1/k.$$

Then consider $B = \cap_k \cup_n A_n^k$, then obviously

$$\mu^*(A) = \mu(B).$$

The results for the inner measure follows similarly.

If A is μ^*-measurable, then obviously $\mu^*(A) = \mu_*(A)$. On the contrary, if $\mu^*(A) = \mu_*(A)$, since the above infimum and supremum are both achieved, then we can choose $A_1, A_2 \in \sigma(\mathscr{A}), A_1 \subset A \subset A_2$ and $\mu(A_1) = \mu(A_2)$. For each $E \subset \Omega$, choose $B \in \sigma(\mathscr{A})$ such that $\mu^*(E) = \mu(B)$. Then

$$\begin{aligned}\mu^*(A \cap E) + \mu^*(A^c \cap E) &\leq \mu(A_2 \cap B) + \mu(A_1^c \cap B) \\ &\leq \mu(B) + \mu(A_2 \backslash A_1) = \mu(B) = \mu^*(E).\end{aligned}$$

So A is μ^*-measurable. □

QUESTION 6.39 *If $(X_n)_{n\geq 1}$ is a random variable sequence on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Assume $\mathscr{G} \subset \mathscr{F}$ and $X_n \xrightarrow{L^p} X$, where $p \geq 1$. Then*

$$\mathbb{P}(X_n|\mathscr{G}) \xrightarrow{L^p} \mathbb{P}(X|\mathscr{G}).$$

Proof. Since we have

$$\begin{aligned}\mathbb{P}[|\mathbb{P}(X_n|\mathscr{G}) - \mathbb{P}(X|\mathscr{G})|^p] &\leq \mathbb{P}[\mathbb{P}(|X_n - X|^p|\mathscr{G})] \\ &= \mathbb{P}(|X_n - X|^p) \to 0, \quad n \to \infty.\end{aligned}$$

So $\mathbb{P}(X_n|\mathscr{G}) \xrightarrow{L^p} \mathbb{P}(X|\mathscr{G})$. □

QUESTION 6.40 *Presume $X \geq 0$ is a random variable on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$, then we have*

$$\mathbb{P}(X) = \int_0^\infty \mathbb{P}(X > x)dx = \int_0^\infty [1 - F(x)]dx.$$

Prove when X is not nonnegative, we have the following expression:

$$\mathbb{P}(X) = \int_0^\infty [1 - F(x)]dx + \int_{-\infty}^0 F(x)dx.$$

Proof. We have

$$\begin{aligned}\mathbb{P}(X) &= \int X d\mathbb{P} = \int \mathbb{1}_{\{X\geq 0\}} X d\mathbb{P} + \int \mathbb{1}_{\{X<0\}} X d\mathbb{P} \\ &= \int \mathbb{1}_{\{X\geq 0\}} \int_0^\infty \mathbb{1}_{\{x<X\}} dx d\mathbb{P} + \int \mathbb{1}_{\{X<0\}} \int_{-\infty}^0 \mathbb{1}_{\{x\geq X\}} d\mathbb{P} \\ &= \int \int_0^\infty \mathbb{1}_{\{x<X\}} dx d\mathbb{P} + \int \int_{-\infty}^0 \mathbb{1}_{\{x\geq X\}} dx d\mathbb{P} \\ &= \int_0^\infty \mathbb{P}(X > x)dx + \int_{-\infty}^0 \mathbb{P}(X \leq x)dx \\ &= \int_0^\infty [1 - F(x)]dx + \int_{-\infty}^0 F(x)dx.\end{aligned}$$

□

QUESTION 6.41 *Consider measurable space* $(\mathbb{R}^n, \mathscr{B}^n)$. *Then* $\mu_k \xrightarrow{v} \mu$ *iff for all the continuous intervals* I *of* μ, $\mu_k(I) \to \mu(I)$.

Proof. (1) Sufficiency. Suppose for all the continuous intervals I of μ, $\mu_k(I) \to \mu(I)$, then for all $f \in C_0(\mathbb{R}^n)$, exist $M > 0$, s.t. $|f(x)| \leq M$. Note $C(\mu)$ is dense in $\mathbb{R}^n$, we can find a continuous interval I of μ, such that $\operatorname{supp}(f) \subset I$. Moreover, notice f is uniformly continuous on I, then $\forall \varepsilon > 0, \exists \delta > 0, \forall |x-y| < \delta$, then $|f(x) - f(y)| < \varepsilon$. Thus we can find a finite partition of I, say $\{I_\ell\}_{\ell=1}^n$, such that every I_ℓ is the continuous interval of μ, the radius of I_k is lease than δ so that $\forall x, y \in I_\ell, |f(x) - f(y)| < \varepsilon$. Define the following simple function:

$$g = \sum_{\ell=1}^n f(\alpha_\ell) \mathbf{1}_{I_\ell},$$

where $\alpha_\ell \in I_\ell$. Then $|f(x) - g(x)| < \varepsilon, \forall x \in I$ and

$$\begin{aligned}
&\left| \int f d\mu_k - \int f d\mu \right| = \left| \int_I f d\mu_k - \int_I f d\mu \right| \\
\leq &\left| \int_I f d\mu_k - \int_I g d\mu_k \right| + \left| \int_I g d\mu_k - \int_I g d\mu \right| + \left| \int_I g d\mu - \int_I f d\mu \right| \\
\leq &\int_I |f - g| d\mu_k + \sum_{\ell=1}^n f(\alpha_\ell) |\mu_k(I_\ell) - \mu(I_\ell)| + \int_I |f - g| d\mu \\
< &\varepsilon \left[\mu_k(I) + \mu(I) \right] + M \sum_{\ell=1}^n |\mu_k(I_\ell) - \mu(I_\ell)|.
\end{aligned}$$

Let $k \to \infty$ and $\varepsilon \to 0$ to obtain

$$\overline{\lim_{k \to \infty}} \left| \int f d\mu_k - \int f d\mu \right| = 0.$$

(2) Necessity. For all continuous interval I of μ, write $I = [a, b]$. Then $\forall \varepsilon > 0, \exists \delta > 0$ small enough, such that $a + \delta < b - \delta$ and $\mu(U) < \varepsilon$, where

$$U = (a - \delta, a + \delta) \cup (b - \delta, b + \delta).$$

We can construct two functions $f, g \in C_0(\mathbb{R}^n)$, s.t. $f \leq \mathbf{1}_I \leq g$ and $|f - g| \leq I_U$. Thus

$$\int f d\mu_k \leq \mu_k(I) \leq \int g d\mu_k$$

and

$$\int f d\mu \leq \mu(I) \leq \int g d\mu.$$

Combine the above two formula to obtain

$$|\mu_k(I) - \mu(I)| \leq \min\left\{\left|\int g d\mu - \int f d\mu_k\right|, \left|\int g d\mu_k - \int f d\mu\right|\right\}.$$

Let $k \to \infty$ to obtain

$$\begin{aligned}\overline{\lim_{k\to\infty}} |\mu_k(I) - \mu(I)| \leq & \min\left\{\left|\int g d\mu - \int f d\mu\right|, \left|\int g d\mu - \int f d\mu\right|\right\} \\ = & \left|\int g d\mu - \int f d\mu\right| \leq \int |f - g| d\mu \\ \leq & \int \mathbf{1}_U d\mu = \mu(U) < \varepsilon.\end{aligned}$$

Let $\varepsilon \to 0$, we will obtain

$$\overline{\lim_{k\to\infty}} |\mu_k(I) - \mu(I)| = 0.$$

□

QUESTION 6.42 *Given finite measure sequence $\{\mu_n\}_{n\in\mathbb{N}}$ and μ on measurable space $(\Omega, \mathscr{F})$. Prove the following assertions.*

1. *The uniformly convergence:*

$$\sup_{A\in\mathscr{F}} |\mu_n(A) - \mu(A)| \to 0, \quad n \to \infty,$$

is equivalent to

$$\sup_{f\in\mathscr{F}, |f|\leq 1} |\mu_n(f) - \mu(f)| \to 0, \quad n \to \infty.$$

2. *The strong convergence: for any $A \in \mathscr{F}$,*

$$|\mu_n(A) - \mu(A)| \to 0, \quad n \to \infty,$$

is equivalent to for any $f \in \mathrm{b}\mathscr{F}$

$$|\mu_n(f) - \mu(f)| \to 0, \quad n \to \infty.$$

Proof. 1. For the sufficiency, since $1_A \in \mathscr{F}$, $|1_A| \leq 1$, then

$$\begin{aligned}\sup_{A\in\mathscr{F}} |\mu_n(A) - \mu(A)| &= \sup_{A\in\mathscr{F}} |\mu_n(1_A) - \mu(1_A)| \\ &\leq \sup_{f\in\mathscr{F}, |f|\leq 1} |\mu_n(f) - \mu(f)| \to 0.\end{aligned}$$

For the necessity, firstly let $A = \Omega$. Then $\mu_n(\Omega) \to \mu(\Omega)$ implies exists $M > 0$ such that

$$\sup_{n\in\mathbb{N}} \mu_n(\Omega) \vee \mu(\Omega) \leq M.$$

For any $f \in \mathscr{F}, |f| \leq 1$, we can find increasing simple functions:

$$f_m = \sum_{k=0}^{2^m} k/2^m \mathbb{1}_{\{k/2^m < f \leq (k+1)/2^m\}} =: \sum_{k=0}^{2^m} k/2^m \mathbb{1}_{A_k} \uparrow f, \quad m \to \infty$$

and $||f_m - f||_\infty \leq 1/2^m$. Therefore

$$\begin{aligned}
&|\mu_n(f) - \mu(f)| \\
&\qquad \leq |\mu_n(f) - \mu_n(f_m)| + |\mu_n(f_m) - \mu(f_m)| + |\mu(f_m) - \mu(f)| \\
&\qquad \leq \mu_n(|f - f_m|) + \sum_{k=0}^{2^m} k/2^m |\mu_n(A_k) - \mu(A_k)| + \mu(|f - f_m|) \\
&\qquad \leq 1/2^m M + \sup_{A\in\mathscr{F}} |\mu_n(A) - \mu(A)| \sum_{k=0}^{2^m} k/2^m + 1/2^m M.
\end{aligned}$$

It follows from the above formula that

$$\sup_{f\in\mathscr{F}, |f|\leq 1} |\mu_n(f) - \mu(f)| \leq M/2^{m-1} + \sup_{A\in\mathscr{F}} |\mu_n(A) - \mu(A)| \sum_{k=0}^{2^m} k/2^m.$$

Let $n \to \infty$ first, and then let $m \to \infty$, we have

$$\sup_{f\in\mathscr{F}, |f|\leq 1} |\mu_n(f) - \mu(f)| \to 0, \quad n \to \infty.$$

2. The proof of the second assertion is nearly a copy of the first one. □

QUESTION 6.43 *Given a measurable space $(E, \mathscr{B}(E))$, where E is a metric space with metric $d(\cdot,\cdot)$, and $\{\mu_n\}_{n\geq 1}, \mu$ are uniformly bounded measure on $(E, \mathscr{B}(E))$. Then the following statements are all equivalent.*

1. $\mu_n \xrightarrow{\text{w}} \mu, n \to \infty$.
2. $\mu_n(B) \to \mu(B)$ *for all μ-continuous closed set $B \in \mathscr{B}(E)$.*
3. $\mu_n(C) \to \mu(C)$ *for all μ-continuous open set $C \in \mathscr{B}(E)$.*

Proof. Way 1. First we show $2 \Rightarrow 3$. Since E is naturally the μ-continuous closed/open set. Then $\mu_n(E) \to \mu(E)$. For any μ-continuous open set $C \in \mathscr{B}(E)$, we have $\mu(\partial C) = 0$. Note $B := C^c$ is closed and $\partial B = \partial C$. So $\mu(\partial B) = \mu(\partial C) = 0$. Therefore

$$\begin{aligned}\mu_n(C) &= \mu_n(E) - \mu_n(C^c) = \mu_n(E) - \mu_n(B)\\ &\to \mu(E) - \mu(B) = \mu(E) - \mu(C^c) = \mu(C).\end{aligned}$$

Similarly we can prove $3 \Rightarrow 2$. Then we show $1 \Leftrightarrow 2(3)$. If $\mu_n \xrightarrow{\text{w}} \mu$, then $\mu_n(A) \to \mu(A)$ for all μ-continuous open/closed set $A \in \mathscr{B}(E)$ naturally holds. We just prove the opposite. If $\mu_n(B) \to \mu(B)$ for all μ-continuous open/closed set $B \in \mathscr{B}(E)$, then for any μ-continuous set A, we have

$$\overline{\lim_{n\to\infty}}\, \mu_n(A) \le \overline{\lim_{n\to\infty}}\, \mu_n(\overline{A}) = \lim_{n\to\infty} \mu_n(\overline{A}) = \mu(\overline{A}) = \mu(A)$$

and

$$\underline{\lim_{n\to\infty}}\, \mu_n(A) \ge \underline{\lim_{n\to\infty}}\, \mu_n(\mathring{A}) = \lim_{n\to\infty} \mu_n(\mathring{A}) = \mu(\mathring{A}) = \mu(A).$$

Thus $\lim_{n\to\infty} \mu_n(A)$ exists and equals to $\mu(A)$. So $\mu_n \xrightarrow{w} \mu, n \to \infty$.

Way 2. First we show $1 \Leftrightarrow 2$. If $\mu_n \xrightarrow{\text{w}} \mu$, then $\mu_n(A) \to \mu(A)$ for all μ-continuous closed set $A \in \mathscr{B}(E)$ naturally holds. We just prove the opposite. For any open set $A \in \mathscr{B}(E)$, define

$$K_r := \{x \in E : d(x, A^c) \ge r\}, \quad r > 0.$$

Then K_r is closed, $K_r \subset A$ for any $r > 0$ and $K_r \uparrow A, r \to 0$. Now we claim there exists $\{r_n\} \subset (0, \infty)$ such that $\{K_{r_n}\}_{n\ge1}$ are all μ-continuous sets. In fact, since we have proved $\partial f^{-1}(A) \subset f^{-1}(\partial A)$ for any $A \in \mathscr{B}(E)$ in the topology part, then

$$(*) \qquad \partial K_r \subset \{x \in E : d(x, A^c) = r\}.$$

We say

$$\{r > 0 : \mu(\partial K_r) > 0\} = \bigcup_{m\ge1} \{r > 0 : \mu(\partial K_r) > 1/m\}$$

is at most countable. If not, there exists $m \ge 1$, such that $\{r > 0 : \mu(\partial K_r) > 1/m\}$ is uncountable. So we can arbitrarily choose a sequence $\{s_n\}_{n\ge1} \subset \{r > 0 : \mu(\partial K_r) > 1/m\}$. Moreover, formula $(*)$ shows for different r, the boundary ∂K_r is disjoint. So

$$\mu\left(\sum_{n\ge1} \partial K_{s_n}\right) = \sum_{n\ge1} \mu(\partial K_{s_n}) > \sum_{n\ge1} 1/m = \infty.$$

This contradicts the fact that μ is bounded. So $\{r > 0 : \mu(\partial K_r) > 0\}$ is at most countable. Therefore we can choose $\{r_k\}_{k\geq 1} \in \{r > 0 : \mu(\partial K_r) = 0\}$ and $r_k \downarrow 0, k \to \infty$. So $\{K_{r_k}\}_{k\geq 1}$ are naturally all μ-continuous sets. Finally,

$$\varliminf_{n\to\infty} \mu_n(A) \geq \varliminf_{n\to\infty} \mu_n(K_{r_k}) = \lim_{n\to\infty} \mu_n(K_{r_k}) = \mu(K_{r_k}).$$

Let $k \to \infty$ in above formula to obtain

$$\varliminf_{n\to\infty} \mu_n(A) \geq \mu\left(\bigcup_{k\geq 1} K_{r_k}\right) = \mu(A).$$

This means $\mu_n \xrightarrow{\text{w}} \mu, n \to \infty$.

So the proof is complete. □

QUESTION 6.44 *Suppose* $\{a_i\}_{i\in\mathbb{N}}$ *and* $\{a_i^{(n)}\}_{i\in\mathbb{N}}, n \in \mathbb{N}$ *are probability distributions. Then*

$$\lim_{n\to\infty} a_i^{(n)} = a_i, \forall i \in \mathbb{N} \Longleftrightarrow \lim_{n\to\infty} \sum_{i=0}^{\infty} s^i a_i^{(n)} = \sum_{i=0}^{\infty} s^i a_i, \quad \forall s \in (0,1).$$

Proof. The necessity part can be simply obtained by using dominance convergence theorem. We just prove the sufficiency part. We use the induction on i to verify this part.

1. When $i = 0$, since

$$\begin{aligned}\sum_{i=0}^{\infty} s^i a_i^{(n)} - \sum_{i=0}^{\infty} s^i a_i &= \sum_{i=1}^{\infty} s^i (a_i^{(n)} - a_i) + (a_0^{(n)} - a_0) \\ &\leq \sum_{i=1}^{\infty} s^i + (a_0^{(n)} - a_0) \\ &= \frac{s}{1-s} + (a_0^{(n)} - a_0).\end{aligned}$$

Then

$$0 = \varliminf_{n\to\infty} \left(\sum_{i=0}^{\infty} s^i a_i^{(n)} - \sum_{i=0}^{\infty} s^i\right) \leq \frac{s}{1-s} + \varliminf_{n\to\infty} (a_0^{(n)} - a_0).$$

Let $s \to 0+$ to obtain

$$0 \leq \varliminf_{n\to\infty} (a_0^{(n)} - a_0).$$

Interchange the position of $a_0^{(n)}$ and a_0, we will get

$$\varliminf_{n\to\infty} (a_0^{(n)} - a_0) = \varlimsup_{n\to\infty} (a_0^{(n)} - a_0) = 0,$$

which is equivalent to $\lim_{n\to\infty} a_0^{(n)} = a_0$.

2. Suppose when $i \le k$, we have $\lim_{n\to\infty} a_i^{(n)} = a_i$. When $i = k+1$, the induction hypothesis ensure

$$\lim_{n\to\infty} \sum_{i=k+1}^{\infty} s^i a_i^{(n)} = \sum_{i=k+1}^{\infty} s^i a_i$$

and furthermore,

$$\lim_{n\to\infty} \sum_{i=k+1}^{\infty} s^{i-(k+1)} a_i^{(n)} = \sum_{i=k+1}^{\infty} s^{i-(k+1)} a_i.$$

Then the similar argument shows $\lim_{n\to\infty} a_{k+1}^{(n)} = a_{k+1}$.

3. Hence the argument is correct for all $i \in \mathbb{N}$. □

QUESTION 6.45 *A random variable X on probability space $(\Omega, \mathscr{F}, \mathbb{P})$ is distributed as a constant c iff $X \overset{\text{a.s.}}{=} c$.*

Proof. Obviously, if $X \overset{\text{a.s.}}{=} c$, then $X \sim c$. On the contrary, if $X \sim c$, then we have for any $n \ge 1$,

$$\mathbb{P}(c - 1/n \le X \le c + 1/n) = \mathbb{P}(c - 1/n \le c \le c + 1/n) = 1.$$

Then let $\varepsilon \to 0+$, we will obtain

$$\mathbb{P}(X = c) = 1,$$

which is equivalent to $X \overset{\text{a.s.}}{=} c$. □

QUESTION 6.46 (The generalization of MCT) *Given measurable functions $\{f_t\}_{t\in I}$ on measure space $(E, \mathscr{E}, \mu)$, where $I = \mathbb{N}, I = \mathbb{R}_+$ (or $I = [a, b]$). If $\{f_t\}_{t\in I}$ is monotone and exists $s \in I$ (or $s \in [a, b)$) such that $f_s \in L^1(\mu)$, then*

$$\int_E f_t d\mu \to \int_E f d\mu, \quad t \to \infty \text{ (or } t \to b).$$

Proof. If $\{f_t\}_{t\in I}$ is increasing, then $f_s \le f_t$ for any $I \ni t \ge s$. Then the normal MCT gives us the desired results. If $\{f_t\}_{t\in I}$ is decreasing, then $f_s \ge f_t \implies -f_s \le -f_t$ for any $I \ni t \ge s$. Then the normal MCT also gives us the desired results. □

QUESTION 6.47 *Suppose $(\Omega, \mathscr{F}, \mathbb{P})$ is a probability space and $A \notin \mathscr{F}$. Let $\mathscr{F}_1 := \sigma(\mathscr{F} \cup \{A\})$. Prove $\mathbb{P}$ can be extended to $(\Omega, \mathscr{F}_1)$.*

Proof. 1. For any $D \in \mathscr{F}$, we define

$$\widetilde{\mathbb{P}}(A \cap D) := \inf\{\mathbb{P}(E) : A \cap D \subset E, E \in \mathscr{F}\}.$$

Next we show exists $E_D \in \mathscr{F}, A \cap D \subset E_D$ such that $\mathbb{P}(E_D) = \widetilde{\mathbb{P}}(A \cap D)$. In fact, by the definition of infimum, exists $\{E_n\}_{n\geq 1} \subset \mathscr{F}$ satisfying $A \cap D \subset E_n$ for any $n \geq 1$ and $\mathbb{P}(E_n) \downarrow \widetilde{\mathbb{P}}(A \cap D)$. Define $E_D := \underline{\lim}_{n\to\infty} E_n \in \mathscr{F}$, then

$$E_D = \underline{\lim}_{n\to\infty} E_n = \bigcup_{n\geq 1} \bigcap_{k\geq n} E_k \supset \bigcup_{n\geq 1} \bigcap_{k\geq n} (A \cap D) = A \cap D$$

and

$$\widetilde{\mathbb{P}}(A \cap D) \leq \mathbb{P}(E_D) = \lim_{n\to\infty} \mathbb{P}\left(\bigcap_{k\geq n} E_k\right) \leq \lim_{n\to\infty} \mathbb{P}(E_n) = \widetilde{\mathbb{P}}(A \cap D).$$

This shows $\mathbb{P}(E_D) = \widetilde{\mathbb{P}}(A \cap D)$. We can additionally presume $E_D \subset D$. In fact, consider $E_D \cap D$ is enough.

2. Moreover, we show for any $(A \cap D_1) \cap (A \cap D_2) = \emptyset$, we have

$$\widetilde{\mathbb{P}}[(A \cap D_1) + (A \cap D_2)] = \widetilde{\mathbb{P}}[A \cap (D_1 \cup D_2)] = \widetilde{\mathbb{P}}(A \cap D_1) + \widetilde{\mathbb{P}}(A \cap D_2).$$

In fact, exist $E_1, E_2 \in \mathscr{F}$ such that $\mathbb{P}(E_1) = \widetilde{\mathbb{P}}(A \cap D_1)$ and $\mathbb{P}(E_2) = \widetilde{\mathbb{P}}(A \cap D_2)$. We claim

$$\mathbb{P}(E_1 \cap E_2) = 0.$$

Otherwise, since $(A \cap D_1) \cap (A \cap D_2) = \emptyset$ and $A \cap D_i \subset E_i, i = 1, 2$, then

$$\mathscr{F} \ni E_1 \cap E_2 \subset [A \cap (D_1 \cup D_2)]^c$$

and $\mathbb{P}(E_1 \cap E_2) > 0$. Therefore $\hat{E}_i := E_i - E_1 \cap E_2$ satisfying

$$A \cap D_i \subset \hat{E}_i, \qquad \mathbb{P}(\hat{E}_i) = \mathbb{P}(E_i) - \mathbb{P}(E_1 \cap E_2) < \mathbb{P}(E_i), \quad i = 1, 2,$$

which contradicts the definition of $E_i, i = 1, 2$. Since we have proved $\mathbb{P}(E_1 \cap E_2) = 0$, then

$$\mathbb{P}(E_1 \cup E_2) = \mathbb{P}(E_1) + \mathbb{P}(E_2) - \mathbb{P}(E_1 \cap E_2) = \mathbb{P}(E_1) + \mathbb{P}(E_2).$$

Since $A \cap D_1 \subset E_1$ and $A \cap D_2 \subset E_2$, then

$$A \cap (D_1 \cup D_2) \subset E_1 \cup E_2.$$

Therefore

$$\begin{aligned}\widetilde{\mathbb{P}}[A\cap(D_1\cup D_2)] &\le \widetilde{\mathbb{P}}(E_1\cup E_2)=\mathbb{P}(E_1)+\mathbb{P}(E_2)\\ &=\widetilde{\mathbb{P}}(A\cap D_1)+\widetilde{\mathbb{P}}(A\cap D_2).\end{aligned}$$

On the contrary, exists $E_{12}\in\mathscr{F}$ such that $A\cap(D_1\cup D_2)\subset E_{12}\subset D_1\cup D_2$ and $\mathbb{P}(E_{12})=\widetilde{\mathbb{P}}[A\cap(D_1\cup D_2)]$. Consider $E_i':=E_{12}\cap D_i, i=1,2$. Since $A\cap(D_1\cup D_2)\subset E_{12}$, then $A\cap D_i\subset E_i', i=1,2$ and $E_{12}=E_1'\cup E_2'$. We claim $\mathbb{P}(E_1'\cap E_2')=0$. In fact, if $\mathbb{P}(E_1'\cap E_2')>0$, since $(A\cap D_1)\cap(A\cap D_2)=\emptyset$, then we must have

$$E_1'\cap E_2'\subset[A\cap(D_1\cup D_2)]^c.$$

Consider $E_{12}'=E_{12}-E_1'\cap E_2'$, then $A\cap(D_1\cup D_2)\subset E_{12}'$ and

$$\mathbb{P}(E_{12}')=\mathbb{P}(E_{12})-\mathbb{P}(E_1'\cap E_2')<\mathbb{P}(E_{12}),$$

which contradicts the definition of E_{12}. So $\mathbb{P}(E_{12})=\mathbb{P}(E_1')+\mathbb{P}(E_2')$ and therefore

$$\begin{aligned}\widetilde{\mathbb{P}}[A\cap(D_1\cup D_2)] &= \mathbb{P}(E_{12})=\mathbb{P}(E_1')+\mathbb{P}(E_2')\\ &\ge \mathbb{P}(E_1)+\mathbb{P}(E_2)=\widetilde{\mathbb{P}}(A\cap D_1)+\widetilde{\mathbb{P}}(A\cap D_2).\end{aligned}$$

The above two formulas show for any $D_1,D_2\in\mathscr{F}$, $(A\cap D_1)\cap(A\cap D_2)=\emptyset$,

$$\widetilde{\mathbb{P}}[A\cap(D_1\cup D_2)]=\widetilde{\mathbb{P}}(A\cap D_1)+\widetilde{\mathbb{P}}(A\cap D_2).$$

3. Next we show for any disjoint sequence $\{A\cap D_n\}_{n\ge1}$, we have

$$\widetilde{\mathbb{P}}\left(\sum_{n\ge1}A\cap D_n\right)=\sum_{n\ge1}\widetilde{\mathbb{P}}(A\cap D_n).$$

We have shown exists $E_{\ge1}\in\mathscr{F}$ such that

$$\sum_{n\ge1}A\cap D_n\subset E_{\ge1}\subset\bigcup_{n\ge1}D_n \text{ and } \mathbb{P}(E_{\ge1})=\widetilde{\mathbb{P}}\left(\sum_{n\ge1}A\cap D_n\right)$$

and exist $\{E_n\}_{n\ge1}\subset\mathscr{F}$ such that $A\cap D_n\subset E_n\subset D_n$,

$$\widetilde{\mathbb{P}}(A\cap D_n)=\mathbb{P}(E_n) \text{ and } \mathbb{P}(E_n\cap E_m)=0,\quad \forall n\ne m\ge1.$$

Observe $\sum_{n\ge1}A\cap D_n\subset\cup_{n\ge1}E_n$, then

$$\widetilde{\mathbb{P}}\left(\sum_{n\ge1}A\cap D_n\right)=\mathbb{P}(E_{\ge1})\le\mathbb{P}\left(\bigcup_{n\ge1}E_n\right)\le\sum_{n\ge1}\mathbb{P}(E_n)=\sum_{n\ge1}\widetilde{\mathbb{P}}(A\cap D_n)$$

Define $E'_n := E_{\geq 1} \cap D_n$, then $A \cap D_n \subset E'_n \subset D_n$ for any $n \geq 1$. We claim $\mathbb{P}(E'_i \cap E'_j) = 0$ for any $i \neq j \geq 1$. In fact, if $\mathbb{P}(E'_i \cap E'_j) > 0$, since $\cap_{n\geq 1} A \cap D_n = \emptyset$, then we must have

$$E'_i \cap E'_j \subset [(A \cap D_i) + (A \cap D_j)]^c \text{ and } E'_i \cap E'_j \subset D_i \cap D_j.$$

This must yields

$$E'_i \cap E'_j \subset \left[\sum_{n\geq 1} A \cap D_n\right]^c.$$

Consider $\hat{E}_{\geq 1} = E_{\geq 1} - E'_i \cap E'_j$, then $\sum_{n\geq 1} A \cap D_n \subset \hat{E}_{\geq 1}$ and

$$\mathbb{P}(\hat{E}_{\geq 1}) = \mathbb{P}(E_{\geq 1}) - \mathbb{P}(E'_i \cap E'_j) < \mathbb{P}(E_{\geq 1}),$$

which contradicts the definition of $E_{\geq 1}$. So we have proved $\mathbb{P}(E'_i \cap E'_j) = 0$ for any $i \neq j \geq 1$.

Define $H_1 = E'_1$ and $H_n = E'_n / \cup_{k=1}^{n-1} E'_k$ for $n \geq 2$. Then $\{H_n\}_{n\geq 1} \subset \mathscr{F}$ is disjoint and $\mathbb{P}(H_n) = \mathbb{P}(E'_n)$. In fact, we have for any $n \geq 2$,

$$\mathbb{P}\left(E'_n \cap \bigcup_{k=1}^{n-1} E'_k\right) = \mathbb{P}\left[\bigcup_{k=1}^{n-1} (E'_n \cap E'_k)\right] \leq \sum_{k=1}^{n-1} \mathbb{P}(E'_n \cap E'_k) = 0.$$

Then for any $n \geq 2$,

$$\mathbb{P}(H_n) = \mathbb{P}\left(E'_n \Big/ \bigcup_{k=1}^{n-1} E'_k\right) = \mathbb{P}(E'_n) - \mathbb{P}\left(E'_n \cap \bigcup_{k=1}^{n-1} E'_k\right) = \mathbb{P}(E'_n).$$

Therefore

$$\begin{aligned}\sum_{n\geq 1} \mathbb{P}(E'_n) &= \sum_{n\geq 1} \mathbb{P}(H_n) = \mathbb{P}\left(\sum_{n\geq 1} H_n\right) = \mathbb{P}\left(\bigcup_{n\geq 1} E'_n\right) \\ &= \mathbb{P}(E_{\geq 1}) = \widetilde{\mathbb{P}}\left(\sum_{n\geq 1} A \cap D_n\right).\end{aligned}$$

Finally, we have

$$\sum_{n\geq 1} \widetilde{\mathbb{P}}(A \cap D_n) = \sum_{n\geq 1} \mathbb{P}(E_n) \leq \sum_{n\geq 1} \mathbb{P}(E'_n) = \widetilde{\mathbb{P}}\left(\sum_{n\geq 1} A \cap D_n\right).$$

Combine with the former formula, we have proved

$$\widetilde{\mathbb{P}}\left(\sum_{n\geq 1} A \cap D_n\right) = \sum_{n\geq 1} \widetilde{\mathbb{P}}(A \cap D_n).$$

4. Define

$$\widetilde{\mathbb{P}}(A^c \cap D) := \sup\left\{\mathbb{P}(F) : A^c \cap D \supset F, F \in \mathscr{F}\right\}.$$

We want to show exists $F_D \in \mathscr{F}$ such that $F_D \subset A^c \cap D$ and $F_D = \widetilde{\mathbb{P}}(A^c \cap D)$. Observe

$$A^c \cap D \supset F \Longleftrightarrow D - (A^c \cap D) \subset D - F \Longleftrightarrow A \cap D \subset D - F$$

and $F = D - (D - F)$ $(F \subset D)$. Then

$$\begin{aligned}
\widetilde{\mathbb{P}}(A^c \cap D) &= \sup\left\{\mathbb{P}(D) - \mathbb{P}(D - F) : A \cap D \subset D - F, F \in \mathscr{F}\right\} \\
&= \sup\left(\mathbb{P}(D) - \{\mathbb{P}(D - F) : A \cap D \subset D - F, F \in \mathscr{F}\}\right) \\
&= \mathbb{P}(D) - \inf\{\mathbb{P}(D - F) : A \cap D \subset D - F, F \in \mathscr{F}\} \\
&= \mathbb{P}(D) - \inf\{\mathbb{P}(E) : A \cap D \subset E, E \in \mathscr{F}\} \\
&= \mathbb{P}(D) - \widetilde{\mathbb{P}}(A \cap D).
\end{aligned}$$

Select $E_D \in \mathscr{F}$ such that $A \cap D \subset E_D \subset D$ and $\mathbb{P}(E_D) = \widetilde{\mathbb{P}}(A \cap D)$. Then

$$\widetilde{\mathbb{P}}(A^c \cap D) = \mathbb{P}(D) - \widetilde{\mathbb{P}}(A \cap D) = \mathbb{P}(D) - \mathbb{P}(E_D) = \mathbb{P}(D - E_D).$$

Then $F_D := D - E_D \subset A^c \cap D$ is what we want. Besides, we still want to show for any $D_1, D_2 \in \mathscr{F}, (A^c \cap D_1) \cap (A^c \cap D_2) = \emptyset$,

$$\widetilde{\mathbb{P}}[A^c \cap (D_1 \cup D_2)] = \widetilde{\mathbb{P}}(A^c \cap D_1) + \widetilde{\mathbb{P}}(A^c \cap D_2).$$

In fact, exists $F_1, F_1 \in \mathscr{F}$ satisfying $F_i \subset A^c \cap D_i$ and $\mathbb{P}(F_i) = \widetilde{\mathbb{P}}(A^c \cap D_i)$ for $i = 1, 2$. Observe $F_1 \cup F_2 \subset A^c \cap (D_1 \cup D_2)$, then $(A^c \cap D_1) \cap (A^c \cap D_2) = \emptyset$ yields

$$\begin{aligned}
\widetilde{\mathbb{P}}[A^c \cap (D_1 \cup D_2)] &\geq \mathbb{P}(F_1 \cup F_2) = \mathbb{P}(F_1) + \mathbb{P}(F_2) \\
&= \widetilde{\mathbb{P}}(A^c \cap D_1) + \widetilde{\mathbb{P}}(A^c \cap D_2).
\end{aligned}$$

On the contrary, exists $F_{12} \in \mathscr{F}$ satisfying $F_{12} \subset A^c \cap (D_1 \cup D_2)$ and $\mathbb{P}(F_{12}) = \widetilde{\mathbb{P}}[A^c \cap (D_1 \cup D_2)]$. Define $F_i' := F_{12} \cap D_i \subset A^c \cap D_i$ for $i = 1, 2$. Then $(A^c \cap D_1) \cap (A^c \cap D_2) = \emptyset$ yields

$$\begin{aligned}
\widetilde{\mathbb{P}}[A^c \cap (D_1 + D_2)] &= \mathbb{P}(F_{12}) = \mathbb{P}(F_1') + \mathbb{P}(F_2') \\
&\leq \mathbb{P}(F_1) + \mathbb{P}(F_2) = \widetilde{\mathbb{P}}(A^c \cap D_1) + \widetilde{\mathbb{P}}(A^c \cap D_2).
\end{aligned}$$

So we have proved for any $D_1, D_2 \in \mathscr{F}, (A^c \cap D_1) \cap (A^c \cap D_2) = \emptyset$,

$$\widetilde{\mathbb{P}}[A^c \cap (D_1 \cup D_2)] = \widetilde{\mathbb{P}}(A^c \cap D_1) + \widetilde{\mathbb{P}}(A^c \cap D_2).$$

And similar argument shows for any disjoint $\{A^c \cap D_n\}_{n\geq 1}$, we have

$$\widetilde{\mathbb{P}}\left(\sum_{n\geq 1} A^c \cap D_n\right) = \sum_{n\geq 1} \widetilde{\mathbb{P}}(A^c \cap D_n).$$

5. Note $\mathscr{F}_1 = \sigma(\mathscr{F} \cup \{A\}) = \{(A \cap D_1) + (A^c \cap D_2) : D_1, D_2 \in \mathscr{F}\}$, then define

$$\widetilde{\mathbb{P}}[(A \cap D_1) + (A^c \cap D_2)] := \widetilde{\mathbb{P}}(A \cap D_1) + \widetilde{\mathbb{P}}(A^c \cap D_2).$$

We show $\widetilde{\mathbb{P}}(\cdot)$ is σ-additive. For any disjoint sets $\{(A \cap D_1^n) + (A^c \cap D_2^n)\}_{n\geq 1}$, it is easy to find $\{A \cap D_1^n\}_{n\geq 1}$ and $\{A \cap D_2^n\}_{n\geq 1}$ are two sequences of disjoint sets. The former discussion yields

$$\begin{aligned}
&\widetilde{\mathbb{P}}\left\{\sum_{n\geq 1}[(A \cap D_1^n) + (A^c \cap D_2^n)]\right\} \\
&\qquad = \widetilde{\mathbb{P}}\left[A \cap \left(\bigcup_{n\geq 1} D_1^n\right) + A^c \cap \left(\bigcup_{n\geq 1} D_2^n\right)\right] \\
&\qquad = \widetilde{\mathbb{P}}\left[A \cap \left(\bigcup_{n\geq 1} D_1^n\right)\right] + \widetilde{\mathbb{P}}\left[A^c \cap \left(\bigcup_{n\geq 1} D_2^n\right)\right] \\
&\qquad = \widetilde{\mathbb{P}}\left[\sum_{n\geq 1}(A \cap D_1^n)\right] + \widetilde{\mathbb{P}}\left[\sum_{n\geq 1}(A^c \cap D_2^n)\right] \\
&\qquad = \sum_{n\geq 1}[\widetilde{\mathbb{P}}(A \cap D_1^n) + \widetilde{\mathbb{P}}(A^c \cap D_2^n)] \\
&\qquad = \sum_{n\geq 1} \widetilde{\mathbb{P}}\left[(A \cap D_1^n) + (A^c \cap D_2^n)\right].
\end{aligned}$$

Till now, $(\Omega, \mathscr{F}, \mathbb{P})$ has been extended to $(\Omega, \mathscr{F}_1, \widetilde{\mathbb{P}})$. □

QUESTION 6.48 *Given a probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Suppose π-system $\mathscr{A}$ satisfies $\Omega \in \mathscr{A}$ and $\mathscr{F} = \sigma(\mathscr{A})$. Define $\mathcal{D}$ the linear span of $\{1_A : A \in \mathscr{A}\}$. Then $\mathcal{D}$ is dense in $L^p(\Omega, \mathscr{F}, \mathbb{P})$ for $0 < p < \infty$.*

Proof. Define

$$\mathscr{L} = \{f \in L^p(\Omega, \mathscr{F}, \mathbb{P}) : \text{exists } \{g_n\}_{n\geq 1} \subset \mathcal{D} \text{ such that } ||g_n - f||_{L^p} \to 0\}.$$

First it is easy to know $\{1_A : A \in \mathscr{A}\} \subset \mathscr{L}$. We next shows $\mathscr{L}$ is a $\mathscr{L}$-system.

1. Since $\Omega \in \mathscr{A}$, then $1 = 1_\Omega \in \mathscr{L}$.

2. For any $f_1, f_2 \in \mathscr{L}$, then there exists $\{g_{i,n}\}_{n\geq 1} \subset \mathcal{D}$ such that $||g_{i,n} - f_i||_{L^p} \to 0, i = 1, 2$. Therefore for any $\alpha, \beta \in \mathbb{R}$, we have $\{\alpha g_{1,n} + \beta g_{2,n}\}_{n\geq 1} \subset \mathcal{D}$ and

$$\begin{aligned} &||\alpha g_{1,n} + \beta g_{2,n} - (\alpha f_1 + \beta f_2)||_{L^p} \\ &\qquad \leq \alpha ||g_{1,n} - f_1||_{L^p} + \beta ||g_{2,n} - f_2||_{L^p} \to 0. \end{aligned}$$

So $\alpha f_1 + \beta f_2 \in \mathscr{L}$.

3. For any $\{f_i\}_{i\geq 1} \subset \mathscr{L}, f \in \mathscr{L}$ and $0 \leq f_i \uparrow f$, there exists $\{g_{i,n}\}_{n\geq 1} \subset \mathcal{D}$ such that $||g_{i,n} - f_i||_{L^p} \to 0, n \to \infty, i \geq 1$. So for any $i \geq 1$, exists $n_i \geq 1$ such that $||g_{i,n_i} - f_i||_{L^p} < 1/i$. Moreover, $0 \leq f_n \uparrow f$ means $0 \leq f_n^p \uparrow f^p$. Monotoncity convergence theorem yields $||f_n||_{L^p} \to ||f||_{L^p}$. So $f_n^p \uparrow f^p$ and $||f_n||_{L^p} \to ||f||_{L^p}$ yield $||f_n - f||_{L^p} \to 0$. Finally, consider $\{g_{i,n_i}\}_{i\geq 1} \subset \mathcal{D}$, we have

$$\begin{aligned} ||g_{i,n_i} - f|| &\leq ||g_{i,n_i} - f_i||_{L^p} + ||f_i - f||_{L^p} \\ &< 1/i + ||f_i - f||_{L^p} \to 0, \quad i \to \infty. \end{aligned}$$

So $f \in \mathscr{L}$.

Therefore $\mathscr{L}$ is a $\mathscr{L}$-system and $\{1_A : A \in \mathscr{A}\} \subset \mathscr{L}$. The function form monotonicity class theorem yields $\mathscr{L} = L^p(\Omega, \mathscr{F}, \mathbb{P})$. □

QUESTION 6.49 *Consider the measure space $(\mathbb{R}^d, \mathscr{B}(\mathbb{R}^d), \mu)$. Define the dual of μ as $\widetilde{\mu}(A) := \mu(-A)$ for $A \in \mathscr{B}(\mathbb{R}^d)$ and the symmetrization of μ as $\mu^\sharp := \mu * \widetilde{\mu}$. Then*

1. *For any $f \in \mathscr{B}(\mathbb{R}^d)$ and $\mu(f)$ exists, we have*

$$\int f(x)\widetilde{\mu}(dx) = \int f(-x)\mu(dx).$$

2. *Consider product measure $\mu \times \mu$. For any $g \in \mathscr{B}(\mathbb{R}^{2d})$ and $\int g d(\mu \times \mu)$ exists, we have*

$$(*) \qquad \iint g(x,y)\mu(dx)\mu(dy) = \iint g(y,x)\mu(dx)\mu(dy).$$

3. *$\mu^\sharp$ is symmetric. Namely, $\mu^\sharp(A) = \mu^\sharp(-A)$ for $A \in \mathscr{B}(\mathbb{R}^d)$.*

Proof. 1. For $f = 1_A$, where $A \in \mathscr{B}(\mathbb{R}^d)$, we can easily find the assertion is correct. For the general case, we can just apply the standard 'simple function', 'nonnegative measurable function' and 'general measurable function' procedure to obtain the desired result.

2. For any $A, B \in \mathscr{B}(\mathbb{R}^d)$, we have

$$\iint \mathbb{1}_{A\times B}(x,y)\mu(dx)\mu(dy) = \mu(A)\mu(B) = \iint \mathbb{1}_{A\times B}(y,x)\mu(dx)\mu(dy).$$

Write $\mathscr{G} := \{C \in \mathscr{B}(\mathbb{R}^{2d}) :, g = 1_C \text{ satisfies}(*)\}$ and $\mathscr{A} := \{A \times B : A, B \in \mathscr{B}(\mathbb{R}^d)\}$. What we have proved shows π system $\mathscr{A} \subset \mathscr{G}$. Apply monotone convergence theorem, we will easily find $\mathscr{G}$ is a λ system. So $\mathscr{G} = \sigma(\mathscr{A}) = \mathscr{B}(\mathbb{R}^{2d})$. Use the standard procedure, we will find the assertion is correct for all $g \in \mathscr{B}(\mathbb{R}^{2d})$.

3. For any $A \in \mathscr{B}(\mathbb{R}^d)$, apply Fubini's theorem and the previous two small questions to obtain

$$\begin{aligned}
\mu^\sharp(A) &= \iint 1_A(x+y)\mu(dx)\tilde{\mu}(dy) \\
&= \int \mu(A-y)\tilde{\mu}(dy) = \int \mu(A+y)\mu(dy) \\
&= \iint 1_A(x-y)\mu(dx)\mu(dy) \\
&= \iint 1_A(y-x)\mu(dx)\mu(dy) \\
&= \iint \mathbb{1}_{-A}(x-y)\mu(dx)\mu(dy) \\
&= \mu^\sharp(-A).
\end{aligned}$$

□

QUESTION 6.50 *Given a distribution μ on measurable space $([0,\infty), \mathscr{B}[0,\infty))$. Write its Laplace transformation as*

$$L_\mu(u) = \int_{[0,\infty)} e^{-ux}\mu(dx), \quad u \geq 0.$$

For two distributions μ_1 and μ_2 on $([0,\infty), \mathscr{B}[0,\infty))$, if $L_{\mu_1}(u) = L_{\mu_2}(u)$, then $\mu_1 = \mu_2$. Namely, distributions and its Laplace transformations are mutually uniquely determined.

Proof. Since μ_1, μ_2 are only determined on $\mathscr{B}[0,\infty)$, then $|e^{-ux}| \leq 1, u \geq 0$ and thus $L_\mu(u)$ is well-defined. Moreover, write the region $D := \{w \in \mathbb{C} : \operatorname{Re} w < 0\}$, then $\overline{D} = \{w \in \mathbb{C} : \operatorname{Re} w \leq 0\}$. We find

$$|e^{zx}| = e^{(\operatorname{Re} z)x} \leq 1, \quad \forall z \in D.$$

So

$$f_j(z) := \int_{[0,\infty)} e^{zx}\mu_j(dx), \quad j = 1, 2$$

is still well-defined. Besides, for any $n \geq 1$, there exists $M_n > 0$ satisfies

$$|x^n e^{zx}| = x^n e^{(\operatorname{Re} z)x} \leq M_n.$$

By the dominated convergence theorem, we know $f_j(z) \in H(D) \cap C(\overline{D})$. Let $z = -u < 0$, we have

$$f_1(-u) = L_{\mu_1}(u) = L_{\mu_2}(u) = f_2(-u), \quad \forall u \geq 0.$$

Hence $f_1(z)$ and $f_2(z)$ coincide on $(-\infty, 0]$. Apply the unique determination theorem for analytic functions, we have $f_1(z) = f_2(z)$ on D. Note $f_j(z) \in C(\overline{D})$, then $f_1(z) = f_2(z)$ on $\overline{D}$. Let $z = it, t \in \mathbb{R}$, we have

$$\varphi_{\mu_1}(t) = f_1(it) = f_2(it) = \varphi_{\mu_2}(t), \quad t \in \mathbb{R},$$

where $\varphi_{\mu_j}(t)$ is the characteristic function of μ_j. So $\mu_1 = \mu_2$. □

QUESTION 6.51 *Given two measures μ, ν on probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Assume $\mu \ll \nu$. Then $\nu \ll \mu$ iff $d\mu/d\nu \neq 0, \nu$-a.e..*

Proof. $\Rightarrow$. If $\nu \ll \mu$, then for any $A \in \mathscr{F}$, we have

$$\mu(A) = \int_A \frac{d\mu}{d\nu} d\nu = \int_A \frac{d\mu}{d\nu}\frac{d\nu}{d\mu} d\mu.$$

This shows

$$\int_A \left(1 - \frac{d\mu}{d\nu}\frac{d\nu}{d\mu}\right) d\mu \equiv 0, \quad \forall A \in \mathscr{F} \Longrightarrow \frac{d\mu}{d\nu}\frac{d\nu}{d\mu} \equiv 1, \quad \mu\text{-a.e..}$$

Thus $d\mu/d\nu \neq 0, \mu$-a.e. and by the mutual absolute continuity between μ and ν, we have $d\mu/d\nu \neq 0, \nu$-a.e..

$\Leftarrow$. If $\nu \not\ll \mu$, then exists $B \in \mathscr{F}$ such that $\mu(B) = 0$ but $\nu(B) > 0$. Moreover, it is easy to find $d\mu/d\nu \geq 0, \nu$-a.e.. So $d\mu/d\nu \neq 0, \nu$-a.e. means $d\mu/d\nu > 0, \nu$-a.e.. Thus

$$0 = \mu(B) = \int_B \frac{d\mu}{d\nu} d\nu > 0.$$

This contradiction yields the desired result. □

QUESTION 6.52 *For finite measures $\{\mu_n\}_{n\geq 1}, \mu$ on metric space $(E, \mathscr{B}(E))$. Assume $\mu_n \xrightarrow{\text{w}} \mu$, then for any μ-continuous set B, we have $\mu_n(B \cap \cdot) \xrightarrow{\text{w}} \mu(B \cap \cdot)$. It is also equivalent to*

$$\int_B f(x)\mu_n(dx) \to \int_B f(x)\mu(dx), \quad n \to \infty.$$

Proof. Since we have proved $\partial(A \cup B) \subset \partial A \cup \partial B$. Then for any μ-continuous set A and the given μ-continuous set B, we have

$$\mu[\partial(A \cup B)] \leq \mu(\partial A \cup \partial B) \leq \mu(\partial A) + \mu(\partial B) = 0.$$

Thus $A \cup B$ is also a μ-continuous set for any μ-continuous set A. Therefore

$$\mu_n(B \cap A) \to \mu(B \cap A), \quad n \to \infty.$$

And the last assertion is obvious. □

QUESTION 6.53 *Given a metric measurable space $(E, \mathscr{B}(E))$ with metric d. If E is Polish, then the diagonal $D = \{(x, x) : x \in E\} \in \mathscr{B}^2(E) = \mathscr{B}(E^2)$.*

Proof. Since E is separable, then there exists a countable dense set $\{q_n\}_{n\geq 1} \subset E$. We claim $\{(q_n, q_n)\}_{n\geq 1} \subset D$ is also a countable dense set of D. In fact, by the definition of the metric on the product metric space, for any $x \in E$ and $\varepsilon > 0$, there exists $q_x \in \{q_n\}_{n\geq 1}$ satisfying $d(x, q_x) < \varepsilon/2$. Then

$$d((x, x), (q_x, q_x)) = 2d(x, q_x) < \varepsilon.$$

And $\{(q_n, q_n)\}_{n\geq 1}$ is obviously countable. So $\{(q_n, q_n)\}_{n\geq 1} \subset D$ is a countable dense set of D. Define

$$A_m := \bigcup_{n\geq 1} B((q_n, q_n), 1/m) \in \mathscr{B}^2(E) = \mathscr{B}(E^2), \quad m \geq 1.$$

We claim $D \subset A_m$ for any $m \geq 1$. In fact, for any $(x, x) \in D$, since $\{(q_n, q_n)\}_{n\geq 1} \subset D$ is a countable dense set of D, then exists $(q_x, q_x) \in \{(q_n, q_n)\}_{n\geq 1}$ such that $d((x, x), (q_x, q_x)) < 1/m$. Namely,

$$(x, x) \in B((q_x, q_x), 1/m) \subset \bigcup_{n\geq 1} B((q_n, q_n), 1/m) = A_m.$$

Thus $D \subset A_m$ for any $m \geq 1$ and therefore $D \subset \cap_{m\geq 1} A_m$. Next we show $\cap_{m\geq 1} A_m \subset D$. In fact, Given any

$$(x, y) \in \bigcap_{m\geq 1} A_m = \bigcap_{m\geq 1} \bigcup_{n\geq 1} B((q_n, q_n), 1/m),$$

then for any $m \geq 1$, there exists $n_m \geq 1$, such that $(x, y) \in B((q_{n_m}, q_{n_m}), 1/m)$. So $d(x, y) \leq 2/m$ for any $m \geq 1$. Letting $m \to \infty$ yields $d(x, y) = 0 \Leftrightarrow x = y$. That is to say, $(x, y) \in D$. In conclusion, $D = \cap_{m\geq 1} A_m \in \mathscr{B}^2(E) = \mathscr{B}(E^2)$.□

QUESTION 6.54 *Given a real valued random variable X on probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Suppose the σ-algebra $\mathscr{G} \subset \mathscr{F}$. Then the following claims are all equivalent (all the equalities below are in the sense of $\mathbb{P}$-a.s).*

1. *X is independent to $\mathscr{G}$.*
2. $\mathbb{P}[f(X)|\mathscr{G}] = \mathbb{P}[f(X)]$ *for any* $f \in \mathrm{b}\mathscr{B}(\mathbb{R})$.
3. $\mathbb{P}(e^{i\theta X}|\mathscr{G}) = \mathbb{P}(e^{i\theta X})$.
4. $\mathbb{P}(e^{i\theta X}1_A) = \mathbb{P}(e^{i\theta X})\mathbb{P}(A)$ *for any* $A \in \mathscr{G}$.

Proof. 1. $1 \Leftrightarrow 2$. For any $f \in \mathrm{b}\mathscr{B}(\mathbb{R})$ and $A \in \mathscr{G}$, note X is independent to $\mathscr{G}$, we have

$$\mathbb{P}[f(X)1_A] = \mathbb{P}[f(X)]\mathbb{P}(A) = \mathbb{P}\{\mathbb{P}[f(X)]1_A\}.$$

This shows $\mathbb{P}[f(X)|\mathscr{G}] = \mathbb{P}[f(X)]$. On the contrary, since $\mathbb{P}[f(X)|\mathscr{G}] = \mathbb{P}[f(X)]$, then

$$\begin{aligned}\mathbb{P}[f(X)1_A] &= \mathbb{P}\{\mathbb{P}[f(X)1_A|\mathscr{G}]\} = \mathbb{P}\{1_A\mathbb{P}[f(X)|\mathscr{G}]\} \\ &= \mathbb{P}\{1_A\mathbb{P}[f(X)]\} = \mathbb{P}[f(X)]\mathbb{P}(A).\end{aligned}$$

So X is independent to $\mathscr{G}$.

2. $3 \Leftrightarrow 4$. The proof is essentially the same as $1 \Leftrightarrow 2$.

3. $1(2) \Leftrightarrow 3(4)$. First $2 \Longrightarrow 3$ is obvious. On the contrary, we show $3 \Rightarrow 1$. Note $\mathbb{R}$ is a polish space, then the regular conditional distribution of X exists. Namely,

$$\mu_X^{\mathscr{G}}(dx) := \mathbb{P}(X \in dx|\mathscr{G})$$

is a measure. And we have

$$\mathbb{P}(e^{i\theta X}|\mathscr{G}) = \int e^{i\theta x}\mu_X^{\mathscr{G}}(dx), .$$

Since we have $\mathbb{P}(e^{i\theta X}|\mathscr{G}) = \mathbb{P}(e^{i\theta X})$, then

$$\int e^{i\theta x}\mu_X^{\mathscr{G}}(dx) = \mathbb{P}(e^{i\theta X}|\mathscr{G}) = \mathbb{P}(e^{i\theta X}) = \int e^{i\theta x}\mu_X(dx).$$

Since the characteristic function and the distribution are mutually uniquely determined, then we have $\mu_X^{\mathscr{G}}(dx) = \mu_X(dx)$. Namely

$$\mathbb{P}(X \in A|\mathscr{G}) = \mathbb{P}(X \in A), \quad \forall A \in \mathscr{B}(\mathbb{R}),$$

which implies X is independent to $\mathscr{G}$. □

QUESTION 6.55 *Given a random variable X on probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and valued on $(\bar{\mathbb{R}}_+, \mathscr{B}(\bar{\mathbb{R}}_+))$, where $\bar{\mathbb{R}}_+ = \mathbb{R}_+ \cup \{+\infty\}$ Consider its Laplace trasformation $\hat{f}(r) = \mathbb{P}(e^{-rX}), r > 0$. Then we have*

$$\lim_{r \downarrow 0} \hat{f}(r) = \mathbb{P}(X < +\infty).$$

Proof. Note

$$\begin{aligned} e^{-rX} &= e^{-rX} \mathbb{1}_{[X<+\infty]} + e^{-r\cdot\infty} \mathbb{1}_{[X=+\infty]} \\ &= e^{-rX} \mathbb{1}_{[X<+\infty]} \uparrow \mathbb{1}_{[X<+\infty]}, \quad r \downarrow 0. \end{aligned}$$

Using the dominant convergence theorem to obtain

$$\lim_{r \downarrow 0} \hat{f}(r) = \lim_{r \downarrow 0} \mathbb{P}(e^{-rX}) = \mathbb{P}(X < +\infty).$$

□

QUESTION 6.56 *Given random variables $(\xi_n)_{n\geq 1}, (\eta_n)_{n\geq 1}$ and ζ on probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Assume $\xi_n - \eta_n \xrightarrow{\mathbb{P}} 0$ and $\xi_n \xrightarrow{\mathrm{w}} \zeta$, then $\eta_n \xrightarrow{\mathrm{w}} \zeta$.*

Proof. For any $x \in C(F_\zeta)$, note $C^c(F_\zeta)$ is at most countable, then $C(F_\zeta)$ is dense in $\mathbb{R}$. Thus we can choose $\varepsilon > 0$ such that $x + \varepsilon \in C(F_\zeta)$. Therefore for ant $k \geq 1$,

$$\begin{aligned} \mathbb{P}(\eta_n \leq x) &= \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| < \varepsilon) + \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| \geq \varepsilon) \\ &\leq \mathbb{P}(\xi_n - \varepsilon \leq x, |\xi_n - \eta_n| < \varepsilon) + \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| \geq \varepsilon) \\ &= \mathbb{P}(\xi_n \leq x + \varepsilon) - \mathbb{P}(\xi_n - \varepsilon \leq x, |\xi_n - \eta_n| \geq \varepsilon) \\ &\quad + \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| \geq \varepsilon). \end{aligned}$$

Let $n \to \infty$ and note $\mathbb{P}(|\xi_n - \eta_n| \geq \varepsilon) \to 0, n \to \infty$ for each $k \geq 1$, then we have

$$\overline{\lim_{n\to\infty}} \mathbb{P}(\eta_n \leq x) \leq \mathbb{P}(\zeta \leq x + \varepsilon).$$

Let $\varepsilon \to 0$, we have

$$\overline{\lim_{n\to\infty}} \mathbb{P}(\eta_n \leq x) \leq \mathbb{P}(\zeta \leq x).$$

On the other side, similarly, we can choose $\delta > 0$ such that $x - \delta \in C(F_\zeta)$, so

$$\begin{aligned} \mathbb{P}(\eta_n \leq x) &= \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| < \delta) + \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| \geq \delta) \\ &\geq \mathbb{P}(\xi_n + \delta \leq x, |\xi_n - \eta_n| < \delta) + \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| \geq \delta) \\ &= \mathbb{P}(\xi_n \leq x - \delta) - \mathbb{P}(\xi_n + \delta \leq x, |\xi_n - \eta_n| \geq \delta) \\ &\quad + \mathbb{P}(\eta_n \leq x, |\xi_n - \eta_n| \geq \delta). \end{aligned}$$

Let $n \to \infty$ to obtain

$$\varliminf_{n\to\infty} \mathbb{P}(\eta_n \le x) \ge \mathbb{P}(\zeta \le x - \delta).$$

Let $\delta \to 0$ and note $x \in C(F_\zeta)$, we have

$$\varliminf_{n\to\infty} \mathbb{P}(\eta_n \le x) \ge \mathbb{P}(\zeta < x) = \mathbb{P}(\zeta \le x).$$

The above argument yields

$$F_{\eta_n}(x) = \mathbb{P}(\eta_n \le x) \to \mathbb{P}(\zeta \le x) = F_\zeta(x), \quad \forall x \in C(F_\zeta).$$

□

QUESTION 6.57 *Given random variables $(X_n)_{n\ge1}, (Y_n)_{n\ge1}, X, Y$ on probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Assume $X_n \xrightarrow{\mathbb{P}} X$ and $Y_n \xrightarrow{\mathrm{w}} Y$, then the following four claims hold:*

1. *$X_n \xrightarrow{\mathbb{P}} X$ implies $X_n \xrightarrow{\mathrm{w}} X$.*
2. *If $X = C \in \mathbb{R}$ is a constant, then $X_n \xrightarrow{\mathbb{P}} C$ is equivalent to $X_n \xrightarrow{\mathrm{w}} C$.*
3. *If $X_n \xrightarrow{\mathbb{P}} C$ is a constant, then $X_n + Y_n \xrightarrow{\mathrm{w}} C + Y$.*
4. *If $X_n \xrightarrow{\mathbb{P}} C$ is a constant, then $X_nY_n \xrightarrow{\mathrm{w}} CY$.*

Proof. 1. Apply the above **QUESTION** with $\xi_n = X, \eta_n = X_n$ and $\zeta = X$.

2. The first sub-question shows that $X_n \xrightarrow{\mathbb{P}} C$ implies $X_n \xrightarrow{\mathrm{w}} C$, we only need to show the converse implication. For any $\varepsilon > 0$, we have

$$\begin{aligned}\mathbb{P}(|X_n - C| \ge \varepsilon) &= \mathbb{P}(X_n \ge C + \varepsilon) + \mathbb{P}(X_n \le C - \varepsilon)\\ &= 1 - \mathbb{P}(X_n < C + \varepsilon) + \mathbb{P}(X_n \le C - \varepsilon).\end{aligned}$$

Note $F_C(x)$ has only one discontinuous point $x = C$, thus $C + \varepsilon, C - \varepsilon \in C(F_C)$. Let $n \to \infty$, we will get

$$\begin{aligned}\varlimsup_{n\to\infty} \mathbb{P}(|X_n - C| \ge \varepsilon) &= 1 - \lim_{n\to\infty} \mathbb{P}(X_n < C + \varepsilon) + \lim_{n\to\infty} \mathbb{P}(X_n \le C - \varepsilon)\\ &= 1 - F_C(C + \varepsilon) + F_C(C - \varepsilon) = 1 - 1 + 0 = 0.\end{aligned}$$

3. Since we have $(X_n + Y_n) - (C + Y_n) = X_n - C \xrightarrow{\mathbb{P}} 0$. Moreover we can get $C + Y_n \xrightarrow{\mathrm{w}} C + Y$ from $Y_n \xrightarrow{\mathrm{w}} Y$. In fact, it can easily be found that $C(F_{C+Y}) = C(F_Y) + C := \{y + C : y \in C(F_Y)\}$. Therefore, for any $x \in C(F_{C+Y})$, we have $x - C \in C(F_Y)$. And

$$\begin{aligned}\mathbb{P}(C + Y_n \le x) &= \mathbb{P}(Y_n \le x - C)\\ &\to \mathbb{P}(Y \le X - C) = \mathbb{P}(C + Y \le x), \quad \forall x \in C(F_{C+Y})\end{aligned}$$

So we can apply the previous **QUESTION** with $\xi_n = C+Y_n, \eta_n = X_n+Y_n$ and $\zeta = C+Y$.

4. Since we have proved in the 2nd sub-question that, convergence in probability to a constant is equivalent to convergence in distribution, we only need to use $\xi_n \xrightarrow{\mathbb{P}} \zeta$ and $\xi_n \xrightarrow{\mathbb{P}} 0$ to show $\xi_n\eta_n \xrightarrow{\mathbb{P}} 0$. However, this is obviously true, since the function $f(x,y) = xy$ is continuous in $\mathbb{R}^2$ and $\mathbb{P}$ is a finite measure. We only need to prove this assertion for $C \neq 0$.

   Suppose $C > 0$. For any $x \in C(F_{CY})$ and $x \geq 0$, note $C(F_Y)$ is dense in $\mathbb{R}$, we can choose $\varepsilon > 0$ such that $C - \varepsilon > 0$ and $\frac{x}{C-\varepsilon} \in C(F_Y)$. Then for any $k \geq 1$,

$$\begin{aligned}
&\mathbb{P}(X_nY_n \leq x) = \mathbb{P}(X_nY_n \leq x, |X_n - C| < \varepsilon) + \mathbb{P}(X_nY_n \leq x, |X_n - C| \geq \varepsilon)\\
=&\mathbb{P}(X_nY_n \leq x, Y_n \geq 0, |X_n - C| < \varepsilon) + \mathbb{P}(X_nY_n \leq x, Y_n < 0, |X_n - C| < \varepsilon)\\
&+ \mathbb{P}(X_nY_n \leq x, |X_n - C| \geq \varepsilon)\\
\leq&\mathbb{P}[(C-\varepsilon)Y_n \leq x, Y_n \geq 0, |X_n - C| < \varepsilon] + \mathbb{P}(Y_n < 0, |X_n - C| < \varepsilon) + o(1)\\
=&\mathbb{P}\left(0 \leq Y_n \leq \frac{x}{C-\varepsilon}\right) + \mathbb{P}(Y_n < 0) + o(1)\\
=&\mathbb{P}\left(Y_n \leq \frac{x}{C-\varepsilon}\right) + o(1).
\end{aligned}$$

   Let $n \to \infty$, we have

$$\overline{\lim_{n\to\infty}}\, \mathbb{P}(X_ny_n \leq x) \leq \mathbb{P}\left(Y \leq \frac{x}{C-\varepsilon}\right).$$

   Let $\varepsilon \to 0$ and we will obtain

$$\overline{\lim_{n\to\infty}}\, \mathbb{P}(X_ny_n \leq x) \leq \mathbb{P}(Y \leq x/C) = \mathbb{P}(CY \leq x).$$

   If $x \in C(F_{CY})$ and $x < 0$, we can choose $\varepsilon > 0$ such that $\frac{x}{C+\varepsilon} \in C(F_Y)$. Then

$$\begin{aligned}
\mathbb{P}(X_nY_n \leq x) &= \mathbb{P}(X_nY_n \leq x, Y_n < 0, |X_n - C| < \varepsilon) + o(1)\\
&\leq \mathbb{P}[(C+\varepsilon)Y_n \leq x, Y_n < 0, |X_n - C| < \varepsilon] + o(1)\\
&= \mathbb{P}\left(Y_n \leq \frac{x}{C+\varepsilon}\right) + o(1).
\end{aligned}$$

   Let $n \to \infty$, we have

$$\overline{\lim_{n\to\infty}}\, \mathbb{P}(X_ny_n \leq x) \leq \mathbb{P}\left(Y \leq \frac{x}{C+\varepsilon}\right).$$

   Let $\varepsilon \to 0$ and note $C(F_{CY}) = C \cdot C(F_Y) = \{Cy : y \in C(F_Y)\}$. Therefore for all $x \in C(F_{CY})$, we have $x/C \in C(F_Y)$. Then we will

again obtain

$$\overline{\lim_{n\to\infty}}\,\mathbb{P}(X_nY_n \le x) \le \mathbb{P}(Y < x/C) = \mathbb{P}(Y \le x/C) = \mathbb{P}(CY \le x).$$

In short, we have $\overline{\lim}_{n\to\infty} F_{X_nY_n}(x) \le F_{CY}(x)$ for any $x \in C(F_{CY})$.

Similarly, if $C > 0$ and $x \ge 0$, we can select $\delta > 0$ such that $\frac{x}{C+\delta} \in C(F_Y)$. So

$$\begin{aligned}\mathbb{P}(X_nY_n \le x) &\ge \mathbb{P}[(C+\delta)Y_n \le x, Y_n \ge 0] + \mathbb{P}(Y_n < 0) + o(1)\\ &= \mathbb{P}\left(Y_n \le \frac{x}{C+\delta}\right) + o(1).\end{aligned}$$

Let $n \to \infty$ to find

$$\underline{\lim_{n\to\infty}}\,\mathbb{P}(X_nY_n \le x) \ge \mathbb{P}\left(Y \le \frac{x}{C+\delta}\right).$$

Let $\delta \to 0$ to obtain

$$\underline{\lim_{n\to\infty}}\,\mathbb{P}(X_nY_n \le x) \ge \mathbb{P}(Y < x/C) = \mathbb{P}(Y \le x/C) = \mathbb{P}(CY \le x).$$

If $x \in C(F_{CY})$ and $x < 0$, we can select $\delta > 0$ such that $\frac{x}{C-\delta} \in C(F_Y)$, therefore

$$\mathbb{P}(X_nY_n \le x) \ge \mathbb{P}\left(Y_n \le \frac{x}{C-\delta}\right) + o(1).$$

Let $n \to \infty$ and then let $\delta \to 0$, we will get

$$\underline{\lim_{n\to\infty}}\,\mathbb{P}(X_nY_n \le x) \ge \mathbb{P}(CY \le x).$$

In short, we have gotten $\underline{\lim}_{n\to\infty} F_{X_nY_n}(x) \ge F_{CY}(x)$ for any $x \in C(F_{CY})$.

Hence when $C > 0$, we have shown $F_{X_nY_n}(x) \to F_{CY}(x)$ for any $x \in C(F_{CY})$.

When $C < 0$, we have $-X_n \xrightarrow{\mathbb{P}} -C > 0$. So the above discuss yields $-X_nY_n \xrightarrow{\mathrm{w}} -CY$. However, using the characteristic function will give us

$$\mathbb{E}e^{i(-t)X_nY_n} = \mathbb{E}e^{it(-X_nY_n)} \to \mathbb{E}e^{it(-CY)} = \mathbb{E}e^{i(-t)CY}, \quad t \in \mathbb{R},$$

which shows $X_nY_n \xrightarrow{\mathrm{w}} CY$. □

QUESTION 6.58 *Given a Poisson process $N = (N_t)_{t\geq 0}$ and i.i.d. random variables $(Y_i)_{i\geq 1}$ on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Then the compound Poisson process*

$$X_t := \sum_{i=1}^{N_t} Y_i, \quad t \geq 0$$

has independent increment.

Proof. For any $0 < s < t$, we have to show the following random variables are independent,

$$X_s := \sum_{i=1}^{N_s} Y_i, \quad X_t - X_s = \sum_{i=N_s+1}^{N_t} Y_i,$$

where $\sum_{i=m}^{n} \cdot = 0$ for any $m > n$. If $N_t = N_s$, we know $X_t - X_s = 0$, and therefore the theorem is trivially true. If $N_t > N_s$, for any $f, g \in \mathscr{B}(\mathbb{R})$, we have

$$\begin{aligned}
\mathbb{P}[f(X_s)g(X_t - X_s)] &= \mathbb{P}\{\mathbb{P}[f(X_s)g(X_t - X_s)|N]\} \\
&= \mathbb{P}\left\{\mathbb{P}\left[f\left(\sum_{i=1}^{k_s} Y_i\right) g\left(\sum_{i=k_s+1}^{k_t} Y_i\right)\right]\Bigg|_{k_s=N_s, k_t=N_t}\right\} \\
&= \mathbb{P}\left\{\mathbb{P}\left[f\left(\sum_{i=1}^{k_s} Y_i\right)\right] \mathbb{P}\left[g\left(\sum_{i=k_s+1}^{k_t} Y_i\right)\right]\Bigg|_{k_s=N_s, k_t=N_t}\right\} \\
&=: \mathbb{P}[F(N_s)G(N_s, N_t)].
\end{aligned}$$

Observe $(Y_i)_{i\geq 1}$ is an i.i.d. random variable sequence, then

$$G(k_s, k_t) = \mathbb{P}\left[g\left(\sum_{i=k_s+1}^{k_t} Y_i\right)\right], \quad k_s \leq k_t$$

only depends on the deference $k_t - k_s \in \mathbb{Z}_+$. Thus we can write $\widetilde{G}(k_t - k_s) := G(k_s, k_t)$. Finally, we can apply the independent increment property of $(N_t)_{t\geq 0}$ to obtain

$$\begin{aligned}
\mathbb{P}[f(X_s)g(X_t - X_s)] &= \mathbb{P}[F(N_s)\widetilde{G}(N_t - N_s)] \\
&= \mathbb{P}[F(N_s)]\mathbb{P}[\widetilde{G}(N_t - N_s)] \\
&= \mathbb{P}\left\{\mathbb{P}\left[f\left(\sum_{i=1}^{k_s} Y_i\right)\right]\Bigg|_{k_s=N_s}\right\} \mathbb{P}\left\{\mathbb{P}\left[g\left(\sum_{i=k_s+1}^{k_t} Y_i\right)\right]\Bigg|_{k_s=N_s, k_t=N_t}\right\} \square \\
&= \mathbb{P}\{\mathbb{P}[f(X_s)|N]\}\mathbb{P}\{\mathbb{P}[g(X_t - X_s)]|N\} \\
&= \mathbb{P}[f(X_s)]\mathbb{P}[g(X_t - X_s)].
\end{aligned}$$

QUESTION 6.59 *Given a probability space* $(\Omega, \mathscr{F}, \mathbb{P})$ *and* $A, B \in \mathscr{F}$. *Then we have* $|\mathbb{P}(AB) - \mathbb{P}(A)\mathbb{P}(B)| \leq 1/4$.

Proof. Note $\mathbb{P}(AB) - \mathbb{P}(A)\mathbb{P}(B) = \mathrm{Cov}(1_A, 1_B)$. So

$$\begin{aligned}|\mathrm{Cov}(1_A, 1_B)| &\leq \sqrt{\mathrm{Var}(1_A)\mathrm{Var}(1_B)} \\ &\leq \sqrt{[\mathbb{P}(A) - \mathbb{P}^2(A)]\,[\mathbb{P}(B) - \mathbb{P}^2(B)]} \\ &\leq \sqrt{(1/4)^2} = 1/4,\end{aligned}$$

where the last inequality comes from the quadratic function $f(x) = x - x^2 \leq 1/4$ for any $x \in \mathbb{R}$. □

QUESTION 6.60 *Given a probability space* $(\Omega, \mathscr{F}, \mathbb{P})$. *Suppose* $A \in \mathscr{F}$ *satisfies* $\mathbb{P}(A) > 0$. *Define* $\mathbb{Q}(B) := \mathbb{P}(B|A)$. *Prove for any* $F \in \mathrm{b}\mathscr{F}$ *and sub-*σ *algebra* $\mathscr{C}$,

$$\mathbb{Q}(F|\mathscr{C}) = \frac{\mathbb{P}(F1_A|\mathscr{C})}{\mathbb{P}(A|\mathscr{C})}, \quad \mathbb{Q}\text{-a.s..}$$

Proof. First we show $\mathbb{P}(A|\mathscr{C}) \neq 0, \mathbb{Q}$-a.s.. Define $\Lambda := \{\mathbb{P}(A|\mathscr{C}) = 0\} \in \mathscr{C}$. We have

$$0 = \mathbb{P}[1_\Lambda \mathbb{P}(A|\mathscr{C})] = \mathbb{P}(1_\Lambda 1_A) = \mathbb{P}(\Lambda \cap A).$$

Thus $\mathbb{Q}(\Lambda) = \mathbb{P}(\Lambda|A) = \mathbb{P}(\Lambda \cap A)/\mathbb{P}(A) = 0$. This implies $\mathbb{P}(A|\mathscr{C}) \neq 0, \mathbb{Q}$-a.s..

Applying the monotonic class theorem, we can show

$$\int F d\mathbb{Q} = \frac{1}{\mathbb{P}(A)} \int F 1_A d\mathbb{P}, \quad F \in \mathrm{b}\mathscr{F}.$$

For any $G \in \mathrm{b}\mathscr{C}$, we have

$$\begin{aligned}\int \frac{\mathbb{P}(F1_A|\mathscr{C})}{\mathbb{P}(A|\mathscr{C})} G d\mathbb{Q} &= \frac{1}{\mathbb{P}(A)} \int \frac{\mathbb{P}(F1_A|\mathscr{C})}{\mathbb{P}(A|\mathscr{C})} G 1_A d\mathbb{P} \\ &= \frac{1}{\mathbb{P}(A)} \int \frac{\mathbb{P}(F1_A|\mathscr{C})}{\mathbb{P}(A|\mathscr{C})} G \mathbb{P}(A|\mathscr{C}) d\mathbb{P} \\ &= \frac{1}{\mathbb{P}(A)} \int G \mathbb{P}(F1_A|\mathscr{C}) d\mathbb{P} \\ &= \frac{1}{\mathbb{P}(A)} \int G F 1_A d\mathbb{P} \\ &= \int G F d\mathbb{Q}.\end{aligned}$$

This implies the statement holds. □

QUESTION 6.61 *Given a probability space* $(\Omega, \mathscr{F}, \mathbb{P})$. *Suppose* $\mathscr{D} = \sigma(A_i, i \geq 1)$, *where* A_i *is a partition of* Ω *and* $\mathscr{C} \vee \mathscr{D} := \sigma(\mathscr{C} \cup \mathscr{D})$. *For any* $B \in \mathscr{F}$,

$$(*) \qquad \mathbb{P}(B|\mathscr{C} \vee \mathscr{D}) = \sum_{n\geq 1} \frac{\mathbb{P}(A_n \cap B|\mathscr{C})}{\mathbb{P}(A_n|\mathscr{C})} \mathbb{1}_{A_n}.$$

Proof. If there exists $m \geq 1$ such that $\mathbb{P}(A_m) = 0$, then $\mathbb{P}[\mathbb{P}(A_m|\mathscr{C})] = \mathbb{P}(A_m) = 0$ and $\mathbb{P}(A_m|\mathscr{C}) \geq 0$ yield $\mathbb{P}(A_m|\mathscr{C}) = 0, \mathbb{P}$-a.s.. So

$$\frac{\mathbb{P}(A_m \cap B|\mathscr{C})}{\mathbb{P}(A_m|\mathscr{C})} = \infty.$$

Noticing $\mathbb{P}(A_m) = 0$, we can define

$$\frac{\mathbb{P}(A_m \cap B|\mathscr{C})}{\mathbb{P}(A_m|\mathscr{C})} \mathbb{1}_{A_m} = c\mathbb{1}_{A_m}, \quad \forall c \in \mathbb{R} \cup \{\infty\}.$$

In short, the right side of $(*)$ is well-defined.

For any $C \in \mathscr{C}$ and $D \in \mathscr{D}$, there exist $\{A_{n_k}\}_{k=1}^m \subset \{A_i\}_{i\geq 1}$ such that $D = \sum_{k=1}^m A_{n_k}$. Suppose without loss of generality $\{A_{n_k}\}_{k=1}^\ell$ are $\mathbb{P}$-null sets, where $1 \leq \ell \leq m$. Applying the result to the previous question, we have

$$\begin{aligned}
\int \sum_{n\geq 1} \frac{\mathbb{P}(A_n \cap B|\mathscr{C})}{\mathbb{P}(A_n|\mathscr{C})} \mathbb{1}_{A_n} \mathbb{1}_{C\cap D} d\mathbb{P} &= \int \sum_{k=1}^m \frac{\mathbb{P}(A_{n_k} \cap B|\mathscr{C})}{\mathbb{P}(A_{n_k}|\mathscr{C})} \mathbb{1}_{A_{n_k}} \mathbb{1}_C d\mathbb{P} \\
&= \sum_{k=\ell+1}^m \int \frac{\mathbb{P}(A_{n_k} \cap B|\mathscr{C})}{\mathbb{P}(A_{n_k}|\mathscr{C})} \mathbb{1}_{A_{n_k}} \mathbb{1}_C d\mathbb{P} \\
&= \sum_{k=\ell+1}^m \int \frac{\mathbb{P}(A_{n_k} \cap B|\mathscr{C})}{\mathbb{P}(A_{n_k}|\mathscr{C})} \mathbb{P}(A_{n_k}|\mathscr{C}) \mathbb{1}_C d\mathbb{P} \\
&= \sum_{k=\ell+1}^m \int \mathbb{P}(A_{n_k} \cap B|\mathscr{C}) \mathbb{1}_C d\mathbb{P} \\
&= \sum_{k=\ell+1}^m \int \mathbb{1}_{A_{n_k}\cap B} \mathbb{1}_C d\mathbb{P} \\
&= \int \sum_{k=1}^m \mathbb{1}_{A_{n_k}} 1_B \mathbb{1}_C d\mathbb{P} \\
&= \int \sum_{n\geq 1} \mathbb{1}_{A_n} 1_B \mathbb{1}_C 1_D d\mathbb{P} \\
&= \int 1_B \mathbb{1}_{C\cap D} d\mathbb{P}.
\end{aligned}$$

This implies the statement holds. □

QUESTION 6.62 *Suppose* X *is a random variable on the probability space* $(\Omega, \mathscr{F}, \mathbb{P})$ *and* $\mathscr{G}, \mathscr{A} \subset \mathscr{F}$ *are two sub* σ*-algebras. Then* $\mathbb{P}(X|\mathscr{G} \vee \mathscr{A}) = \mathbb{P}(X|\mathscr{G})$ *iff* X *and* $\mathscr{A}$ *are conditional independent with respect to* $\mathscr{G}$.

Proof. For any $G \in \mathscr{G}, A \in \mathscr{A}$, we have

$$\begin{aligned}\mathbb{P}(1_G 1_A X) &= \mathbb{P}[1_G \mathbb{P}(1_A X|\mathscr{G})] = \mathbb{P}[1_G \mathbb{P}(A|\mathscr{G})\mathbb{P}(X|\mathscr{G})] \\ &= \mathbb{P}[1_G 1_A \mathbb{P}(X|\mathscr{G})].\end{aligned}$$

where the second equality holds because X and $\mathscr{A}$ are conditional independent with respect to $\mathscr{G}$. The monotone class theorem yields $\mathbb{P}(1_B X) = \mathbb{P}[1_B \mathbb{P}(X|\mathscr{G})]$, where $B \in \mathscr{G} \vee \mathscr{A}$. So $\mathbb{P}(X|\mathscr{G}) = \mathbb{P}(X|\mathscr{G} \vee \mathscr{A})$. On the contrary, if $\mathbb{P}(X|\mathscr{G} \vee \mathscr{A}) = \mathbb{P}(X|\mathscr{G})$, then for any $G \in \mathscr{G}, A \in \mathscr{A}$,

$$(*) \qquad \mathbb{P}(1_G 1_A X) = \mathbb{P}[1_G 1_A \mathbb{P}(X|\mathscr{G})] = \mathbb{P}[1_G \mathbb{P}(A|\mathscr{G})\mathbb{P}(X|\mathscr{G})].$$

So $(*)$ shows $\mathbb{P}(A|\mathscr{G})\mathbb{P}(X|\mathscr{G}) = \mathbb{P}(1_A X|\mathscr{G})$, which means X and $\mathscr{A}$ are conditional independent with respect to $\mathscr{G}$. □

QUESTION 6.63 *There is a random variable X on $(\Omega, \mathscr{F}, \mathbb{P})$ such that $\mathbb{P}(|X|^\alpha) = \infty$ for any $\alpha > 0$.*

Proof. Consider a continuous random variable X with density

$$f(x) = \begin{cases} \frac{1}{2|x|\log^2|x|}, & \text{if } |x| > e, \\ 0, & \text{otherewise.} \end{cases}$$

Because

$$\begin{aligned}\int_{-\infty}^{\infty} f(x)dx &= \int_{[|x|>e]} \frac{dx}{2|x|\log^2|x|} = \int_e^\infty \frac{dx}{x\log^2 x} \\ &= \int_e^\infty \frac{d\log x}{\log^2 x} = \frac{1}{\log x}\bigg|_\infty^e = 1,\end{aligned}$$

$f(x)$ is indeed a density. And for any $\alpha > 0$, we have $\frac{x^\alpha}{\log^2 x} \to \infty, x \to \infty$. So there exists $C > 0$ such that for any $\frac{x^\alpha}{\log^2 x} \geq 1$ for any $x > C$. Whence

$$\begin{aligned}\mathbb{P}(|X|^\alpha) &= \int_{[|x|>e]} \frac{|x|^\alpha}{2|x|\log^2|x|}dx = \int_e^\infty \frac{x^\alpha}{x\log^2 x}dx \\ &\geq \int_e^\infty x^{-1}dx = \infty.\end{aligned}$$

□

QUESTION 6.64 *There is two random variables X, Y on $(\Omega, \mathscr{F}, \mathbb{P})$ such that X, Y are related but there exists some $f \in \mathrm{b}\mathscr{B}$ such that $\mathbb{P}[f(X)|Y] = \mathbb{P}[f(X)]$.*

Proof. Consider $\mathbb{P}(Y = \pm 1) = 1/2$ and $X = -Y$. Obviously, X, Y are related. Let $f(x) = x^2$. We have $\mathbb{P}[f(X)|Y] = \mathbb{P}(X^2|Y) = 1 = \mathbb{P}(X^2)$. □

QUESTION 6.65 *Assume X, Y, D are three random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and $\mathbb{P}|Y| < \infty$. If $\mathbb{P}[f(X)|D] \neq \mathbb{P}[f(X)]$ for any non-constant $f \in \mathscr{B}$ satisfying $\mathbb{P}[|f(X)|] < \infty$, and Y, D are independent, then $\mathbb{P}(Y|X, D) = \mathbb{P}(Y|X)$ implies $\mathbb{P}(Y|X) = \mathbb{P}Y$.*

Proof. Noting $\mathbb{P}[\mathbb{P}(Y|X, D)|D] = \mathbb{P}(Y|D)$, combining the fact $\mathbb{P}(Y|X, D) = \mathbb{P}(Y|X)$ and Y, D are independent, we have

$$\mathbb{P}[\mathbb{P}(Y|X)|D] = \mathbb{P}[\mathbb{P}(Y|X, D)|D] = \mathbb{P}(Y|D) = \mathbb{P}Y.$$

Since $\mathbb{P}[f(X)|D] \neq \mathbb{P}[f(X)]$ for any non-constant $f \in \mathscr{B}$ satisfying $\mathbb{P}[|f(X)|] < \infty$, and $\mathbb{P}Y = \mathbb{P}[\mathbb{P}(Y|X)]$, we have $\mathbb{P}(Y|X) = \mathbb{P}Y$ is a constant. □

QUESTION 6.66 *Assume X, Y are two random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and $A \in \mathscr{B}$. Prove*

$$\mathbb{P}[\mathbb{P}(Y|X)|X \in A] = \mathbb{P}(Y|X \in A).$$

Proof. We just need to note

$$\begin{aligned}
\mathbb{P}[\mathbb{P}(Y|X)|X \in A] &= \frac{1}{\mathbb{P}(X \in A)} \int \mathbb{P}(Y|X) \mathbb{1}_{[X \in A]} d\mathbb{P} \\
&= \frac{1}{\mathbb{P}(X \in A)} \int Y \mathbb{1}_{[X \in A]} d\mathbb{P} \\
&= \mathbb{P}(Y|X \in A).
\end{aligned}$$

□

QUESTION 6.67 *Assume X, Y, Z are three random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and $A \in \mathscr{B}$. Prove*

$$\mathbb{P}[\mathbb{P}(X|Y, Z \in A)|Z \in A] = \mathbb{P}(X|Z \in A).$$

Proof. Since

$$\mathbb{P}(X|Y, Z \in A) = \frac{\mathbb{P}(X \mathbb{1}_{[Z \in A]}|Y)}{\mathbb{P}(Z \in A|Y)}.$$

The definition of conditional expectation yields

$$\begin{aligned}
\mathbb{P}[\mathbb{P}(X|Y, Z \in A)|Z \in A] &= \frac{1}{\mathbb{P}(Z \in A)} \int \mathbb{P}(X|Y, Z \in A) \mathbb{1}_{[Z \in A]} d\mathbb{P} \\
&= \frac{1}{\mathbb{P}(Z \in A)} \int \frac{\mathbb{P}(X \mathbb{1}_{[Z \in A]}|Y)}{\mathbb{P}(Z \in A|Y)} \mathbb{1}_{[Z \in A]} d\mathbb{P} \\
&= \frac{1}{\mathbb{P}(Z \in A)} \int \frac{\mathbb{P}(X \mathbb{1}_{[Z \in A]}|Y)}{\mathbb{P}(Z \in A|Y)} \mathbb{P}(Z \in A|Y) d\mathbb{P}
\end{aligned}$$

$$
\begin{aligned}
&= \frac{1}{\mathbb{P}(Z \in A)} \int \mathbb{P}(X \mathbb{1}_{[Z \in A]} | Y) d\mathbb{P} \\
&= \frac{1}{\mathbb{P}(Z \in A)} \mathbb{P}[\mathbb{P}(X \mathbb{1}_{[Z \in A]} | Y)] \\
&= \frac{1}{\mathbb{P}(Z \in A)} \mathbb{P}(X \mathbb{1}_{[Z \in A]}) \\
&= \mathbb{P}(X | Z \in A).
\end{aligned}
$$

□

QUESTION 6.68 *Suppose $(\Omega, \mathscr{F})$ is a measurable space. Assume $\mathscr{G} \subset \mathscr{F}$ is a sub σ algerba and $A \in \mathscr{F}$. Prove*

$$\mathscr{G} \vee \{A\} := \sigma(\mathscr{G} \cup \{A\}) = \{(B_1 \cap A) \cup (B_2 \cap A^c) : B_1, B_2 \in \mathscr{G}\}.$$

Proof. Write $\mathscr{A} := \{(B_1 \cap A) \cup (B_2 \cap A^c) : B_1, B_2 \in \mathscr{G}\}$. Obviously $\mathscr{A} \subset \mathscr{G} \vee \{A\}$. We only need to prove $\mathscr{A}$ is a σ algebra. First, $\Omega \in \mathscr{A}$ is obvious. Next, for any $(B_1 \cap A) \cup (B_2 \cap A^c) \in \mathscr{A}$, we have

$$
\begin{aligned}
[(B_1 \cap A) \cup (B_2 \cap A^c)]^c &= (B_1^c \cup A^c) \cap (B_2^c \cup A) \\
&= [(B_1^c \cup A^c) \cap B_2^c] \cup [(B_1^c \cup A^c) \cap A] \\
&= [(B_1^c \cap B_2^c) \cup (A^c \cap B_2^c)] \cup (B_1^c \cap A).
\end{aligned}
$$

We claim $B_1^c \cap B_2^c \subset (A^c \cap B_2^c) \cup (B_1^c \cap A)$. In fact, since A and A^c is a partition of Ω, for any $\omega \in B_1^c \cap B_2^c$, we have either $w \in A$ or $w \in A^c$. Without loss of generality, we assume $\omega \in A^c$. Because $\omega \in B_1^c \cap B_2^c$, we have $\omega \in A^c \cap B_1^c \cap B_2^c \subset A^c \cap B_2^c \subset (A^c \cap B_2^c) \cup (B_1^c \cap A)$. So, the claim is true. The formula above turns out to be

$$[(B_1 \cap A) \cup (B_2 \cap A^c)]^c = (A^c \cap B_2^c) \cup (B_1^c \cap A) \in \mathscr{A}.$$

Finally, for any sequence $\{(B_{1,k} \cap A) \cup (B_{2,k} \cap A^c)\}_{k \geq 1} \subset \mathscr{A}$, we have

$$
\begin{aligned}
\bigcup_{k \geq 1} [(B_{1,k} \cap A) \cup (B_{2,k} \cap A^c)] &= \left[\bigcup_{k \geq 1} (B_{1,k} \cap A)\right] \cup \left[\bigcup_{k \geq 1} (B_{2,k} \cap A^c)\right] \\
&= \left[\left(\bigcup_{k \geq 1} B_{1,k}\right) \cap A\right] \cup \left[\left(\bigcup_{k \geq 1} B_{2,k}\right) \cap A^c\right] \in \mathscr{A}.
\end{aligned}
$$

In conclusion, $\mathscr{A}$ is a σ algebra and therefore $\mathscr{G} \vee \{A\} = \mathscr{A}$. □

QUESTION 6.69 *Suppose $(\Omega, \mathscr{F})$ is a measurable space. Assume $\mathscr{G} \subset \mathscr{F}$ is a sub σ algerba and $A \in \mathscr{F}$. For any measurable function $H \in \mathscr{G} \vee \{A\}$, there exist $F_1, F_2 \in \mathscr{G}$ such that $H = F_1 \mathbb{1}_A + F_2 \mathbb{1}_{A^c}$.*

Proof. Define

$$\Lambda := \{H \in \mathscr{G} \vee \{A\} : \text{there exist } F_1, F_2 \in \mathscr{G} \text{ such that } H = F_1 \mathbb{1}_A + F_2 \mathbb{1}_{A^c}\}.$$

Based on the previous QUESTION, we know all the indicator functions of $\mathscr{G} \vee \{A\}$ are in Λ. For any $H, H' \in \Lambda$, there exist $F_1, F_1', F_2, F_2' \in \mathscr{G}$ such that

$$H = F_1 \mathbb{1}_A + F_2 \mathbb{1}_{A^c}, \quad H' = F_1' \mathbb{1}_A + F_2' \mathbb{1}_{A^c}.$$

Therefore, for any $\alpha, \beta \in \mathbb{R}$,

$$\alpha H + \beta H' = (\alpha F_1 + \beta F_1')\mathbb{1}_A + (\alpha F_2 + \beta F_2')\mathbb{1}_{A^c} \in \Lambda.$$

If $H \in \mathscr{G} \vee \{A\}$ and there exists a sequence $\{H_n\}_{n\geq 1} \subset \Lambda$ satisfying $0 \leq H_n = F_{1,n}\mathbb{1}_A + F_{2,n}\mathbb{1}_{A^c} \uparrow H$, defining $F_1 := \overline{\lim} F_{1,n}, F_2 := \overline{\lim} F_{2,n} \in \mathscr{G}$, we have $H = F_1 \mathbb{1}_A + F_2 \mathbb{1}_{A^c}$. Actually, noticing A and A^c are a partition of Ω, then the pointwise convergence implies $F_{1,n}\mathbb{1}_A \uparrow F_1 \mathbb{1}_A$ and $F_{2,n}\mathbb{1}_A \uparrow F_2 \mathbb{1}_A$.

Based on the *monotonic class theorem of functions*, we know $\Lambda = \mathscr{G} \vee \{A\}$.□

QUESTION 6.70 *Suppose X and Y are two random variables on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$, and $A \in \mathscr{F}$. We have*

$$\mathbb{E}[X|\sigma(Y) \vee \{A\}] = \mathbb{E}(X|Y, A)\mathbb{1}_A + \mathbb{E}(X|Y, A^c)\mathbb{1}_{A^c},$$

where $\mathbb{E}(X|Y, A) := \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]}$.

Proof. We only need to prove $\mathbb{E}[X|\sigma(Y) \vee \{A\}]\mathbb{1}_A = \mathbb{E}(X|Y, A)\mathbb{1}_A$. Because $\mathbb{E}[X|\sigma(Y) \vee \{A\}] \in \sigma(Y) \vee \{A\}$ and $A \in \sigma(Y) \vee \{A\}$, we have $\mathbb{E}[X|\sigma(Y) \vee \{A\}]\mathbb{1}_A \in \sigma(Y) \vee \{A\}$. Similarly, since $\mathbb{E}(X|Y, A) \in \sigma(Y) \subset \sigma(Y) \vee \{A\}$, we also have $\mathbb{E}(X|Y, A)\mathbb{1}_A \in \sigma(Y) \vee \{A\}$. Next, we only need to show

$$\int \mathbb{E}[X|\sigma(Y) \vee \{A\}]\mathbb{1}_A H d\mathbb{P} = \int \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]}\mathbb{1}_A H d\mathbb{P},$$

for any $H \in \sigma(Y) \vee \{A\}$. Based on the previous QUESTION, we can find $H_1, H_2 \in \sigma(Y)$ such that $H = H_1\mathbb{1}_A + H_2\mathbb{1}_{A^c}$. Therefore

$$\begin{aligned}
&\int \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]}\mathbb{1}_A H d\mathbb{P} \\
=&\int \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]}\mathbb{1}_A (H_1\mathbb{1}_A + H_2\mathbb{1}_{A^c}) d\mathbb{P} \\
=&\int \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]} H_1\mathbb{1}_A d\mathbb{P}
\end{aligned}$$

$$
\begin{aligned}
&= \int \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]}H_1\mathbb{P}(A|Y)d\mathbb{P} \\
&= \int \mathbb{E}(X\mathbb{1}_A|Y)\mathbb{1}_{[\mathbb{P}(A|Y)>0]}H_1 d\mathbb{P} \\
&= \int X\mathbb{1}_A\mathbb{1}_{[\mathbb{P}(A|Y)>0]}H_1 d\mathbb{P} \\
&= \int X\mathbb{1}_A\mathbb{1}_{[\mathbb{P}(A|Y)>0]}H d\mathbb{P} \\
&= \int \mathbb{E}[X|\sigma(Y)\vee\{A\}]\mathbb{1}_A\mathbb{1}_{[\mathbb{P}(A|Y)>0]}H d\mathbb{P}.
\end{aligned}
$$

Next, we show $\{\mathbb{P}(A|Y)=0\}\subset A^c, \mathbb{P}$-a.s.. In fact,

$$
\begin{aligned}
\mathbb{P}(A\cap\{\mathbb{P}(A|Y)=0\}) &= \mathbb{E}[\mathbb{P}(A\cap\{\mathbb{P}(A|Y)=0\}|Y)] \\
&= \mathbb{E}[\mathbb{1}_{[\mathbb{P}(A|Y)=0]}\mathbb{P}(A|Y)] = 0,
\end{aligned}
$$

which means $\{\mathbb{P}(A|Y)=0\}\subset A^c, \mathbb{P}$-a.s.. So, $A\subset\{\mathbb{P}(A|Y)>0\}, \mathbb{P}$-a.s. and therefore

$$
\begin{aligned}
&\int \frac{\mathbb{E}(X\mathbb{1}_A|Y)}{\mathbb{P}(A|Y)}\mathbb{1}_{[\mathbb{P}(A|Y)>0]}\mathbb{1}_A H d\mathbb{P} \\
&= \int \mathbb{E}[X|\sigma(Y)\vee\{A\}]\mathbb{1}_A\mathbb{1}_{[\mathbb{P}(A|Y)>0]}H d\mathbb{P} \\
&= \int \mathbb{E}[X|\sigma(Y)\vee\{A\}]\mathbb{1}_A H d\mathbb{P}.
\end{aligned}
$$

□

QUESTION 6.71 *Suppose X is a random variable on $(\Omega,\mathscr{F},\mathbb{P})$. Given sub σ-field $\mathscr{G}\subset\mathscr{F}$, and measurable set $A\in\mathscr{F}$. The following conditional law of iterated expectation*

$$(*)\qquad \mathbb{P}[\mathbb{P}(X|\mathscr{G})|A] = \mathbb{P}(X|A)$$

holds if at least one of the following conditions holds:

(1) $X\in\mathscr{G}$.

(2) $A\in\mathscr{G}$.

(3) X and A are independent conditional on $\mathscr{G}$.

Proof. 1. If (1) holds, then $\mathbb{P}(X|\mathscr{G})=X$. So, $(*)$ naturally holds.

2. If (2) holds, we have

$$
\begin{aligned}
\mathbb{P}[\mathbb{P}(X|\mathscr{G})|A] &= \frac{\mathbb{P}[\mathbb{P}(X|\mathscr{G})\mathbb{1}_A]}{\mathbb{P}(A)} = \frac{\mathbb{P}[\mathbb{P}(X\mathbb{1}_A|\mathscr{G})]}{\mathbb{P}(A)} \\
&= \frac{\mathbb{P}(X\mathbb{1}_A)}{\mathbb{P}(A)} = \mathbb{P}(X|A).
\end{aligned}
$$

3. If (3) holds, we have

$$\begin{aligned}\mathbb{P}[\mathbb{P}(X|\mathscr{G})|A] &= \frac{\mathbb{P}[\mathbb{P}(X|\mathscr{G})\mathbb{1}_A]}{\mathbb{P}(A)} = \frac{\mathbb{P}\{\mathbb{P}[\mathbb{P}(X|\mathscr{G})\mathbb{1}_A|\mathscr{G}]\}}{\mathbb{P}(A)} \\ &= \frac{\mathbb{P}[\mathbb{P}(X|\mathscr{G})\mathbb{P}(A|\mathscr{G})]}{\mathbb{P}(A)} = \frac{\mathbb{P}[\mathbb{P}(X\mathbb{1}_A|\mathscr{G})]}{\mathbb{P}(A)} \\ &= \frac{\mathbb{P}(X\mathbb{1}_A)}{\mathbb{P}(A)} = \mathbb{P}(X|A).\end{aligned}$$

□

QUESTION 6.72 *Given random variables X, Y on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$, a measurable function $f \in \mathscr{F}$ and set $A \in \mathscr{F}$. Suppose A and X are independent conditional on Y. Then, A and X are independent conditional on $f(Y)$ if $\mathbb{P}(A|Y) \in \sigma(f(Y))$.*

Proof. A and X are independent conditional on Y is equivalent to

$$\mathbb{P}(A|X, Y) = \mathbb{P}(A|Y).$$

Since $\mathbb{P}(A|Y) \in \sigma(f(Y))$ and $\sigma(f(Y)) \subset \sigma(Y)$, we have

$$\begin{aligned}\mathbb{P}[A|X, f(Y)] &= \mathbb{P}[\mathbb{P}(A|X, Y)|X, f(Y)] \\ &= \mathbb{P}[\mathbb{P}(A|Y)|X, f(Y)] = \mathbb{P}(A|Y) \\ &= \mathbb{P}[\mathbb{P}(A|Y)|f(Y)] = \mathbb{P}[A|f(Y)].\end{aligned}$$

Thus, A and X are independent conditional on $f(Y)$. □

QUESTION 6.73 *Suppose $(X_n)_{n\geq 1}$ is a r.v. sequence on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and valued in $(E, \mathscr{E})$. Prove one part of the Pormanteau theorem: $\mathbb{P}f(X_n) \to \mathbb{P}f(X)$ for any $f \in C_b(E)$ iff $\underline{\lim}_{n\to\infty} \mathbb{P}f(X_n) \geq \mathbb{P}f(X)$ for any $f \in C_+(E)$, where $C_+(E)$ is the set of all the non-negative and continuous functions.*

Proof. (1) If $\underline{\lim}_{n\to\infty} \mathbb{P}f(X_n) \geq \mathbb{P}f(X)$ for any $f \in C_+(E)$, for any $f \in C_b(E)$, there exists $m, M \in \mathbb{R}$ such that $m \leq f \leq M$. So, $M - f \geq 0, f - m \geq 0$ and therefore

$$\begin{aligned} & \varliminf_{n\to\infty} \mathbb{P}[M - f(X_n)] \geq \mathbb{P}[M - f(X)] \\ \iff\quad & M - \varlimsup_{n\to\infty} \mathbb{P}f(X_n) \geq M - \mathbb{P}f(X) \\ \iff\quad & \varlimsup_{n\to\infty} \mathbb{P}f(X_n) \leq \mathbb{P}f(X).\end{aligned}$$

Similarly,

$$\begin{aligned} & \varliminf_{n\to\infty} \mathbb{P}[f(X_n) - m] \geq \mathbb{P}[f(X) - m] \\ \iff\quad & \varliminf_{n\to\infty} \mathbb{P}f(X_n) \geq \mathbb{P}f(X).\end{aligned}$$

Combine the above two equalities to obtain

$$Pf(X) \leq \varliminf_{n\to\infty} \mathbb{P}f(X_n) \leq \varlimsup_{n\to\infty} \mathbb{P}f(X_n) \leq \mathbb{P}f(X),$$

which means $\mathbb{P}f(X_n) \to \mathbb{P}f(X)$.

(2) If $\mathbb{P}f(X_n) \to \mathbb{P}f(X)$ for any $f \in C_b(E)$, for any $f \in C_+(E)$, we know $f \wedge M \in C_b(E)$, where $M > 0$. So,

$$\lim_{n\to\infty} \mathbb{P}[f(X_n) \wedge M] = \mathbb{P}[f(X) \wedge M].$$

Observe $f \geq f \wedge M$, therefore

$$\varliminf_{n\to\infty} \mathbb{P}f(X_n) \geq \lim_{n\to\infty} \mathbb{P}[f(X_n) \wedge M] = \mathbb{P}[f(X) \wedge M].$$

Let $M \to \infty$ and apply the monotone convergence theorem to obtain

$$\varliminf_{n\to\infty} \mathbb{P}f(X_n) \geq \mathbb{P}f(X).$$

□

QUESTION 6.74 *Suppose $(X_n)_{n\geq 1}$ is a non-negative r.v. sequence on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. If $X_n \xrightarrow{\mathrm{d}} X$, then X is also non-negative.*

Proof. Since X_n is non-negative, we have $\mathbb{P}(X_n < 0) = 0$. By the Portmanteau theorem, we know

$$0 = \varliminf_{n\to\infty} \mathbb{P}(X_n < 0) \geq \mathbb{P}(X < 0).$$

This implies $\mathbb{P}(X < 0) = 0$. □

QUESTION 6.75 *Suppose $(X_n)_{n\geq 1}$ and X are r.v.s in the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. If $X_n \xrightarrow{L^r} X, n \to \infty$, then $X_n^r \xrightarrow{L^1} X^r, n \to \infty$.*

Proof. Since $X_n \xrightarrow{L^r} X, n \to \infty$, we know that $X_n \xrightarrow{\mathbb{P}} X, n \to \infty$ and

$$\int_A |X_n|^r d\mathbb{P} \text{ is integral absolutely continuous.}$$

Since $x \mapsto |x|^r$ is a continuous function, $|X_n|^r \xrightarrow{\mathbb{P}} |X|^r, n \to \infty$. Moreover,

$$\int_A ||X_n|^r| d\mathbb{P} = \int_A |X_n|^r d\mathbb{P} \text{ is integral absolutely continuous.}$$

So, $X_n^r \xrightarrow{L^1} X^r, n \to \infty$. □

QUESTION 6.76 *If $(X_n)_{n\geq 1}$ and $(Y_n)_{n\geq 1}$ are two tight r.v. sequences in the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and valued in $(E, \mathscr{B}(E))$, then $\{(X_n, Y_n)\}_{n\geq 1}$ is also a tight r.v. sequences in the product probability space $(\Omega^2, \mathscr{F}^2, \mathbb{P}^2)$.*

Proof. For any $\varepsilon > 0$, there exist compact sets $M_\varepsilon, V_\varepsilon \subset E$ such that

$$\sup_{n\geq 1} \mathbb{P}(X_n \in M_\varepsilon^c) < \varepsilon, \quad \sup_{n\geq 1} \mathbb{P}(X_n \in V_\varepsilon^c) < \varepsilon.$$

Therefore

$$\begin{aligned}\sup_{n\geq 1} \mathbb{P}[(X_n, Y_n) \in (M_\varepsilon \times V_\varepsilon)^c] &= \sup_{n\geq 1} \mathbb{P}(X_n \in M_\varepsilon^c \text{ or } Y_n \in V_\varepsilon^c) \\ &\leq \sup_{n\geq 1} \mathbb{P}(X_n \in M_\varepsilon^c) + \sup_{n\geq 1} \mathbb{P}(X_n \in V_\varepsilon^c) < 2\varepsilon,\end{aligned}$$

where $M_\varepsilon \times V_\varepsilon \subset E^2$ is also compact in the topology of product topology. So, $\{(X_n, Y_n)\}_{n\geq 1}$ is also a compact r.v. sequences in the product probability space $(\Omega^2, \mathscr{F}^2, \mathbb{P}^2)$. □

QUESTION 6.77 *Suppose μ, ν are two measures on the measurable space $(\Omega, \mathscr{F})$ satisfying $\mu \ll \nu$, and $\lambda, \#$ are two measures on the measurable space $(E, \mathscr{E})$ satisfying $\lambda \ll \#$. Then $\mu \times \lambda \ll \nu \times \#$ on the measurable space $(\Omega \times E, \mathscr{F} \times \mathscr{E})$.*

Proof. For any $A \in \mathscr{F} \times \mathscr{E}$ such that $(\nu \times \#)(A) = 0$, we have

$$\int_E \nu(A_x)\#(dx) = 0 \Longrightarrow \nu(A_x) = 0, \quad \# \text{ - a.s.},$$

where A_x is the section of A at x. Since $\mu \ll \nu$ and $\lambda \ll \#$, it follows from the above expression that

$$\mu(A_x) = 0, \quad \lambda \text{ - a.s..}$$

Therefore

$$(\mu \times \lambda)(A) = \int_E \mu(A_x)\lambda(dx) = 0,$$

which implies $\mu \times \lambda \ll \nu \times \#$ on the measurable space $(\Omega \times E, \mathscr{F} \times \mathscr{E})$. □

QUESTION 6.78 *Suppose $X_n \xrightarrow{\mathbb{P}} X$. There exists a sequence $\{\varepsilon_n\}_{n\geq 1}$ satisfying $\varepsilon_n \downarrow 0, n \to \infty$, such that $\mathbb{P}(|X_n - X| > \varepsilon_n) \to 0, n \to \infty$.*

Proof. Since $X_n \xrightarrow{\mathbb{P}} X$, we have

$$\mathbb{P}(|X_n - X| > \varepsilon) \to 0, n \to \infty, \quad \text{for any } \varepsilon > 0.$$

Given any decreasing sequence $\gamma_n \downarrow 0, n \to \infty$, we can construct another decreasing sequence $\{\varepsilon_n\}_{n\geq 1}$ as follows:

- Let $\varepsilon_1 := \gamma_1$ and $N_1 := 1$.
- For γ_k, where $k \geq 2$, since $\mathbb{P}(|X_n - X| > \gamma_k) \to 0, n \to \infty$, there exists $\ell_k \in \mathbb{Z}_+$ such that

$$\mathbb{P}(|X_n - X| > \gamma_k) \leq 1/k, \quad \forall n \geq \ell_k.$$

  Define $N_k := \max\{N_{k-1}, \ell_k\} + 1$ and $\varepsilon_n := \gamma_{k-1}, \forall N_{k-1} \leq n < N_k - 1$.

Based on this construction, the sequence $\{\varepsilon_n\}_{n\geq 1}$ satisfies $\varepsilon_n \to 0, n \to \infty$. Moreover, for any $n \in \mathbb{Z}_+$, there exists $k(n) \in \mathbb{Z}_+$ such that $N_{k(n)} \leq n < N_{k(n)} - 1$ and

$$\mathbb{P}(|X_n - X| > \varepsilon_n) \leq 1/k(n). \tag{5}$$

Noting the integer sequence $\{N_k\}$ is strictly increasing to ∞, we know $k(n) \uparrow \infty, n \to \infty$. Therefore, letting $n \to \infty$ in (5), we will get the desired result. □

QUESTION 6.79 *Suppose X_n and Y_n are two r.v.s on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Then $X_n \leq Y_n + o_p(1)$ is equivalent to*

$$\mathbb{P}(X_n > Y_n + \varepsilon) \to 0, \quad n \to \infty, \quad \forall \varepsilon > 0. \tag{$*$}$$

Proof. Suppose $X_n \leq Y_n + o_p(1)$, there exists $Z_n = o_p(1)$, such that $X_n \leq Y_n + Z_n$. Then, for any $\varepsilon > 0$,

$$\mathbb{P}(X_n > Y_n + \varepsilon) \leq \mathbb{P}(Z_n > \varepsilon) \leq \mathbb{P}(|Z_n| > \varepsilon) \to 0, \quad n \to \infty.$$

On the contrary, if $(*)$ holds, then define $Z_n := (X_n - Y_n)^+$, we have

$$X_n = Y_n + (X_n - Y_n) \leq Y_n + (X_n - Y_n)^+ = Y_n + Z_n$$

and

$$\begin{aligned}\mathbb{P}(|Z_n| > \varepsilon) &= \mathbb{P}[(X_n - Y_n)^+ > \varepsilon] = \mathbb{P}(X_n - Y_n > \varepsilon) \\ &= \mathbb{P}(X_n > Y_n + \varepsilon) \to 0, \quad n \to \infty, \quad \forall \varepsilon > 0.\end{aligned}$$

So, $Z_n = o_p(1)$ and $X_n \leq Y_n + Z_n$. □

QUESTION 6.80 *Suppose X_n and Y_n are two r.v.s on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. If $\mathbb{P}(X_n \leq Y_n) \to 1, n \to \infty$, then $X_n \leq Y_n + o_p(1)$.*

Proof. Define $Z_n = (X_n - Y_n)^+$, then

$$\begin{aligned}\mathbb{P}(|Z_n| > \varepsilon) &= \mathbb{P}[(X_n - Y_n)^+ > \varepsilon] = \mathbb{P}(X_n - Y_n > \varepsilon) \\ &= \mathbb{P}(X_n > Y_n + \varepsilon) \leq \mathbb{P}(X_n > Y_n) \\ &= 1 - \mathbb{P}(X_n \leq Y_n) \to 0, \quad n \to \infty.\end{aligned}$$

Thus $Z_n = o_p(1)$ and obviously $X_n \leq Y_n + Z_n$. □

QUESTION 6.81 *Suppose X_n, Y_n and Z_n are three positive r.v.s on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Suppose $X_n = O_p(Y_n) + O_p(Z_n)$, then $X_n = O_p(Y_n + Z_n)$.*

Proof. For any $\forall \varepsilon > 0$, there exists $M_1^\varepsilon > 0$ and $M_2^\varepsilon > 0$ such that

$$\sup_{n\geq 1} \mathbb{P}\left(\frac{O_p(Y_n)}{Y_n} > M_1^\varepsilon\right) < \varepsilon/2 \quad \text{and} \quad \sup_{n\geq 1} \mathbb{P}\left(\frac{O_p(Z_n)}{Z_n} > M_2^\varepsilon\right) < \varepsilon/2.$$

Define $M := 2(M_1^\varepsilon \vee M_2^\varepsilon) > 0$, then

$$\begin{aligned}
\mathbb{P}\left(\frac{X_n}{Y_n + Z_n} > M\right) &= \mathbb{P}\left(\frac{O_p(Y_n) + O_p(Z_n)}{Y_n + Z_n} > M\right) \\
&= \mathbb{P}\left(\frac{O_p(Y_n)}{Y_n + Z_n} + \frac{O_p(Z_n)}{Y_n + Z_n} > M\right) \\
&\leq \mathbb{P}\left(\frac{O_p(Y_n)}{Y_n} + \frac{O_p(Z_n)}{Z_n} > M\right) \\
&\leq \mathbb{P}\left(\frac{O_p(Y_n)}{Y_n} > \frac{M}{2}\right) + \mathbb{P}\left(\frac{O_p(Z_n)}{Z_n} > \frac{M}{2}\right) < \varepsilon.
\end{aligned}$$

So, $X_n = O_p(Y_n + Z_n)$ □

QUESTION 6.82 *Suppose X, Y and T are three r.v.s on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and valued on the measurable space $(E, \mathscr{E})$. If $Y \perp T|X$ and $T \perp X$, then $Y \perp X$.*

Proof. Since $Y \perp T|X$, for any $f, g \in \mathrm{b}\mathscr{E}$,

$$\mathbb{E}[f(Y)g(T)|X] = \mathbb{E}[f(Y)|X]\mathbb{E}[g(T)|X] = \mathbb{E}[f(Y)|X]\mathbb{E}[g(T)].$$

Taking the expectation on both sides, we will get

$$\mathbb{E}[f(Y)g(T)] = \mathbb{E}[f(Y)]\mathbb{E}[g(T)],$$

which implies $Y \perp X$. □

QUESTION 6.83 *Let X and Y be independent and identically distributed random variables with $\mathbb{E}|X| < \infty$. Show that $\mathbb{E}[|X + Y|] \geq \mathbb{E}[|X - Y|]$, where equality holds if and only if the distribution of X is symmetric around 0.*

Proof. The proof uses the Dirichlet integral identity

$$\int_{-\infty}^{+\infty} \frac{1 - \cos(ax)}{x^2}\, dx = \pi|a|.$$

Therefore, by Tonelli's theorem,

$$\begin{aligned}
&\mathbb{E}\big[|X+Y|-|X-Y|\big] \\
= \;& \frac{1}{\pi}\,\mathbb{E}\left[\int_{-\infty}^{+\infty}\frac{1-\cos\big((X+Y)t\big)}{t^2}\,dt \;-\; \int_{-\infty}^{+\infty}\frac{1-\cos\big((X-Y)t\big)}{t^2}\,dt\right] \\
= \;& \frac{1}{\pi}\int_{-\infty}^{+\infty}\mathbb{E}\left[\frac{\cos\big((X-Y)t\big)-\cos\big((X+Y)t\big)}{t^2}\right]dt.
\end{aligned}$$

Using the trigonometric identity $\cos u - \cos v = -2\sin\frac{u+v}{2}\,\sin\frac{u-v}{2}$ with $u = (X-Y)t$ and $v = (X+Y)t$, we obtain

$$\cos\big((X-Y)t\big)-\cos\big((X+Y)t\big) = 2\sin(Xt)\sin(Yt).$$

Hence, by independence and identical distribution of X and Y,

$$\begin{aligned}
\mathbb{E}\big[|X+Y|-|X-Y|\big] &= \frac{2}{\pi}\int_{-\infty}^{+\infty}\mathbb{E}\left[\frac{\sin(Xt)\sin(Yt)}{t^2}\right]dt \\
&= \frac{2}{\pi}\int_{-\infty}^{+\infty}\frac{1}{t^2}\,\mathbb{E}[\sin(Xt)]\,\mathbb{E}[\sin(Yt)]\,dt \\
&= \frac{2}{\pi}\int_{-\infty}^{+\infty}\left(\frac{\mathbb{E}[\sin(Xt)]}{t}\right)^2 dt \;\geq\; 0.
\end{aligned}$$

This proves $\mathbb{E}[|X+Y|] \geq \mathbb{E}[|X-Y|]$. If equality holds, then necessarily $\mathbb{E}[\sin(Xt)] = 0$ for all $t \in \mathbb{R}$. Since the characteristic function is $\varphi_X(t) = \mathbb{E}[e^{itX}] = \mathbb{E}[\cos(tX)] + i\,\mathbb{E}[\sin(tX)]$, this means φ_X is real for all t, hence the distribution of X is symmetric about 0. The converse is immediate. □

QUESTION 6.84 (Equivalences for stochastic boundedness) *Let $\{X_n\}_{n\geq 1}$ be random variables taking values in $\mathbb{R}^d$ (for any fixed norm $\|\cdot\|$ on $\mathbb{R}^d$). The following statements are equivalent:*

(1) For every $\varepsilon > 0$ there exists $M < \infty$ such that

$$\sup_{n\geq 1}\mathbb{P}\big(\|X_n\| > M\big) \leq \varepsilon.$$

Equivalently, $\lim_{M\to\infty}\sup_n \mathbb{P}(\|X_n\| > M) = 0$.

(2) For every deterministic sequence $\{\ell_n\}_{n\geq 1}$ with $\ell_n \to \infty$,

$$\mathbb{P}\big(\|X_n\| > \ell_n\big) \;\longrightarrow\; 0.$$

(3) For every $\varepsilon > 0$ there exists $M < \infty$ such that

$$\limsup_{n\to\infty}\mathbb{P}\big(\|X_n\| > M\big) \leq \varepsilon.$$

In particular, these are all equivalent to saying that $\{X_n\}$ is bounded in probability (stochastically bounded, i.e. $X_n = O_{\mathbb{P}}(1)$).

Proof. **(1) ⇒ (2).** Fix $\varepsilon > 0$ and a sequence $\ell_n \to \infty$. By (1), choose $M < \infty$ with

$$\sup_n \mathbb{P}(\|X_n\| > M) \le \varepsilon.$$

For all large n, $\ell_n \ge M$, hence

$$\mathbb{P}(\|X_n\| > \ell_n) \le \mathbb{P}(\|X_n\| > M) \le \varepsilon,$$

so $\mathbb{P}(\|X_n\| > \ell_n) \to 0$.

(2) ⇒ (1). Argue by contradiction. Suppose (1) fails. Then there exists $\varepsilon_0 > 0$ such that

$$\lim_{M\to\infty} \sup_n \mathbb{P}(\|X_n\| > M) \;\ge\; \varepsilon_0.$$

Define $g_N(M) := \sup_{n\ge N} \mathbb{P}(\|X_n\| > M)$ for $N \in \mathbb{N}$. If for some N we had $\lim_{M\to\infty} g_N(M) = 0$, then the tail family $\{X_n : n \ge N\}$ would be tight; combined with the finiteness of $\{X_1, \dots, X_{N-1}\}$ this would yield (1), a contradiction. Hence for every N,

$$\limsup_{M\to\infty} g_N(M) \;\ge\; \varepsilon_0.$$

Let $\eta := \varepsilon_0/2$. We now construct strictly increasing sequences $M_k \uparrow \infty$ and $n_k \uparrow \infty$ recursively. Set $N_1 := 1$. Choose M_1 large with $g_{N_1}(M_1) \ge \eta$, and then pick $n_1 \ge N_1$ such that $\mathbb{P}(\|X_{n_1}\| > M_1) \ge \eta$. Given (n_{k-1}, M_{k-1}), set $N_k := n_{k-1}+1$. Choose $M_k > M_{k-1}$ with $g_{N_k}(M_k) \ge \eta$, and then pick $n_k \ge N_k$ so that $\mathbb{P}(\|X_{n_k}\| > M_k) \ge \eta$.

Define a deterministic, piecewise-constant sequence

$$\ell_n := \begin{cases} M_1, & n < n_1, \\ M_k, & n_k \le n < n_{k+1} \quad (k \ge 1). \end{cases}$$

Then $\ell_n \to \infty$ and, for all k,

$$\mathbb{P}(\|X_{n_k}\| > \ell_{n_k}) = \mathbb{P}(\|X_{n_k}\| > M_k) \;\ge\; \eta,$$

which contradicts (2). Thus (1) must hold.

(1) ⇒ (3). Immediate, since for any M,

$$\limsup_{n\to\infty} \mathbb{P}(\|X_n\| > M) \;\le\; \sup_n \mathbb{P}(\|X_n\| > M).$$

(3) ⇒ (1). Fix $\varepsilon > 0$. By (3), choose $M_\infty < \infty$ with

$$\limsup_{n\to\infty} \mathbb{P}(\|X_n\| > M_\infty) \le \varepsilon/2,$$

so there exists N such that

$$\sup_{n\ge N} \mathbb{P}(\|X_n\| > M_\infty) \le \varepsilon/2.$$

For each fixed $n < N$, the sets $\{\|X_n\| > t\}$ decrease to the empty set as $t \to \infty$, hence $\mathbb{P}(\|X_n\| > t) \downarrow 0$. Thus we can choose $M_0 < \infty$ with

$$\max_{1\le n<N} \mathbb{P}(\|X_n\| > M_0) \le \varepsilon/2.$$

Let $M := \max\{M_\infty, M_0\}$. Then

$$\begin{aligned}\sup_n \mathbb{P}(\|X_n\| > M) &= \max\Big\{\max_{n<N} \mathbb{P}(\|X_n\| > M),\ \sup_{n\ge N} \mathbb{P}(\|X_n\| > M)\Big\}\\ &\le \varepsilon/2 + \varepsilon/2 = \varepsilon,\end{aligned}$$

which proves (1).

Combining the implications, we obtain the equivalence of (1), (2), and (3).□

QUESTION 6.85 (Uniform integrability vs. tightness under moment bounds) *Let $(X_n)_{n\ge1}$ be real-valued random variables on a common probability space. Assume that for some $r > 0$,*

$$M_r := \sup_{n\ge1} \mathbb{E}|X_n|^r < \infty.$$

Prove that:

(a) If $r > 0$, then (X_n) is tight.

(b) If $r > 1$, then (X_n) is uniformly integrable.

(c) Uniform integrability implies tightness.

(d) The condition $r = 1$ alone does not *imply uniform integrability: construct a counterexample.*

Proof. **(a)** $r > 0 \Rightarrow$ **tight.** Applying Markov inquality with $r > 0$ and taking the supremum over n gives

$$\sup_n \mathbb{P}(|X_n| > K) \le \frac{\sup_n \mathbb{E}|X_n|^r}{K^r} \le \frac{M_r}{K^r} \xrightarrow[K\to\infty]{} 0,$$

which is the definition of tightness.

(b) $r > 1 \Rightarrow$ **UI.** Let $r > 1$ and $r^* := \frac{r}{r-1}$ be the Hölder conjugate. For each n and $K > 0$,

$$\mathbb{E}\big[|X_n|\mathbb{1}_{\{|X_n|>K\}}\big] \le \big(\mathbb{E}|X_n|^r\big)^{1/r}\big(\mathbb{P}(|X_n|>K)\big)^{1/r^*} \le M_r^{1/r}\left(\frac{M_r}{K^r}\right)^{1/r^*} = \frac{M_r}{K^{r-1}}.$$

Taking $\sup_n$ and then $K \to \infty$ yields

$$\lim_{K\to\infty} \sup_n \mathbb{E}\big[|X_n|\mathbb{1}_{\{|X_n|>K\}}\big] = 0,$$

so (X_n) is uniformly integrable.

Alternative (de la Vallée Poussin). For $r > 1$, the convex function $\Phi(t) = t^r$ satisfies $\Phi(t)/t = t^{r-1} \to \infty$ as $t \to \infty$, and

$$\sup_n \mathbb{E}\Phi(|X_n|) = \sup_n \mathbb{E}|X_n|^r \le M_r < \infty,$$

hence (X_n) is UI.

(c) UI $\Rightarrow$ tight. For any $K > 0$ and any n,

$$K\,\mathbb{1}_{\{|X_n|>K\}} \le |X_n|\,\mathbb{1}_{\{|X_n|>K\}},$$

so taking expectations and then $\sup_n$,

$$K \sup_n \mathbb{P}(|X_n| > K) \;\le\; \sup_n \mathbb{E}\big[|X_n|\mathbb{1}_{\{|X_n|>K\}}\big].$$

Letting $K \to \infty$ and using UI, the right-hand side tends to 0, hence $\sup_n \mathbb{P}(|X_n| > K) \to 0$, i.e. tightness.

(d) Counterexample for $r = 1$. Let $X_n := n\,\mathbb{1}_{A_n}$ where $\mathbb{P}(A_n) = 1/n$ (the events A_n can be arbitrary). Then

$$\mathbb{E}|X_n| = n \cdot \mathbb{P}(A_n) = 1 \quad \text{for all } n,$$

but for any $K \ge 0$,

$$\sup_n \mathbb{P}(|X_n| > K) = \sup_{n>K} \frac{1}{n} = \frac{1}{\lfloor K \rfloor + 1} \xrightarrow[K\to\infty]{} 0,$$

so (X_n) is tight. However,

$$\sup_n \mathbb{E}\big[|X_n|\mathbb{1}_{\{|X_n|>K\}}\big] = \sup_{n>K} n \cdot \mathbb{P}(A_n) = \sup_{n>K} 1 = 1,$$

which does not converge to 0 as $K \to \infty$. Hence (X_n) is *not* uniformly integrable. This shows that $r = 1$ alone is insufficient for UI. $\square$

QUESTION 6.86 (Weak convergence implies $O_P(1)$ (tightness)) *Let $X_n \Rightarrow X$ in $\mathbb{R}^d$. Prove that (X_n) is tight, i.e., for every $\varepsilon > 0$ there exists $M < \infty$ such that*

$$\sup_{n\geq 1} \mathbb{P}(\|X_n\| > M) \leq \varepsilon.$$

Equivalently, $X_n = O_P(1)$.

Proof. For $M > 0$, define the closed set

$$F_M := \{x \in \mathbb{R}^d : \|x\| \geq M\}.$$

By the Portmanteau theorem,

$$\limsup_{n\to\infty} \mathbb{P}(X_n \in F_M) \leq \mathbb{P}(X \in F_M) = \mathbb{P}(\|X\| \geq M).$$

Since $\|X\| < \infty$ almost surely, we have $\mathbb{P}(\|X\| \geq M) \to 0$ as $M \to \infty$. Hence, given any $\varepsilon > 0$, choose M large enough so that $\mathbb{P}(\|X\| \geq M) \leq \varepsilon/2$. Then there exists N such that for all $n \geq N$,

$$\mathbb{P}(\|X_n\| \geq M) = \mathbb{P}(X_n \in F_M) \leq \mathbb{P}(X \in F_M) + \varepsilon/2 \leq \varepsilon.$$

Therefore $\sup_{n\geq N} \mathbb{P}(\|X_n\| > M) \leq \varepsilon$, which is tightness. In particular, $X_n = O_P(1)$. □

QUESTION 6.87 *Let X and Y be (marginally) normal random variables.*

(a) Must the sum $X + Y$ be normal?

(b) Can $\mathrm{Var}(X + Y)$ be smaller than both $\mathrm{Var}(X)$ and $\mathrm{Var}(Y)$?

Proof. (a) Not necessarily. It is not enough that X and Y are each normal; dependence can make $X + Y$ non-normal. Construct the following counterexample: let $X \sim N(0, 1)$ and let $Z \in \{-1, +1\}$ be independent of X with $\mathbb{P}(Z = 1) = \mathbb{P}(Z = -1) = \frac{1}{2}$. Define $Y := ZX$. Then $Y \mid \{Z = 1\} \sim N(0, 1)$ and $Y \mid \{Z = -1\} \sim N(0, 1)$, hence $Y \sim N(0, 1)$ marginally. However,

$$X + Y = \begin{cases} 2X, & Z = 1 \ \text{(prob. } 1/2), \\ 0, & Z = -1 \ \text{(prob. } 1/2), \end{cases}$$

so $X + Y$ has an atom at 0 of mass $1/2$ and is therefore not normal.

On the other hand, a sufficient (indeed, necessary and sufficient) condition for any linear combination $aX+bY$ to be normal is that (X, Y) be jointly normal. In particular, if X and Y are independent normals, then $X + Y$ is normal with

$$\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y), \qquad \mathrm{Var}(X + Y) = \mathrm{Var}(X) + \mathrm{Var}(Y),$$

which can be verified, for instance, from the product of moment generating functions.

(b) Yes, it can be smaller under negative correlation (but not if X and Y are independent). In general,

$$\mathrm{Var}(X+Y) = \mathrm{Var}(X) + \mathrm{Var}(Y) + 2\,\mathrm{Cov}(X,Y).$$

If X and Y are independent, then $\mathrm{Cov}(X,Y) = 0$ and hence

$$\mathrm{Var}(X+Y) = \mathrm{Var}(X) + \mathrm{Var}(Y) \geq \max\{\mathrm{Var}(X), \mathrm{Var}(Y)\},$$

with equality only if one variance is zero.

If X and Y are (say, jointly normal and) negatively correlated with correlation $\rho \in [-1, 0)$, writing $\mathrm{Var}(X) = \sigma_X^2$, $\mathrm{Var}(Y) = \sigma_Y^2$, we have

$$\mathrm{Var}(X+Y) = \sigma_X^2 + \sigma_Y^2 + 2\rho\,\sigma_X\sigma_Y.$$

Choosing ρ sufficiently negative can make $\mathrm{Var}(X+Y)$ smaller than both σ_X^2 and σ_Y^2. For example, with $\sigma_X^2 = \sigma_Y^2 = 1$ and $\rho = -0.9$,

$$\mathrm{Var}(X+Y) = 1 + 1 + 2(-0.9) = 0.2 < 1.$$

In the extreme case $\rho = -1$ and $\sigma_X = \sigma_Y$, we have $Y = -X$ almost surely and hence $\mathrm{Var}(X+Y) = 0$. □

Proposition 6.1 (Conditional Strong Law of Large Numbers) *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $\mathcal{G} \subset \mathcal{F}$ be a sub-σ-algebra. Suppose $(Y_i)_{i\geq 1}$ are integrable random variables such that* conditionally on $\mathcal{G}$ *they are i.i.d., and*

$$\mathbb{E}[\,|Y_1|\,|\,\mathcal{G}] < \infty \quad \text{a.s.}$$

Then

$$\frac{1}{n}\sum_{i=1}^{n} Y_i \xrightarrow{\text{a.s.}} \mathbb{E}[Y_1\,|\,\mathcal{G}]\,.$$

Proof. (Disintegration + Kolmogorov SLLN.) Assume a regular conditional probability $\mathbb{P}(\cdot\,|\,\mathcal{G})(\omega)$ exists (e.g., on standard Borel spaces). For ω in a full-measure set, define $\mathbb{P}_\omega(\cdot) := \mathbb{P}(\cdot\,|\,\mathcal{G})(\omega)$ and

$$m(\omega) := \mathbb{E}[Y_1\,|\,\mathcal{G}]\,(\omega).$$

By the hypothesis "i.i.d. given $\mathcal{G}$", under $\mathbb{P}_\omega$ the sequence (Y_i) is i.i.d. with mean $m(\omega)$ and is integrable. Hence, by Kolmogorov's SLLN,

$$\frac{1}{n}\sum_{i=1}^{n} Y_i \xrightarrow{\text{a.s. under } \mathbb{P}_\omega} m(\omega).$$

Therefore $\mathbb{P}_\omega(\lim_{n\to\infty} n^{-1}\sum_{i=1}^n Y_i = m(\omega)) = 1$ for a.e. ω. Integrating both sides (Fubini/Tonelli) yields

$$\mathbb{P}\left(\lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^n Y_i = \mathbb{E}[Y_1 \,|\, \mathcal{G}]\right) = 1,$$

which is the claim. □

Example 6.1 (Random environment: $Y_i \mid \Lambda \overset{iid}{\sim} N(\Lambda, 1)$) Let Λ be a real random variable with $\mathbb{E}|\Lambda| < \infty$. Conditional on Λ, let $(Y_i)_{i\geq 1}$ be i.i.d. with distribution $N(\Lambda, 1)$. Equivalently, write

$$Y_i = \Lambda + \varepsilon_i, \qquad \varepsilon_i \overset{iid}{\sim} N(0,1), \;\; \varepsilon_i \text{ independent of } \Lambda.$$

So, Y_i is not independent if $\Lambda \neq \text{const.}$ $\mathbb{P}$-a.s.. And,

$$\frac{1}{n}\sum_{i=1}^n Y_i \xrightarrow{\text{a.s.}} \Lambda.$$

Proof. By construction, given Λ the Y_i are independent and $Y_i \mid \Lambda \sim N(\Lambda, 1)$, so they are i.i.d. given $\mathcal{G} = \sigma(\Lambda)$. Unconditionally, for $i \neq j$,

$$\text{Cov}(Y_i, Y_j) = \mathbb{E}[\text{Cov}(Y_i, Y_j \,|\, \Lambda)] + \text{Cov}(\mathbb{E}[Y_i \mid \Lambda], \mathbb{E}[Y_j \mid \Lambda]) = 0 + \text{Var}(\Lambda).$$

If $\text{Var}(\Lambda) > 0$, then $\text{Cov}(Y_i, Y_j) > 0$ and hence Y_i, Y_j cannot be independent.

For the last statement,

- Proof 1 (decomposition). Write $\bar{Y}_n = \frac{1}{n}\sum_{i=1}^n Y_i = \Lambda + \bar{\varepsilon}_n$ with $\bar{\varepsilon}_n = \frac{1}{n}\sum_{i=1}^n \varepsilon_i$. By Kolmogorov's SLLN, $\bar{\varepsilon}_n \to 0$ almost surely (indeed, both conditionally on Λ and unconditionally). Hence $\bar{Y}_n \to \Lambda$ almost surely.

- Proof 2 (conditional SLLN). Let $\mathcal{G} = \sigma(\Lambda)$. Then Y_i are i.i.d. given $\mathcal{G}$ with

$$\mathbb{E}[Y_1 \mid \mathcal{G}] = \Lambda, \qquad \mathbb{E}[\,|Y_1| \,|\, \mathcal{G}] < \infty.$$

  Apply the Conditional SLLN (the proposition) to deduce $\bar{Y}_n \to \mathbb{E}[Y_1 \mid \mathcal{G}] = \Lambda$ almost surely. □

QUESTION 6.88 *Suppose X is a random variable in the probability space $(\Omega, \mathscr{F}, \mathbb{P})$, $\mathscr{A} \subset \mathscr{F}$ is a sub σ-algebra, and $X \in L^r$ for any $r \in (0, \infty)$. Then $||\mathbb{E}(X|\mathscr{A})||_r \leq ||X||_r$. When $r = 2$, the inequality is the well-known* triangle inequality.

Proof. The Hölder's inequality yields

$$\begin{aligned} ||\mathbb{E}(X|\mathscr{A})||_r &= \mathbb{E}[|\mathbb{E}(X|\mathscr{A})|^r]^{1/r} \leq \mathbb{E}[\mathbb{E}(|X||\mathscr{A})^r]^{1/r} \\ &\leq \mathbb{E}[\mathbb{E}(|X|^r|\mathscr{A})]^{1/r} = \mathbb{E}(|X|^r)^{1/r} = ||X||_r. \end{aligned}$$

□

QUESTION 6.89 *Let (X_n) and (Y_n) be $\mathbb{R}^d$-valued random variables. If*

$$Y_n - X_n \xrightarrow{\mathbb{P}} 0 \quad \textit{and} \quad X_n \xrightarrow{\mathrm{d}} X,$$

then

$$Y_n \xrightarrow{\mathrm{d}} X.$$

Proof. By the Portmanteau lemma, it is enough to show that for every bounded Lipschitz function $f : \mathbb{R} \to \mathbb{R}$,

$$\mathbb{E}[f(Y_n)] \to \mathbb{E}[f(X)].$$

Fix such an f. There exist $M < \infty$ and $K < \infty$ such that $|f(u)| \leq M$ for all u and

$$|f(u) - f(v)| \leq K\,|u - v| \quad \text{for all } u, v \in \mathbb{R}.$$

For any $\varepsilon > 0$,

$$\begin{aligned}
&|\mathbb{E}[f(Y_n)] - \mathbb{E}[f(X_n)]| \leq \mathbb{E}[\,|f(Y_n) - f(X_n)|\,] \\
=&\mathbb{E}[\,|f(Y_n) - f(X_n)|\,\mathbb{1}_{[|Y_n - X_n| \leq \varepsilon]}] + \mathbb{E}[\,|f(Y_n) - f(X_n)|\,\mathbb{1}\{|Y_n - X_n| > \varepsilon\}] \\
\leq& K\,\varepsilon + 2M\,\mathbb{P}(|Y_n - X_n| > \varepsilon).
\end{aligned}$$

Hence, by the triangle inequality,

$$\begin{aligned}
|\mathbb{E}[f(Y_n)] - \mathbb{E}[f(X)]| &\leq |\mathbb{E}[f(Y_n)] - \mathbb{E}[f(X_n)]| + |\mathbb{E}[f(X_n)] - \mathbb{E}[f(X)]| \\
&\leq K\varepsilon + 2M\,\mathbb{P}(|Y_n - X_n| > \varepsilon) + |\mathbb{E}[f(X_n)] - \mathbb{E}[f(X)]|.
\end{aligned}$$

Taking $n \to \infty$, the second term vanishes because $Y_n - X_n \xrightarrow{\mathbb{P}} 0$, and the third term vanishes by the Portmanteau lemma since $X_n \xrightarrow{\mathrm{d}} X$. Therefore,

$$\limsup_{n\to\infty} \Big|\mathbb{E}[f(Y_n)] - \mathbb{E}[f(X)]\Big| \leq K\varepsilon.$$

Because $\varepsilon > 0$ was arbitrary, the limit is 0, so $\mathbb{E}[f(Y_n)] \to \mathbb{E}[f(X)]$ for all bounded Lipschitz f. Another application of the Portmanteau lemma yields $Y_n \xrightarrow{\mathrm{d}} X$, as claimed. □

Theorem 6.1 *Let (X_n) and (Y_n) be $\mathbb{R}^d$-valued random variables. Suppose $X_n \xrightarrow{\mathrm{d}} X$ and $Y_n \xrightarrow{\mathbb{P}} c$, where c is a constant. Then*

$$(X_n,\, Y_n) \xrightarrow{\mathrm{d}} (X,\, c).$$

Proof. Show that $(X_n,\, c) \xrightarrow{\mathrm{d}} (X,\, c)$. By the portmanteau lemma, it suffices to verify that for every bounded continuous $f : \mathbb{R}^2 \to \mathbb{R}$,

$$\mathbb{E}[f(X_n,\, c)] \longrightarrow \mathbb{E}[f(X,\, c)]\,.$$

Fix such f and define the bounded continuous function $g : \mathbb{R} \to \mathbb{R}$ by $g(x) := f(x,\, c)$. Since $X_n \xrightarrow{\mathrm{d}} X$, another application of the portmanteau lemma yields

$$\mathbb{E}[f(X_n,\, c)] = \mathbb{E}[g(X_n)] \longrightarrow \mathbb{E}[g(X)] = \mathbb{E}[f(X,\, c)]\,,$$

Therefore $(X_n,\, c) \xrightarrow{\mathrm{d}} (X,\, c)$. Since

$$\|(X_n,\, Y_n) - (X_n,\, c)\| \asymp \|Y_n - c\| \xrightarrow{\mathbb{P}} 0,$$

applying the previous QUESTION, we have

$$(X_n,\, Y_n) \xrightarrow{\mathrm{d}} (X,\, c).$$

□

§ 7 Stochastic Processes

QUESTION 7.1 *Suppose $(X_n)_{n\geq 0}$ is a Markov chain. If the state space E has at least two states, then the following statement is not true:*

$$\mathbb{P}(X_{n+1} = j | X_n \in A, X_{n-1} = i_{n-1}, \cdots, X_0 = i_0) = \mathbb{P}(X_{n+1} = j | X_n \in A),$$

where $A \subset E$. Use an example to show this.

Proof. Consider the (p, q)-biased random walk $\{X_n\}$, where $0 < q < p < 1$ and the initial distribution is $\mathbb{P}(X_0 = 0) = \mathbb{P}(X_0 = 2) = 1/2$. Let $A = \{1, 3\}$. Then we have

$$\mathbb{P}(X_2 = 2 | X_1 \in A, X_0 = 0) = \mathbb{P}(X_2 = 2 | X_1 = 1) = p,$$

and

$$\begin{aligned}
\mathbb{P}(X_1 = 1) &= \mathbb{P}_0(X_1 = 1)\mathbb{P}(X_0 = 0) + \mathbb{P}_2(X_1 = 1)\mathbb{P}(X_0 = 2) \\
&= p/2 + q/2 = 1/2, \\
\mathbb{P}(X_1 = 3) &= \mathbb{P}_0(X_1 = 3)\mathbb{P}(X_0 = 0) + \mathbb{P}_2(X_1 = 3)\mathbb{P}(X_0 = 2) = p/2.
\end{aligned}$$

So

$$\begin{aligned}
\mathbb{P}(X_2 = 2|X_1 \in A) &= \frac{\mathbb{P}\left[\{X_2 = 2\} \cap (\{X_1 = 1\} + \{X_1 = 3\})\right]}{\mathbb{P}(\{X_1 = 1\} + \{X_1 = 3\})} \\
&= \frac{\mathbb{P}(X_2 = 2, X_1 = 1) + \mathbb{P}(X_2 = 2, X_1 = 3)}{\mathbb{P}(X_1 = 1) + \mathbb{P}(X_1 = 3)} \\
&= \frac{\mathbb{P}(X_2 = 2|X_1 = 1)\mathbb{P}(X_1 = 1) + \mathbb{P}(X_2 = 2|X_1 = 3)\mathbb{P}(X_1 = 3)}{\mathbb{P}(X_1 = 1) + \mathbb{P}(X_1 = 3)} \\
&= \frac{p/2 + qp/2}{1/2 + p/2} = \frac{p(1+q)}{1+p}.
\end{aligned}$$

Hence

$$\mathbb{P}(X_2 = 2|X_1 \in A, X_0 = 0) \neq \mathbb{P}(X_2 = 2|X_1 \in A).$$

□

QUESTION 7.2 *Use an example to show exists an adapted stochastic process $(X_t : t \in I)$ with respect to $(\mathscr{F}_t)$ which is not measurable. Namely, $I \times \Omega \to E, (t, \omega) \mapsto X_t(\omega)$ is not measurable.*

Proof. Let $I = \mathbb{R}_+$. Choose an unmeasurable subset $A \subset [0, 1]$ and a measurable set $F \in \mathscr{F}$. Consider the following stochastic process

$$(\mathbb{R}_+ \times \Omega, \mathscr{B}(\mathbb{R}_+) \times \mathscr{F}) \to (\mathbb{R}, \mathscr{B}(\mathbb{R})), \quad (t, \omega) \mapsto \mathbf{1}_{A\times F}(t, \omega) =: X_t(\omega)$$

and the filtration $(\mathscr{F}_t) = (\mathscr{F})$. For any $t \geq 0$, if $t \in A$, then $X_t(\omega) = \mathbf{1}_F(\omega) \in \mathscr{F}$. If $t \notin A$, then $X_t(\omega) = 0 \in \mathscr{F}$. So X is an adapted stochastic process. However, $\{(t, \omega) : X_t(\omega) = 1\} = A \times F \notin \mathscr{B}(\mathbb{R}_+) \times \mathscr{F}$. So X is not measurable. □

QUESTION 7.3 *Use an example to show exists a measurable stochastic process $(X_t : t \in I)$ which is not adapted to $(\mathscr{F}_t)$.*

Proof. Let $I = \mathbb{Z}_+$. Choose a measurable set $F \in \mathscr{F}$ $(F \neq \Omega, \emptyset)$. Consider the following stochastic process

$$(\mathbb{Z}_+ \times \Omega, \mathscr{B}(\mathbb{Z}_+) \times \mathscr{F}) \to (\mathbb{R}, \mathscr{B}(\mathbb{R})), \quad (t, \omega) \mapsto \mathbf{1}_{\{0\}\times F}(t, \omega) =: X_t(\omega)$$

and the filtration $(\mathscr{F}_t) = (\{\Omega, \emptyset\})$. Since

$$\sigma((t, \omega) \mapsto X_t(\omega)) = \{\mathbb{Z}_+ \times \Omega, \emptyset, \{0\} \times F\} \subset \mathscr{B}(\mathbb{Z}_+) \times \mathscr{F}.$$

Thus X is measurable. However for $t = 0$, we know $X_0(\omega) = \mathbf{1}_F(\omega)$ and

$$\sigma(\omega \mapsto X_0(\omega)) = \{\Omega, \emptyset, F\} \not\subset \mathscr{F}_0 = \{\Omega, \emptyset\}.$$

So X is not adapted to $(\mathscr{F}_t)$. □

QUESTION 7.4 *Use an example to show exists an adapted stochastic process $(X_t : t \in I)$ with respect to $(\mathscr{F}_t)$ which is not progressively measurable.*

Proof. We still use the example in **QUESTION 4.2**. We have proved X is adapted to $(\mathscr{F}_t) = (\mathscr{F})$. However, for any $t > 1$, we have

$$\{(t, \omega) : X_t(\omega) = 1\} \cap ([0, t] \times \Omega) = A \times F \notin \mathscr{B}[0, t] \times \mathscr{F}.$$

So we have finished the proof. □

QUESTION 7.5 *Use an example to show exists a measurable stochastic process $(X_t : t \in I)$ which is not progressively measurable.*

Proof. We still use the example in **QUESTION 4.3**. We have proved X is measurable. However for $t = 0$, we know

$$\begin{aligned}&\sigma((t, \omega) \mapsto X_t(\omega)) \cap (\{0\} \times \Omega)\\ =&\{\{0\} \times \Omega, \emptyset, \{0\} \times F\} \not\subset \mathscr{B}(\{0\}) \times \mathscr{F}_0.\end{aligned}$$

So X is not progressively measurable. □

QUESTION 7.6 *Use an example to show exists a measurable and adapted stochastic process $(X_t : t \in I)$ with respect to $(\mathscr{F}_t)$ which is not progressively measurable.*

Proof. I still can not find a proper example!!! □

QUESTION 7.7 *Suppose $(\mathscr{F}_t)$ is a filtration. Prove $(\mathscr{F}_{t+})$ is right continuous.*

Proof. Clearly, we have $\mathscr{F}_{t+} \subset \mathscr{F}_{t++}$. On the contrary, for all $A \in \mathscr{F}_{t++} = \cap_{\ell > t} \mathscr{F}_{\ell+} = \cap_{\ell > t} \cap_{s > \ell} \mathscr{F}_s$, then $\forall s, \ell$ satisfying $s > \ell > t$, we have $A \in \mathscr{F}_s$. Hence for each $\gamma > t$, exists ℓ_γ, such that $\gamma > \ell_\gamma > t$. So $A \in \mathscr{F}_\gamma$. By the arbitrariness of γ, we obtain $A \in \cap_{\gamma > t} \mathscr{F}_\gamma = \mathscr{F}_{t+}$. In conclusion, $\mathscr{F}_{t++} \subset \mathscr{F}_{t+}$. □

QUESTION 7.8 *Suppose $(E, \mathscr{E})$ is measurable space and isomorphic to a complete separable metric space $(F, \mathscr{B}(F))$. Given a Poisson random measure $X : (\Omega, \mathscr{F}, \mathbb{P}) \to (E, \mathscr{E})$ with intensity μ. Then for each measurable functions $\{f_i\}_{i=1}^n$ and disjoint sets $\{A_i\} \subset \mathscr{E}$, if $\{\mu(f_i)\}_{i=1}^n$ all make sense, then $X(f_1 \mathbf{1}_{A_1}), \cdots X(f_n \mathbf{1}_{A_n})$ is independent.*

Proof. When $f_i = \sum_{j=1}^{n_i} \alpha_{ij} \mathbf{1}_{U_{ij}}$, where $\{U_{ij}\}_{j=1}^{n_i} \subset \mathscr{E}$ is disjoint for each $i = 1, \cdots, n$, then we have

$$X(f_i; A_i) = \int_{A_i} \sum_{j=1}^{n_i} \alpha_{ij} \mathbf{1}_{U_{ij}}(x) X(dx) = \sum_{j=1}^{n_i} \alpha_{ij} X(A_i \cap U_{ij}).$$

By the definition of Poisson random measure, we know

$$\{X(A_i \cap U_{ij}) : \forall i = 1, \cdots, n, j = 1, \cdots, n_i\}$$

are independent. So the random vectors

$$\{(X(A_i \cap U_{i1}), \cdots, X(A_i \cap U_{in_i})) : \forall i = 1, \cdots, n\}$$

are independent. Furthermore, we know

$$\left\{X(f_i; A_i) = \sum_{j=1}^{n_i} \alpha_{ij} X(A_i \cap U_{ij}) : i = 1, \cdots, n\right\}$$

are independent. When $\{f_i\}_{i=1}^n \subset \mathrm{p}\mathscr{E}$, we can find simple functions $\{g_{ij}\}$ such that $g_{ij} \uparrow f, j \to \infty$. Then for $\{\theta_i\}_{i=1}^n > 0$ and $A \in \mathscr{E}$, by Monotonicity convergence theorem, we know $X(f_{ij}\mathbf{1}_{A_i}) \uparrow X(f_i\mathbf{1}_{A_i})$. Then by Dominance convergence theorem, we have

$$\begin{aligned}
\mathbb{P}\exp\left\{-\sum_{i=1}^n \theta_i X(f_i\mathbf{1}_{A_i})\right\} &= \lim_{j\to\infty} \mathbb{P}\exp\left\{-\sum_{i=1}^n \theta_i X(f_{ij}\mathbf{1}_{A_i})\right\} \\
&= \lim_{j\to\infty} \mathbb{P}\left(\prod_{i=1}^n \exp\{-\theta_i X(f_{ij}\mathbf{1}_{A_i})\}\right) \\
&= \lim_{j\to\infty} \prod_{i=1}^n \mathbb{P}\exp\{-\theta_i X(f_{ij}\mathbf{1}_{A_i})\} \\
&= \prod_{i=1}^n \mathbb{P}\exp\{-\theta_i X(f_i\mathbf{1}_{A_i})\}
\end{aligned}$$

By the property of generating function, we know $\{X(f_i\mathbf{1}_{A_i})\}_{i=1}^n$ are independent. When $\{f_i\}_{i=1}^n \in \mathscr{E}$, then

$$\begin{aligned}
X(f_i\mathbf{1}_{A_i}) &= X(f_i^+\mathbf{1}_{A_i}) - X(f_i^-\mathbf{1}_{A_i}) \\
&= X(f\mathbf{1}_{\{f_i>0\}\cap A_i}) - X((-f)\mathbf{1}_{\{f_i<0\}\cap A_i}).
\end{aligned}$$

Utilize what we have proved, we know

$$\{X(f_i^\pm\mathbf{1}_{A_i}) : i = 1, \cdots, n\}$$

are independent. So the random vectors

$$\{(X(f_i^+\mathbf{1}_{A_i}), X(f_i^-\mathbf{1}_{A_i})) : i = 1, \cdots, n\}$$

are independent. Furthermore, we know

$$\{X(f_i\mathbf{1}_{A_i}) : i = 1, \cdots, n\}$$

are independent. □

QUESTION 7.9 *Suppose T is a $(\mathscr{F}_t)_{t\in I}$ stopping time. Then $\{T=\infty\}\in\mathscr{F}_\infty$.*

Proof. Observe for each $n\geq 1$, $\{T>n\}=\{T\leq n\}^c\in\mathscr{F}_n\subset\mathscr{F}_\infty$, then

$$\{T=\infty\}=\bigcap_{n=1}^{\infty}\{T>n\}\in\mathscr{F}_\infty.$$

The proof is completed. □

QUESTION 7.10 *Suppose $(\mathscr{G}_t)_{t\in I}$ is the enlargement of the filtration $(\mathscr{F}_t)_{t\in I}$. Then $\mathscr{G}_\infty=\sigma(\mathscr{F}_\infty\cup\mathscr{N})$.*

Proof. Obviously $\mathscr{F}_\infty\subset\mathscr{G}_\infty$ and $\mathscr{N}\subset\mathscr{G}_0\subset\mathscr{G}_\infty$, thus $\sigma(\mathscr{F}_\infty\cup\mathscr{N})\subset\mathscr{G}_\infty$. On the contrary, for each $t\in I$, we have $\mathscr{G}_t=\sigma(\mathscr{F}_t\cup\mathscr{N})\subset\sigma(\mathscr{F}_\infty\cup\mathscr{N})$. Thus $\mathscr{G}_\infty=\sigma(\cup_{t\in I}\mathscr{G}_t)\subset\sigma(\mathscr{F}_\infty\cup\mathscr{N})$. □

QUESTION 7.11 *Suppose $(B_t)_{t\geq 0}$ is the Brownian motion on the probability space $(\Omega,\mathscr{F},\mathbb{P})$. Then $B_t/t\to 0,\mathbb{P}$-a.s.. Therefore*

$$X_t=\begin{cases}B_t/t & ,\text{if } t>0,\\ 0 & ,\text{if } t=0.\end{cases}$$

is a.s. continuous.

Proof. Since $(B_t)_{t\geq 0}$ is a continuous time martingale and $B_t\sim N(0,t)$, then for each $\varepsilon>0$ and $0<\delta<\gamma$,

$$\begin{aligned}\mathbb{P}\left(\sup_{\delta\leq t\leq\gamma}|B_t/t|\geq\varepsilon\right)&\leq\mathbb{P}\left(\sup_{\delta\leq t\leq\gamma}|B_t|/\delta\geq\varepsilon\right)\\&\leq\varepsilon^{-2}\mathbb{P}\left(B_\gamma^2/\delta^2\right)=\frac{\gamma}{\varepsilon^2\delta^2}.\end{aligned}$$

Especially, we take $\delta=2^n,\gamma=2^{n+1}$ for each $n\geq 0$. Then we have

$$\sum_n\mathbb{P}\left(\sup_{2^n\leq t\leq 2^{n+1}}|B_t/t|\geq\varepsilon\right)\leq\sum_n\frac{1}{2^{n-1}\varepsilon^2}<\infty.$$

The Borel-Cantelli's lemma shows

$$\mathbb{P}\left(\bigcap_{m=0}^{\infty}\bigcup_{n=m}^{\infty}\sup_{2^n\leq t\leq 2^{n+1}}|B_t/t|\geq\varepsilon\right)=0,\quad\forall\varepsilon>0.$$

This yields $B_t/t\to 0,\mathbb{P}$-a.s.. □

QUESTION 7.12 *Suppose $(X_t)_{t\in I}$ is a Markov chain on the discrete state space E. For each $i \in E$, define $S_i^0 = 0$ and successively define $S_i^n = \inf\{t > S_i^{n-1} : X_t = i\}$ for $n \geq 1$. Write $T_i^n = S_i^n - S_i^{n-1}$ for $n \geq 1$. If $\mathbb{P}_i(T_i^1 < \infty) = \mathbb{P}_i(S_i^1 < \infty) = 1$, then $\{T_i^n\}_{n\geq 1}$ is a i.i.d.random variables sequence under the probability measure $\mathbb{P}^i$ and $T_i^k \sim T_i^1$ for each $k \geq 1$.*

Proof. We use the notation of the Markov system. Then $S_i^{n+1} = S_i^n + S_i \circ \theta_{S_i^n}$ and $T_i^{n+1} = S_i \circ \theta_{S_i^n}$ for each $n \geq 0$. For each □

QUESTION 7.13 *Suppose $(X_n)_{n\geq 0}$ is a Markov chain on the discrete state space E. For each $i \in E$, define*

$$N_n(i) = \{\text{the number of visits to } i \text{ at times} \leq n\},$$
$$N(i) = \{\text{the number of visits to } i\}.$$

Then $N(i) = \infty, \mathbb{P}_i$-a.s. iff i is recurrent. Moreover, when $i \in E$ is recurrent, then $N_n(i) \to \infty (n \to \infty), \mathbb{P}_i$-a.s.

Proof. Define $S_i^0 = 0$ and successively define $S_i^n = \inf\{n > S_i^{n-1} : X_n = i\}$ for $n \geq 1$. By the strong Markov property, we have

$$\mathbb{P}_i(S_i^n < \infty) = [\mathbb{P}_i(S_i^1 < \infty)]^n = (f_{ii})^n, \quad n \geq 1$$

and

$$\mathbb{P}_i[N(i) = \infty] = \lim_{n\to\infty} \mathbb{P}_i[N(i) \geq n] = \lim_{n\to\infty} \mathbb{P}_i(S_i^n < \infty) = \lim_{n\to\infty} (f_{ii})^n.$$

Therefore $\mathbb{P}_i[N(i) = \infty] = 1$ iff $f_{ii} = 1$. Namely, $N(i) = \infty, \mathbb{P}_i$-a.s. iff i is recurrent. Moreover, notice

$$N_n(i) = \sum_{k=1}^{n} \mathbb{1}_{\{X_k = i\}}.$$

When i is recurrent, we have $N_n(i) \uparrow N(i) = \infty, \mathbb{P}_i$-a.s.. □

QUESTION 7.14 *Suppose (N_t) is a Poisson process with parameter $\lambda > 0$ on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$. Define $T_1 = \inf\{t \geq 0 : N_t = 1\}$, then we have*

$$\mathbb{P}(T_1 \leq s | N_t = 1) = s/t, \quad \forall 0 \leq s \leq t.$$

Proof. Utilize the independence of N_s and $N_t - N_s$, then we have

$$\begin{aligned}\mathbb{P}(T_1 \le s | N_t = 1) = \mathbb{P}(N_s = 1 | N_t = 1) &= \frac{\mathbb{P}(N_s = 1, N_t = 1)}{\mathbb{P}(N_t = 1)} \\ &= \frac{\mathbb{P}(N_s = 1, N_t - N_s = 0)}{\mathbb{P}(N_t = 1)} \\ &= \frac{\mathbb{P}(N_s = 1)\mathbb{P}(N_t - N_s = 0)}{\mathbb{P}(N_t = 1)} \\ &= \frac{\lambda s e^{-\lambda s} e^{-\lambda(t-s)}}{\lambda t e^{-\lambda t}} = \frac{s}{t}.\end{aligned}$$

This is what we desire. □

QUESTION 7.15 *Suppose T is a $(\mathscr{F}_n)_{n\ge 0}$ stopping time on the filtered probability space $(\Omega, \mathscr{F}, (\mathscr{F}_n), \mathbb{P})$. If the random variable $X \in \mathscr{F}$ is integrable, then*

$$\mathbb{P}(X|\mathscr{F}_T) = \sum_{n=0}^{\infty} \mathbb{P}(X|\mathscr{F}_n)\mathbb{1}_{\{T=n\}} + \mathbb{P}(X|\mathscr{F}_\infty)\mathbb{1}_{\{T=\infty\}}.$$

Proof. Write F for the right side. For each $k \ge 0$, we have

$$F\mathbb{1}_{\{T\le k\}} = \sum_{n=0}^{k} \mathbb{P}(X|\mathscr{F}_n)\mathbb{1}_{\{T=n\}} \in \mathscr{F}_k,$$

so $F \in \mathscr{F}_T$. Moreover

$$\begin{aligned}&\mathbb{P}\left[\left|\sum_{n=0}^{\infty} \mathbb{P}(X|\mathscr{F}_n)\mathbb{1}_{\{T=n\}} + \mathbb{P}(X|\mathscr{F}_\infty)\mathbb{1}_{\{T=\infty\}}\right|\right] \\ &\quad\le \mathbb{P}\left[\sum_{n=0}^{\infty} \mathbb{P}(|X||\mathscr{F}_n)\mathbb{1}_{\{T=n\}} + \mathbb{P}(|X||\mathscr{F}_\infty)\mathbb{1}_{\{T=\infty\}}\right] \\ &\quad= \sum_{n=0}^{\infty} \mathbb{P}[\mathbb{P}(|X||\mathscr{F}_n)\mathbb{1}_{\{T=n\}}] + \mathbb{P}[\mathbb{P}(|X||\mathscr{F}_\infty)\mathbb{1}_{\{T=\infty\}}] \\ &\quad= \sum_{n=0}^{\infty} \mathbb{P}(|X|; T = n) + \mathbb{P}(|X|; T = \infty) = \mathbb{P}(|X|) < \infty.\end{aligned}$$

Thus F is also integrable. For each $A \in \mathscr{F}_T$, by the Fubini's theorem,

$$\begin{aligned}\mathbb{P}(F; A) &= \sum_{n=0}^{\infty} \mathbb{P}[\mathbb{P}(X|\mathscr{F}_n)\mathbb{1}_{\{T=n\}\cap A}] + \mathbb{P}[\mathbb{P}(X|\mathscr{F}_\infty)\mathbb{1}_{\{T=\infty\}\cap A}] \\ &= \sum_{n=0}^{\infty} \mathbb{P}(X; \{T = n\} \cap A) + \mathbb{P}(X; \{T = \infty\} \cap A] \\ &= \mathbb{P}(X; A).\end{aligned}$$

So $\mathbb{P}(X|\mathscr{F}_T) = F$. □

QUESTION 7.16 *Let $M = (M_n)_{n\geq 0}$ be a $(\mathscr{F}_n)$ uniformly integrable martingale and let M_∞ be the a.s. and L^1 limit of M, where $(\mathscr{F}_n)$ is reinforced. Then for any stopping time T, we have*

$$\mathbb{P}(M_\infty|\mathscr{F}_T) = M_T,$$

where with convention that $M_T = M_\infty$ on the event $\{T = \infty\}$. Therefore,

1. *For each stopping times S, T and $S \leq T$, we have*

$$\mathbb{P}(M_T|\mathscr{F}_S) = M_S.$$

2. *For each stopping time T, we have $\mathbb{P}(|M_T|) < \infty$ and $\mathbb{P}(M_T) = \mathbb{P}(M_0)$.*

Proof. Way 1. Since M is a uniformly integrable martingale, then exists M_∞ such that $M_n \xrightarrow[\text{a.s.}]{L^1} M_\infty$ and $\mathbb{P}(|M_\infty|) < \infty$. Moreover, for each $n \geq 0$,

$$\mathbb{P}(M_\infty|\mathscr{F}_n) = M_n.$$

Furthermore, for the given $n \geq 0$, by the optional stopping theorem, we have

$$\mathbb{P}(M_n|\mathscr{F}_{T\wedge n}) = M_{T\wedge n}.$$

Combine the above two conditions, note $\mathscr{F}_{T\wedge n} \subset \mathscr{F}_n$, we have

$$(*) \qquad \mathbb{P}(M_\infty|\mathscr{F}_{T\wedge n}) = \mathbb{P}[\mathbb{P}(M_\infty|\mathscr{F}_n)|\mathscr{F}_{T\wedge n}] = \mathbb{P}(M_n|\mathscr{F}_{T\wedge n}) = M_{T\wedge n}.$$

It follows from $(*)$ that

$$|M_{T\wedge n}| \leq \mathbb{P}(|M_\infty||\mathscr{F}_{T\wedge n}).$$

Since $\{T \leq n\} \in \mathscr{F}_{T\wedge n}$, then

$$\begin{aligned}\mathbb{P}(|M_T|;\{T \leq n\}) &= \mathbb{P}(|M_{T\wedge n}|; \{T \leq n\}) \\ &\leq \mathbb{P}[\mathbb{P}(|M_\infty||\mathscr{F}_{T\wedge n}); \{T \leq n\}] = \mathbb{P}(|M_\infty|; \{T \leq n\}).\end{aligned}$$

Let $n \to \infty$, the Monotone convergence theorem yields

$$\mathbb{P}(|M_T|; \{T < \infty\}) \leq \mathbb{P}(|M_\infty|; \{T < \infty\}).$$

Moreover, observe $\mathbb{P}(|M_T|; \{T = \infty\}) = \mathbb{P}(|M_\infty|; \{T = \infty\})$, then

$$\begin{aligned}\mathbb{P}(|M_T|) &= \mathbb{P}(|M_T|; \{T < \infty\}) + \mathbb{P}(|M_T|; \{T = \infty\}) \\ &\leq \mathbb{P}(|M_\infty|; \{T < \infty\}) + \mathbb{P}(|M_\infty|; \{T = \infty\}) \\ &= \mathbb{P}(|M_\infty|) < \infty.\end{aligned}$$

Therefore $\mathbb{P}(|M_T|) < \infty$.

If $A \in \mathscr{F}_T$, we claim $A \cap \{T \le n\} \in \mathscr{F}_{T\wedge n}$. In fact, $A \cap \{T \le n\} \in \mathscr{F}_n$ is obvious. Moreover $A \in \mathscr{F}_T$ and $\{T \le n\} \in \mathscr{F}_T$ implies $A \cap \{T \le n\} \in \mathscr{F}_T$. So $A \cap \{T \le n\} \in \mathscr{F}_T \cap \mathscr{F}_n = \mathscr{F}_{T\wedge n}$. So $(*)$ implies

$$\mathbb{P}(M_\infty; A \cap \{T \le n\}) = \mathbb{P}(M_{T\wedge n}; A \cap \{T \le n\}) = \mathbb{P}(M_T; A \cap \{T \le n\}).$$

Let $n \to \infty$ and utilize the Dominance convergence theorem,

$$\mathbb{P}(M_\infty; A \cap \{T < \infty\}) = \mathbb{P}(M_T; A \cap \{T < \infty\}).$$

Observe $\mathbb{P}(M_\infty; A \cap \{T = \infty\}) = \mathbb{P}(M_T; A \cap \{T = \infty\})$. then

$$\mathbb{P}(M_\infty; A) = \mathbb{P}(M_T; A), \quad \forall A \in \mathscr{F}_T.$$

Since $M_T \in \mathscr{F}_T$, then $\mathbb{P}(M_\infty|\mathscr{F}_T) = M_T$. Therefore

1. For each stopping times S, T and $S \le T$, observe $\mathscr{F}_S \subset \mathscr{F}_T$, then

$$(**) \qquad \mathbb{P}(M_T|\mathscr{F}_S) = \mathbb{P}[\mathbb{P}(M_\infty|\mathscr{F}_T)|\mathscr{F}_S] = \mathbb{P}(M_\infty|\mathscr{F}_S) = M_S.$$

2. For each stopping time T, we have shown $\mathbb{P}(|M_T|) < \infty$. Use $(**)$, we have

$$\mathbb{P}(M_T) = \mathbb{P}[\mathbb{P}(M_T|\mathscr{F}_0)] = \mathbb{P}(M_0).$$

Way 2. We just prove $\mathbb{P}(M_\infty|\mathscr{F}_T) = M_T$ since the rest is similar. Observe $(\mathscr{F}_n)$ is reinforced, then $M_\infty \in \mathscr{F}_\infty$. Because M_∞ is integrable, then the previous **QUESTION** shows

$$\begin{aligned}\mathbb{P}(M_\infty|\mathscr{F}_T) &= \sum_{n=0}^{\infty} \mathbb{P}(M_\infty|\mathscr{F}_n)\mathbb{1}_{\{T=n\}} + \mathbb{P}(M_\infty|\mathscr{F}_\infty)\mathbb{1}_{\{T=\infty\}} \\ &= \sum_{n=0}^{\infty} M_n\mathbb{1}_{\{T=n\}} + M_\infty\mathbb{1}_{\{T=\infty\}} = M_T.\end{aligned}$$

The result is completed. □

QUESTION 7.17 *Suppose $X = (X_n)$ and $Y = (Y_n)$ are two irreducible and aperiodic Markov chains on the state space E with common transition probability P and initial distributions λ and π, where π is the stationary distribution of P. Consider the independent coupling $W_n = (X_n, Y_n)$, which has the transition probability $\tilde{P}$, where $\tilde{p}_{(i,j)(\gamma,\ell)} = p_{i\gamma}p_{j,\ell}$, and the initial distribution $\tilde{\mu}_i = \lambda_i\pi_i$. Fix $i \in E$ and define the following stopping time,*

$$T := \inf\{n \ge 1 : W_n = (i, i)\}.$$

The we define

$$Z_n := \begin{cases} (X_n, Y_n) & \text{if } n < T, \\ (Y_n, X_n) & \text{if } n \geq T. \end{cases}$$

Prove $Z = (Z_n)$ is a Markov chain with transition probability $\tilde{P}$ and the initial distribution $\tilde{\mu}$.

Proof. For each $n \geq 1$ and $\{i_k\}_{k=1}^n, \{j_k\}_{k=1}^n \subset E$, without loss of generality, we assume exists $1 \leq m \leq n$, such that

$$\{Z_0 = (i_0, j_0), \cdots, Z_n = (i_n, j_n)\} \subset \{T = m\} \subset \{n \geq T\}.$$

Therefore, utilize the Markov property of $W_n = (X_n, Y_n)$,

$$\begin{aligned}
&\mathbb{P}[Z_0 = (i_0, j_0), \cdots, Z_T = (i_T, j_T), \cdots, Z_n = (i_n, j_n)] \\
=&\mathbb{P}(X_0 = i_0, Y_0 = j_0, \cdots, Y_T = i_T, X_T = j_T, \cdots, Y_n = i_n, X_n = j_n) \\
=&\mathbb{P}(X_0 = i_0, Y_0 = j_0, \cdots, Y_m = i, X_m = i, \cdots, Y_n = i_n, X_n = j_n) \\
=&\mathbb{P}[W_0 = (i_0, j_0), \cdots, W_m = (i, i), \cdots, W_n = (j_n, i_n)] \\
=&\tilde{\mu}_{(i_0,j_0)}\tilde{p}_{(i_0,j_0)(i_1,j_1)} \cdots \tilde{p}_{(i_{m-1},j_{m-1})(i,i)}\tilde{p}_{(i,i)(j_{m+1},i_{m+1})} \cdots \tilde{p}_{(j_{n-1},i_{n-1})(j_n,i_n)} \\
=&\lambda_{i_0}\pi_{i_0}p_{i_0i_1}p_{j_0j_1} \cdots p_{i_{m-1}i}p_{j_{m-1}i}p_{ij_{m+1}}p_{ii_{m+1}} \cdots p_{j_{n-1}j_n}p_{i_{n-1}i_n} \\
=&\lambda_{i_0}\pi_{i_0}p_{i_0i_1}p_{j_0j_1} \cdots p_{i_{m-1}i}p_{j_{m-1}i}p_{ii_{m+1}}p_{ij_{m+1}} \cdots p_{i_{n-1}i_n}p_{j_{n-1}j_n} \\
=&\tilde{\mu}_{(i_0,j_0)}\tilde{p}_{(i_0,j_0)(i_1,j_1)} \cdots \tilde{p}_{(i_{m-1},j_{m-1})(i,i)}\tilde{p}_{(i,i)(i_{m+1},j_{m+1})} \cdots \tilde{p}_{(i_{n-1},j_{n-1})(i_n,j_n)}.
\end{aligned}$$

This shows $Z = (Z_n)$ is a Markov chain with transition probability $\tilde{P}$ and the initial distribution $\tilde{\mu}$. □

QUESTION 7.18 *Suppose $\{S_n\}_{n\geq 1}$ is a sequence of i.i.d. exponential distribution with parameter λ. Fix $i \geq 1$, define $J_n := \sum_{k=1}^n S_k$ and*

$$\tilde{S}_1 = S_{i+1} - (s - J_i), \qquad \tilde{S}_n = S_{i+n} \quad \forall n \geq 2.$$

Then conditional on the event $J_i \leq s < J_{i+1}$, we have $\{\tilde{S}_n\}_{n\geq 1}$ are i.i.d. exponential distribution with parameter λ.

Proof. Given $k \geq 1$, for each $\{t_\ell\}_{\ell=1}^k \subset \mathbb{R}_+$, utilize the independence between

$\{S_n\}_{n\geq 1}$ and **QUESTION 4.31**, we have

$$\begin{aligned}
&\mathbb{P}(\tilde{S}_1 > t_1, \cdots, \tilde{S}_k > t_n | J_i \leq s < J_{i+1}) \\
&\quad = \mathbb{P}[S_{i+1} - (s - J_i) > t_1, \cdots, S_{i+k} > t_k | s - J_i \geq 0, S_{i+1} > s - J_i] \\
&\quad = \frac{\mathbb{P}[S_{i+1} - (s - J_i) > t_1, \cdots, S_{i+k} > t_k, s - J_i \geq 0, S_{i+1} > s - J_i]}{\mathbb{P}(s - J_i \geq 0, S_{i+1} > s - J_i)} \\
&\quad = \frac{\mathbb{P}[S_{i+1} - (s - J_i) > t_1, s - J_i \geq 0, S_{i+1} > s - J_i]}{\mathbb{P}(s - J_i \geq 0, S_{i+1} > s - J_i)} \\
&\qquad \times \mathbb{P}(S_{i+2} > t_2, \cdots, S_{i+k} > t) \\
&\quad = \mathbb{P}[S_{i+1} > t_1 + (s - J_i) | s - J_i \geq 0, S_{i+1} > s - J_i] \times \prod_{\ell=2}^{k} \mathbb{P}(S_{i+\ell} > t_\ell) \\
&\quad = \prod_{\ell=1}^{k} \mathbb{P}(S_{i+\ell} > t_\ell) = e^{-\lambda \sum_{\ell=1}^{k} t_\ell}.
\end{aligned}$$

So the desired result follows. □

QUESTION 7.19 *Suppose (X_n) is an irreducible Markov chain with transition matrix P and state space E. If μ is a non-trivial invariant measure of P, then μ is strictly positive.*

Proof. Since μ is a non-trivial invariant measure of P, then exists $i \in E$, such that $\mu_i > 0$. Observe P is irreducible, then for any $j \in E$, exists $n \geq 1$ satisfying $p_{ij}^{(n)} > 0$. Then utilize $\mu = \mu P^{(n)}$, we have

$$\mu_j = \sum_{k \in E} \mu_k p_{kj}^{(n)} \geq \mu_i p_{ij}^{(n)} > 0.$$

So μ is strictly positive. □

QUESTION 7.20 *Suppose $(X_t)_{t\geq 0}$ is a stochastic process in the filtered probability space $(\Omega, \mathscr{F}, (\mathscr{F}_t)_{t\geq 0}, \mathbb{P})$ and valued on a metric space (E, d). Define $T = \inf\{t \geq 0 : X_t \in A\}$, where $A \in \mathscr{B}(E)$ ensuring T is a $(\mathscr{F}_t)_{t\geq 0}$ optional time.*

1. *If $(X_t)_{t\geq 0}$ has continuous-path, then $X_T \in \partial A$.*
2. *If $(X_t)_{t\geq 0} \in D_E[0, \infty)$, then $X_T \notin \mathrm{Ext}A$. Especially, when A is a skeleton, say $A = \{x\}$, then $X_T = x$.*
3. *Assume $(X_t)_{t\geq 0} \in D_E[0, \infty)$ and A is a open set. If $X_s \in A$, then exists $\delta > 0$ satisfying $X_t \in A, \forall t \in [s, s + \delta)$. Especially, if $A = \{x\}$ is an open set, then exists $\delta > 0$ satisfying $X_t = x, \forall t \in [s, s + \delta)$*

Proof. 1. Firstly, if $X_T \in \mathring{A}$, notice $\mathring{A}$ is open and $(X_t)_{t\geq 0}$ is left continuous, then exists $0 \leq s < T$ such that $X_s \in \mathring{A}$, which contradicts to definition of T. Besides, if $X_T \in \mathrm{Ext}A$, notice $\mathrm{Ext}A$ is open and $(X_t)_{t\geq 0}$ is right continuous, then exists $\delta > 0$ such that $X_s \in \mathrm{Ext}A, \forall T \leq s < T + \delta$. Namely, $X_s \notin A$ for any $T \leq s < T + \delta$, which also contradicts to definition of T.

2. If $(X_t)_{t\geq 0} \in D_E[0, \infty)$, the similar argument concerning the exterior of A in the first question gives the desired results. Finally, since when $A = \{x\}$, we have $A = \mathring{A} \cup \partial A$. Then $X_T \notin \mathrm{Ext}A \Longrightarrow X_T = x$.

3. The last assertion is just the definition. □

QUESTION 7.21 *Discuss the the relationship between the three statements.*

1. *For any $i, j \in E$, exist $n \in \mathbb{N}_+$ and distinctive states $i = i_1, \cdots, i_n = j$ such that $q_{i_1 i_2} \cdots q_{i_{n-1} i_n} > 0$.*

2. *For any $i, j \in E$, exist $n \in \mathbb{N}_+$ and states $i = i_1, \cdots, i_n = j$ such that $q_{i_1 i_2} \cdots q_{i_{n-1} i_n} \neq 0$.*

3. *For any $i, j \in E$, exist $n \in \mathbb{N}_+$ such that $q_{ij}^{(n)} \neq 0$.*

Proof. 1. $1 \Longleftrightarrow 2$. First, $1 \Longrightarrow 2$ is obvious.

$2 \Longrightarrow 1$. If $i_1, \cdots, i_{n-1}$ are distinct, then all are done and it is clear that $q_{i_1 i_2} \cdots q_{i_{n-1} i_n} > 0$. Otherwise, presume $i_{n_1} < \cdots < i_{n_{m_1}}$ are the same, $i_{n_2} < \cdots < i_{n_{m_2}}$ are the same, $\cdots$, and $i_{n_3} < \cdots < i_{n_{m_3}}$ are the same. We can cut all the loops, then the rest part satisfies our requirement.

2. $3 \Longrightarrow 2$. Observe we have the following expression:

$$\sum_{i_1, \cdot, i_{n-1} \in E} q_{i i_1} q_{i_1 i_2} \cdots q_{i_{n-1} i_n} \neq 0,$$

which implies exist states $i = i_1, \cdots, i_n = j$ such that $q_{i_1 i_2} \cdots q_{i_{n-1} i_n} \neq 0$.

3. However $2 \not\Longrightarrow 3$. Consider the following example.

$$Q = \begin{pmatrix} -3 & 2 & 1 \\ 2 & -3 & 1 \\ 3 & 12 & -15 \end{pmatrix}.$$

It is trivial to text Q is indeed a Q-matrix. We have

$$q_{12}^{(2)} = q_{11} q_{12} + q_{12} q_{22} + q_{13} q_{32} = -6 - 6 + 12 = 0.$$

However, clearly $q_{11} q_{12} \neq 0$, $q_{12} q_{22} \neq 0$ and $q_{13} q_{32} \neq 0$. □

QUESTION 7.22 *Suppose P satisfyies $\exists i_0 \in E$ such that $\sum_{j\in E} p_{i_0 j} < 1$. Then*

1. *For all $n \in \mathbb{N}$, we have $\sum_{j\in E} p_{i_0 j}^{(n)} < 1$.*
2. *If exists $m \in \mathbb{N}$ and $i_1 \in E$ such that $p_{i_1 i_0}^{(m)} > 0$, then for any $n > m$, we have $\sum_{j\in E} p_{i_1 j}^{(n)} < 1$.*

Proof. 1. For any $n \geq 2$, we have

$$\begin{aligned}\sum_{j\in E} p_{i_0 j}^{(n)} &= \sum_{j\in E}\sum_{k\in E} p_{i_0 k} p_{kj}^{(n-1)} = \sum_{k\in E} p_{i_0 k} \sum_{j\in E} p_{kj}^{(n-1)} \\ &\leq \sum_{k\in E} p_{i_0 k} < 1.\end{aligned}$$

2. For any $n > m$, by the previous question, we have $\sum_{j\in E} p_{i_0 j}^{(n-m)} < 1$. Moreover, notice $p_{i_1 i_0}^{(m)} > 0$, then

$$\begin{aligned}\sum_{j\in E} p_{i_1 j}^{(n)} &= \sum_{j\in E}\sum_{k\in E} p_{i_1 k}^{(m)} p_{kj}^{(n-m)} = \sum_{j\in E}\sum_{k\neq i_0} p_{i_1 k}^{(m)} p_{kj}^{(n-m)} + \sum_{j\in E} p_{i_1 i_0}^{(m)} p_{i_0 j}^{(n-m)} \\ &< \sum_{k\neq i_0} p_{i_0 k}^{(m)} + p_{i_0 i_0}^{(m)} \leq 1.\end{aligned}$$

□

Theorem 7.1 (Perron-Frobenius theorem of Q) *Presume Q is irreducible. Then*

1. *Q has a principal eigenvalue $-\lambda \leq 0$ and two eigenvectors u and v satisfying*

$$uQ = -\lambda u, \quad Qv = -\lambda v.$$

2. *If we restrict $\sum_{i\in E} u_i = 1$ and $\sum_{i\in E} u_i v_i = 1$, then u and v is unique.*
3. *For any other eigenvalue λ' of Q, we have $|\lambda'| > \lambda$.*
4. *Let $P(t) = e^{tQ}$, then*

$$p_{ij}(t) \sim e^{-\lambda t} v_i u_j, \quad t \to \infty.$$

Proof. Let $c = \max_{i\in E} q_i$ and consider $Q^c =: Q + cI \geq 0$. The P-F theorem of nonnegative irreducible matrix yields

1. Q^c has a principal eigenvalue $\lambda^c > 0$, a positive left eigenvector u and a positive right eigenvector v satisfying

$$(*) \qquad uQ^c = \lambda^c u, \quad Q^c v = \lambda^c v.$$

2. If we restrict $\sum_{i\in E} u_i = 1, \sum_{i\in E} u_i v_i = 1$, the eigenvectors u, v are unique.
3. For any other eigenvalues λ_0^c of Q^c, we have $|\lambda_0^c| < \lambda^c$.

From $(*)$ we have

$$u\,(Q + cI) = \lambda^c u \Longrightarrow uQ = -(c - \lambda^c)u,$$
$$(Q + cI)\,v^c = \lambda^c v \Longrightarrow Qv = -(c - \lambda^c)v.$$

Define $\lambda =: c - \lambda^c$. Hence

$$uQ = -\lambda u, \quad Qv = -\lambda v,$$

and u, v are unique with the restriction $\sum_{i\in E} u_i = 1$ and $\sum_{i\in E} u_i v_i = 1$. Observe $uQ^c = \lambda^c u$ and $\sum_{j\in E} q_{ij}^c \le c$, then

$$\sum_{i\in E} u_i q_{ij}^c = \lambda^c u_j \Rightarrow \sum_{j\in E}\sum_{i\in E} u_i q_{ij}^c = \lambda^c \sum_{j\in E} u_j \Rightarrow c \ge \lambda^c.$$

Therefore $\lambda > 0$. Note $|\lambda_0^c| < \lambda^c$, then

$$(**) \qquad \lambda = c - \lambda^c < c - |\lambda_0^c| \le |\lambda_0^c - c|.$$

Since there is a one-to-one correspondence

$$(.) \qquad \varphi : \sigma\,(Q^c) \to \sigma\,(Q) : \mu \mapsto \mu - c.$$

Hence $(**)$ and $(.)$ means for any other eigenvalue λ' of Q, we have $|\lambda'| > \lambda$. Furthermore, for any **fixed** $t > 0$,

$$\begin{aligned} uP(t) &= u\sum_{n=0}^{\infty} \frac{Q^n t^n}{n!} = \sum_{n=0}^{\infty} \frac{uQ^n t^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{(-\lambda)^n u t^n}{n!} = \left(\sum_{n=0}^{\infty} \frac{(-\lambda)^n t^n}{n!}\right) u = \mathrm{e}^{-\lambda t} u. \end{aligned}$$

And symmetrically $P(t)v = \mathrm{e}^{-\lambda t} v$. Besides, we can also construct the following one-to-one correspondence

$$\phi : \sigma\,(Q) \to \sigma\,(P(t)) : \mu \mapsto \mathrm{e}^{\mu t}.$$

Consider $\phi \circ \varphi$, then for any other eigenvalue of $P(t)$, we can write it as $e^{(\lambda_0^c - c)t}$, where $\lambda_0^c \in \sigma(Q^c)$ and $\lambda_0^c \ne \lambda^c$. Thus

$$(..) \qquad |e^{(\lambda_0^c - c)t}| = \mathrm{e}^{(\mathrm{Re}\lambda_0^c - c)t} \le e^{(|\lambda_0^c| - c)t} < e^{(\lambda^c - c)t} = \mathrm{e}^{-\lambda t}.$$

Formula $(..)$ mean $\mathrm{e}^{-\lambda t}$ is the principal eigenvalue of $P(t)$. In conclusion, $P(t)$ has principal eigenpairs $(\mathrm{e}^{-\lambda t}, u, v)$.

For any **fixed** $t > 0$, the P-F theorem and C-K equation yield

$$(...) \qquad p_{ij}(nt) = p_{ij}^{(n)}(t) \sim \mathrm{e}^{-\lambda nt} v_i u_j, \quad n \to \infty.$$

Finally, we just need to show $p_{ij}(t) \sim \mathrm{e}^{-\lambda t} v_i u_j (t \to \infty)$. Or equivalently, we just need to prove $\forall i, j \in E$,

$$e^{\lambda t} p_{ij}(t) \to v_i u_j, \quad t \to \infty.$$

Define $\varphi_{ij}(t) = e^{\lambda t} p_{ij}(t), \forall t > 0$. Then the following properties naturally follow by $(...)$ and $P(t)v = e^{-\lambda t} v$:

$$\forall h > 0, \varphi_{ij}(nh) \to v_i u_j (n \to \infty), \qquad \forall t > 0, \sum_{j \in E} \varphi_{ij}(t) v_j = v_i.$$

Consider the following decomposition:

$$\varphi_{ij}(t) = \sum_{k \in E} \varphi_{ik}(t - nh) \varphi_{kj}(nh),$$

where $n = [t/h]$. Then $t - nh \in [0, h]$ and $t \to \infty$ implies $n \to \infty$. Moreover, observe $\varphi_{ij}(t)$ in bounded in $[0, h]$. Thus for any $i, j \in E$,

$$\begin{aligned} |\varphi_{ij}(t) - v_i u_j| &= \left| \sum_{k \in E} \varphi_{ik}(t - nh) \left[\varphi_{kj}(nh) - v_k u_j \right] \right| \\ &\leq \sum_{k \in E} \max_{s \in [0,h]} \varphi_{ik}(s) |\varphi_{kj}(nh) - v_k u_j| \to 0, \quad t \to \infty. \end{aligned}$$

Till now, the proof is completed. □

QUESTION 7.23 *Suppose the Markov chain P is irreducible and $i \neq j$.*

1. *If $f_{jj} = 1$, then $\sum_{n \in \mathbb{N}} p_{ij}^{(n)} = \infty$ and $f_{ij} = 1$.*
2. *If $f_{jj} < 1$, then $\sum_{n \in \mathbb{N}} p_{ij}^{(n)} < \infty$. However, we can not determine f_{ij}.*

Proof. 1. If $f_{jj} = 1$, the results from the following two equalities

$$\sum_{n \in \mathbb{N}} p_{ij}^{(n)} = \frac{f_{ij}}{1 - f_{jj}} = \infty$$

and

$$f_{jj} = p_{jj} + \sum_{k \neq j} p_{jk} f_{kj} = \sum_{k \in E} p_{jk} f_{kj} \Longrightarrow F_j = PF_j,$$

where $F_j = (f_{kj})_{k\in E}$. For any $i \neq j$, the irreducibility implies exists $m \in \mathbb{N}$ such that $p_{ij}^{(m)} > 0$. Therefore,

$$F_j = PF_j \Longrightarrow F_j = P^m F_j \Longrightarrow 1 = f_{jj} = \sum_{k\in E} p_{jk}^{(m)} f_{kj}.$$

However $\sum_{k\in E} p_{jk}^{(m)} = 1$, so $f_{ij} = 1$.

2. If $f_{jj} < 1$, The first equality shows $\sum_{n\in\mathbb{N}} p_{ij}^{(n)} < \infty$. We use two example to show $f_{ij} = 1$ and $f_{ij} < 1$ are both possible. Consider $E = \{1, 2, 3\}$ and $i = 1, j = 2$.

   (a) Let

$$P = \begin{pmatrix} 0 & 1 & 0 \\ 1/2 & 0 & 1/2 \\ 0 & 0 & 1 \end{pmatrix}$$

   Observe 3 is an absorb point, then $f_{12} = p_{12} = 1, f_{12}^{(n)} = 0 (n \geq 2)$. Moreover, $f_{22}^{(2)} = 1/2, f_{22}^{(n)} = 0 (n \neq 2)$. Therefore

$$\begin{aligned} f_{12} &= \sum_{n\in\mathbb{N}} f_{12}^{(n)} = p_{12} + \sum_{n=2}^{\infty} f_{12}^{(n)} = 1 + 0 = 1, \\ f_{22} &= \sum_{n\in\mathbb{N}} f_{12}^{(n)} = f_{22}^{(2)} = 1/2 \in (0, 1). \end{aligned}$$

   (b) Let

$$P = \begin{pmatrix} 0 & 1/2 & 1/2 \\ 1/2 & 0 & 1/2 \\ 0 & 0 & 1 \end{pmatrix}$$

   Observe 3 is an absorb point, then $f_{12} = p_{12} = 1/2, p_{12}^{(n)} = 0 (n \geq 2)$. Moreover $f_{22}^{(2)} = 1/4, f_{22}^{(n)} = 0 (n \neq 2)$ and 3 is an absorb point. Therefore

$$\begin{aligned} f_{12} &= \sum_{n\in\mathbb{N}} f_{12}^{(n)} = p_{12} + \sum_{n=2}^{\infty} f_{12}^{(n)} = 1/2 + 0 = 1/2, \\ f_{22} &= \sum_{n\in\mathbb{N}} f_{12}^{(n)} = f_{22}^{(2)} = 1/4 \in (0, 1). \end{aligned}$$

□

QUESTION 7.24 *Suppose E is a countable state space and $P = (p_{ij})_{i,j\in E}$ is an irreducible and aperiodic transition matrix. Suppose P is null-recurrent, then $p_{ij}^{(n)} \to 0, n \to \infty$.*

Proof. The diagonal method provides a sub-sequence $\{n_m\}$ such that

$$v_k := \lim_{m\to\infty} p_{ik}^{(n_m)}, \quad \forall k \in E.$$

If exists $j \in E$ such that $v_j > 0$, then the Fatou's lemma yields:

$$\sum_k v_k \le \lim_{m\to\infty} \sum_k p_{ik}^{(n_m)} = 1.$$

Moreover, the dominated convergence theorem yields:

$$\lim_{m\to\infty} p_{ik}^{(n_m+1)} = \lim_{m\to\infty} \sum_\ell p_{i\ell} p_{\ell k}^{(n_m)} = \sum_\ell p_{i\ell} v_k = v_k.$$

Therefore, use Fatou's lemma again to obtain

$$v_k = \lim_{m\to\infty} p_{ik}^{(n_m+1)} = \lim_{m\to\infty} \sum_\ell p_{i\ell}^{(n_m)} p_{\ell k} \ge \sum_\ell v_\ell p_{\ell k}.$$

Add all the k together, we will find the inequality above is actually equality. The irreducibility of P ensures $v_k > 0, \forall k \in E$. Observe $\sum_k v_k \le 1$, then P has a stationary distribution, which is equivalent to the positive recurrence of P. The contradiction arise.

If $v_k = 0, \forall k \in E$, then the dominated convergence theorem yields: $\forall r \ge 1$,

$$\lim_{m\to\infty} p_{ik}^{(n_m+r)} = \lim_{m\to\infty} \sum_\ell p_{i\ell}^{(r)} p_{\ell k}^{(n_m)} = \sum_\ell p_{i\ell}^{(r)} \lim_{m\to\infty} p_{\ell k}^{(n_m)} = 0.$$

Next we shows the limit $\lim_{m\to\infty} p_{ik}^{(n_m+r)}$ is uniform with respect to the variable r. In fact, the coupling inequality

$$|p_{ik}^{(n)} - p_{jk}^{(n)}| \le \mathbb{P}(T \ge n), \quad \forall i, j \in E$$

shows the limit $\lim_{m\to\infty} p_{\ell k}^{(n_m)}$ is uniform with respect to the state ℓ. Namely, for any $\varepsilon > 0$, exists $N \in \mathbb{N}$, for any $m > N$ and $\ell \in E$, we have $p_{\ell k}^{(n_m)} < \varepsilon$. This result together with the C-K equation yield: for any $r \in \mathbb{N}$,

$$p_{ik}^{(n_m+r)} = \sum_\ell p_{i\ell}^{(r)} p_{\ell k}^{(n_m)} < \varepsilon \sum_\ell p_{i\ell}^{(r)} = \varepsilon.$$

Observe the N mentioned above has nothing to do with $\ell \in E$ and $r \in \mathbb{N}$, the uniformity of r in $\lim_{m\to\infty} p_{ik}^{(n_m+r)}$ naturally follows.

For any $n \in \mathbb{N}$, exists unique $m \in \mathbb{N}$ such that $n_m \le n < m_{m+1}$. Therefore we can write $p_{ik}^{(n)} = p_{ik}^{(n_m+r_m)}$, where $r_n = n - n_m$. Observe $n \to \infty$ implies $m \to \infty$, then the uniformity of r in $\lim_{m\to\infty} p_{ik}^{(n_m+r)}$ implies

$$\lim_{n\to\infty} p_{ik}^{(n)} = \lim_{m\to\infty} p_{ik}^{(n_m+r_n)} = 0.$$

□

QUESTION 7.25 *If the state space E is finite, and P is random. Then there is at least one recurrent state and there is no null recurrence state.*

Proof. Since the asymptotic property of non-recurrence has nothing to do with the irreducibility, then if all the states are non-recurrent, then $p_{ij}^{(n)} \to 0, i, j \in E$. Therefore,

$$1 = \lim_{n\to\infty} \sum_j p_{ij}^{(n)} = \sum_j \lim_{n\to\infty} p_{ij}^{(n)} = 0,$$

which is a contradiction. We have two ways to handle the next assertion.

1. If $k \in E$ is null-recurrent, consider the irreducible class containing C_k such that $k \in C_k$. So all the states in C_k is null-recurrent and therefore C_k is closed. We have $p_{ij}^{(n)} \to 0, i, j \in C_k$. The same argument provides the same contradiction.

2. If $k \in E$ is recurrent, we want to shows k is positive recurrent. In fact, in the closed irreducible class C_k, exists invariant measure μ such that $\mu P|_{C_k} = \mu$. Observe E therefore C_k is finite, so the existence of invariant measure is equivalent to the existence of stationary distribution. Therefore k is positive recurrent. □

QUESTION 7.26 *Suppose P is a Markov chain and μ is its invariant measure. Define $\widetilde{p}_{ij} = \mu_j p_{ji}/\mu_i$. Prove $\widetilde{P} = (\widetilde{p}_{ij})$ is a Markov chain. $\widetilde{P}$ is called the dual process of P. Please prove:*

1. *P is irreducible if and only if $\widetilde{P}$ is irreducible.*
2. *P is recurrent if and only if $\widetilde{P}$ is recurrent.*
3. *P is positive recurrent if and only if $\widetilde{P}$ is positive recurrent.*

Proof. 1. First, we use induction on n to verify $\widetilde{p}_{ij}^{(n)} = \mu_j p_{ji}^{(n)}/\mu_i$. When $n = 1$, it is exactly the definition. Suppose when $n = k - 1$, our assertion is true. When $n = k$,

$$\begin{aligned}\widetilde{p}_{ij}^{(k)} &= \sum_{\tau\in E} \widetilde{p}_{i\tau}^{(k-1)} \widetilde{p}_{\tau j} = \sum_{\tau\in E} \frac{\mu_\tau p_{\tau i}^{(k-1)}}{\mu_i} \frac{\mu_j p_{j\tau}}{\mu_\tau} \\ &= \frac{\mu_j}{\mu_i} \sum_{\tau\in E} p_{j\tau} p_{\tau i}^{(k-1)} = \frac{\mu_j}{\mu_i} p_{ji}^{(k)}.\end{aligned}$$

Therefore the assertion is correct. For any $i, j \in E$, exist $n_{ij} \in \mathbb{N}$ such that $p_{ij}^{(n_{ij})} > 0$ iff $\widetilde{p}_{ij}^{(n_{ij})} > 0$. So the irreducibility is equivalent.

2. For any $j \in E$,

$$\sum_{n=0}^{\infty} \widetilde{p}_{jj}^{(n)} = \sum_{n=0}^{\infty} \frac{\mu_j}{\mu_j} p_{jj}^{(n)} = \sum_{n=0}^{\infty} p_{jj}^{(n)}.$$

So P is recurrent iff $\widetilde{P}$ is recurrent.

3. There are three ways to prove this assertion.

   (a) Notice $\forall j \in E, \widetilde{p}_{jj}^{(n)} = p_{jj}^{(n)}$, so

$$\widetilde{P}_{jj}(1) = \sum_{n=0}^{\infty} \widetilde{p}_{jj}^{(n)} = \sum_{n=0}^{\infty} p_{jj}^{(n)} = P_{jj}(1)$$

   $\forall \{s_n\}_{n\in\mathbb{N}}$, and $s_n \uparrow 1$, we have

$$\begin{aligned}\widetilde{m}_{jj} &= \lim_{n\to\infty} \left(\frac{\mathrm{d}}{\mathrm{d}s} \widetilde{F}_{jj}(s) \bigg|_{s=s_n} \right) = \lim_{n\to\infty} \left[\frac{\mathrm{d}}{\mathrm{d}s} \left(1 - \frac{1}{\widetilde{P}_{jj}(s)} \right) \bigg|_{s=s_n} \right] \\ &= \lim_{n\to\infty} \left[\frac{\mathrm{d}}{\mathrm{d}s} \left(1 - \frac{1}{P_{jj}(s)} \right) \bigg|_{s=s_n} \right] = \lim_{n\to\infty} \left(\frac{\mathrm{d}}{\mathrm{d}s} F_{jj}(s) \bigg|_{s=s_n} \right) = m_{jj}\end{aligned}$$

   So P is positive recurrent if and only if $\widetilde{P}$ is positive recurrent.

   (b) The existence of the dual chain $\widetilde{P}$ implies the existence of invariant measure μ of P. Moreover,

$$\sum_{i\in E} \mu_i \widetilde{p}_{ij} = \sum_{i\in E} \mu_i \frac{\mu_j p_{ji}}{\mu_i} = \sum_{i\in E} \mu_j p_{ji} = \mu_i.$$

   Therefore μ is also a invariant measure of $\widetilde{P}$. *By a general conclusion, P is positive recurrent iff P admit a stationary distribution (even P is not irreducible, and μ is not unique at the same time)*, then

$$P \text{ is positive recurrent} \Leftrightarrow \sum_{i\in E} \mu_i < \infty \Leftrightarrow \widetilde{P} \text{ is positive recurrent}$$

   (c) Observe

$$\begin{aligned}\widetilde{f}_{jj}^{(n)} &= \sum_{j_1,\cdots,j_{n-1}\neq j} \widetilde{p}_{jj_1}\widetilde{p}_{j_1j_2}\cdots\widetilde{p}_{j_{n-1}j} \\ &= \sum_{j_1,\cdots,j_{n-1}\neq j} \frac{\mu_{j_1}p_{j_1j}}{\mu_j}\frac{\mu_{j_2}p_{j_2j_1}}{\mu_{j_1}}\cdots\frac{\mu_j p_{jj_{n-1}}}{\mu_{j_{n-1}}} \\ &= \sum_{j_1,\cdots,j_{n-1}\neq j} p_{jj_{n-1}}p_{j_{n-1}j_{n-2}}\cdots p_{j_1j}^{(n)} \\ &= f_{jj}.\end{aligned}$$

   Then $\widetilde{m}_{jj} = m_{jj}, \forall j \in E$. □

QUESTION 7.27 *Markov chain (X_n) is invertible if and only if any times $n, m \in \mathbb{N}$ and any states $i, j \in E$, then*

$$\mathbb{P}(X_n = i, X_m = j) = \mathbb{P}(X_n = j, X_m = i)$$

1. *The above formula is equivalent to any times $n_1 < \cdots < n_m$, which suffice $n_{j+1} - n_j = n_{m+1-j} - n_{m-j}, j = 1, \cdots, m-1$ and any state $i_1, \cdots, i_m \in E$, then*

$$\mathbb{P}(X_{n_1} = i_1, \cdots, X_{n_m} = i_m) = \mathbb{P}(X_{n_1} = i_m, \cdots, X_{n_m} = i_1)$$

2. *(X_n) is invertible if and only if $\exists$ stationary distribution (π_i), s.t. $\pi_i p_{ij} = \pi_j p_{ji}$ and the initial distribution $\pi_0 = \pi$.*

Proof. 1. We just need to prove the sufficiency part. Let $n = 0, m = 1$, then we have

$$\begin{aligned}\mathbb{P}(X_0 = i)p_{ij} &= \mathbb{P}(X_0 = i, X_1 = j) \\ &= \mathbb{P}(X_0 = j, X_1 = i) = \mathbb{P}(X_0 = j)p_{ji}.\end{aligned}$$

Write $\pi_i := \mathbb{P}(X_0 = i)$ for $i \in E$, then the above formula yields

$$(*) \qquad \pi_i p_{ij} = \pi_j p_{ji} \Longrightarrow \pi P^n = \pi, \quad \forall n \geq 1.$$

Thus $X_0 \sim \pi$ is the reversible (stationary) distribution and thus for any $n \geq 1$,

$$\mathbb{P}(X_n = j) = \mathbb{P}(X_0 = j) = \pi_j, \quad \forall j \in E.$$

Therefore

$$\begin{aligned}&\mathbb{P}(X_{n_1} = i_1, \cdots, X_{n_m} = i_m) \\ =&\mathbb{P}(X_{n_1} = i_1)p_{i_1 i_2}^{(n_2-n_1)} p_{i_2 i_3}^{(n_3-n_2)} \cdots p_{i_{m-1} i_m}^{(n_m-n_{m-1})} \\ =&\pi_{i_1} p_{i_1 i_2}^{(n_2-n_1)} p_{i_2 i_3}^{(n_3-n_2)} \cdots p_{i_{m-1} i_m}^{(n_m-n_{m-1})} \\ =&p_{i_2 i_1}^{(n_2-n_1)} \pi_{i_2} p_{i_2 i_3}^{(n_3-n_2)} \cdots p_{i_{m-1} i_m}^{(n_m-n_{m-1})} \\ =&\cdots = p_{i_2 i_1}^{(n_2-n_1)} p_{i_3 i_2}^{(n_3-n_2)} \cdots p_{i_m i_{m-1}}^{(n_m-n_{m-1})} \pi_{i_m} \\ =&\pi_{i_m} p_{i_m i_{m-1}}^{(n_m-n_{m-1})} \cdots p_{i_3 i_2}^{(n_3-n_2)} p_{i_2 i_1}^{(n_2-n_1)} \\ =&\mathbb{P}(X_{n_1} = i_m, \cdots, X_{n_m} = i_1).\end{aligned}$$

2. Formula $(*)$ has proven the necessity part. For then sufficiency, we use induction on n to prove $\forall n \in \mathbb{N}_+, \pi_i p_{ij}^{(n)} = \pi_j p_{ji}^{(n)}$. When $n = 1$, this is

the requirement of this question. Suppose when $n = k - 1$, the assertion is right. When $n = k$,

$$\begin{aligned}\pi_i p_{ij}^{(k)} &= \pi_i \sum_{\tau\in E} p_{i\tau}^{(k-1)} p_{\tau j} = \sum_{\tau\in E} \pi_i p_{i\tau}^{(k-1)} p_{\tau j} = \sum_{\tau\in E} \pi_\tau p_{\tau i}^{(k-1)} p_{\tau j} \\ &= \sum_{\tau\in E} p_{\tau i}^{(k-1)} \pi_\tau p_{\tau j} = \sum_{\tau\in E} p_{\tau i}^{(k-1)} \pi_j p_{j\tau} = \pi_j \sum_{\tau\in E} p_{j\tau} p_{\tau i}^{(k-1)} \\ &= \pi_j p_{ji}^{(k)}.\end{aligned}$$

In conclusion, our assertion holds for $\forall n \in \mathbb{N}$. Since π is also a stationary distribution, then $\forall m \in \mathbb{N}, X_m \sim \pi$. So for any $n < m$,

$$\begin{aligned}\mathbb{P}(X_n = i, X_m = j) &= \mathbb{P}(X_n = i) p_{ij}^{(m-n)} = \pi_i p_{ij}^{(m-n)} \\ &= \pi_j p_{ji}^{(m-n)} = \mathbb{P}(X_n = j, X_m = i).\end{aligned}$$

□

QUESTION 7.28 *Suppose $(X_n)_{n\in\mathbb{N}}$ is an irreducible, aperiodic Markov chain on $\mathbb{Z}_+$. Then for any initial distribution μ,*

1. *$(X_n)_{n\in\mathbb{N}}$ is non-recurrent iff $X_n \xrightarrow{\text{a.s.}} \infty$*
2. *$(X_n)_{n\in\mathbb{N}}$ is non-recurrent iff $X_n \xrightarrow{\mathbb{P}} \infty$.*

Proof. 1. Since for any $N \in \mathbb{N}$ and any initial distribution $\mu = (\mu_k)_{k\in E}$, we have

$$\begin{aligned}\sum_{n=0}^{\infty} \mathbb{P}^\mu(X_n \le N) &= \sum_{n=0}^{\infty}\sum_{j=0}^{N} \mathbb{P}^\mu(X_n = j) = \sum_{n=0}^{\infty}\sum_{j=0}^{N}\sum_{k\in E} \mu_k p_{kj}^{(n)} \\ &= \sum_{j=0}^{N}\sum_{k\in E} \mu_k \sum_{n=0}^{\infty} p_{kj}^{(n)} = \sum_{j=0}^{N}\sum_{n=0}^{\infty} p_{\mu j}^{(n)}\end{aligned}$$

□

QUESTION 7.29 *Suppose there are two kinds of genes: A and B. And the number of the genes is $2N$. The composition of the next generation's genes is determined by $2N$ times of independent binomial experiments. Namely, if the numbers of A and B are i and $2N - i$, then the probabilities of each experiment with outcome A and B are*

$$p_i = \frac{i}{2N}, \quad q_i = 1 - p_i.$$

Let X_n be the number of A in the nth generation. Prove when $0 < j < 2N$,

$$\lim_{n\to\infty} p_{ij}^{(n)} = 0, \quad \lim_{n\to\infty} p_{i0}^{(n)} = 1 - \frac{i}{2N}, \quad \lim_{n\to\infty} p_{i,2N}^{(n)} = \frac{i}{2N}.$$

Proof. Define $T = \inf\{n \in \mathbb{N}_+ : X_n = 0 \text{ or } 2N\} = \tau_0^+ \wedge \tau_{2N}^+$. Observe $A := \{1, \cdots, 2N-1\}$ is a irreducible class and $P|_A$ is sub-random, then $p_{ij}^{(n)} \to 0, \forall i, j \in A$. Moreover, note $0, 2N$ are two absorbing points, then

$$p_{i0}^{(n)} = \sum_{m=1}^{n} f_{i0}^m p_{00}^{(n-m)} = \sum_{m=1}^{n} f_{i0}^m.$$

So $f_{i0} = \lim_{n\to\infty} p_{i0}^{(n)}$ and similarly $f_{i,2N} = \lim_{n\to\infty} p_{i,2N}^{(n)}$. Therefore for any $i \in A$,

$$\begin{aligned}
\mathbb{P}^i(T < \infty) &= \mathbb{P}^i(\tau_0^+ < \infty) + \mathbb{P}^i(\tau_{2N}^+ < \infty) = f_{i0} + f_{i,2N} \\
&= \lim_{n\to\infty} \left(p_{i0}^{(n)} + p_{i,2N}^{(n)}\right) = \lim_{n\to\infty} \left(1 - \sum_{j=1}^{2N} p_{ij}^{(n)}\right) \\
&= 1 - \sum_{j=1}^{2N} \lim_{n\to\infty} p_{ij}^{(n)} = 1.
\end{aligned}$$

We claim $(X_n)_{n\in\mathbb{N}}$ is a martingale. In fact, given $X_n = i$, we know $X_{n+1} \sim B(2N, i/2N)$, then the property of binomial distribution yields

$$\mathbb{P}(X_{n+1}|X_n = i) = i.$$

Hence $\mathbb{P}(X_{n+1}|X_n) = X_n$ and $(X_n)_{n\in\mathbb{N}}$ is a martingale. Although T is not bounded, we have $|X_n| \le 2N$. Therefore the *Doob's stopping theorem* still yields

$$0 f_{i0} + 2N f_{i,2N} = \mathbb{P}^i(X_T) = \mathbb{P}^i(X_0) = i.$$

This formula together with $f_{i0} + f_{i,2N} = 1$ show us

$$\lim_{n\to\infty} p_{i0}^{(n)} = f_{i0} = \frac{i}{2N}, \quad \lim_{n\to\infty} p_{i,2N}^{(n)} = f_{i,2N} = \frac{i}{2N}.$$

□

QUESTION 7.30 *Suppose $(M_t)_{t\ge0}$ is a $(\mathscr{F}_t)_{t\ge0}$ martingale. Given $0 \le r < u$ and $Z \in \mathscr{F}_r$, define*

$$N_t := Z(M_{u\wedge t} - M_{r\wedge t}).$$

Then $(N_t)_{t\ge0}$ is also a $(\mathscr{F}_t)_{t\ge0}$ martingale.

Proof. Observe for any $s, t > 0$, we have $\mathbb{P}(M_t|\mathscr{F}_s) = M_{s\wedge t}$ and $N_t = 0, \forall 0 \le t \le r$. Then for any $0 \le s < t$,

1. When $s < r$, we have

$$\begin{aligned}\mathbb{P}(N_t|\mathscr{F}_s) &= \mathbb{P}[Z(M_{u\wedge t} - M_{r\wedge t})|\mathscr{F}_s] \\ &= \mathbb{P}\{\mathbb{P}[Z(M_{u\wedge t} - M_{r\wedge t})|\mathscr{F}_r]\mathscr{F}_s\} \\ &= \mathbb{P}[Z\mathbb{P}(M_{u\wedge t} - M_{r\wedge t}|\mathscr{F}_r)\mathscr{F}_s] \\ &= \mathbb{P}[Z(M_{u\wedge t\wedge r} - M_{r\wedge t\wedge r})|\mathscr{F}_s] \\ &= \mathbb{P}[Z(M_{r\wedge t} - M_{r\wedge t})|\mathscr{F}_s] \\ &= 0 = N_s.\end{aligned}$$

2. When $s \geq r$, we have

$$\begin{aligned}\mathbb{P}(N_t|\mathscr{F}_s) &= \mathbb{P}[Z(M_{u\wedge t} - M_{r\wedge t})|\mathscr{F}_s] \\ &= Z\mathbb{P}(M_{u\wedge t} - M_{r\wedge t}|\mathscr{F}_s) \\ &= Z(M_{u\wedge t\wedge s} - M_{r\wedge t\wedge s}) \\ &= Z(M_{u\wedge t} - M_{r\wedge s}) \\ &= N_s.\end{aligned}$$

Thus the proof is completed. □

QUESTION 7.31 *Prove the minimum solution of the forward equation and the backward equation are identical.*

Proof. Consider the following expression of the forward equation and the backward equation:

$$\begin{aligned}x_{ij} &= \sum_{k\neq i} \frac{q_{ik}}{\lambda + q_i} x_{kj} + \frac{\delta_{ij}}{\lambda + q_i}, \\ y_{ij} &= \sum_{k\neq j} y_{ik} \frac{q_{kj}}{\lambda + q_j} + \frac{\delta_{ij}}{\lambda + q_j}.\end{aligned}$$

Consider the second iteration. Write the outcomes of the each step as $\{x_{ij}^{(n)}\}_{i,j\in E}$ and $\{y_{ij}^{(n)}\}_{i,j\in E}$ for the forward equation and the backward equation respectively. We use induction on n to prove this.

1. When $n = 1$, then

$$x_{ij}^{(1)} = \frac{\delta_{ij}}{\lambda + q_i} = \frac{\delta_{ij}}{\lambda + q_j} = y_{ij}^{(1)}.$$

2. When $n \le \ell$, assume the statement is correct. When $n = \ell + 1$, we have

$$\begin{aligned}
x_{ij}^{(\ell+1)} &= \sum_{k\neq i} \frac{q_{ik}}{\lambda + q_i} x_{kj}^{(\ell)} = \sum_{k\neq i} \frac{q_{ik}}{\lambda + q_i} y_{kj}^{(\ell)} \\
&= \sum_{k\neq i} \frac{q_{ik}}{\lambda + q_i} \sum_{s\neq j} y_{ks}^{(\ell-1)} \frac{q_{sj}}{\lambda + q_j} = \sum_{s\neq j}\sum_{k\neq i} \frac{q_{ik}}{\lambda + q_i} x_{ks}^{(\ell-1)} \frac{q_{sj}}{\lambda + q_j} \\
&= \sum_{s\neq j} x_{is}^{(\ell)} \frac{q_{sj}}{\lambda + q_j} = \sum_{s\neq j} y_{is}^{(k)} \frac{q_{sj}}{\lambda + q_j} \\
&= y_{ij}^{(\ell+1)}.
\end{aligned}$$

3. So $x_{ij}^{(n)} = y_{ij}^{(n)}$ for any $n \ge 1$.

Therefore

$$p_{ij}^{\min} = x_{ij} = \sum_{n\ge 1} x_{ij}^{(n)} = \sum_{n\ge 1} y_{ij}^{(n)}.$$

□

QUESTION 7.32 *Given a continuous process $(X_t)_{t\ge 0}$ on the probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and valued on metric space (E, d). Define the stopping time*

$$T_a := \inf\{t \ge 0 : X_t = a\}.$$

Then

$$\begin{aligned}
\{T_a \le t\} =& \{\exists 0 \le s \le t, X_s = a\} \\
=& \left\{\inf_{0\le s\le t} d(X_s, a) = 0\right\} \\
=& \left\{\inf_{s\in[0,t]\cap\mathbb{Q}} d(X_s, a) = 0\right\}
\end{aligned}$$

Proof. 1. We have

$$\{T_a \le t\} = \{T_a < t\} \cup \{T_a = t\} = \{\exists 0 \le s < t, X_s = a\} \cup \{T_a = t\}.$$

Since $T_a = t$ is equivalent to either of the following statement:

(a) $X_s \neq 0$ for any $0 \le s < t$ and $X_t = a$.

(b) there is a sequence $\{t_n\} \subset (t, \infty)$ such that $t_n \downarrow t$ and $X_{t_n} = a$ for any $n \ge 1$. Then the continuity of (X_t) implies

$$X_t = \lim_{n\to\infty} X_{t_n} = a.$$

Therefore $\{T_a = t\} \subset \{\forall 0 \le s < t, X_s \ne a, X_t = a\}$. Besides, another direction is obvious. So $\{T_a = t\} = \{\forall 0 \le s < t, X_s \ne a, X_t = a\}$ and it is easy to verify

$$\begin{aligned}&\{T_a \le t\}\\ =&\{\exists 0 \le s < t, X_s = a\} \cup \{\forall 0 \le s < t, X_s \ne a, X_t = a\}\\ =&\{\exists 0 \le s \le t, X_s = a\}.\end{aligned}$$

2. Now we start to prove the second equality. First,

$$\{\exists 0 \le s \le t, X_s = a\} \subset \left\{\inf_{0\le s\le t} d(X_s, a) = 0\right\}$$

is trivial. Ob the contrary, if $\inf_{0\le s\le t} d(X_s, a) = 0$, then we can find a sequence $\{s_n\} \subset [0, t]$ such that $d(X_{s_n}, a) \to 0$. Knowing $[0, t]$ is compact, then we can find a convergent subsequence $\{s_{n_k}\}$, say $s_{n_k} \to s_0 \in [0, t]$. Then the continuity of (X_t) and $d(\cdot, \cdot)$ yields

$$d(X_{s_0}, a) = \lim_{k\to\infty} d(X_{s_{n_k}}, a) = 0,$$

which means $X_{s_0} = a$. Therefore

$$\{\exists 0 \le s \le t, X_s = a\} \supset \left\{\inf_{0\le s\le t} d(X_s, a) = 0\right\}.$$

3. Finally, we started to prove the third equality. We only need to show

$$\left\{\inf_{0\le s\le t} d(X_s, a) = 0\right\} \subset \left\{\inf_{s\in[0,t]\cap\mathbb{Q}} d(X_s, a) = 0\right\}.$$

In fact, if $\inf_{0\le s\le t} d(X_s, a) = 0$, then we can find a sequence $\{s_n\} \subset [0, t]$ such that $d(X_{s_n}, a) \to 0$. Utilize the density of $\mathbb{Q}$ in $\mathbb{R}$ and the continuity of (X_t), for any $n \ge 1$, we can find $q_n \in \mathbb{Q}$ such that $d(X_{q_n}, X_{s_n}) < 1/n$. Thus

$$\begin{aligned}\inf_{s\in[0,t]\cap\mathbb{Q}} d(X_s, a) &\le \overline{\lim_{n\to\infty}}\, d(X_{q_n}, a)\\ &\le \overline{\lim_{n\to\infty}}\, d(X_{q_n}, X_{s_n}) + \overline{\lim_{n\to\infty}}\, d(X_{s_n}, a) = 0.\end{aligned}$$

Thus the whole statement is completed. □

QUESTION 7.33 *Consider a Markov chain $(X_n)_{n\ge0}$ on probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and valued on countable state space E. For each $a \in E$, consider the following stopping times*

$$T_a = \inf\{n \ge 0 : X_n = a\}, \quad T_a^+ = \inf\{n \ge 1 : X_n = a\}.$$

Then

$$T_a^+(\omega) = \begin{cases} (T_a \circ \theta_1)(\omega) + 1 = (T_a^+ \circ \theta_1)(\omega) + 1 & , \textit{if } \omega \in \{X_1 \neq a\}, \\ (T_a \circ \theta_1)(\omega) + 1 & , \textit{if } \omega \in \{X_1 = a\}. \end{cases}$$

Moreover, if additionally $\omega \in \{X_0 \neq a\}$, then $T_a^+(\omega) = T_a(\omega)$. Hence T_a also has the above expression.

Proof. Since if $\omega \in \{X_1 \neq a\}$, then obviously $(T_a \circ \theta_1)(\omega) = (T_a^+ \circ \theta_1)(\omega)$. Then we just need to show for any $\forall \omega \in \Omega$,

$$T_a^+(\omega) = (T_a \circ \theta_1)(\omega) + 1.$$

If $\omega \in \{X_1 = a\}$, then $T_a^+(\omega) = 1$ and $(T_a \circ \theta_1)(\omega) = 0$. So the above formula holds. If $\omega \in \{X_1 \neq a\}$, by the definition of shift operator θ_1, we have the following equality

$$T_a^+(\omega) - (T_a^+ \circ \theta_1)(\omega) = 1.$$

Moreover, we know $(T_a \circ \theta_1)(\omega) = (T_a^+ \circ \theta_1)(\omega), \forall \omega \in \{X_1 \neq a\}$. So the proof is completed. □

QUESTION 7.34 *Suppose Q is a Q matrix on the state space E and $\sup_{i \in E} |q_i| < \infty$. Then we have the following estimation:*

$$\sup_{i,j \in E} |q_{ij}^{(n)}| \leq \frac{1}{2} \left(2 \sup_{i \in E} |q_i| \right)^n, \quad \forall n \geq 1.$$

Especially, when $|E| = m < \infty$, for any $A = (a_{ij}) \in M_m(\mathbb{R})$, define $||A|| := m \sup_{i,j \in E} |a_{ij}|$, then $(M_m(\mathbb{R}), ||\cdot||)$ is a Banach algebra.

Proof. We use induction on n to prove the first estimation.

1. When $n = 1$, obviously $|q_{ij}| \leq \sup_{i \in E} |q_i|$.
2. Suppose $n = k$, the estimation is correct. When $n = k + 1$, we have

$$\begin{aligned} |q_{ij}^{(k+1)}| = \left| \sum_k q_{ik} q_{kj}^{(k)} \right| &\leq \sum_{k \neq i} q_{ik} |q_{kj}^{(k)}| + q_i |q_{ij}^{(k)}| \\ &\leq \frac{1}{2} \left(2 \sup_{i \in E} |q_i| \right)^k \left(\sum_{k \neq i} q_{ik} + q_i \right) \\ &= \frac{1}{2} \left(2 \sup_{i \in E} |q_i| \right)^k 2 q_i \\ &\leq \frac{1}{2} \left(2 \sup_{i \in E} |q_i| \right)^{k+1}. \end{aligned}$$

3. In conclusion, the estimation is correct for all $n \geq 1$.

Furthermore, we can infer from this conclusion that e^{tQ} is well-defined.

Obviously $||\cdot||$ is a norm. For any $A = (a_{ij}), B = (b_{ij}) \in M_n(\mathbb{R})$, we have

$$\begin{aligned} ||AB|| &= m \sup_{i,j\in E} \left| \sum_{k\in E} a_{ik} b_{kj} \right| \leq m \sum_{k\in E} \sup_{i,j\in E} a_{ij} \sup_{i,j\in E} b_{ij} \\ &= m^2 \sup_{i,j\in E} a_{ij} \sup_{i,j\in E} b_{ij} = ||A|| \cdot ||B||. \end{aligned}$$

Thus $(M_m(\mathbb{R}), ||\cdot||)$ is a Banach algebra. Moreover, it provide us a less strict estimation for $|q_{ij}^{(n)}|$ when $|E| = m < \infty$. In fact,

$$m \sup_{i,j\in E} |q_{ij}^{(n)}| = ||Q^n|| \leq ||Q||^n = \left(m \sup_{i,j\in E} q_{ij} \right)^n,$$

which is equivalent to

$$\sup_{i,j\in E} |q_{ij}^{(n)}| \leq \frac{1}{m} \left(m \sup_{i,j\in E} q_{ij} \right)^n.$$

□

QUESTION 7.35 *Consider a symmetry matrix $A \in M_n(\mathbb{R})$ and write all its eigenpairs as $(\lambda_j, x_j)_{j=1}^n$. Then*

$$A = \sum_{j=1}^n \lambda_j x_j x_j^{\mathrm{T}} = \sum_{j=1}^n \lambda_j \langle \cdot, x_j \rangle x_j.$$

Proof. Since $A \in M_n(\mathbb{R})$ is symmetry, then indeed it has n eigenvalues which form a unit standard orthogonal basis, denoted as $\{x_j\}_{j=1}^n$ (column vectors). And we write their corresponding eigenvalues as $\{\lambda_j\}_{j=1}^n$ (note some eigenvalues in $\{\lambda_j\}$ might be repeated). Then exists a orthogonal matrix P such that

$$PAP^{\mathrm{T}} = \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix},$$

where $P = \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix}$. Therefore

$$
\begin{aligned}
A &= P^{\mathrm{T}} \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix} P \\
(*) \qquad &= \begin{pmatrix} x_1^{\mathrm{T}} \\ x_2^{\mathrm{T}} \\ \vdots \\ x_n^{\mathrm{T}} \end{pmatrix} \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix} \begin{pmatrix} x_1 & x_2 & \cdots & x_n \end{pmatrix} \\
&= \sum_{j=1}^{n} \lambda_j x_j x_j^{\mathrm{T}}.
\end{aligned}
$$

Moreover, since $\{x_j\}_{j=1}^n$ is a unit standard orthogonal basis on $\mathbb{R}^n$, then for any $y \in \mathbb{R}^n$, exists $\{y_j\}_{j=1}^n$ such that

$$
y = \sum_{j=1}^{n} y_j x_j.
$$

Therefore

$$
Ay = \sum_{j=1}^{n} y_j A x_j = \sum_{j=1}^{n} \lambda_j y_j x_j = \sum_{j=1}^{n} \lambda_j \langle y, x_j \rangle x_j,
$$

which is equivalent to

$$
(**) \qquad A = \sum_{j=1}^{n} \lambda_j \langle \cdot, x_j \rangle x_j.
$$

Compare $(*)$ and $(**)$, we can concludes that

$$
x_j x_j^{\mathrm{T}} = \langle \cdot, x_j \rangle x_j.
$$

□

QUESTION 7.36 *Suppose $N = (N_t, t \geq 0)$ is a Poisson process with parameter $\lambda > 0$, then the following statement is true:*

1. *N is increasing.*
2. *For any $t \geq 0$, there exists $\delta > 0$ such that N is constant in the interval $(t, t + \delta]$.*
3. *Define stopping times recursively by $T_0 = 0$ and $T_n = \inf\{t > T_{n-1} : N_t - N_{T_{n-1}}\}$ for $n \geq 1$. Then $N_{T_n} = n$ for $n \in \mathbb{Z}_+$.*

4. $(\Delta N(t), t \geq 0)$ *takes values in* $\{0, 1\}$ *and for any* $t \geq 0$.

Proof. 1. For any $0 \leq s < t$, we have $N_t = N_s + (N_t - N_s)$ and $N_t - N_s \sim P(\lambda(t-s))$. So $N_t \geq N_s$ and therefore N is increasing.

2. If there exists $t \geq 0$ satisfying for any $\delta > 0$, N_t is not constant in the interval $(t, t+\delta]$. Note N is increasing, then there exist $t < t_1 < t+\delta$ satisfying $N_{t+\delta} > N_{t_1}$. Since $N_{t+\delta} - N_{t_1}$ is poisson distributed, then $N_{t+\delta} - N_{t_1} \geq 1$. Consider $(t, t+t_1]$, since N is not constant in $(t, t+t_1]$, we can find $t < t_2 < t_1$ and $N_{t_1} - N_{t_2} \geq 1$. Do this repeatedly, we will obtain a sequence $\{t_n\}_{n\geq 1}$ satisfying

$$t < \cdots < t_n < t_{n-1} < \cdots < t_1 < t_0 := t+\delta \text{ and } N_{t_n} - N_{t_{n+1}} \geq 1.$$

So for any $n \geq 1$,

$$N_{t+\delta} - N_t \geq \sum_{k=0}^{n} (N_{t_k} - N_{t_{k+1}}) \geq n.$$

Observe n is arbitrary, then $N_{t+\delta} - N_t = \infty$. However we know $N_{t+\delta} - N_t$ is Poisson distributed, so $\mathbb{P}(N_{t+\delta} - N_t = \infty) = 0$.

3. First we show $\Delta N_t \in \mathbb{Z}_+$. Apply the increasing property of N, we have

$$\Delta N_t = N_t - N_{t-} = \lim_{n\to\infty} (N_t - N_{t-1/n}).$$

Observe $N_t - N_{t-1/n} \sim P(\lambda/n)$, then $N_t - N_{t-1/n}$ is integer valued and therefore ΔN_t is integer valued. □

QUESTION 7.37 *Suppose* $(X_n)_{n\geq 1}$ *is an irreducible Markov chain on a countable state space* E.

1. *For* $x \neq y \in E$, *if there exists* $n \geq 1$ *satisfies* $p_{ij}^{(n)} > 0$, *then there exists* $m \leq n$ *such that* $\mathbb{P}_i(X_m = j, \tau_i^+ > m) > 0$.

2. *Apply this property to show: if* $f_{ii} = 1$ *for any* $i \in E$, *then* $f_{ij} = 1$ *for any* $i, j \in E$.

Proof. 1. If $\mathbb{P}_i(X_n = j, \tau_i^+ > n) > 0$, then all is done. If $\mathbb{P}(X_n = j, \tau_i^+ > n) = 0$, then $\mathbb{P}_i(X_n = j) = \mathbb{P}_i(X_n = j, \tau_i^+ \leq n)$ and

$$\begin{aligned}
\mathbb{P}_i(X_n = j, \tau_i^+ \leq n) &= \mathbb{P}_i(X_n = j, \tau_i^+ < n) \\
&= \sum_{k=1}^{n-1} \mathbb{P}_i(X_n = j, \tau_i^+ = k) = \sum_{k=1}^{n-1} \mathbb{P}_i(X_n = j | \tau_i^+ = k)\mathbb{P}_i(\tau_i^+ = k) \\
&= \sum_{k=1}^{n-1} \mathbb{P}_i(X_{n-k} = j)\mathbb{P}_i(\tau_i^+ = k) > 0.
\end{aligned}$$

So there exists $n_1 < n$ such that $p_{ij}^{(n_1)} = \mathbb{P}_i(X_{n_1} = j) > 0$. Again if $\mathbb{P}_i(X_{n_1} = j, \tau_i^+ > n_1) > 0$, then all is done. If $\mathbb{P}_i(X_{n_1} = j, \tau_i^+ > n_1) = 0$, we can repeat the above procedure to find there $n_2 < n_1$ such that $p_{ij}^{(n_2)} = \mathbb{P}_i(X_{n_2} = j) > 0$. If there exists $1 < m < n$ such that $\mathbb{P}_i(X_m = j, \tau_i^+ > m) = 0$, then all is done. If not, we will find

$$\mathbb{P}_i(X_1 = j) = \mathbb{P}_i(X_1 = j, \tau_i^+ > 1) > 0.$$

2. If exists $i \neq j \in E$ such that $f_{ij} < 1$, then $\mathbb{P}_i(\tau_j^+ = \infty) > 0$. Since $(X_n)_{n\geq 1}$ is irreducible, then there exists $n \geq 1$ satisfies $p_{ji}^{(n)} > 0$. So what we have proved yields $\mathbb{P}_j(X_m = i, \tau_j^+ > m) > 0$. Thus

$$\begin{aligned}
\mathbb{P}_j(\tau_j^+ = \infty) &= \sum_{k\in E} \mathbb{P}_j(X_m = k, \tau_j^+ > m, \tau_j^+ = \infty) \\
&\geq \mathbb{P}_j(X_m = i, \tau_j^+ > m, \tau_j^+ = \infty) \\
&= \mathbb{P}_j(X_m = i, \tau_j^+ > m, \tau_j^+ \circ \theta_m + m = \infty) \\
&= \mathbb{P}_j(\tau_j^+ \circ \theta_m + m = \infty | X_m = i, \tau_j^+ > m)\mathbb{P}_j(X_m = i, \tau_j^+ > m) \\
&= \mathbb{P}_j(\tau_j^+ \circ \theta_m = \infty | X_m = i, \tau_j^+ > m)\mathbb{P}_j(X_m = i, \tau_j^+ > m) \\
&= \mathbb{P}_i(\tau_j^+ = \infty)\mathbb{P}_j(X_m = i, \tau_j^+ > m) > 0.
\end{aligned}$$

This contradict the fact $f_{jj} = 1$. □

QUESTION 7.38 *Given a nonnegative supermartingale $(X_n)_{n\geq 0}$ with respect to $(\mathscr{F}_n)_{n\geq 1}$. Then $X_\infty := \lim_{n\to\infty} X_n$ exists almost surely. Moreover we have $\mathbb{P}(X_\infty|\mathscr{F}_n) \leq X_n$ for any $n \geq 0$ and $\{X_\infty < \infty\} \subset \{X_0 < \infty\}$ almost surely.*

Proof. $X_\infty := \lim_{n\to\infty} X_n$ exists almost surely is a well-known result. To show $\mathbb{P}(X_\infty|\mathscr{F}_n) \leq X_n$, we use the nonnegativity of $(X_n)_{n\geq 1}$ and the Fatou's lemma. For any $A \in \mathscr{F}_n$ and $m \geq n$,

$$\mathbb{P}(X_\infty 1_A) \leq \varliminf_{m\to\infty} \mathbb{P}(X_m 1_A) \leq \varliminf_{m\to\infty} \mathbb{P}(X_n 1_A) = \mathbb{P}(X_n 1_A).$$

This yields $\mathbb{P}(X_\infty|\mathscr{F}_n) \leq X_n$. Let $n = 0$ and $A = \{X_0 \leq M\}$, where $M > 0$. Then

$$\mathbb{P}(X_\infty \mathbb{1}_{\{X_0\leq M\}}) \leq \mathbb{P}(X_0 \mathbb{1}_{\{X_n\leq M\}}) \leq M < \infty.$$

This means $\{X_\infty = \infty\} \cap \{X_0 \leq M\} = \emptyset, \mathbb{P}$-a.s. for any $M > 0$. Hence □

$$\{X_\infty = \infty\} \subset \bigcap_{M=1}^{\infty} \{X_0 > M\} = \{X_0 = \infty\}, \quad \mathbb{P}\text{-a.s.}.$$

QUESTION 7.39 *Suppose $(\xi_n, n \geq 0)$ is a discrete-time Markov process with transition kernel $P(x, dy)$ and initial distribution $\mu(dx)$ on $(E, \mathscr{E})$. Besides, $(N_t, t \geq 0)$ is an independent Poisson process with parameter $\alpha > 0$. Let $X_t := \xi_{N_t}$. Then the continuous-time process $(X_t, t \geq 0)$ is a Markov process with transition kernel as follows:*

$$Q_t(x, A) := e^{-\alpha t} \sum_{k \geq 0} \frac{(\alpha t)^k}{k!} P^k(x, A), \quad A \in \mathscr{E}.$$

Proof. We just need to show that

$$\mathbb{P}[f(X_{s+t}) | X_{s_1}, \cdots, X_{s_n}] = Q_t f(X_s)$$

for any $n \geq 1, t \geq 0, 0 = s_0 \leq s_1 \leq \cdots \leq s_n = s$ and $f \in \mathrm{b}\mathscr{E}$. And this is again equivalent to showing

$$\mathbb{P}[f(X_{s+t}) g(X_{s_1}, \cdots, X_{s_n})] = \mathbb{P}[Q_t f(X_s) g(X_{s_1}, \cdots, X_{s_n})]$$

for any $n \geq 1, t \geq 0, 0 = s_0 \leq s_1 \leq \cdots \leq s_n = s$ and $g \in \mathrm{b}\mathscr{E}^n$. First note by the independency of $(\xi_n, n \geq 0)$ and $(N_t, t \geq 0)$,

$$\begin{aligned}
&\mathbb{P}[g(X_{s_1}, \cdots, X_{s_n})] = \mathbb{P}[g(\xi_{N_{s_1}}, \cdots, \xi_{N_{s_n}})] \\
&= \sum_{k_1, \cdots, k_n \geq 0} \mathbb{P}\left[g(\xi_{N_{s_1}}, \cdots, \xi_{N_{s_n}}); N_{s_r} - N_{s_{r-1}} = k_r, 1 \leq r \leq n\right] \\
&= \sum_{k_1, \cdots, k_n \geq 0} \mathbb{P}\left[g(\xi_{k_1}, \cdots, \xi_{k_1 + \cdots + k_n}); N_{s_r} - N_{s_{r-1}} = k_r, 1 \leq r \leq n\right] \\
&= \sum_{k_1, \cdots, k_n \geq 0} \mathbb{P}[g(\xi_{k_1}, \cdots, \xi_{k_1 + \cdots + k_n})] \mathbb{P}(N_{s_r} - N_{s_{r-1}} = k_r, 1 \leq r \leq n) \\
&= \sum_{k_1, \cdots, k_n \geq 0} \int_{E^n} g(x_1, \cdots, x_n) \mu P_{J_n}(dx_1, \cdots, dx_n) \mathbb{P}(N_{s_r} - N_{s_{r-1}} = k_r, 1 \leq r \leq n),
\end{aligned}$$

where $J_n = \{k_1, k_1 + k_2, \cdots, k_1 + \cdots + k_n\}$ and

$$P_{J_n}(dx_1, \cdots, dx_n) = P_{k_1}(x, dx_1) P_{k_2}(x_1, dx_2) \cdots P_{k_n}(x_{n-1}, dx_n).$$

Thus let $s_{n+1} = s + t$, we have

$$\begin{aligned}
&\mathbb{P}[f(X_{s+t}) g(X_{s_1}, \cdots, X_{s_n})] \\
&= \sum_{k_1, \cdots, k_n, k_{n+1} \geq 0} \int_{E^{n+1}} f(x_{n+1}) g(x_1, \cdots, x_n) \mu P_{J_{n+1}}(dx_1, \cdots, dx_{n+1}) \\
&\qquad \times \mathbb{P}(N_{s_r} - N_{s_{r-1}} = k_r, 1 \leq r \leq n+1) \\
&= \sum_{k_1, \cdots, k_n, k_{n+1} \geq 0} \int_{E^n} P_{k_{n+1}} f(x_n) g(x_1, \cdots, x_n) \mu P_{J_n}(dx_1, \cdots, dx_n)
\end{aligned}$$

$$
\begin{aligned}
&\times \mathbb{P}(N_{s_r} - N_{s_{r-1}} = k_r, 1 \le r \le n)\mathbb{P}(N_{s_{n+1}} - N_{s_n} = k_{n+1}) \\
= &\sum_{k_1,\cdots,k_n \ge 0} \int_{E^n} \Big[\sum_{k_{n+1} \ge 0} \mathbb{P}(N_{s_{n+1}} - N_{s_n} = k_{n+1}) P_{k_{n+1}} f(x_n) \\
&\times g(x_1, \cdots, x_n) \mu P_{J_n}(dx_1, \cdots, dx_n) \Big] \mathbb{P}(N_{s_r} - N_{s_{r-1}} = k_r, 1 \le r \le n) \\
= &\sum_{k_1,\cdots,k_n \ge 0} \int_{E^n} e^{-\alpha t} \sum_{k \ge 0} \frac{(\alpha t)^k}{k!} P_k f(x_n) g(x_1, \cdots, x_n) \mu P_{J_n}(dx_1, \cdots, dx_n) \\
&\times \mathbb{P}(N_{s_r} - N_{s_{r-1}} = k_r, 1 \le r \le n) \\
= &\mathbb{P}[Q_t f(X_s) g(X_{s_1}, \cdots, X_{s_n})].
\end{aligned}
$$

Till now, we have completed the proof. □

QUESTION 7.40 *For a $\mathbb{R}^d$-valued Lévy process $(\Omega, \mathscr{F}, \mathscr{F}_t, X_t, t \ge 0)$, write its transition and convolution semigroups as $(P_t)_{t \ge 0}$ and $(\mu_t)_{t \ge 0}$ respectively. Then the following statements are both equivalent: for any $(\mathscr{F}_t)_{t \ge 0}$ stopping time T and $f \in \mathrm{b}\mathscr{B}(\mathbb{R}^d)$,*

1. $\mathbb{P}[f(X_{T+t}) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T] = P_t f(X_T) \mathbb{1}_{\{T<\infty\}}$.
2. $\mathbb{P}[f(X_{T+t} - X_T) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T] = P_t f(0) \mathbb{1}_{\{T<\infty\}}$.

Proof. If the first formula is satisfied, write $g_y(x) := f(x-y)$, then

$$
\begin{aligned}
&\mathbb{P}[f(X_{T+t} - X_T) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T] \\
&= \mathbb{P}[f(X_{T+t} - y) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T]|_{y=X_T} = \mathbb{P}[g_y(X_{T+t}) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T]|_{y=X_T} \\
&= P_t g_y(X_T) \mathbb{1}_{\{T<\infty\}}|_{y=X_T} = \mathbb{1}_{\{T<\infty\}} \int g_y(x) P_t(X_T, dx) \Big|_{y=X_T} \\
&= \mathbb{1}_{\{T<\infty\}} \int g_y(X_T + x) \mu_t(dx) \Big|_{y=X_T} \\
&= \mathbb{1}_{\{T<\infty\}} \int f(X_T + x - y) \mu_t(dx) \Big|_{y=X_T} = \mathbb{1}_{\{T<\infty\}} \int f(x) \mu_t(dx) \\
&= \mathbb{1}_{\{T<\infty\}} \int f(x) P_t(0, dx) = P_t f(0) \mathbb{1}_{\{T<\infty\}}.
\end{aligned}
$$

On the contrary, if the second formula is satisfied, write $h_y(x) := f(x+y)$, then

$$
\begin{aligned}
&\mathbb{P}[f(X_{T+t}) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T] \\
&= \mathbb{P}[f(X_{T+t} - X_T + X_T) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T] \\
&= \mathbb{P}[f(X_{T+t} - X_T + y) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T]|_{y=X_T} \\
&= \mathbb{P}[h_y(X_{T+t} - X_T) \mathbb{1}_{\{T<\infty\}} | \mathscr{F}_T]|_{y=X_T} = P_t h_y(0) \mathbb{1}_{\{T<\infty\}}|_{y=X_T} \\
&= \mathbb{1}_{\{T<\infty\}} \int h_y(x) P_t(0, dx) \Big|_{y=X_T} = \mathbb{1}_{\{T<\infty\}} \int f(x + X_T) \mu_t(dx) \\
&= \mathbb{1}_{\{T<\infty\}} \int f(x) P_t(X_T, dx) = P_t f(X_T) \mathbb{1}_{\{T<\infty\}}.
\end{aligned}
$$

□

§ 8 Functional Analysis

QUESTION 8.1 *Suppose (M, d) is a compact metric space, then $C(M)$ with the supremum norm $||\cdot||$ is a Polish space.*

Proof. Obviously, $C(M)$ is a Banach space, so we just need to verify $C(M)$ is separable. In fact, since M is a compact metric space, then M is totally bounded and therefore is separable. Since M is a separable metric space, then M is a C_2 topology space, namely, M has a countable topology basis, denoted as $\{U_n\}$. Consider $f_n(x) = d(x, U_n)$, we claim $\{f_n\} \subset C(M)$ can separate points in M. In fact, note M is a Hausdorff topology space, then for each $x, y \in M$, exists $U_m, U_k \in \{U_n\}$ such that $x \in U_m, y \in U_k$ and $U_m \cap U_k = \emptyset$. Therefore $f_k(x) = d(x, U_k) = 0$ and $f_k(y) = d(y, U_k) > 0$. So we have proved $\{f_n\}$ can separate points in M. Let

$$\mathscr{G} := \left\{ \sum_{n=1}^{m} q_n f_n : m \in \mathbb{N}, q_n \in \mathbb{Q}, f_n \in \{f_n\} \right\}.$$

Next, we define

$$\mathscr{F} := \left\{ \prod_{n=1}^{m} g_n : m \in \mathbb{N}, g_n \in \mathscr{G} \right\} \cup \{1\},$$

where 1 denoted $f(x) \equiv 1$. We can find $\mathscr{F}$ is countable. Then $\bar{\mathscr{F}}$ has the unit element 1 and can separate points in E, where the closure is in the sense of $||\cdot||$. Moreover, it can be found that $\bar{\mathscr{F}}$ is an algebra. The Stone-Weirestrass theorem yields $\overline{\mathscr{F}} = C(M)$. So the assertion is proved. □

QUESTION 8.2 *Suppose N is a normal operator on Hilbert space H. For all $\Psi \in B(\sigma(N))$, define $\Psi(N)$ as follows: for every $x, y \in H$,*

$$\langle \Psi(N)x, y \rangle = \int_{\sigma(N)} \Phi(z) m_{x,y}(dz),$$

where $m_{x,y}(dz)$ is the spectral measure of N corresponding to x, y. Moreover, suppose $(\mathbb{C}, \mathscr{B}(\mathbb{C}), E)$ is the spectral family of N, and $\Phi : B_E(\mathbb{C}) \to \mathscr{N}(H)$ is the $*$-isometry corresponding to the spectral family $(\mathbb{C}, \mathscr{B}(\mathbb{C}), E)$. Then

$$\Phi|_{B(\sigma(N))} : \Psi \mapsto \Psi(N).$$

Proof. For all $\Psi \in B(\sigma(N))$, define

$$\widetilde{\Psi}(z) := \begin{cases} \Psi(z) & \text{, if } z \in \sigma(N) \\ 0 & \text{, if } z \notin \sigma(N) \end{cases}.$$

Since the support of E is $\sigma(N)$, then $\forall f \in B(\mathbb{C}), f = \widetilde{f|_{\sigma(N)}}, E-a.e.$. As a result, $\forall \Psi \in B(\sigma(N))$, we identify it with $\widetilde{\Psi} \in B(\mathbb{C})$. When

$$\Psi = \sum_{k=1}^{n} \alpha_k \mathbf{1}_{A_k} \in \mathscr{S}(\sigma(N)), \quad \alpha_k \subset \mathbb{C}, A_k \in \mathscr{B}(\mathbb{C}) \cap \sigma(N),$$

For every $x, y \in H$, observe $(\sigma(N), \mathscr{B}(\mathbb{C}) \cap \sigma(N), m_{x,y}) = (\sigma(N), \mathscr{B}(\mathbb{C}) \cap \sigma(N), E)$, namely, $m_{x,y} = E|_{\mathscr{B}(\mathbb{C})\cap\sigma(N)}$ and use the definition of Φ in [1, Theorem 2.5.29], then

$$\begin{aligned}
\langle \Psi(N)x, y\rangle &= \int_{\sigma(N)} \Psi(z) m_{x,y}(dz) = \int_{\sigma(N)} \Psi(z) d\langle E(z)x, y\rangle \\
&= \int_{\sigma(N)} \sum_{k=1}^{n} \alpha_k \mathbf{1}_{A_k} d\langle E(z)x, y\rangle = \sum_{k=1}^{n} \alpha_k \int_{\sigma(N)} \mathbf{1}_{A_k} d\langle E(z)x, y\rangle \\
&= \sum_{k=1}^{n} \alpha_k \langle E(A_k)x, y\rangle = \left\langle \sum_{k=1}^{n} \alpha_k E(A_k)x, y \right\rangle \\
&= \langle \Phi(\Psi)x, y\rangle .
\end{aligned}$$

Since this formula holds for every $x, y \in H$, then $\Phi(\Psi) = \Psi(N), \forall \Psi \in \mathscr{S}(\sigma(N))$. When $\Psi \in B(\sigma(N))$, exist $\{\Psi_n\} \subset \mathscr{S}(\sigma(N))$, s.t. $\Psi_n \to \Psi$ and $\exists M > 0$, s.t. $||\Psi_n|| \leq M$. By the definition of Φ in [1, Theorem 2.5.29], then $\Phi(\Psi) := \lim_{n\to\infty} \Phi(\Psi_n)$. By [1, Theorem 2.5.20 (iii)], we have

$$\langle \Psi(N)x, y\rangle = \lim_{n\to\infty} \langle \Psi_n(N)x, y\rangle = \lim_{n\to\infty} \langle \Phi(\Psi_n)x, y\rangle = \langle \Phi(\Psi)x, y\rangle.$$

Since the above formula holds for every $x, y \in H$, then $\Phi(\Psi) = \Psi(N), \forall \Psi \in B(\sigma(N))$. Namely, we have proved

$$\Phi|_{B(\sigma(N))} : \Psi \mapsto \Psi(N).$$

□

QUESTION 8.3 *Suppose N is a normal operator on Hilbert space H. If $\varphi \in C(\sigma(N)), \Psi \in B(\varphi(\sigma(N)))$, then*

$$(\Psi \circ \varphi)(N) = \Psi(\varphi(N)).$$

Proof. We have proved: $\forall \varphi \in C(\sigma(N)), \Psi \in C(\varphi(\sigma(N)))$, then $(\Psi \circ \varphi)(N) = \Psi(\varphi(N))$. When $\Psi \in B(\varphi(\sigma(N)))$, we can find a sequence $\{\Psi_n\} \in C(\varphi(\sigma(N)))$, s.t. $\Psi_n \to \Psi, n \to \infty$ and $\exists M > 0$, s.t. $||\Psi_n|| \leq M$. Hence $\Psi_n \circ \varphi \to \Psi \circ \varphi, ||\Psi_n \circ \varphi|| \leq M$. Notice $\varphi(N)$ is also a normal operator, by [1, Theorem 2.5.20 (iii)], $\forall x \in H$,

$$\begin{aligned}
(\Psi \circ \varphi)(N)x &= \lim_{n\to\infty} (\Psi_n \circ \varphi)(N)x = \lim_{n\to\infty} \Psi_n(\varphi(N))x \\
&= \lim_{n\to\infty} \Psi_n(\varphi(N))x = \Psi(\varphi(N))x,
\end{aligned}$$

which means $(\Psi \circ \varphi)(N) = \Psi(\varphi(N))$. □

QUESTION 8.4 *Suppose $\mathscr{X}$ is a Banach space and $T : \mathscr{X} \supset \mathscr{D}(T) \to \mathscr{X}$ is a closed operator. For every $\lambda \in \sigma_c(T)$, we have $(\lambda I - T)^{-1}$ is not bounded.*

Proof. Suppose $\exists \lambda \in \sigma_c(T)$, s.t. $(\lambda I - T)^{-1}$ is bounded. By the definition of $\sigma_c(T)$, we have $R(\lambda I - T) \neq \mathscr{X}$ and $\overline{R(\lambda I - T)} = \mathscr{X}$. Hence $\forall y \in \mathscr{X}, \exists \{y_n\} \subset R(\lambda I - T)$, such that $y_n \to y, n \to \infty$. Write $x_n := (\lambda I - T)^{-1} y_n$. Since $\{y_n\}$ is a Cauchy sequence and $(\lambda I - T)^{-1}$ is bounded, $\{x_n\}$ is also a Cauchy sequence. Because $\mathscr{X}$ is complete, then $\exists x \in \mathscr{X}$, s.t. $x_n \to x, n \to \infty$. Till now, we have

$$x_n \to x, (\lambda I - T)x_n = y_n \to y, n \to \infty$$

Moreover, $\lambda I - T$ is also a closed operator, whence $y = (\lambda I - T)x \in R(\lambda I - T)$. This implies $\mathscr{X} = R(\lambda I - T)$, which contradicts to the fact that $R(\lambda I - T) \neq \mathscr{X}$. □

QUESTION 8.5 *Suppose H is a Hilbert space and $u_k \rightharpoonup u$. Then $||u|| \leq \underline{\lim}_{k\to\infty} ||u_k||$.*

Proof. Since for each $f \in H^*$, we have

$$|f(x)| = \lim_{k\to\infty} |f(x_k)| \leq ||f|| \varliminf_{k\to\infty} ||x_k||.$$

Moreover, we have

$$||x|| = \sup_{||f||=1} |f(x)| \leq \sup_{||f||=1} \left(||f|| \varliminf_{k\to\infty} ||x_k|| \right) = \varliminf_{k\to\infty} ||x_k||.$$

This is what we want. □

QUESTION 8.6 *Given a Banach space $(B, ||\cdot||)$ and a separable subset $E \subset B$. Write C for its countable dense subset of E and $\widetilde{C}$ to be the set of rational multiples of elements of C. Then for any $x \in E$ and $\varepsilon > 0$ there exists $y \in \widetilde{C}$ satisfying $||y|| \leq ||x||$ and $||y - x|| < \varepsilon$.*

Proof. For the given $x \in E$ and $0 < \delta < \varepsilon/2$, there exists $y' \in C$ such that $||y' - x|| < \delta$. If $||y'|| \leq ||x||$, then we let $y := y'$ and all we need is completed. Otherwise, if $||y'|| > ||x||$, consider a rational $r \in \mathbb{Q} \cap [0, 1]$, then we have the following estimation

$$\begin{aligned} ||ry' - x|| &\leq ||ry' - rx|| + ||rx - x|| \\ &= |r|\, ||y' - x|| + (1 - r)||x|| \\ &\leq \delta + (1 - r)||x||. \end{aligned}$$

Let $(1-r)||x|| < \varepsilon/2$, then we have $r > 1-\varepsilon/(2||x||)$. Choose a $r \in (1-\varepsilon/(2||x||), 1) \cap \mathbb{Q}$, and define $y := ry' \in \widetilde{C}$. Hence the above estimation yields

$$||y-x|| \leq \delta + (1-r)||x|| < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

□

QUESTION 8.7 *Any finite-dimensional metric space (E, d) is compact.*

Proof. Since every norm on a finite-dimensional space is equivalent, everything boils down to the finite-dimensional Euclidean space. Obviously, it is compact.□

QUESTION 8.8 *Any finite-dimensional subspace of a metric space (E, d) is closed.*

Proof. Let $A \subset E$ be a finite-dimensional subspace with dimension $m < \infty$ and basis $\{x_k\}_{k=1}^n$. Any element $x \in A$ can be uniquely expressed as $x = \sum_{k=1}^m \alpha_k x_k$. Given a convergent sequence $\{x^n\}_{n\in\mathbb{Z}_+}$, say

$$x^n = \sum_{k=1}^m \alpha_k^n x_k \to x \in E, \quad n \to \infty,$$

we want to prove $x \in A$. Firstly, the convergence of $\{x^n\}_{n\in\mathbb{Z}_+}$ ensures that $\{x^n\}_{n\in\mathbb{Z}_+}$ is bounded. The above QUESTION shows that A is compact. So, there exists a subsequence $\{x^{n_\ell}\}_{\ell\in\mathbb{Z}_+}$ such that $x^{n_\ell} \to y \in A, \ell \to \infty$. Note $x^{n_\ell} \to x \in E, \ell \to \infty$, the uniqueness of the limit yields that $x = y \in A$. In conclusion, A is closed. □

QUESTION 8.9 *Given two subspaces V, W of the Hilbert space $\mathscr{X}$. Suppose $V \perp W$, then $\overline{V \oplus W} = \overline{V} \oplus \overline{W}$. Besides, give an example to show $\overline{V \oplus W} = \overline{V} \oplus \overline{W}$ is not always true if $V \not\perp W$.*

Proof. Since $V \perp W$, the notation $V \oplus W$ is reasonable.

Given a convergent sequence $x_n = v_n + w_n, n \geq 1$, where $(v_n)_{n\geq 1} \subset V, (w_n)_{n\geq 1} \subset W$ and $x_n \to x \in \mathscr{X}$, since $\overline{V}$ and $\overline{W}$ are both closed subspace of $\mathscr{X}$, we can get the following decomposition:

$$x = P_{\overline{V}} x + P_{\overline{V}^\perp} x,$$

where $P_{\overline{V}}$ is the projection operator of $\overline{V}$. Since $V \perp W$, the continuity of the inner product yields $\overline{V} \perp \overline{W}$. Therefore $\overline{W} \subset \overline{V}^\perp$. Since $v_n - P_{\overline{V}} x \in \overline{V}$ and $w_n - P_{\overline{V}^\perp} x \in \overline{V}^\perp$, we have

$$\begin{aligned} ||x_n - x|| &= ||(v_n - P_{\overline{V}} x) + (w_n - P_{\overline{V}^\perp} x)|| \\ &= ||v_n - P_{\overline{V}} x|| + ||w_n - P_{\overline{V}^\perp} x|| \to 0, \quad n \to \infty. \end{aligned}$$

This is equivalent to

$$||v_n - P_{\overline{V}}x|| \to 0, \quad ||w_n - P_{\overline{V}^\perp}x|| \to 0, \quad n \to \infty.$$

So, $(v_n)_{n\geq 1}$ and $(w_n)_{n\geq 1}$ are both convergent sequences and thus

$$v_n \to P_{\overline{V}}x \in \overline{V}, \quad w_n \to P_{\overline{V}^\perp}x \in \overline{W}, \Longrightarrow x \in \overline{V} \oplus \overline{W}.$$

So, $\overline{V \oplus W} \subset \overline{V} \oplus \overline{W}$.

On the contrary, for any $v \in \overline{V}$ and $w \in \overline{W}$, there exist $(v_n)_{n\geq 1} \subset V$ and $(w_n)_{n\geq 1} \subset W$ such that $v_n \to v$ and $w_n \to w$. Therefore, $V \oplus W \ni v_n + w_n \to v + w$. This implies $\overline{V} \oplus \overline{W} \subset \overline{V \oplus W}$.

Example: Consider three subspaces of ℓ^2: $V = \text{span}\{e_1\}, U = \text{span}\{e_1 + e_n\}_{n\geq 1}$ and $W = \overline{U}$. It is easy to show

$$\begin{aligned} V &= \{(a_1, 0, \cdots) : a_1 \in \mathbb{R}\}, \\ W &= \left\{ \left(\sum_{k\geq 2} a_k, a_2, a_3, \cdots \right) : \sum_{k\geq 2} a_k < \infty, \sum_{k\geq 2} a_k^2 < \infty \right\}. \end{aligned}$$

And from the expression above, we can find $V \cap W = \{0\}$. So the notation $V \oplus W$ is reasonable. Obviously, V is closed and U is closed by its definition, so $\overline{V} \oplus \overline{W} = V \oplus W$. However, consider

$$w_n = \left(\sum_{k=2}^{n} \frac{1}{k}, \frac{1}{2}, \cdots, \frac{1}{n}, 0, \cdots \right) \in W, \quad v_n = \left(-\sum_{k=2}^{n} \frac{1}{k}, 0, \cdots \right) \in V.$$

Noting $\sum_k \frac{1}{k^2} < \infty$, we have

$$\begin{aligned} v_n + w_n &= \left(0, \frac{1}{2}, \cdots, \frac{1}{n}, 0, \cdots \right) \in V \oplus W \\ \text{and } v_n + w_n \to x &= \left(0, \frac{1}{2}, \frac{1}{3}, \cdots \right) \in \ell^2. \end{aligned}$$

Since $\sum_k \frac{1}{k} = \infty$, there is no $v \in V$ and $w \in W$ such that $x = v + w$. This means $\overline{V} \oplus \overline{W} = V \oplus W \subsetneq \overline{V \oplus W}$.

This partial inclusion is because $V \not\perp W$. □

QUESTION 8.10 *Given a subspace A in the Hilbert space H. Suppose $\Lambda \subset A$ is a subspace of A. Suppose $\mathscr{A}, \mathscr{B}, \lambda$ are three subspaces of the Hilbert space $\mathscr{H}$. Then*

1. $\text{Proj}_{\Lambda^\perp}(A) \subset A$.

2. *If Λ is a q-replicating space $\Lambda = \lambda \times \cdots \times \lambda = \lambda^q \subset \mathscr{H}^q = H$ $(q \geq 2)$, we have*

$$\Lambda^\perp = \lambda^\perp \times \cdots \times \lambda^\perp.$$

3. $\mathrm{Proj}_{\Lambda^\perp}(A) = A \cap \Lambda^\perp$. *Furthermore, if $A = \mathscr{A} \times \cdots \times \mathscr{A} = \mathscr{A}^q, \Lambda = \lambda \times \cdots \times \lambda = \lambda^q$, we have*

$$\mathrm{Proj}_{\Lambda^\perp}(A) = (\mathscr{A} \cap \lambda) \times \cdots \times (\mathscr{A} \cap \lambda) = (\mathscr{A} \cap \lambda)^q,$$

which is also a q-replicating space.

Proof. 1. Note $\mathrm{Proj}_{\Lambda^\perp}(A) = \{x - \mathrm{Proj}_\Lambda(x) : \forall x \in A\}$ and $\mathrm{Proj}_\Lambda(x) \in \Lambda \subset A$, the fact that A is closed under addition yields

$$x - \mathrm{Proj}_\Lambda(x) \in A, \quad \forall x \in A.$$

Therefore, $\mathrm{Proj}_{\Lambda^\perp}(A) \subset A$.

2. We can find that $x = (x_1, \cdots, x_q) \in \Lambda^\perp \subset \mathscr{H}^q = H$ iff

$$(*) \qquad \langle x, y\rangle = \sum_{k=1}^{q} \langle x_k, y_k\rangle = 0, \quad \forall y = (y_1, \cdots, y_q) \in \Lambda.$$

Letting $x_{-k} \equiv 0 \in \mathscr{H}^{q-1}$ and x_k to vary, we can find $(*)$ is equivalent to $x_k \perp \lambda$ for any $k = 1, \cdots, q$. So,

$$\begin{aligned}\Lambda^\perp &= \{x = (x_1, \cdots, x_q) \in H : x_k \in \mathscr{A}^\perp, \forall k = 1, \cdots, q\} \\ &= \lambda^\perp \times \cdots \times \lambda^\perp.\end{aligned}$$

3. Since we have proved in 1. that $\mathrm{Proj}_{\Lambda^\perp}(A) \subset A$ and obviously $\mathrm{Proj}_{\Lambda^\perp}(A) \subset \Lambda^\perp$, we have $\mathrm{Proj}_{\Lambda^\perp}(A) \subset A \cap \Lambda^\perp$. On the contrary, since $A \cap \Lambda^\perp \subset \Lambda^\perp$, obviously we have $A \cap \Lambda^\perp = \mathrm{Proj}_{\Lambda^\perp}(A \cap \Lambda^\perp) \subset \mathrm{Proj}_{\Lambda^\perp}(A)$. Thus, $\mathrm{Proj}_{\Lambda^\perp}(A) = A \cap \Lambda^\perp$. The second assertion is obvious. □

QUESTION 8.11 (Variance identity iff the norm is Hilbertian) *Let V be a finite-dimensional real or complex vector space (e.g. $V = \mathbb{R}^{m\times n}$ with matrices viewed as vectors), equipped with a norm $\|\cdot\|$. The following are equivalent:*

(i) There exists an inner product $\langle\cdot,\cdot\rangle$ on V such that $\|x\|^2 = \langle x, x\rangle$ for all $x \in V$.

(ii) For every V-valued random element X with $\mathbb{E}\|X\|^2 < \infty$,

$$\mathbb{E}\|X - \mathbb{E}X\|^2 = \mathbb{E}\|X\|^2 - \|\mathbb{E}X\|^2.$$

Proof. **(i) ⇒ (ii).** Write $m := \mathbb{E}X$. Using the inner product that induces the norm and the bilinearity (sesquilinearity in the complex case) of $\langle \cdot, \cdot \rangle$,

$$\|X - m\|^2 = \langle X - m, X - m \rangle = \langle X, X \rangle - 2\,\mathrm{Re}\langle X, m \rangle + \langle m, m \rangle.$$

Taking expectations,

$$\mathbb{E}\|X - m\|^2 = \mathbb{E}\langle X, X \rangle - 2\,\mathrm{Re}\langle \mathbb{E}X, m \rangle + \langle m, m \rangle = \mathbb{E}\|X\|^2 - \|m\|^2,$$

which is the desired identity.

(ii) ⇒ (i). We derive the parallelogram law. Fix arbitrary $u, v \in V$ and let X take the two values u and v with probability $1/2$ each. Then $\mathbb{E}X = (u+v)/2$ and

$$\mathbb{E}\|X - \mathbb{E}X\|^2 = \frac{1}{2}\left\|\frac{u-v}{2}\right\|^2 + \frac{1}{2}\left\|\frac{v-u}{2}\right\|^2 = \frac{1}{4}\|u - v\|^2,$$

while

$$\mathbb{E}\|X\|^2 - \|\mathbb{E}X\|^2 = \frac{1}{2}(\|u\|^2 + \|v\|^2) - \left\|\frac{u+v}{2}\right\|^2 = \frac{1}{2}(\|u\|^2 + \|v\|^2) - \frac{1}{4}\|u+v\|^2.$$

By assumption (ii) these two quantities are equal, hence for all u, v,

$$\|u + v\|^2 + \|u - v\|^2 = 2(\|u\|^2 + \|v\|^2),$$

the parallelogram identity. By the Jordan–von Neumann theorem, a norm satisfies the parallelogram law if and only if it is induced by an inner product. Concretely, one may define, for real V,

$$\langle u, v \rangle := \frac{1}{4}\big(\|u + v\|^2 - \|u - v\|^2\big),$$

and for complex V,

$$\langle u, v \rangle := \frac{1}{4}\sum_{k=0}^{3} i^k \, \|u + i^k v\|^2,$$

which are inner products satisfying $\|x\|^2 = \langle x, x \rangle$ for all $x \in V$. This proves (i). □

§ 9 Partial Differential Equations

QUESTION 9.1 *Suppose $\{u_n\} \subset L^\infty(\mathbb{R}^n)$ is bounded, then exists $u \in L^\infty(\mathbb{R}^n)$ such that $u_n \rightharpoonup v$ in $L^2_{\mathrm{loc}}(\mathbb{R}^n)$.*

Proof. Observe $L^\infty(\mathbb{R}^n) = [L^1(\mathbb{R}^n)]^*$ and $L^1(\mathbb{R}^n)$ is separable. Since $\{u_n\} \subset L^\infty(\mathbb{R}^n)$ is bounded then exists $u \in L^\infty(\mathbb{R}^n)$, such that $w^*\text{-}\lim_{n\to\infty} u_n = u$. Namely, for each $v \in L^1(\mathbb{R}^n)$, we have

$$\int_{\mathbb{R}^n} u_n v dx \to \int_{\mathbb{R}^n} uv dx.$$

For any $V \subset\subset \mathbb{R}^n$ and each $w \in L^2(V)$, by Hölder inequality, we know $w \in L^1(V)$ as well. Moreover, we can extend w to $\mathbb{R}^n$ as follows:

$$v(x) = \begin{cases} w(x) & \text{, if } x \in V \\ 0 & \text{, if } x \in V^c. \end{cases}$$

Then $v \in L^1(\mathbb{R}^n)$. Thus, the above limit means

$$\int_V u_n w dx = \int_{\mathbb{R}^n} u_n v dx \to \int_{\mathbb{R}^n} uv dx = \int_V uw dx.$$

This shows $u_n \rightharpoonup v$ in $L^2_{\text{loc}}(\mathbb{R}^n)$. □

QUESTION 9.2 *Assume $U \subset \mathbb{R}^n$ is a bounded open set. Prove $W^{1,p}(U) \subset\subset L^p(U)$ for each $1 \le p \le \infty$.*

Proof.

1. When $1 \le p < n$, this is exactly the conclusion of Rellich-Kondrachov Compactness Theorem.

2. When $p = n$. Observe U is bounded, then the Hölder inequality tells us $u \in W^{1,n}(U)$ implies $u \in W^{1,p}(U)$ for each $1 \le p < n$. Observe $p* = \frac{np}{n-p} \uparrow \infty (p \uparrow n)$. So we can find a $p \in [1, n)$ such that $p^* > n$. Utilize Rellich-Kondrachov Compactness Theorem, we know $W^{1,p}(U) \subset\subset L^p(U)$.

3. When $n < p \le \infty$.

   (a) Way 1. *However, this way is suitable only for $n < p < \infty$.* It is similar to the previous situation. For each $u \in W^{1,p}(U)$, we can find $1 \le q < n$ such that $u \in W^{1,q}(U)$. Besides, observe $q* = \frac{nq}{n-q} \uparrow \infty (q \uparrow n)$, so we assume the given q is closed enough to n such that $q^* > p$. Thus the Rellich-Kondrachov Compactness Theorem gives $W^{1,q}(U) \subset\subset L^p(U)$. However, by Hölder inequality, we know the embedding mapping $F : W^{1,p}(U) \to W^{1,q}(U), u \mapsto u$ is continuous. So we have $W^{1,p}(U) \subset\subset L^p(U)$.

   (b) Way 2. Given a bounded sequence $\{u_m\} \subset W^{1,p}(U)$. By Morrey's inequality, we have

$$||u_m^*||_{C(\bar{U})} + [u_m^*]_{C^{0,1-n/p}(\bar{U})} \le C||u_m^*||_{W^{1,p}(U)},$$

where u_m^* is the continuous version of u_m. Next we verify u_m^* is uniformly and equicontinuous. In fact, the above inequality implies

$$\sup_{n\geq 1}||u_m^*||_{C(\bar{U})} \leq C\sup_{m\geq 1}||u_m^*||_{W^{1,p}(U)} = C'$$

and

$$\sup_{m\geq 1}[u_m^*]_{C^{0,1-n/p}(\bar{U})} \leq C\sup_{m\geq 1}||u_m^*||_{W^{1,p}(U)} = C'.$$

The first inequality show $\{u_m^*\}$ is uniformly bounded. The second inequality shows

$$\sup_{m\geq 1}|u_m^*(x) - u_m^*(y)| \leq C'||x-y||^{1-n/p},$$

which means $\{u_m^*\}$ is equicontinuous. So exists $u \in C(\bar{U})$ and subsequence $\{m_k\}_{k\geq 1}$ such that $||u_{m_k}^* - u||_{C(\bar{U})} \to 0, k \to \infty$. However, in the bounded open set, $u_{m_k}^* \to u$ in $C(\bar{U})$ implies $u_{m_k}^* \to u$ in $L^p(U)$. This assertion holds for each $n < p \leq \infty$. Thus $W^{1,p}(U) \subset\subset L^p(U)$. □

QUESTION 9.3 (A corollary of Weak maximum principle for $c \geq 0$) *Assume $u \in C^2(U)\cap C(\bar{U})$ and $Lu = 0$ in U, where $L = -a^{ij}\partial_{ij} + b^i\partial_i + c$ is the elliptic partial differential operators of second order. Assume $c \geq 0$ in U, then we have*

$$\max_{\bar{U}}|u| = \max_{\partial U}|u|.$$

Proof. Since $Lu = 0$, then we have

$$\max_{\bar{U}} u \leq \max_{\partial U} u^+ \leq \max_{\partial U}|u|, \quad \max_{\bar{U}}(-u) \leq \max_{\partial U} u^- \leq \max_{\partial U}|u|.$$

Hence,

$$\max_{\bar{U}}|u| = \max_{\bar{U}} u \vee \max_{\bar{U}}(-u) \leq \max_{\partial U}|u|$$

The converse inequality is obvious. □

QUESTION 9.4 *Assume $u \in C^2(U) \cap C(\bar{U})$ and U is connected, open and bounded. Write $M := \max_{\bar{U}} u$ and assume $C := \{x \in U | u(x) = M\} \neq \emptyset$. If $u \not\equiv M$ in U, so $V := \{x \in U | u(x) < M\} \neq \emptyset$. Then exists a point $y \in V$ satisfying $d(y, C) < d(y, \partial U)$.*

Proof. Choose and fix $x \in C$ and $z \in V$. Since U is connected and bounded, then exists a path $\gamma : [0,1] \to U$ satisfying $\gamma(0) = x, \gamma(1) = z$ and $\gamma[0,1]$ is a compact subset of $\mathbb{R}^n$. Observe $r := d(\gamma[0,1], \partial U) > 0$, we can find finite open balls $\{B(x_i, r/3)\}_{i=1}^n$ covering $\gamma[0,1]$, where $x_i \in \gamma[0,1]$ and $B(x_i, r/3) \cap B(x_{i+1}, r/3) \neq \emptyset$ for each i. There must exists $B(x_k, r/3)$ such that $B(x_k, r/3) \cap C \neq \emptyset$ and $B(x_k, r/3) \cap V \neq \emptyset$. We can choose $y \in B(x_k, r/3) \cap V$. So

$$d(y, \partial U) \geq d(\gamma[0,1], \partial U) - d(y, \gamma[0,1]) \geq r - \frac{r}{3} = \frac{2r}{3} > d(y, C).$$

□

QUESTION 9.5 *We write $\widetilde{\partial}_i$ for the weak derivative operator with respect to the i-th component. Suppose $u \in W^{1,p}(U)$, then $\widetilde{\partial}^2_{ij} u := \widetilde{\partial}_i(\widetilde{\partial}_j u)$ exists iff $\widetilde{\partial}^2_{ji} u$ exists. In this case, we still have $\widetilde{\partial}^2_{ij} u = \widetilde{\partial}^2_{ji} u$, where $U \subset \mathbb{R}^n$ is an open subset and $1 \leq i, j \leq n$.*

Proof. Suppose $\widetilde{\partial}^2_{ij} u$ exists, then for each $\phi \in C_c^\infty(U)$, we have

$$\int_U \widetilde{\partial}^2_{ij} u \phi dx = -\int_U \widetilde{\partial}_j u \partial_i \phi dx = \int_U u \partial^2_{ji} \phi dx = \int_U u \partial^2_{ij} \phi dx.$$

By the definition of the weak derivative, we have $\widetilde{\partial}^2_{ij} u = \widetilde{\partial}^2_{ji} u$. □

QUESTION 9.6 *Suppose $u \in W^{k,n}(U)$ and $k > 2$, where $U \subset \mathbb{R}^n$. Then for each $|\alpha| \leq k-2$, the weak derivative $D^\alpha u = \partial^\alpha u$ (the strong derivative). Therefore $u \in C^{k-2,\gamma}(\bar{U})$ for each $0 < \gamma < 1$.*

Proof. Since $u \in W^{k,n}(U)$, then the Gagliardo-Nirenberg-Sobolev inequality and the Hölder inequality shows: for each $n \leq q < \infty$, exists $p \in [1, n)$ such that for each $|\beta| \leq k-1$,

$$||D^\beta u||_{L^q(U)} \leq C||D^\beta u||_{W^{1,p}(U)} \leq C||D^\beta u||_{W^{1,n}(U)} \leq C||u||_{W^{k,n}(U)}.$$

Thus we have $u \in W^{k-1,q}(U)$ for each $n \leq q < \infty$. As a consequence, the Morrey's inequality shows $D^\alpha u \in C^{0,1-n/q}(\bar{U})$ for all $n < q < \infty$ and $|\alpha| \leq k-2$. Next we show $D^\alpha u = \partial^\alpha u$. For each $x \in U$, select a close ball $B(x, \epsilon) \subset U$. Consider the standard modifier η and the convolution $D^\alpha u^\varepsilon = \eta_\varepsilon * D^\alpha u$ in U_ε. When ε small enough, we know $B(x, \epsilon) \subset U_\varepsilon$. Observe $D^\alpha u \in C^{0,1-n/q}(\bar{U}) \subset C(\bar{U})$, so $D^\alpha u^\varepsilon \to D^\alpha u$ uniformly in $B(x, \epsilon)$. So for any sequence $\{\varepsilon_m\}$ satisfying $\varepsilon_m \to 0$, we have $D^\alpha u^{\varepsilon_m} \to D^\alpha u$ uniformly in $B(x, \epsilon)$. Based on the previous discussion, $\{u^{\varepsilon_m}\}$ is a Cauchy sequence in the Banach space $C^{k-2}(B(x, \epsilon))$, so we can find $v \in C^{k-2}(B(x, \epsilon))$ such that $u^{\varepsilon_m} \to v$ in $C^{k-2}(B(x, \epsilon))$. The uniqueness of the limit shows $u = v \in C^{k-2}(B(x, \epsilon))$. Thus $D^\alpha u = \partial^\alpha u$ for each $|\alpha| \leq k-2$. □

QUESTION 9.7 *Suppose $U \subset \mathbb{R}^n$ and $\partial U \cap \mathbb{R}_n^+ \neq \emptyset$. Assume $f \in C^1(\bar{U})$, then for each $x \in \partial U \cap \mathbb{R}_n^+$, we have*

$$\partial_{x_n+} f(x) := \lim_{h\to 0} \frac{f(x+he_n)-f(x)}{h} = \lim_{h\to 0} \partial_{x_n} f(x+he_n).$$

Proof. Write $x = (x', x_n) \in \mathbb{R}^n$ and fix x'. Then define $g(x_n) := f(x', x_n)$. We know $g(x_n)$ is continuous on $[0, h]$ and differentiable in $(0, h)$. So the Differential mean value theorem yields exists $\theta(h) \in (0, 1)$ such that

$$\begin{aligned} f(x+he_n) - f(x) &= g(h) - g(0) \\ &= g'(\theta(h)h)h = \partial_{x_n} f(x+\theta(h)he_n)h. \end{aligned}$$

Observe $f \in C^1(\bar{U})$, then we have

$$\begin{aligned} \partial_{x_n+} f(x) &= \lim_{h\to 0} \frac{f(x+he_n)-f(x)}{h} = \lim_{h\to 0} \frac{\partial_{x_n} f(x+\theta(h)he_n)h}{h} \\ &= \lim_{h\to 0} \partial_{x_n} f(x+\theta(h)he_n) = \lim_{h\to 0} \partial_{x_n} f(x+he_n). \end{aligned}$$

□

QUESTION 9.8 *Suppose $U \subset \mathbb{R}^n$ is an open, bounded subset. Consider $u(x) = C$, a.e. $x \in U$, where C is a constant. Then we have $u \in W^{1,p}(U)$ for each $1 \le p < \infty$ and $Tu = C$, a.e. $x \in \partial U$ provided $1 \le p < \infty$.*

Proof. We can easily verify $u \in W^{1,p}(U)$. Consider another function $v(x) \equiv C, \forall x \in \bar{U}$. Since $v \in W^{1,p}(U) \cap C(\bar{U})$ for each $1 \le p < \infty$, then the Trace theorem shows $Tv = v|_{\partial U} = C$. Thus

$$||C - Tu||_{L^p(\partial U)} = ||Tv - Tu||_{L^p(\partial U)} \le C'||v-u||_{W^{1,p}(U)} = 0.$$

This yields $Tu = C$, a.e. $x \in \partial U$. □

§ 10 Econometrics

QUESTION 10.1 *Consider a univariate regression model*

$$y_i = \alpha + \beta x_i + \varepsilon_i.$$

If we want to do a weighted OLS with weight $\sqrt{w_i}$:

$$\sqrt{w_i} y_i = \sqrt{w_i}\alpha + \sqrt{w_i}\beta x_i + \sqrt{w_i}\varepsilon_i,$$

we can write it as

$$\begin{aligned} & V^{1/2}Y = \alpha V^{1/2}\mathbf{1} + V^{1/2}X\gamma + V^{1/2}\varepsilon \\ \iff \quad & \widetilde{Y} = \alpha\widetilde{\mathbf{1}} + \beta\widetilde{x} + \widetilde{\varepsilon}. \end{aligned}$$

Then, the weighted OLS estimator has the following expression:

$$\widetilde{\gamma} = \frac{\widehat{\mathrm{Cov}}_w(\widetilde{x}, \widetilde{Y})}{\widehat{\mathrm{Var}}_w(\widetilde{x})},$$

where the subscript w means taking the sample covariance, variance, and mean with respect to the measure $w = (w_i)_{i=1}^n$.

Proof. Write $\widetilde{X} := (\widetilde{\mathbf{1}}, \widetilde{x})$. Then, the weighted OLS estimate $\widetilde{\gamma}$ is

$$\begin{aligned}
\widetilde{\gamma} &= (\widetilde{X}'X)^{-1}\widetilde{X}'\widetilde{Y} \\
&= \left[\begin{pmatrix} \widetilde{\mathbf{1}}' \\ \widetilde{x}' \end{pmatrix} \begin{pmatrix} \widetilde{\mathbf{1}} & \widetilde{x} \end{pmatrix}\right]^{-1} \begin{pmatrix} \widetilde{\mathbf{1}}' \\ \widetilde{x}' \end{pmatrix} \widetilde{Y} \\
&= \left[\begin{pmatrix} \mathbf{1}'V^{1/2} \\ x'V^{1/2} \end{pmatrix} \begin{pmatrix} V^{1/2}\mathbf{1} & V^{1/2}x \end{pmatrix}\right]^{-1} \begin{pmatrix} \mathbf{1}'V^{1/2} \\ x'V^{1/2} \end{pmatrix} V^{1/2}Y \\
&= \begin{pmatrix} \mathbf{1}'V\mathbf{1} & \mathbf{1}'Vx \\ x'V\mathbf{1} & x'Vx \end{pmatrix}^{-1} \begin{pmatrix} \mathbf{1}'VY \\ x'VY \end{pmatrix} \\
&= \begin{pmatrix} \sum_i w_i & \sum_i w_i x_i \\ \sum_i w_i x_i & \sum_i w_i x_i^2 \end{pmatrix}^{-1} \begin{pmatrix} \sum_i w_i y_i \\ \sum_i w_i x_i y_i \end{pmatrix} \\
&= \frac{1}{\sum_i w_i \sum_i w_i x_i^2 - \left(\sum_i w_i x_i\right)^2} \begin{pmatrix} \sum_i w_i x_i^2 & -\sum_i w_i x_i \\ -\sum_i w_i x_i & \sum_i w_i \end{pmatrix} \begin{pmatrix} \sum_i w_i y_i \\ \sum_i w_i x_i y_i \end{pmatrix}.
\end{aligned}$$

Therefore,

$$\widetilde{\beta} = \frac{\sum_i w_i \sum_i w_i x_i y_i - \sum_i w_i x_i \sum_i w_i y_i}{\sum_i w_i \sum_i w_i x_i^2 - \left(\sum_i w_i x_i\right)^2}.$$

Note

$$\begin{aligned}
\widehat{\mathrm{Cov}}_w(\widetilde{x}, \widetilde{Y}) &= \frac{1}{\sum_i w_i} \sum_{i=1}^n w_i \left(x_i - \frac{1}{\sum_i w_i} \sum_{j=1}^n w_j x_j\right)\left(y_i - \frac{1}{\sum_i w_i} \sum_{j=1}^n w_j y_j\right) \\
&= \frac{1}{\sum_i w_i} \left[\sum_{i=1}^n w_i x_i y_i - \sum_{i=1}^n w_i x_i \left(\frac{1}{\sum_i w_i} \sum_{j=1}^n w_j y_j\right)\right. \\
&\quad \left. - \sum_{i=1}^n w_i y_i \left(\frac{1}{\sum_i w_i} \sum_{j=1}^n w_j x_j\right) + \frac{1}{\sum_i w_i} \left(\sum_{j=1}^n w_j x_j\right)\left(\sum_{j=1}^n w_j y_j\right)\right] \\
&= \frac{1}{\sum_i w_i} \sum_{i=1}^n w_i x_i y_i - \left(\frac{1}{\sum_i w_i}\right)^2 \left(\sum_{i=1}^n w_i x_i\right)\left(\sum_{j=1}^n w_j y_j\right).
\end{aligned}$$

So,

$$\widetilde{\beta} = \frac{\left(\sum_i w_i\right)^2 \widehat{\mathrm{Cov}}_w(\widetilde{x}, \widetilde{Y})}{\left(\sum_i w_i\right)^2 \widehat{\mathrm{Var}}_w(\widetilde{x})} = \frac{\widehat{\mathrm{Cov}}_w(\widetilde{x}, \widetilde{Y})}{\widehat{\mathrm{Var}}_w(\widetilde{x})}.$$

□

QUESTION 10.2 *Given a parameter space Θ and observable distribution family $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$. Define $G : \Theta \to \mathcal{P}, \theta \mapsto P_\theta$. Then*

- *Model is point identified at true value θ_0 if*

$$P_\theta = P_{\theta_0} \Longrightarrow \theta = \theta_0,$$

  which is equivalent to $G^{-1}(\{P_{\theta_0}\}) = \{\theta\}$

- *Model is globally identified if*

$$P_\theta = P_{\theta'} \Longrightarrow \theta = \theta', \quad \forall \theta, \theta' \in \Theta,$$

  which is equivalent to saying G is an injection.

- *Model is locally identified if G is an injection in some neighborhood of the true value θ_0.*

- *Model is partially identified if $G^{-1}(\{P_{\theta_0}\}) \subsetneq \Theta$ and $G^{-1}(\{P_{\theta_0}\})$ is not a singleton.*

- *For a functional parameter $\psi()$*

When there is a functional F such that $\theta = F(P_\theta)$, θ is globally identified.

§ 11 Mathemetical Statistics

QUESTION 11.1 *Let $X_1, \ldots, X_n$ be i.i.d. with distribution $U(\theta, \theta + |\theta|)$, where $\theta \neq 0$. Find the MLE of θ.*

Proof. Write the order statistics $X_{(1)} = \min_i X_i$ and $X_{(n)} = \max_i X_i$. The density is

$$f(x;\theta) = \frac{1}{|\theta|}\mathbf{1}\{\theta \le x \le \theta + |\theta|\},$$

hence the likelihood is

$$L(\theta) = \frac{1}{|\theta|^n}\mathbf{1}\left\{\theta \le X_{(1)} \text{ and } X_{(n)} \le \theta + |\theta|\right\}.$$

Case 1: $\theta > 0$. Then the support is $[\theta, 2\theta]$ and feasibility requires

$$\theta \le X_{(1)} \quad \text{and} \quad 2\theta \ge X_{(n)} \iff \theta \in \left[X_{(n)}/2,\ X_{(1)}\right].$$

On this interval $L(\theta) = \theta^{-n}$ is strictly decreasing in θ, so the maximizer is

$$\hat{\theta}_+ = \frac{X_{(n)}}{2}, \qquad L\left(\tfrac{X_{(n)}}{2}\right) = \left(\frac{2}{X_{(n)}}\right)^n.$$

Case 2: $\theta < 0$**.** Then the support is $[\theta, 0]$ and feasibility requires

$$\theta \le X_{(1)} \quad \text{and} \quad X_{(n)} \le 0 \iff \theta \in (-\infty,\, X_{(1)}].$$

Here $L(\theta) = (-\theta)^{-n}$ is strictly increasing in θ, so the maximizer is

$$\hat{\theta}_- = X_{(1)}, \qquad L\big(X_{(1)}\big) = \frac{1}{\big(-X_{(1)}\big)^n}.$$

Since the sample must come from one of the two feasible regimes, the MLE is

$$\hat{\theta}_{\text{MLE}} = \begin{cases} X_{(1)}, & \text{if } X_{(n)} \le 0 \quad \text{(negative–support case)}, \\ \dfrac{X_{(n)}}{2}, & \text{if } \dfrac{X_{(n)}}{2} \le X_{(1)} \quad \text{(positive–support case)}. \end{cases}$$

Equivalently, when both candidates are admissible, select the one with the larger likelihood:

$$\hat{\theta}_{\text{MLE}} = \begin{cases} X_{(1)}, & \text{if } \left(\dfrac{2}{X_{(n)}}\right)^n \le \dfrac{1}{\big(-X_{(1)}\big)^n}, \\ \dfrac{X_{(n)}}{2}, & \text{if } \left(\dfrac{2}{X_{(n)}}\right)^n > \dfrac{1}{\big(-X_{(1)}\big)^n}. \end{cases}$$

□

§ 12 Semiparametric Model

QUESTION 12.1 *Given a model $\mathcal{P}$ and its dominating measure ν. Assume $P_0 \in \mathcal{P}$ is the true model and observation $Z \sim P_0$. Define its n-dimensional regular submodels as follows*

$$\mathcal{P}^{(n)} := \{P_\gamma = P_{(\gamma_1, \cdots, \gamma_n)} : \gamma_i \in (-\varepsilon_i, \varepsilon_i), 1 \le i \le n\} \subset \mathcal{P}.$$

Write $s_{\gamma, \mathcal{P}^{(n)}}$ as the score of submodels $\mathcal{P}^{(n)}$ in the sense of DQM. Then the following definition of tangent space does not depend on n:

$$\mathcal{T}_n(\mathcal{P}) := \overline{\left\{h' s_{\gamma, \mathcal{P}^{(n)}}(Z) : \mathcal{P}^{(n)} \subset \mathcal{P}, h \in \mathbb{R}^n\right\}}^{L_0^2}.$$

Proof. We just need to prove $\mathcal{T}_1(\mathcal{P}) = \mathcal{T}_n(\mathbb{P}), \forall n \ge 2$.

For any $s_{\ell, \overline{\mathcal{P}}^{(1)}}(Z)$, where

$$\overline{\mathcal{P}}^{(1)} = \{\overline{P}_\ell : \ell \in (-\varepsilon_1, \varepsilon_1)\} \subset \mathcal{P},$$

we can define the n-dimensional submodel

$$\mathcal{P}^{(n)} := \{P_\gamma = P_{(\ell,\gamma_2,\cdots,\gamma_n)} = \overline{P}_\ell : \gamma_i \in (-\varepsilon_i, \varepsilon_i), 1 \le i \le n\} \subset \mathcal{P},$$

which means that the model $\mathcal{P}^{(n)}$ will not change with respect to parameters $(\gamma_2, \cdots, \gamma_n)$. We now show $\mathcal{P}^{(n)}$ is regular, i.e., $\mathcal{P}^{(n)}$ is DQM at $\gamma = 0$ with score $s_{\gamma,\mathcal{P}^{(n)}}(Z) := (s_{\ell,\overline{\mathcal{P}}^{(1)}}(Z), 0, \cdots, 0)'$:

$$\begin{aligned}
&||\sqrt{p_{\gamma,\mathcal{P}^{(n)}}} - \sqrt{p_0} - \frac{1}{2}\sqrt{p_0}\gamma' s_{\gamma,\mathcal{P}^{(n)}}||_{L_0^2(\nu)} \\
&\quad = ||\sqrt{p_{\ell,\overline{\mathcal{P}}^{(1)}}} - \sqrt{p_0} - \frac{1}{2}\sqrt{p_0}\ell s_{\gamma,\overline{\mathcal{P}}^{(1)}}||_{L_0^2(\nu)} = o_{L_0^2(\nu)}(|\ell|),
\end{aligned}$$

where the last equality is because $\overline{\mathcal{P}}^{(1)}$ is DQM at $\ell = 0$ with score $s_{\ell,\overline{\mathcal{P}}^{(1)}}(Z)$. Since $|\ell| \le ||\gamma||$, we have $o_{L_0^2(\nu)}(|\ell|) = o_{L_0^2(\nu)}(||\gamma||)$. Therefore, for any $h = (h_1, \cdots, h_n) \in \mathbb{R}^n$, $h_1 s_{\gamma,\overline{\mathcal{P}}^{(1)}} = h' s_{\gamma,\mathcal{P}^{(n)}}$, which implies $\mathcal{P}^{(1)} \subset \mathcal{P}^{(n)}$.

On the contrary, for any $\widetilde{h} = (\widetilde{h}_1, \cdots, \widetilde{h}_n)$ and $\widetilde{h}' s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z)$, where

$$\overline{\mathcal{P}}^{(n)} = \{\overline{P}_\gamma = \overline{P}_{(\gamma_1,\cdots,\gamma_n)} : \gamma_i \in (-\varepsilon_i, \varepsilon_i), 1 \le i \le n\},$$

We can let $M > 0$ big enough such that $h := \widetilde{h}/M$ satisfies $h = \widetilde{h}_i/M \in (-\varepsilon_i, \varepsilon_i)$. The model

$$\widetilde{\mathcal{P}}^{(n)} := \{\overline{P}_h = \overline{P}_{(h_1,\cdots,h_n)} : h_i \in (-\varepsilon_i, \varepsilon_i), 1 \le i \le n\}$$

Then we can define the one-dimensional submodel

$$\mathcal{P}^{(1)} := \{P_\ell = \overline{P}_{\ell h} = \overline{P}_{(\ell h_1,\cdots,\ell h_n)} : \ell \in (-1, 1)\}.$$

Write

$$\widetilde{\mathcal{P}}^{(n)} := \{\overline{P}_{\ell h} = \overline{P}_{(\ell h_1,\cdots,\ell h_n)} : \ell h_i \in (-h_i, h_i), 1 \le i \le n\}.$$

Obviously, using the DQM property of $\overline{\mathcal{P}}^{(n)}$ (treat ℓh_i as a new parameter $\tilde{\gamma}_i$), we know $\widetilde{\mathcal{P}}^{(n)}$ is also DQM at $\ell h = 0$ with score $s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z)$. We now show $\mathcal{P}^{(1)}$ is regular, i.e., $\mathcal{P}^{(1)}$ is DQM at $\ell = 0$ with score $s_{\ell,\mathcal{P}^{(1)}}(Z) = h' s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z)$:

$$\begin{aligned}
&||\sqrt{p_{\ell,\mathcal{P}^{(1)}}} - \sqrt{p_0} - \frac{1}{2}\sqrt{p_0}\ell h' s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z)||_{L_0^2(\nu)} \\
&\quad = ||\sqrt{p_{\ell h,\widetilde{\mathcal{P}}^{(n)}}} - \sqrt{p_0} - \frac{1}{2}\sqrt{p_0}\ell h' s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z)||_{L_0^2(\nu)} = o_{L^2(\nu)}(||\ell h||).
\end{aligned}$$

Since h is a constant, $o_{L^2(\nu)}(||\ell h||) = o_{L^2(\nu)}(|\ell|)$. Therefore, $\widetilde{h}' s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z) = M h' s_{\gamma,\overline{\mathcal{P}}^{(n)}}(Z) = M s_{\ell,\mathcal{P}^{(1)}}(Z)$, which implies $\mathcal{P}^{(n)} \subset \mathcal{P}^{(1)}$. $\square$

Definition 12.1 (Two notions of Neyman orthogonality) Let $\mathcal{P}$ be a DQM (Hellinger differentiable) model at P_0, with tangent space $\mathcal{T} \subset L_0^2(P_0)$ and nuisance tangent space $\mathcal{T}_\eta \subset \mathcal{T}$. Let $\psi(W;\theta,\eta)$ be a moment/score map with $\mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0) = 0$.
(i) Score/Geometric orthogonality. We say ψ is *score-orthogonal to the nuisance* if

$$\mathbb{E}_{P_0}\big[\psi(W;\theta_0,\eta_0)\, s(W)\big] = 0 \qquad \text{for all } s \in \mathcal{T}_\eta.$$

(ii) Functional (DML) orthogonality. Fix the true law P_0. We say ψ is *(functionally) Neyman orthogonal* if the Fréchet/Gateaux derivative w.r.t. η vanishes at truth:

$$\partial_\eta\, \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)[h] = 0 \qquad \text{for all admissible directions } h.$$

In general, (i) and (ii) are independent: either one can hold without the other.

Definition 12.2 (EIF and global selection of IF) Let $\theta : \mathcal{P} \to \mathbb{R}^d$ be pathwise differentiable at P_0, with pathwise derivative $\dot\theta_{P_0} : \mathcal{T} \to \mathbb{R}^d$. An $L_0^2(P_0)$ element $\phi^\star_{P_0} \in \overline{\mathcal{T}}$ is the *efficient influence function (EIF)* if

$$\dot\theta_{P_0}(s) = \mathbb{E}_{P_0}\big[\phi^\star_{P_0}(W)\, s(W)\big] \qquad (\forall s \in \mathcal{T}),$$

and $\mathrm{Var}_{P_0}(\phi^\star_{P_0})$ is minimal among all IFs. A *global selection of IFs* is a measurable map $P \mapsto \phi_P$ such that, for each $P \in \mathcal{P}$, ϕ_P is an influence function for θ at P and

$$\mathbb{E}_P\phi_P(W) = 0, \qquad \dot\theta_P(s) = \mathbb{E}_P\big[\phi_P(W)\, s(W)\big] \quad (\forall s \in \mathcal{T}_P).$$

The special case $\phi_P = \phi^\star_P$ is the *global EIF selection*.

Proposition 12.1 (Any IF is score-orthogonal to $\mathcal{T}_\eta$ at P_0) *If ϕ is an influence function at P_0, then for every $s \in \mathcal{T}_\eta$,*

$$\mathbb{E}_{P_0}\big[\phi(W)\, s(W)\big] = \dot\theta_{P_0}(s) = 0,$$

because nuisance-only paths keep θ fixed.

Theorem 12.1 (When both orthogonalities hold) *Assume the model is DQM at P_0.*
(A) Global EIF selection. *Let $\psi(W;\theta,\eta) = \phi^\star_{P_{\theta,\eta}}(W)$, i.e. the moment function equals the EIF at the law $P_{\theta,\eta}$. Then both*

$$\mathbb{E}_{P_0}\big[\psi(W;\theta_0,\eta_0)\, s(W)\big] = 0 \quad (\forall s \in \mathcal{T}_\eta), \qquad \partial_\eta\, \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)[h] = 0 \quad (\forall h)$$

hold simultaneously.
(B) Full-rank linear rescaling of EIF. *More generally, if* $\psi(W;\theta,\eta) = M\,\phi^\star_{P_{\theta,\eta}}(W)$ *with a fixed full-rank* $d \times d$ *matrix* M*, then the same two orthogonality properties hold.*

Proof. (A) Score orthogonality. Fix a nuisance-only regular path $\varepsilon \mapsto P_\varepsilon = P_{\theta_0,\eta_\varepsilon}$ with score $s \in \mathcal{T}_\eta$. At P_0, the EIF identity gives

$$\mathbb{E}_{P_0}\big[\psi(W;\theta_0,\eta_0)\,s(W)\big] = \mathbb{E}_{P_0}\big[\phi^\star_{P_0}(W)\,s(W)\big] = \dot\theta_{P_0}(s) = 0.$$

Functional orthogonality. Define

$$H(\varepsilon) := \mathbb{E}_{P_\varepsilon}\big[\psi(W;\theta_0,\eta_\varepsilon)\big] = \mathbb{E}_{P_{\theta_0,\eta_\varepsilon}}\big[\phi^\star_{P_{\theta_0,\eta_\varepsilon}}(W)\big] \equiv 0,$$

since for every P the EIF has zero mean under P. By DQM, H is differentiable at 0 and

$$0 = H'(0) = \underbrace{\mathbb{E}_{P_0}\big[\psi(W;\theta_0,\eta_0)\,s(W)\big]}_{=0 \text{ by the previous part}} + \partial_\eta\,\mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)[\dot\eta].$$

Hence $\partial_\eta\,\mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)[\dot\eta] = 0$ for every admissible nuisance direction $\dot\eta$, proving functional orthogonality.

(B) Replace ψ by $M\phi^\star_{P_{\theta,\eta}}$. The two conditions are preserved because multiplication by a fixed full-rank matrix does not affect zero inner products nor zero derivatives. □

Proposition 12.2 (Calibration matrix A when using (rescaled) EIF) *Let*

$$A := \partial_\theta\,\mathbb{E}_{P_0}\psi(W;\theta,\eta_0)\big|_{\theta=\theta_0}.$$

If $\psi(W;\theta,\eta) = \phi^\star_{P_{\theta,\eta}}(W)$*, then* $A = -I_d$*. If* $\psi(W;\theta,\eta) = M\,\phi^\star_{P_{\theta,\eta}}(W)$ *with full-rank* M*, then* $A = -M$*.*

Proof. Consider a θ-only regular path $\varepsilon \mapsto P_\varepsilon = P_{\theta_\varepsilon,\eta_0}$ with score s_θ and $\dot\theta := \frac{d}{d\varepsilon}\theta_\varepsilon\big|_0 \in \mathbb{R}^d$ arbitrary (by local identification). Define

$$G(\varepsilon) := \mathbb{E}_{P_\varepsilon}\big[\phi^\star_{P_{\theta_\varepsilon,\eta_0}}(W)\big] \equiv 0.$$

DQM implies

$$0 = G'(0) = \underbrace{\mathbb{E}_{P_0}\big[\phi^\star_{P_0}(W)\,s_\theta(W)\big]}_{=\,\dot\theta \text{ by EIF identity}} + \partial_\theta\,\mathbb{E}_{P_0}\phi^\star_{P_{\theta,\eta_0}}(W)\big|_{\theta_0}[\dot\theta].$$

The local identification guarantees $\dot\theta \in \mathbb{R}^d$. Therefore $A = \partial_\theta\mathbb{E}_{P_0}\phi^\star_{P_{\theta,\eta_0}}|_{\theta_0} = -I_d$. For $\psi = M\phi^\star$, linearity gives $A = M\,(-I_d) = -M$. □

Corollary 12.1 (IF of Z-estimators and efficiency) *Suppose* $(\hat{\theta}, \hat{\eta})$ *solves* $\mathbb{P}_n \psi(W; \theta, \eta) = 0$ *with* $\mathbb{E}_{P_0} \psi(W; \theta_0, \eta_0) = 0$*, and regularity (e.g. cross-fitting, stochastic equicontinuity) ensures the linearization*

$$\sqrt{n}(\hat{\theta} - \theta_0) = -A^{-1} \frac{1}{\sqrt{n}} \sum_{i=1}^{n} \psi(W_i; \theta_0, \eta_0) + o_P(1).$$

If $\psi = \phi^\star_{P_{\theta,\eta}}$*, then by Proposition 12.2,* $A = -I_d$ *and*

$$\mathrm{IF}_{\hat{\theta}}(W) = \phi^\star_{P_0}(W), \qquad \mathrm{Var}\big(\sqrt{n}(\hat{\theta} - \theta_0)\big) = \mathrm{Var}_{P_0}\big(\phi^\star_{P_0}\big),$$

which is the semiparametric efficiency bound. If $\psi = M\phi^\star$*, then* $A = -M$ *and the IF is still* $\phi^\star_{P_0}$*; efficiency is unchanged.*

Remark 12.1 (What each orthogonality controls)

- *Score/geometric orthogonality* $\mathbb{E}_{P_0}[\psi_0 s] = 0$ for $s \in \mathcal{T}_\eta$ kills the "distribution channel" part of the pathwise derivative (variations of P along nuisance directions).
- *Functional (DML) orthogonality* $\partial_\eta \mathbb{E}_{P_0} \psi = 0$ kills the "function channel" that arises when plugging $\hat{\eta}$ for η_0 under a fixed law P_0.
- In general they are independent; when ψ is the (rescaled) *EIF map*, both hold simultaneously, and moreover $A = -I$ (or $-M$), yielding efficiency.
- Any IF at P_0 (not necessarily efficient) is automatically score-orthogonal to $\mathcal{T}_\eta$, but may fail functional (DML) orthogonality under a fixed P_0 unless it is chosen as a global selection and the above argument applies.

Definition 12.3 (Model, tangent space, and pathwise derivative) Let $\mathcal{P}$ be a statistical model and $P_0 \in \mathcal{P}$ the true distribution. For any regular parametric submodel $\{P_\varepsilon : \varepsilon \in (-\delta, \delta)\} \subset \mathcal{P}$ with density p_ε at P_0, the score at $\varepsilon = 0$ is

$$s(W) = \frac{\partial}{\partial \varepsilon} \log p_\varepsilon(W)\bigg|_{\varepsilon=0} \in L^2_0(P_0) := \{f \in L^2(P_0) : \mathbb{E}_{P_0} f = 0\}.$$

The (closure of the) linear span of all such scores is the *tangent space* $\mathcal{T} \subset L^2_0(P_0)$. For a parameter (functional) $\theta : \mathcal{P} \to \mathbb{R}^d$ that is Gateaux differentiable at P_0, its pathwise derivative is the continuous linear map

$$\dot{\theta}_{P_0} : \mathcal{T} \to \mathbb{R}^d, \qquad \dot{\theta}_{P_0}(s) = \frac{\partial}{\partial \varepsilon} \theta(P_\varepsilon)\bigg|_{\varepsilon=0}.$$

Definition 12.4 (Influence function (IF) and efficient influence function (EIF)) An element $\phi \in L_0^2(P_0)$ is an *influence function* for θ at P_0 if

$$\dot{\theta}_{P_0}(s) = \mathbb{E}_{P_0}\big[\phi(W)\, s(W)\big] \qquad \text{for all } s \in \mathcal{T}.$$

Among all such representations, the unique representative that belongs to the closure $\overline{\mathcal{T}}$ is called the *efficient influence function* (EIF) and is denoted $\phi^\star$. By the Riesz representation theorem on $(\overline{\mathcal{T}}, \langle\cdot,\cdot\rangle_{L^2(P_0)})$,

$$\mathrm{Var}_{P_0}(\phi^\star) = \inf\Big\{\mathrm{Var}_{P_0}(\psi) : \ \psi \in L_0^2(P_0),\ \dot{\theta}_{P_0}(s) = \mathbb{E}_{P_0}[\psi s]\ \forall s \in \mathcal{T}\Big\}.$$

Definition 12.5 (Nuisance tangent space) Fix the true value $\theta_0 = \theta(P_0)$. Let $\mathcal{P}_\eta$ be the collection of paths that perturb only the nuisance while keeping θ fixed at θ_0:

$$\mathcal{P}_\eta := \{P_\varepsilon \in \mathcal{P} : \ \theta(P_\varepsilon) \equiv \theta_0\}.$$

The *nuisance tangent space* is

$$\mathcal{T}_\eta := \overline{\{s : \ s \text{ is the score of some } P_\varepsilon \in \mathcal{P}_\eta\}} \ \subset \ \mathcal{T}.$$

Definition 12.6 (Neyman orthogonality and near-orthogonality) Let $\psi(W;\theta,\eta)$ be a score/moment map with $\mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0) = 0$. Its (Fréchet/Gateaux) derivative in the nuisance direction at truth is the linear map

$$\partial_\eta \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)[h], \qquad h \in \mathcal{H}.$$

We say ψ is *Neyman orthogonal* iff

$$\partial_\eta \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)[h] = 0 \quad \text{for all } h \in \mathcal{H}.$$

We say it is *near-orthogonal* if $\|\partial_\eta \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0)\| \le \delta_N$ with $\delta_N \to 0$.

Proposition 12.3 (Uniqueness in $\overline{\mathcal{T}}$ and efficiency) *If $\psi \in \overline{\mathcal{T}}$ also satisfies*

$$\dot{\theta}_{P_0}(s) = \mathbb{E}_{P_0}[\psi s] \qquad \forall\, s \in \mathcal{T},$$

then $\psi = \phi^\star$ in $L^2(P_0)$, and for any influence function $\tilde{\phi}$ one has

$$\mathrm{Var}_{P_0}(\tilde{\phi}) \ \ge \ \mathrm{Var}_{P_0}(\phi^\star),$$

with equality iff $\tilde{\phi} = \phi^\star$ a.s.

Proof. From the two representations, $\mathbb{E}[(\psi - \phi^\star)s] = 0$ for all $s \in \mathcal{T}$, i.e. $\psi - \phi^\star \perp \mathcal{T}$. Since both lie in $\overline{\mathcal{T}}$, orthogonality forces $\psi = \phi^\star$. For efficiency, write any IF as $\tilde{\phi} = \phi^\star + g$ with $g \perp \mathcal{T}$ (this is the orthogonal decomposition relative to $\overline{\mathcal{T}}$). Then

$$\mathrm{Var}(\tilde{\phi}) = \mathrm{Var}(\phi^\star) + \mathrm{Var}(g) \;\geq\; \mathrm{Var}(\phi^\star),$$

with equality iff $g = 0$. □

Proposition 12.4 (Near-orthogonal scores yield root-n linearization; EIF gives efficiency) *Suppose $(\hat{\theta}, \hat{\eta})$ solves the empirical moment equation $\mathbb{P}_n\psi(W;\theta,\eta) = 0$ with $\mathbb{E}\psi(W;\theta_0,\eta_0) = 0$. Let*

$$A := \partial_\theta \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0), \qquad B := \partial_\eta \mathbb{E}_{P_0}\psi(W;\theta_0,\eta_0).$$

Assume A is nonsingular, $\|B\| \leq \delta_N \to 0$ (near orthogonality), $\|\hat{\eta} - \eta_0\| = a_N = o(1)$, and $\delta_N a_N = o(n^{-1/2})$. Then

$$\sqrt{n}(\hat{\theta} - \theta_0) = -A^{-1}\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\psi(W_i;\theta_0,\eta_0) + o_P(1).$$

If, furthermore, $\psi(W;\theta_0,\eta_0) = \phi^\star(W)$, then $A = -I_d$ and

$$\sqrt{n}(\hat{\theta} - \theta_0) \;\rightsquigarrow\; N\Big(0,\; A^{-1}\,\mathbb{E}[\phi^\star\phi^{\star\top}]\,(A^{-1})^\top\Big),$$

achieving the semiparametric efficiency bound.

Proof. A first-order Taylor expansion of the sample moment at (θ_0, η_0) yields

$$0 = \mathbb{P}_n\psi(\hat{\theta},\hat{\eta}) = \mathbb{P}_n\psi(\theta_0,\eta_0) + A(\hat{\theta} - \theta_0) + B(\hat{\eta} - \eta_0) + r_n,$$

with remainder $r_n = o_P(n^{-1/2})$ under standard regularity (e.g., cross-fitting and stochastic equicontinuity). Multiply by $\sqrt{n}$ and rearrange:

$$\sqrt{n}(\hat{\theta} - \theta_0) = -A^{-1}\Big\{\sqrt{n}(\mathbb{P}_n - \mathbb{P})\psi(\theta_0,\eta_0) + \sqrt{n}\,B(\hat{\eta} - \eta_0)\Big\} + o_P(1).$$

By the near-orthogonality and rate assumptions, $\|\sqrt{n}\,B(\hat{\eta} - \eta_0)\| \leq \sqrt{n}\,\delta_N a_N = o(1)$, which gives the asserted linearization. If $\psi = \phi^\star$, the central limit theorem yields the stated Gaussian limit with covariance $A^{-1}\mathbb{E}(\phi^\star\phi^{\star\top})(A^{-1})^\top$. □

Corollary 12.2 (Relationship between IF/EIF and (near) Neyman orthogonality)

- *The EIF is orthogonal to the nuisance tangent space: $\phi^\star \perp \mathcal{T}_\eta$. Thus, using the EIF as a score produces exact Neyman orthogonality in the sense $\partial_\eta \mathbb{E}\psi = 0$ at the truth.*

- *Any IF in $\overline{\mathcal{T}}$ that represents the pathwise derivative coincides with the EIF and attains the minimal asymptotic variance.*
- *Near-orthogonal scores behave like the EIF up to $o(1)$ in $L^2(P_0)$; if $\delta_N a_N = o(n^{-1/2})$, they deliver root-$n$ asymptotics, and if the score equals the EIF at the truth, they achieve the semiparametric efficiency bound.*

References

[1] Dachun Yang, Wen Yuan, *Selected lectures on functional analysis* [M]. Beijing Normal University Press, 2016.