

Problem Set 1

1. 简答题

1. $|\frac{d\vec{v}}{dt}| = 0$ 表示加速度大小为0, 速度的大小和方向均不变
即静止或匀速直线运动

$\frac{d|\vec{v}|}{dt} = 0$ 表示速度大小不变而方向可能改变

2. 可以, 当物体的加速度方向与初速度方向不共线时, 物体运动方向即会改变


3. (1) 错误, 如匀速圆周运动

(2) 正确

(3) 错误, 如匀速圆周运动

4. 因为篮球在出手前已经具有一定初速度
应该在瞄准篮所需的出手速度上减去初速度再出手

2. 教材习题



(1) 无穷多次

(2) $t = \frac{x}{v_{\text{甲乙}}} = 1 \text{ h}$, $S = v_0 \cdot t = 60 \text{ km}$

1.3 (1) $t_1 = 3.0 \text{ s}$ 时, $x_1 = 96 \text{ m}$

$t_2 = 3.10 \text{ s}$ 时, $x_2 = 102.1 \text{ m}$

$t_3 = 3.01 \text{ s}$ 时, $x_3 = 96.601 \text{ m}$

$t_4 = 3.001 \text{ s}$ 时, $x_4 = 96.06001$

$$\bar{v}_1 = \frac{x_2 - x_1}{t_2 - t_1} = 61 \text{ m/s} \quad \bar{v}_2 = \frac{x_3 - x_1}{t_3 - t_1} = 60.1 \text{ m/s}$$

$$\bar{v}_3 = \frac{x_4 - x_1}{t_4 - t_1} = 60.01 \text{ m/s} \quad \vec{v}_1, \vec{v}_2, \vec{v}_3 \text{ 方向均沿 } x \text{ 正向}$$


$$(2) v = \dot{x} = \frac{dx}{dt} = 20t \quad t = 3.0s \text{ 时, } v = 60m/s$$

\vec{v} 方向沿 x 正向

$$(3) v = \dot{x} = \frac{dx}{dt} = \frac{20t dt}{dt} = 20t \text{ m/s}$$

$$a = \dot{v} = \ddot{x} = 20 \text{ m/s}^2$$

1.25



$$t = \frac{L}{\cos\beta \cdot v_0}$$

$$\frac{1}{2} g t^2 = L (\tan\beta - \tan\alpha)$$

$$\frac{gL^2}{2v_0^2 \cos^2\beta} = L \cdot \left(\frac{\sin\beta}{\cos\beta} - \frac{\sin\alpha}{\cos\alpha} \right)$$

$$\frac{gL}{2v_0^2} = \frac{\sin(\beta - \alpha) \cos\beta}{\cos\alpha}$$

$$v_0^2 = \frac{gL \cos\alpha}{2 \sin(\beta - \alpha) \cos\beta}$$

$$\therefore v_0 = \sqrt{\frac{gL \cos\alpha}{2 \sin(\beta - \alpha) \cos\beta}}$$


1.31 (1) 即 $a_n = a_t = 3m/s^2$

$$a_n = \frac{v^2}{R} = 3m/s^2 \quad \therefore v = 3m/s$$

$$t = \frac{v}{a_n} = 1s$$

$$(2) s = \frac{1}{2} a_n \cdot t^2 = 1.5m$$

1.33



以 O 为原点, 杆所在射线为径向建立极坐标系

$$\therefore \vec{v} = \vec{v}_n + \vec{v}_t = a_0 t \hat{r} + \frac{1}{2} a_0 t^2 \omega_0 \hat{\theta}$$

$$\vec{a} = \dot{\vec{v}} = a_0 \hat{r} + a_0 t \frac{d\hat{r}}{dt} + a_0 \omega_0 t \hat{\theta} + \frac{1}{2} a_0 \omega_0 t^2 \frac{d\hat{\theta}}{dt}$$

$$\frac{d\hat{\theta}}{dt} = \omega_0 \cdot \hat{r} \quad \frac{d\hat{r}}{dt} = \omega_0 \cdot \hat{\theta}$$

$$\therefore \vec{a} = \left(a_0 + \frac{1}{2} a_0 \omega_0^2 t^2 \right) \hat{r} + 2a_0 \omega_0 t \hat{\theta}$$

3. 补充习题

1. $\vec{A} = (3, 4, -4)$

(a) 设 $\vec{B} = (a, b, 0)$ $\begin{cases} \vec{B} \cdot \vec{A} = 3a + 4b = 0 \\ a^2 + b^2 = 1 \end{cases}$

$\therefore a = \frac{4}{5}, b = -\frac{3}{5} \quad \therefore \vec{B} = (\frac{4}{5}, -\frac{3}{5}, 0)$

(b) $\vec{C} = (x, y, z)$

$\begin{cases} \vec{C} \cdot \vec{A} = 3x + 4y - 4z = 0 \\ \vec{C} \cdot \vec{B} = \frac{4}{5}x - \frac{3}{5}y = 0 \end{cases}$

$3y = 4x$

取 $\vec{C} = (3, 4, \frac{25}{4})$

(c) $\vec{A} \perp \vec{B}, \vec{A} \perp \vec{C}, \vec{B}, \vec{C}$ 起点均为 O 点,

$\therefore \vec{A}$ 垂直于 \vec{B}, \vec{C} 所在的平面

2. ① $\vec{a} = (1, -2, 1), \vec{b} = (1, -1, 3), \vec{c} = (2, 5, -3)$

$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{e}_x & \vec{e}_y & \vec{e}_z \\ 1 & -2 & 1 \\ 1 & -1 & 3 \end{vmatrix} = \begin{vmatrix} -2 & 1 \\ 1 & 3 \end{vmatrix} \vec{e}_x - \begin{vmatrix} 1 & 1 \\ 1 & 3 \end{vmatrix} \vec{e}_y + \begin{vmatrix} 1 & -2 \\ 1 & -1 \end{vmatrix} \vec{e}_z$

$= (-5, -2, 1)$

② $(\vec{a} \times \vec{b}) \cdot \vec{c} = (-5, -2, 1) \cdot (2, 5, -3) = -23$

③ $\vec{a} \times (\vec{b} \times \vec{c}) = \vec{b}(\vec{a} \cdot \vec{c}) - \vec{c}(\vec{a} \cdot \vec{b})$

$= -11 \times (1, -1, 3) - 6 \times (2, 5, -3)$

$= (-11, 11, -33) - (12, 30, -18) = (-23, -19, -15)$

3. (a) $y' = 24x^2 + 1$

(b) $y' = (x-1)\tan x + (x+1)\tan x + (x^2-1)\frac{1}{\cos^2 x} = 2x\tan x + \frac{x^2-1}{\cos^2 x}$

(c) $y' = \frac{(2x+9)(5x+6) - 5(9x+x^2)}{(5x+6)^2} = \frac{10x^2 + 57x + 54 - 45x - 5x^2}{(5x+6)^2}$

$= \frac{5x^2 + 12x + 54}{25x^2 + 60x + 36}$

(d) $y' = \cos x - x \sin x + \frac{x \cos x - \sin x}{x^2}$

$f'(x) dx = df(x)$

4. (a) $\int (3x^2 + \sin x + \frac{5}{x}) dx = \int 3x^2 dx + \int \sin x dx + \int \frac{5}{x} dx$
 $= x^3 - \cos x + 5 \ln|x|$

(b) $\int \sqrt{a^2 - x^2} dx = a \int \sqrt{1 - \sin^2 x} d \sin x = a \int \sqrt{1 - \sin^2 x} a \cos x dx$
 $(a > 0)$

$$= a^2 \int \cos^2 x dx =$$

5. 代入: $y' = -C_1 \omega \sin \omega t - C_2 \omega \sin \omega t$

$$y'' = -C_1 \omega^2 \cos \omega t - C_2 \omega^2 \sin \omega t$$

$$\therefore y'' + \omega^2 y = 0$$

6. (a) $V_m = \int_0^T a(t) dt = \frac{a_m}{2} \int_0^T (1 - \cos \frac{2\pi}{T} t) dt = \frac{a_m}{2} (t - \frac{T}{2\pi} \sin \frac{2\pi}{T} t) \Big|_0^T = \frac{a_m T}{2}$

(b) 在 $T \ll T$ 时间内, 电梯可视为做匀加速运动

$$\therefore V_T = \int_0^T a(t) dt = \frac{a_m}{2} (t - \frac{T}{2\pi} \sin \frac{2\pi}{T} t) \Big|_0^T = \frac{a_m}{2} (T - \frac{T}{2\pi} \sin \frac{2\pi}{T} \cdot T)$$

$$\sin \frac{2\pi}{T} T = \sin 2\pi = 0 \quad \therefore V_T = 0$$


(c) $V(t) = \int_0^t a(t) dt = \frac{a_m}{2} (t - \frac{T}{2\pi} \sin \frac{2\pi}{T} t)$

$$X(t) = \int_0^t V(t) dt = \frac{a_m}{2} (\frac{1}{2} t^2 + \frac{T^2}{4\pi^2} \cos \frac{2\pi}{T} t)$$


则 0-T 时间内电梯运行距离为 $X(T) = \frac{a_m}{2} (\frac{1}{2} T^2 - \frac{T^2}{4\pi^2})$

$$\text{则 } D = X_{\text{总}} = 2X(T) = \frac{2\pi^2 - 1}{4\pi^2} a_m T^2 \quad \therefore T = \sqrt{\frac{4\pi^2 D}{(2\pi^2 - 1) a_m}}$$

$$\therefore t_1 = 2T = 4\pi \sqrt{\frac{D}{(2\pi^2 - 1) a_m}}$$



设初速度为 V_0 , 运动时间为 t



$$S = \frac{1}{2} g t^2 V_0 \cos \theta$$

$$V_0 t \cos \theta = \frac{2S}{g}$$

S 取最大值时, 位移最远

$$\theta + \varphi = \frac{1}{2} (\pi - (\frac{\pi}{2} - \varphi)) = \frac{1}{2} (\frac{\pi}{2} + \varphi) = \frac{\pi}{4} + \frac{\varphi}{2}$$

$$\therefore \theta = \frac{\pi}{4} - \frac{\varphi}{2}$$

8. (a) $r = \frac{r_0}{1 + e \cos \theta}$, $\theta = 0, r_1 = \frac{r_0}{1 + e}$; $\theta = \pi, r_2 = \frac{r_0}{1 - e}$

$$r_1 + r_2 = r_0 (\frac{1}{1 + e} + \frac{1}{1 - e}) = \frac{2r_0}{1 - e^2} = 2A \quad \therefore r_0 = (1 - e^2) A, e = \frac{\sqrt{A^2 - B^2}}{A}$$

(b) $\vec{r}_{10} = \frac{r_0}{1 + e \cos \theta} \hat{r}$

$$\vec{v}_\theta = \dot{\vec{r}}_{10} = \frac{(1 + e \cos \theta + e \sin \theta r_0) \omega}{(1 + e \cos \theta)^2} \hat{r} + \frac{r_0}{1 + e \cos \theta} \dot{\theta} \hat{\theta}$$

$$= \frac{(1 + e \cos \theta + e \sin \theta r_0) \omega}{(1 + e \cos \theta)^2} \hat{r} + \frac{r_0 \omega}{1 + e \cos \theta} \hat{\theta}$$

$$V_\theta = \sqrt{\left(\frac{(1 + e \cos \theta + e \sin \theta r_0) \omega}{(1 + e \cos \theta)^2} \right)^2 + \left(\frac{r_0 \omega}{1 + e \cos \theta} \right)^2}$$

$$\vec{a}_\theta = \dot{\vec{v}}_\theta =$$

$$9. \quad \vec{r} = r_0 e^{\alpha\theta} \hat{r}$$

$$\begin{aligned} \vec{v} = \dot{\vec{r}} &= r_0 \alpha e^{\alpha\theta} \cdot \dot{\theta} \hat{r} + r_0 e^{\alpha\theta} \dot{\hat{r}} \\ &= r_0 \alpha e^{\alpha\theta} \dot{\theta} \hat{r} + r_0 e^{\alpha\theta} \dot{\theta} \hat{\theta} \end{aligned}$$

$$\hat{v} = \frac{\alpha \hat{r} + \hat{\theta}}{\sqrt{\alpha^2 + 1}} \quad \hat{n} = \frac{\hat{r} - \alpha \hat{\theta}}{\sqrt{\alpha^2 + 1}}$$

$$\begin{aligned} \vec{a} = \dot{\vec{v}} &= (\ddot{r} - r\dot{\theta}^2) \hat{r} + (r\ddot{\theta} + 2\dot{r}\dot{\theta}) \hat{\theta} \\ &= [r\dot{\theta}^2(\alpha^2 - 1) + \alpha r\ddot{\theta}] \hat{r} + [r\ddot{\theta} + 2\alpha r\dot{\theta}^2] \hat{\theta} \end{aligned}$$

$$a_n = \vec{a} \cdot \hat{n} = \frac{1}{\sqrt{\alpha^2 + 1}} \{ [r\dot{\theta}^2(\alpha^2 - 1) + \alpha r\ddot{\theta}] - \alpha [r\ddot{\theta} + 2\alpha r\dot{\theta}^2] \}$$

$$= \frac{1}{\sqrt{\alpha^2 + 1}} [-r\dot{\theta}^2(\alpha^2 + 1)] \quad \therefore |a_n| = r\dot{\theta}^2 \sqrt{\alpha^2 + 1}$$

$$v^2 = r^2 \dot{\theta}^2 (\alpha^2 + 1)$$

$$\rho = \frac{v^2}{|a_n|} = \frac{r^2 \dot{\theta}^2 (\alpha^2 + 1)}{r \dot{\theta}^2 \sqrt{\alpha^2 + 1}} = r \sqrt{1 + \alpha^2} = \sqrt{1 + \alpha^2} r_0 e^{\alpha\theta}$$