

A. Refined module-selection homepage

Knee OA Risk Calculator

Home

Early OA Risk

Arthroplasty Risk

Model Information

About

Knee OA Stage-Specific Risk Calculator

Estimate early radiographic progression and future knee arthroplasty risk using stage-matched clinical prediction models.

Select the module that matches the patient's current radiographic stage.

中文名称：膝骨关节炎阶段特异性风险计算器

Disease-stage framework

Radiographic progression to arthroplasty risk

KL 0-1
symptomatic
knee

→

KL ≥2
definite
OA

→

KL 3/4
advanced
OA

→

TKA/KR
endpoint



→



KL 0-1 → KL ≥2

EARLY OA

24-Month Radiographic Progression

Estimate the probability of progression from KL grade 0-1 to definite radiographic OA (KL ≥2).

CHECK model

Symptomatic knees

Baseline KL 0-1



→



KL 0-4 → TKA/KR

ESTABLISHED OA

60-Month Knee Arthroplasty Risk

Estimate the probability of target-knee arthroplasty during the 60 months after landmark assessment.

OAI model

24-month landmark

KL 0-4

B. Early radiographic progression module

Knee OA Risk Calculator

Home

Early OA Risk

Arthroplasty Risk

Model Information

About

24-Month Early Radiographic Progression Risk

This model estimates the probability that a symptomatic knee with baseline KL grade 0 or 1 will progress to KL grade ≥2 within 24 months.

Patient characteristics

Age, years
56
Typical development cohort range: 30-85 years.

Sex
Female
Typical development cohort range: 30-85 years.

Body mass index, kg/m2
26.1
Typical development cohort range: 15-60 kg/m2.

Knee side
Left

Symptoms and radiographic status

Baseline KL grade
KL0

WOMAC pain score, 0-20
4.9
Use the original CHECK WOMAC pain score from 0 to 20.

WOMAC function score, 0-68
15.5

Baseline knee pain
Yes

CHECK early OA model

Estimated 24-month progression risk

5.6%

Probability of incident KL grade ≥2 within 24 months.

Lower estimated risk

Main contributors

- Presence of baseline knee pain

Clinical interpretation

This estimate may help identify symptomatic knees that are more likely to develop definite radiographic OA over the next 24 months. It may support risk communication, preventive management, and selection of patients for structured follow-up.

Download clinical summary

C. Knee arthroplasty risk module

Knee OA Risk Calculator

Home

Early OA Risk

Arthroplasty Risk

Model Information

About

60-Month Knee Arthroplasty Risk

This model estimates the probability of target-knee TKA/KR during the 60 months following the 24-month landmark assessment.

Patient characteristics

Age at landmark, years
63

Sex
Female

Knee side
Left

Current disease status at landmark

Current pain scores, 0-10
3.8

Current KL grade
KL2

Change during the previous 24 months

Pain score 24 months earlier, 0-10
2.8

KL grade 24 months earlier
KL2

Computed change pain 1.5, KL 0. Positive values indicate worsening.

OAI arthroplasty model

Estimated 60-month TKA/KR risk

11.1%

Estimated probability of target-knee arthroplasty during the 60 months after landmark assessment.

Risk group: Middle predicted-risk tertile

Main contributors

- Current pain burden on the harmonized 0-10 scale
- Recent pain worsening, with modest added information after current pain

Research thresholds

15% threshold
Estimate is above the research threshold.
High-risk knees: n= 61

25% threshold
Estimate is below the research threshold.
High-risk knees: n= 14

35% threshold
Estimate is below the research threshold.
High-risk knees: n= 6