

Knee OA Risk Calculator

[Home](#)
[Early OA Risk](#)
[Arthroplasty Risk](#)
[Model Information](#)
[About](#)

Knee OA Stage-Specific Risk Calculator

Estimate early radiographic progression and future knee arthroplasty risk using stage-matched clinical prediction models.

Select the module that matches the patient's current radiographic stage.

中文名称：膝骨关节炎阶段特异性风险计算器

Disease-stage framework

Radiographic progression to arthroplasty risk

KL 0-1 → KL >=2

EARLY OA

24-Month Radiographic Progression

Estimate the probability of progression from KL grade 0-1 to definite radiographic OA (KL >=2).

CHECK model

Symptomatic knees

Baseline KL 0-1

KL 0-4 → TKA/IR

ESTABLISHED OA

60-Month Knee Arthroplasty Risk

Estimate the probability of target-knee arthroplasty during the 60 months after landmark assessment.

OAI model

24-month landmark

KL 0-4

24-Month Early Radiographic Progression Risk

This model estimates the probability that a symptomatic knee with baseline KL grade 0 or 1 will progress to KL grade ≥ 2 within 24 months.

[Back to home](#)

Patient characteristics

Age, years

55

Typical development cohort range: 30-60 years.

Sex

Female

Body mass index, kg/m²

26.1

Typical development cohort range: 15-40 kg/m².

Knee side

Left

Symptoms and radiographic status

Baseline KL grade

KL0

WOMAC pain score, 0-20

4.9

Use the original Chicago WOMAC pain scale from 0 to 20.

WOMAC function score, 0-48

15.5

Baseline knee pain

Yes

[CHECK early OA model](#)

Estimated 24-month progression risk

5.6%

Probability of incident KL grade ≥ 2 within 24 months.

[Lower estimated risk](#)

Main contributors

- Presence of baseline knee pain

Clinical interpretation

This estimate may help identify symptomatic knees that are more likely to develop definite radiographic OA over the next 24 months. It may support risk communication, preventive management, and selection of patients for structured follow-up.

[Download clinical summary](#)

60-Month Knee Arthroplasty Risk

This model estimates the probability of target-knee TKA/AR during the 60 months following the 24-month landmark assessment.

Patient characteristics

Age at landmark, years

63

Sex

Female

Knee side

Left

Current disease status at landmark

Current pain score, 0-10

3.8

Current KJ, grade

KL2

Change during the previous 24 months

Pain score change, pain 0-10, 0-10

2.5

KJ, grade 24-month earlier

KL2

Computed change: pain 1.3, KL2. Positive values indicate worsening.

[Calculate risk](#)
[Start new calculation](#)

[Old arthroplasty model](#)

Estimated 60-month TKA/AR risk

11.1%

Estimated probability of target-knee arthroplasty during the 60 months after landmark assessment.

Risk group: Middle predicted-risk tertile

Main contributors

- Current pain worsening on the harmonized 0-10 scale
- Recent pain worsening, with model added information after current pain

Research thresholds

10% threshold	25% threshold	35% threshold
Estimate is above this research threshold	Estimate is below this research threshold	Estimate is below this research threshold
High-risk knee: 6.4%	High-risk knee: 6.4%	High-risk knee: 6.4%