



Contents lists available at ScienceDirect

Journal of Algebra

[www.elsevier.com/locate/jalgebra](http://www.elsevier.com/locate/jalgebra)



# On finite $p$ -groups of supersoluble type

A. Ballester-Bolinches<sup>a,b</sup>, R. Esteban-Romero<sup>b,c</sup>, H. Meng<sup>d</sup>,  
N. Su<sup>e,\*</sup>

<sup>a</sup> Department of Mathematics, Guangdong University of Education, 510310, Guangzhou, People's Republic of China

<sup>b</sup> Departament de Matemàtiques, Universitat de València, Dr. Moliner, 50, 46100 Burjassot, València, Spain

<sup>c</sup> Institut Universitari de Matemàtica Pura i Aplicada, Universitat Politècnica de València, Camí de Vera, s/n, 46022 València, Spain<sup>1</sup>

<sup>d</sup> Department of Mathematics, Shanghai University, Shanghai, 200444, People's Republic of China

<sup>e</sup> Department of Mathematics, Sun Yat-sen University, Guangzhou, 510275, People's Republic of China



## ARTICLE INFO

### Article history:

Received 23 July 2019

Available online 15 September 2020

Communicated by Markus Linckelmann

### MSC:

20D20

20D15

20D05

### Keywords:

Finite group

Fusion system

Supersolubility

## ABSTRACT

A finite  $p$ -group  $S$  is said to be of supersoluble type if every fusion system over  $S$  is supersoluble. The main aim of this paper is to characterise the finite  $p$ -groups of supersoluble type. Abelian and metacyclic  $p$ -groups of supersoluble type are completely described. Furthermore, we show that the Sylow  $p$ -subgroups of supersoluble type of a finite simple group must be cyclic.

© 2020 Published by Elsevier Inc.

\* Corresponding author.

E-mail addresses: [Adolfo.Ballester@uv.es](mailto:Adolfo.Ballester@uv.es) (A. Ballester-Bolinches), [Ramon.Esteban@uv.es](mailto:Ramon.Esteban@uv.es), [resteban@mat.upv.es](mailto:resteban@mat.upv.es) (R. Esteban-Romero), [hangyangmenges@gmail.com](mailto:hangyangmenges@gmail.com) (H. Meng), [suning3@mail.sysu.edu.cn](mailto:suning3@mail.sysu.edu.cn) (N. Su).

<sup>1</sup> Permanent address.

## 1. Introduction

All groups considered in this paper will be finite.

A saturated fusion system  $\mathcal{F}$  over a  $p$ -group  $S$ ,  $p$  a prime, is a category whose objects are the subgroups of  $S$ , whose morphisms are monomorphisms between subgroups of  $S$ , and whose morphism sets satisfy certain axioms motivated by properties of conjugacy relations between  $p$ -subgroups of a group. If  $S$  is a Sylow  $p$ -subgroup of a group  $G$ , we can associate the saturated fusion system  $\mathcal{F}_S(G)$  over  $S$ , called the *fusion system* of  $G$ , whose morphisms are those homomorphisms induced by conjugation in  $G$ . We refer to [1] for a detailed introduction to the theory of saturated fusion systems: this book will be our reference for the notation, terminology and results.

In this paper, we continue the study, started in [14], of supersoluble saturated fusion systems.

**Definition 1** ([14, Definition 1.2]). Let  $S$  be a  $p$ -group and let  $\mathcal{F}$  be a saturated fusion system over  $S$ . We say that  $\mathcal{F}$  is *supersoluble* if there exists a series  $1 = S_0 \leq \cdots \leq S_m = S$  of subgroups of  $S$  such that  $S_i$  is strongly closed in  $S$  with respect to  $\mathcal{F}$  and  $S_{i+1}/S_i$  is cyclic for each  $i \in \{0, \dots, m-1\}$ .

It has been shown that supersoluble fusion systems are precisely the fusion systems of supersoluble groups (see [14, Proposition 1.3(b)]).

It is clear that every nilpotent saturated fusion system in the sense of [11] is supersoluble and every supersoluble saturated fusion system is soluble in the sense of [1, Part II, Definition 12.1].

**Definition 2.** A  $p$ -group  $S$  is said to be of *supersoluble type* if for every saturated fusion system  $\mathcal{F}$  over  $S$ ,  $\mathcal{F}$  is supersoluble.

We characterise the  $p$ -groups of supersoluble type in the following theorem.

**Theorem A.** *Let  $S$  be a  $p$ -group. Then  $S$  is of supersoluble type if and only if  $S$  is resistant and every  $p'$ -subgroup of  $\text{Aut}(S)$  is abelian of exponent dividing  $p-1$ .*

We will give several applications of the above characterisation theorem. One of the consequences is the following.

**Corollary 3.** *If  $S$  is a  $p$ -group of supersoluble type, then  $\text{Aut}(S)$  is soluble.*

Theorem A and Corollary 3 will be proved in the next section.

We then apply this characterisation to describe the abelian and metacyclic  $p$ -groups of supersoluble type in Theorems B and C. With all these results at hand, we can show that the Sylow  $p$ -subgroups of supersoluble type of a simple group must be cyclic

(Theorem D), and that the structure of metacyclic Sylow  $p$ -subgroups of a simple group is quite limited (Theorem 12).

## 2. Proof of Theorem A

Recall that a  $p$ -group  $S$  is called *resistant* if  $S$  is normal in every saturated fusion system over  $S$  (see [13]).

**Proof of Theorem A.** We prove first the necessity of the condition. Assume that  $S$  is of supersoluble type. Let  $\mathcal{F}$  be a saturated fusion system over  $S$ . By [14, Proposition 1.3(b)], there exists a supersoluble group  $K$  such that  $S$  is a Sylow  $p$ -subgroup of  $K$  and  $\mathcal{F} = \mathcal{F}_S(K)$ . Without loss of generality, we may assume that  $O_{p'}(K) = 1$ . Since  $K$  is supersoluble with  $O_{p'}(K) = 1$ , we have  $S \trianglelefteq K$  and thus  $S \trianglelefteq \mathcal{F}_S(K) = \mathcal{F}$ . Hence  $S$  is resistant.

Let  $H$  be a  $p'$ -subgroup of  $\text{Aut}(S)$ . We will show that  $H$  is abelian of exponent dividing  $p - 1$ . Set  $G = S \rtimes H$ , the natural semidirect product of  $S$  and  $H$ . Clearly  $C_G(S) \leq S$  since  $C_H(S) = 1$ . Write  $\mathcal{F} = \mathcal{F}_S(G)$ . As  $\mathcal{F}$  is a saturated fusion system over  $S$ , it follows that  $\mathcal{F}$  is supersoluble. By [14, Proposition 1.3(d)],  $\text{Aut}_{\mathcal{F}}(S)$  is  $p$ -closed and a Hall  $p'$ -subgroup of  $\text{Aut}_{\mathcal{F}}(S)$  is abelian of exponent dividing  $p - 1$ . Note that  $\text{Aut}_{\mathcal{F}}(S) = \text{Aut}_G(S) \cong N_G(S)/C_G(S) = HS/C_G(S)$ . Then  $HC_G(S)/C_G(S)$  is a Hall  $p'$ -subgroup of  $G/C_G(S)$  and so  $H \cong HC_G(S)/C_G(S)$  is abelian of exponent dividing  $p - 1$ .

We prove now the sufficiency of the condition. Assume that  $S$  is resistant and every  $p'$ -subgroup of  $\text{Aut}(S)$  is abelian of exponent dividing  $p - 1$ . Let  $\mathcal{F}$  be a saturated fusion system over  $S$ . We shall show that  $\mathcal{F}$  is supersoluble. As  $S$  is resistant,  $S \trianglelefteq \mathcal{F}$  and  $\mathcal{F}$  is constrained. Since  $S \trianglelefteq \mathcal{F}$ , it is clear that  $\mathcal{F}$  is the fusion system of a finite group  $G = S \rtimes H$  for some  $p'$ -subgroup  $H$  of  $\text{Aut}(S)$ . It then follows from the assumption that  $H$  is abelian of exponent dividing  $p - 1$ . Thus  $G$  is soluble and every  $p$ -chief factor of  $G$  is cyclic by [5, Chapter B, Theorem 9.8] and every  $p'$ -chief factor is central. Consequently,  $G$  is supersoluble. By [14, Proposition 1.3(b)],  $\mathcal{F} = \mathcal{F}_S(G)$  is supersoluble. We conclude then that  $S$  is of supersoluble type.  $\square$

**Corollary 4.** *Let  $S$  be a 2-group. Then  $S$  is of supersoluble type if and only if  $S$  is resistant and  $\text{Aut}(S)$  is a 2-group.*

We can then obtain Corollary 3 by combining Theorem A and the following result.

**Theorem 5.** *If every  $p'$ -subgroup of a group  $G$  is abelian of exponent dividing  $p - 1$ , then  $G$  is soluble.*

The proof of Theorem 5 requires the following lemma.

**Lemma 6.** Let  $G = \text{PSL}_2(r^f)$ , where  $r$  is a prime and  $f \geq 1$ . Set  $u = (r^f - 1)/k$  and  $s = (r^f + 1)/k$ , where  $k = (r^f - 1, 2)$ . Then  $G$  has two cyclic subgroups  $U$  and  $S$  of orders  $u$  and  $s$ , respectively. Moreover  $N_G(U)$  is dihedral of order  $2u$  and  $N_G(S)$  is dihedral of order  $2s$ .

**Proof.** It is a consequence of [9, Kapitel II, Satz 8.3 and 8.4].  $\square$

**Proof of Theorem 5.** Let  $G$  be a counterexample of minimal order. Then  $p$  is the largest prime dividing the order of  $G$ . By [5, Chapter I, Section 2],  $p \neq 2$ , and  $p \neq 3$ . The minimal choice of  $G$  implies that every proper subgroup and every nontrivial epimorphic image of  $G$  are soluble. Hence  $G$  is a minimal simple group.

By a result of Thompson (see [9, Kapitel II, Bemerkung 7.5]),  $G$  is isomorphic to one of the following groups:

1.  $\text{PSL}_2(q)$ , where  $q > 3$  is a prime and  $5 \nmid q^2 - 1$ ;
2.  $\text{PSL}_2(2^q)$ , where  $q$  is a prime;
3.  $\text{PSL}_2(3^q)$ , where  $q$  is an odd prime;
4.  $\text{PSL}_3(3)$ ;
5. the Suzuki group  $\text{Sz}(2^q)$ , where  $q$  is an odd prime.

If  $G \cong \text{PSL}_3(3)$ , then  $|G| = 13 \cdot 3^3 \cdot 2^4$  and  $p = 13$ . Observe that  $\text{PSL}_3(3) \cong \text{SL}_3(3)$  has a subgroup isomorphic to  $\text{SL}_2(3)$ , which is a nonabelian  $13'$ -group, contrary to assumption. Hence  $G$  cannot be isomorphic to  $\text{PSL}_3(3)$ . If  $G \cong \text{Sz}(2^q)$ , where  $q$  is an odd prime, we can apply [10, Chapter XI, Lemma 3.1(a) and Theorem 3.3] to conclude that  $G$  has a nonabelian Sylow 2-subgroup. This contradiction shows that  $G$  is not isomorphic to  $\text{Sz}(2^q)$  for any odd prime  $q$ .

Assume that  $G = \text{PSL}_2(r^f)$  for some prime  $r$  and integer  $f$  such that  $r^f \geq 4$ . Then  $|G| = r^f(r^f - 1)(r^f + 1)k^{-1}$ , where  $k = (2, r^f - 1)$ . Observe that  $(r^f + 1, r^f - 1) = 1$  or  $2$ . As  $p \neq 2$ , we can conclude that  $r^f - 1$  or  $r^f + 1$  is a  $p'$ -number. Suppose that  $r^f = 4$  or  $5$ . By [9, Kapitel II, Satz 6.14],  $G \cong \text{PSL}_2(4) \cong \text{PSL}_2(5)$  is isomorphic to the alternating group of degree 5 which has a nonabelian  $p'$ -subgroup isomorphic to the alternating group of degree 4. This contradiction yields  $r^f \geq 6$ . Then  $2(r^f + 1/k) > 2(r^f - 1/k) > 4$ ; by Lemma 6,  $G$  has dihedral  $p'$ -subgroups. This final contradiction proves the theorem.  $\square$

### 3. Abelian and metacyclic $p$ -groups of supersoluble type

Our aim in this section is to characterise the abelian and metacyclic  $p$ -groups of supersoluble type. Such characterizations will be used later in Section 4 to investigate the structure of Sylow  $p$ -subgroups of supersoluble type of finite simple groups and the structure of metacyclic Sylow  $p$ -subgroups of finite simple groups.

We need two preliminary lemmas. The first one is elementary.

**Lemma 7.** *Let  $P$  be a group isomorphic to  $C_{p^n} \times C_{p^n}$  for some positive integer  $n$ . Then  $\text{Aut}(P)$  has a quotient isomorphic to  $\text{GL}_2(p)$ .*

The second lemma is a consequence of Theorem A.

**Lemma 8.** *Let  $S$  be a resistant  $p$ -group. Assume that  $S$  has a series of characteristic subgroups  $\Phi(S) = D_0 \leq D_1 \leq \cdots \leq D_n = S$  such that  $D_i/D_{i-1}$  is cyclic for each  $0 < i \leq n$ . Then  $S$  is of supersoluble type.*

**Proof.** Without loss of generality, we may assume that  $D_i/D_{i-1}$  is of order  $p$  for  $i = 1, \dots, n$ . Let  $H$  be a  $p'$ -subgroup of  $\text{Aut}(S)$ , and let  $H^*$  be the smallest normal subgroup of  $H$  such that  $H/H^*$  is an abelian group of exponent dividing  $p - 1$ . We want to show that  $H^* = 1$ . Let  $1 \leq i \leq n$ . Then  $H/C_H(D_i/D_{i-1})$  is isomorphic to a subgroup of  $\text{Aut}(D_i/D_{i-1}) \cong \text{Aut}(C_p)$  and so  $H/C_H(D_i/D_{i-1})$  is abelian of exponent dividing  $p - 1$ . Thus  $H^* \leq C_H(D_i/D_{i-1})$  for all  $i$ . Therefore  $H^*$  stabilises the chain  $S = D_n \geq D_{n-1} \geq \cdots \geq D_0 = \Phi(S)$ . By [5, Chapter I, Lemma 1.5],  $H^*$  acts trivially on  $D = S/\Phi(S)$ . It follows from [5, Chapter I, Proposition 1.7] that  $H^* = 1$ . Consequently,  $S$  is of supersoluble type by Theorem A.  $\square$

**Theorem B.** *Let  $S$  be an abelian  $p$ -group of type  $(m_1, \dots, m_t)$ . Then  $S$  is of supersoluble type if and only if  $m_1, \dots, m_t$  are all distinct.*

**Proof.** We can assume, arguing by contradiction, that  $S$  is of supersoluble type and  $m_1, \dots, m_t$  are not all distinct. Without loss of generality we may suppose that  $m_1 = m_2 = n$ . Then  $S = P \times H$ , where  $P, H \leq S$  and  $P \cong C_{p^n} \times C_{p^n}$ . By Lemma 7,  $\text{Aut}(P)$  has a quotient isomorphic to  $\text{GL}_2(p)$ . Observe that  $\text{Aut}(P) \times \text{Aut}(H)$  is a subgroup of  $\text{Aut}(S)$ . Thus  $\text{Aut}(S)$  has a section isomorphic to  $\text{GL}_2(p)$ . Suppose that  $p = 2$ . Since  $\text{GL}_2(2) \cong S_3$ , it follows that  $\text{Aut}(S)$  is not a 2-group. Hence  $p > 2$  by Corollary 4. If  $p$  is odd, the Sylow 2-subgroups of  $\text{SL}_2(p)$  are nonabelian by [9, Kapitel II, Hauptsatz 8.27]. This contradicts Theorem A. Consequently,  $m_1, \dots, m_t$  are distinct.

Assume that  $m_1, \dots, m_t$  are all distinct and  $m_1 < m_2 < \cdots < m_t$ . We shall show that  $S$  is of supersoluble type. By [1, Part I, Corollary 4.7],  $S$  is resistant. Let  $D_i = \Omega_i(S)\Phi(S)$ , where  $\Omega_i(S)$  is the subgroup generated by all elements of  $S$  of order dividing  $p^i$ . Then there exists a positive integer  $n$  such that

$$\Phi(S) = D_0 \leq D_1 \leq \cdots \leq D_n = S. \quad (1)$$

Then (1) is a characteristic series of  $S$  such that  $D_i/D_{i-1}$  is cyclic for each  $0 < i \leq n$ . By Lemma 8,  $S$  is of supersoluble type.  $\square$

**Theorem C.** *Let  $S$  be a metacyclic  $p$ -group. Then  $S$  is of supersoluble type if and only if  $S$  is none of the following groups:*

1. the abelian group  $C_{p^n} \times C_{p^n}$  for some positive integer  $n$ ,
2. dihedral, semidihedral or generalised quaternion if  $p = 2$ .

**Proof.** First assume that  $p = 2$  and let  $S$  be a metacyclic 2-group. Applying [4, Theorems 1.1] and Corollary 4, we have that  $S$  is of supersoluble type if and only if  $S$  is none of the groups listed in the statement of the theorem.

Now assume that  $p$  is odd. If  $S$  is an abelian  $p$ -group, then by Theorem B,  $S$  is of supersoluble type if and only if  $S$  is not isomorphic to  $C_{p^n} \times C_{p^n}$  for any positive integer  $n$ .

Suppose that  $S$  is a nonabelian metacyclic  $p$ -group. We prove that  $S$  is of supersoluble type. By [13, Proposition 5.4],  $S$  is resistant. Since  $S'$  is a nontrivial cyclic subgroup of  $S$ , we can apply [9, Kapitel III, Satz 10.2(c)] to conclude that  $S$  is regular.

Assume that the exponent of  $S$  is  $p^m$ . Since  $S$  is regular, we can apply [9, Kapitel III, Hauptsatz 10.5(b)] to conclude that

$$\mathcal{U}_{m-1}(S) = \langle x^{p^{m-1}} : x \in S \rangle = \{x^{p^{m-1}} : x \in S\}.$$

Moreover, by [9, Kapitel III, Satz 10.6],  $\mathcal{U}_{m-1}(S)$  is elementary abelian. Since  $\mathcal{U}_{m-1}(S)$  is metacyclic, we have that  $\mathcal{U}_{m-1}(S) \cong C_p$  or  $\mathcal{U}_{m-1}(S) \cong C_p \times C_p$ .

Suppose that  $\mathcal{U}_{m-1}(S) \cong C_p$ . By [9, Kapitel III, Satz 10.7 (a)], we have that  $|S/\Omega_{m-1}(S)| = |\mathcal{U}_{m-1}(S)| = p$ . Since  $S$  is 2-generated,  $|S/\Phi(S)| = p^2$ . Hence  $\Phi(S) \leq \Omega_{m-1}(S) \leq S$  is a characteristic series of  $S$  with cyclic factors. Applying Lemma 8, we conclude that  $S$  is of supersoluble type.

Suppose that  $\mathcal{U}_{m-1}(S) \cong C_p \times C_p$ . Since  $S'$  is a nontrivial cyclic subgroup of  $S$ , we have  $\mathcal{U}_{m-1}(S)$  is not contained in  $S'$ . Moreover  $\mathcal{U}_{m-1}(S) \cap S' \neq 1$ , because otherwise  $\mathcal{U}_{m-1}(S)S' = \mathcal{U}_{m-1}(S) \times S' \cong (C_p \times C_p) \times C_{p^t}$ ,  $t > 0$ , would not be metacyclic. Hence  $|\mathcal{U}_{m-1}(S) \cap S'| = p$ .

Let  $D$  be the subgroup of  $S$  such that  $\Omega_{m-1}(S/S') = D/S'$ . Clearly  $D$  is a characteristic subgroup of  $G$ , and

$$\begin{aligned} |S : D| &= |S/S' : D/S'| = |S/S' : \Omega_{m-1}(S/S')| = |\mathcal{U}_{m-1}(S/S')| \\ &= |\mathcal{U}_{m-1}(S)S'/S'| = |\mathcal{U}_{m-1}(S) : \mathcal{U}_{m-1}(S) \cap S'| = p. \end{aligned}$$

It follows that  $\Phi(S) \leq D \leq S$  is a characteristic series of  $S$  with  $|S/D| = |D/\Phi(S)| = p$ . By Lemma 8,  $S$  is of supersoluble type.  $\square$

#### 4. Simple groups with Sylow $p$ -subgroups of supersoluble type

The aim of this section is to determine the Sylow  $p$ -subgroups of simple groups that are of supersoluble type. As an application we also determine the metacyclic Sylow  $p$ -subgroups of simple groups. This is achieved in the last two theorems in the section and requires some preliminary results. The first lemma is well known.

**Lemma 9.** *If  $p$  is an odd prime, then the group  $\mathrm{SL}_2(p)$  has an element of order  $p + 1$ .*

**Lemma 10.** *If  $S$  is an extraspecial group of order  $p^3$  and exponent  $p$ , with  $p$  an odd prime, then  $S$  is not of supersoluble type.*

**Proof.** It is enough to prove the existence of a  $p'$ -automorphism of  $S$  with order not dividing  $p - 1$ . Since every  $p'$ -automorphism of the elementary abelian quotient of  $S$  lifts to  $S$ , Lemma 9 yields that  $\mathrm{Aut}(S)$  has an element of order divisible by  $p + 1$ . Since  $p + 1$  is not a divisor of  $p - 1$ , the result follows as a consequence of Theorem A.  $\square$

**Lemma 11.** *The Sylow  $p$ -subgroups of  $G = \mathrm{PSU}_3(q)$  for  $q$  a power of the prime  $p$  are not of supersoluble type.*

**Proof.** As in [9, Kapitel II, Satz 10.12], we consider  $\mathrm{GU}_3(q)$  as the group of matrices  $M \in \mathrm{GL}_3(q^2)$  such that  $M^q J M = J$ , where

$$J = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

and  $\varphi$  acts on each entry of the matrix as the field automorphism  $x \mapsto x^q$ . Then the Sylow  $p$ -subgroups of  $\mathrm{PSU}_3(q)$  are isomorphic to the Sylow  $p$ -subgroups of  $\mathrm{GU}_3(q)$ , and there is a Sylow  $p$ -subgroup  $U$  of  $\mathrm{GU}_3(q)$  composed of matrices of the form

$$M(c, d) = \begin{bmatrix} 1 & c & d \\ 0 & 1 & -c^q \\ 0 & 0 & 1 \end{bmatrix},$$

where  $d \in \mathrm{GF}(q^2)$  and  $c \in \mathrm{GF}(q^2)$  satisfy  $cc^q = -(d + d^q)$  and multiplication given by  $M(c, d)M(e, f) = M(c + e, d + f - ce^q)$ . Let  $U$  be the set composed of all these matrices with  $c, d \in \mathrm{GF}(q^2)$  and  $cc^q = -(d + d^q)$ .

Let  $\zeta$  be a generator of the multiplicative group of  $\mathrm{GF}(q) \subseteq \mathrm{GF}(q^2)$ . Then the matrix

$$D = \begin{bmatrix} \zeta^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \zeta \end{bmatrix}$$

is an element of  $\mathrm{GU}_3(q)$  that induces by conjugation an automorphism  $\delta$  of  $\mathrm{GU}_3(q)$  such that  $D^{-1}M(c, d)D = M(\zeta c, \zeta^2 d)$ , and since  $\zeta^q = \zeta$ ,  $(\zeta c)(\zeta c)^q = \zeta^2 cc^q = -\zeta^2(d + d^q) = -(\zeta^2 d + (\zeta^2 d)^q)$  and  $M(\zeta c, \zeta^2 d) \in U$ . We note that this automorphism has order  $q - 1$ , because if  $\xi$  is a generator of the multiplicative group of  $\mathrm{GF}(q^2)$ , then  $A = M(1, -\xi/(\xi + \xi^q)) \in U$  and  $A^{\delta^t} = M(\zeta^t, -\zeta^{2t}\xi/(\xi + \xi^q))$ . Hence  $A^{\delta^t} = A$  if and only if  $q - 1 \mid t$ , and so the order of  $\delta$  is divisible by  $q - 1$ . If  $U$  is of supersoluble type, then by Theorem A we have that  $q = p$ , a prime number.

Suppose that  $G = \text{PSU}_3(p)$ , with  $p$  prime, then  $p > 2$  since  $\text{PSU}_3(2)$  is soluble. Let  $A = M(1, -\xi/(\xi + \xi^p))$ ,  $B = M(\xi, -\xi^2\xi^p/(\xi + \xi^p))$ , where  $\xi$  is a generator of  $\text{GF}(p^2)^\times$ . Since  $\xi - \xi^p \neq 0$ , these elements do not commute, because  $AB = M(\xi + 1, -\xi - \xi^2\xi^q/(\xi + \xi^q))$  and  $BA = M(\xi + 1, -\xi^q - \xi^2\xi^q/(\xi + \xi^q))$ . Moreover, the elements of  $U$  have order  $p$ , since  $M(c, d)^r = M(rc, rd - (r(r-1)/2)cc^p)$ . We conclude that  $U$  is an extraspecial group of order  $p^3$  and exponent  $p$ . By Lemma 10, this case is also ruled out.  $\square$

**Theorem D.** *Let  $S$  be a Sylow  $p$ -subgroup of a finite simple group  $G$ . If  $S$  is of supersoluble type, then  $S$  is cyclic.*

**Proof.** Since  $S$  is resistant by Theorem A, we have that  $S \trianglelefteq \mathcal{F}_S(G)$ . Then  $G$  is a  $p$ -Goldschmidt group (see [1, Part II, Definition 12.9]). According to results of Foote and Flores and Foote ([7, 6], see also [1, Part II, Theorem 12.10]),  $G$  is a  $p$ -Goldschmidt group if and only if one of the following conditions holds:

1.  $S$  is abelian.
2.  $G$  is of Lie type in characteristic  $p$  of Lie rank 1.
3.  $p = 5$  and  $G \cong \text{McL}$ .
4.  $p = 11$  and  $G \cong J_4$ .
5.  $p = 3$  and  $G \cong J_2$ .
6.  $p = 5$  and  $G \cong \text{HS}$ ,  $\text{Co}_2$ , or  $\text{Co}_3$ .
7.  $p = 3$  and  $G \cong G_2(q)$  for some prime power  $q$  prime to 3 such that  $q$  is not congruent to  $\pm 1$  modulo 9.
8.  $p = 3$  and  $G \cong J_3$ .

First assume that  $S$  is not abelian. In Cases 3–7, according to the Atlas [3], the Sylow  $p$ -subgroup is extraspecial of order  $p^3$  and exponent  $p$ . These cases are ruled out by Lemma 10. In Case 8, if  $p = 3$  and  $G = J_3$ , then  $|S| = 3^5$  and we can check with GAP [8] that  $\text{Aut}(S)$  is a  $\{2, 3\}$ -group whose Sylow 2-subgroup is isomorphic to a semidihedral group  $\text{QD}_{16}$  of order 16, therefore the Sylow 3-subgroup of  $J_3$  is not of supersoluble type by Theorem A. Consequently we can assume  $G$  is of Lie type in characteristic  $p$  and  $G$  has Lie rank 1. Since the Sylow  $p$ -subgroups of  $\text{PSL}_2(q)$  for  $q = p^f$  are isomorphic to the multiplicative group of the field  $\text{GF}(q)$ , that is abelian, we have that  $G$  is either isomorphic to  $\text{PSU}_3(q)$  for  $q = p^f$  a prime power, or to a Suzuki group  $\text{Sz}(2^{2m+1})$  for  $p = 2$ , or to a Ree group  ${}^2G_2(3^{2m+1})$  for  $p = 3$ . By Lemma 11, the Sylow  $p$ -subgroups of  $\text{PSU}_3(q)$  are not of supersoluble type. In the Suzuki and Ree cases, the field automorphism  $x \mapsto x^p$  induces an automorphism of the Sylow subgroup  $S$  of order  $2m+1 \geq 3$ , that cannot be a divisor of  $p-1$  and thus  $S$  is not of supersoluble type by Theorem A.

Therefore we can suppose that  $S$  is abelian. Assume that  $S$  is of type  $(m_1, \dots, m_t)$ . Now, by Theorem B, we know that  $m_1, \dots, m_t$  are all distinct. Moreover, it is shown in



[12] that  $S$  is isomorphic to a direct product of copies of a cyclic group. Hence  $S$  must be cyclic. This completes the proof of the theorem.  $\square$

By combining Theorem C, Theorem D, and [2, Theorem 1], we can determine the structure the metacyclic Sylow  $p$ -subgroups of finite simple groups.

**Theorem 12.** *Let  $S$  be a Sylow  $p$ -subgroup of a finite simple group  $G$ . If  $S$  is metacyclic, then  $S$  is one of the following:*

1.  $C_{p^n} \times C_{p^n}$  for some positive integer  $n$ ,
2. cyclic if  $p \neq 2$ ,
3. dihedral or semidihedral if  $p = 2$ .

**Remark 13.** It is clear that the classes of metacyclic  $p$ -group listed in Theorem 12 do occur as Sylow  $p$ -subgroups of some finite simple groups. For instance,  $A_7$ , the alternating group of degree 7, has dihedral Sylow 2-subgroups, has cyclic Sylow 7-subgroups, and has elementary Sylow 3-subgroups of order 9. And the linear group  $\mathrm{PSL}_3(7)$  has semidihedral Sylow 2-subgroups.

## Acknowledgments

The research of the first and the second author has been supported by the grants PGC2018-095140-B-I00 from the *Ministerio de Ciencia, Innovación y Universidades* and the *Agencia Estatal de Investigación*, Spain, and FEDER, European Union, and PROMETEO/2017/057 from the Generalitat, Valencian Community, Spain. The third author is supported by Shanghai Sailing Program (20YF1413400) and the National Science Foundation of China (12001359). The fourth author is supported by the National Science Foundation of China (11401597 and 12071092).

The authors would like to thank professor Yanjun Liu for helpful conversations.

The authors are grateful to the referee for his/her valuable suggestions.

## References

- [1] M. Aschbacher, R. Kessar, B. Oliver, *Fusion Systems in Algebra and Topology*, London Mathematical Society Lecture Note Series, vol. 391, Cambridge University Press, Cambridge, 2011.
- [2] R. Brauer, M. Suzuki, On finite groups of even order whose 2-Sylow group is a quaternion group, *Proc. Natl. Acad. Sci. USA* 45 (12) (December 1959) 1757–1759.
- [3] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, R.A. Wilson, *Atlas of Finite Groups*, Oxford Univ. Press, London, 1985.
- [4] D.A. Craven, A. Glessner, Fusion systems on small  $p$ -groups, *Trans. Am. Math. Soc.* 364 (11) (2012) 5945–5967.
- [5] K. Doerk, T. Hawkes, *Finite Soluble Groups*, De Gruyter Expositions in Mathematics, vol. 4, Walter de Gruyter & Co., Berlin, 1992.
- [6] R. Flores, R. Foote, Strongly closed subgroups of finite groups, *Adv. Math.* 222 (2) (2009) 453–484.
- [7] R. Foote, A characterization of finite groups containing a strongly closed 2-subgroup, *Commun. Algebra* 25 (2) (1997) 593–606.

- [8] The GAP Group, GAP – groups, algorithms, and programming, version 4.10.1, <http://www.gap-system.org>, 2019.
- [9] B. Huppert, Endliche Gruppen I, Grund. Math. Wiss., vol. 134, Springer Verlag, Berlin, Heidelberg, New York, 1967.
- [10] B. Huppert, N. Blackburn, Finite Groups III, Grund. Math. Wiss., vol. 243, Springer-Verlag, Berlin, Heidelberg, New York, 1982.
- [11] J. Liao, J. Zhang, Nilpotent fusion systems, *J. Algebra* 442 (2015) 438–454.
- [12] R. Shen, Y. Zhou, Finite simple groups with some abelian Sylow subgroups, *Kuwait J. Sci.* 43 (2) (2016) 1–15.
- [13] R. Stancu, Control of fusion in fusion systems, *J. Algebra Appl.* 5 (6) (2006) 817–837.
- [14] N. Su, On supersolvable saturated fusion systems, *Monatshefte Math.* 187 (1) (2018) 171–179.