## **Course Objectives**



- To learn DOE techniques based on probability theory and statistical tools
- Understand proper methods for data collection, defining quality parameters, diagnostic tools, quality analysis, and interpretation for process/product improvement.
- Statistical DOE enables understanding of the relationship between multiple input variables/factors and the key responses, or product/process performance.

In-depth knowledge of these tools is quintessential while conducting scientific experiments

## **List of Topics\***



#### **Fundamental Concepts and Methods**

- Quality philosophy and conceptual framework
- Statistical Methods and Probability Concepts for Data Characterization

#### **Classical Design of Experiments**

- Nature of variability, probability distributions
- Empirical models (regression, hypothesis testing, confidence intervals, applications)
- Two-level factorial designs (factor effects, ANOVA, residual analysis, interactions)
- General 2<sup>k</sup> factorial designs
- Two-level fractional factorial designs

#### **Response Surface Methodology**

- First and second-order models and surfaces
- Central composite designs
- Multiple response analysis, Design rotatability, Box-Behnken design

#### **Robust Design Method**

- Quality loss function, signal, noise and control factors, product life cycle
- Matrix experiments using orthogonal arrays, analysis of means and variance, error prediction
- Steps in robust design: noise factors and testing, signal to noise ratio, degrees of freedom, selection of orthogonal arrays
- Conducting matrix experiments: randomization, confounding, result interpretation, verification
- Factor interactions, dynamic problems

#### Spring 2025

### **ME** 794 Statistical Design of Experiments

### **Chapter 1**

# **Fundamental Concepts**

**Prof. Soham Mujumdar** 

Email: sohammujumdar@iitb.ac.in



**NOTE:** Some of the course material is adopted from 'Design and Analysis of Experiments' by D. C. Montgomery, and similar courses taught by Prof. S. G. Kapoor at the University of Illinois at Urbana-Champaign and Prof. S. S. Joshi at IIT Bombay. You do NOT have permission to share this file or any of its contents with anyone else, and/or upload it on the internet or any of the platforms where it can be accessed by others.



Question: Is Sachin Tendulkar a 'good' batsman?

How would you answer that?

- You need some data to decide ...
- In general, when we collect data, we are interested in,
  - How the process/product is behaving in terms of an output quality characteristic(s)
  - Perhaps, in terms of (a) average value of the characteristic, (b) the variation in individual measurements of the characteristic






#### How would you collect the data?

In general, when we collect data, we have three choices,

 Observe the process ONCE, and use that observation as the absolute reflection of the process behavior

'Fatal Error!'



Observing **entire population/universe**, i.e. all possible realizations of the process – a very large number. **Neither practical nor necessary** 

 Observe PART OF the output of the process and use it to infer something about the process behavior

More 'practical'! We are 'sampling' the process (population)





#### Sachin Tendulkar stats

Batsmar

Career Batting Stats
Right-Handed Batsman

ormat	M	Inn	NO	Runs	HS	Avg	BF	BF	SR	100s	50s
T <b>est</b> 989–13	200	329	33	15921	248*	53.8	29437	379	-	51	68
<b>DDI</b> 989–12	463	452	41	18426	200*	44.8	21367	147	86.2	49	96
Γ <b>20Ι</b> 2006	1	1	0	10	10	10.0	12	12	83.3	0	0
PL 2008–13	78	78	11	2334	100*	34.8	1948	66	119.8	1	13



We need an efficient and adequate method to collect and analyze the data

#### **Critical Issues in Sampling**

- How much do we sample?
- How do we sample?
- When do we sample?
- How/what can we say something about the process from the information contained in the sample?

We will keep exploring these answers throughout the course...

For now, it is important to understand the difference between a Sample vs. the Population






#### Sachin Tendulkar stats


Batsmar

Career Batting Stats

ormat	M	lnn	NO	Runs	HS	Avg	BF	BF	SR	100s	50s
Test 1989–13	200	329	33	15921	248*	53.8	29437	379	-	51	68
<b>ODI</b> 1989–12	463	452	41	18426	200*	44.8	21367	147	86.2	49	96
<b>T20I</b> 2006	1	1	0	10	10	10.0	12	12	83.3	0	0
<b>PL</b> 2008–13	78	78	11	2334	100*	34.8	1948	66	119.8	1	13







## **Sources of Variability**



Taguchi suggests that variation in product and process function arises from three basic sources:

- Outer Noise: Sources of noise which influence performance as measured during field use under actual operating conditions, e.g., temp, humidity, supply voltage, vibration
- Inner Noise: Internal change in product characteristics such as drift from the nominal over time due to deterioration, e.g., mechanical wear, aging
- Variational Noise: Variation in the product parameters from one unit to another as a result of the manufacturing process, e.g., manufacturing imperfection

## **Example 1: Refrigerator**



#### Outer Noise

#### [Operating Conditions]

- The number of times the door is opened and closed
- The amount of food kept and the initial temperature of the food
- Variation in the ambient temperature
- Supply voltage variation

#### • Inner Noise

#### [Deterioration]

- The leakage of Refrigerant
- Mechanical Wear of Compressor parts

#### Variational Noise [Mfg/Use Imperfection]

- The tightness of door closure
- The amount of refrigerant used



## **Example 2: Braking Distance of a Car**



#### Outer Noise

#### [Operating Conditions]

- Wet or dry road
- Concrete or Asphalt pavement
- Number of passengers in the car

#### • Inner Noise

#### [Deterioration]

- The leakage of brake fluid
- Wear of brake drums and brake pads
- Variational Noise [Mfg/Use Imperfection]
  - Variation in friction coefficient of pads and drums
  - The amount of brake fluid



### **Data Characterization**




Match No.	Runs	Balls
463	52	48
462	114	147
461	6	19
460	39	30
459	14	15
458	22	23
457	3	12
456	15	24
455	48	63
454	2	6
152	1Ω	1/
285	18	16
284	87	67




#### **THREE Important Characteristics of Data**

- Central Tendency
- Variability or Dispersion
- Shape of Frequency Distribution



**NOTE:** You do NOT have permission to share this file or any of its contents with anyone else, and/or upload it on internet or any of the platforms where it can be accessed by others.

## **Central Tendency (Mean)**





• Given a sample of n pieces of data  $(y_1, y_2, y_3, ..., y_n)$  taken from a given population of size N, the arithmetic mean of the sample, denoted by  $\overline{y}$  is

$$\bar{y} = \sum_{k=1}^{n} \frac{y_i}{n}$$

• Population mean  $\mu$  given as,

$$\mu = \sum_{k=1}^{N} \frac{y_i}{N}$$

• Note: True mean  $(\mu)$  of the population of size N could be different than sample mean  $\overline{y}$ .

DIY

Can you show that as n -> N, the sample mean -> population mean?

## **Dispersion/Variability**





Range (R): Difference between the largest value and the smallest value of the data

$$R = \max(y_i) - \min(y_i)$$

• Variance ( $s^2$ ): Sample variance is given by

$$s^{2} = \sum_{i=1}^{n} \frac{(y_{i} - \bar{y})^{2}}{n - 1}$$

DIY
Why do we use (n-1) in the denominator for sample variance?

• Note, that the true variance ( $\sigma^2$ ) of the population of size N could be different

$$\sigma^2 = \sum_{i=1}^N \frac{(y_i - \mu)^2}{N}$$


- Notice (n-1) in the denominator for sample variance, while N for true variance
- Standard Deviation = Square root of Variance

## **Dispersion/Variability**




#### Why is variance defined as follows is a GOOD way to assess variability?

• What if we just add the deviations from the mean and take the average?




What if we just take absolute values of deviations from the mean?



• But, when we square...


$$\sqrt{\left(\frac{4^2+4^2+(-4)^2+(-4)^2}{4}\right)} = \sqrt{\left(\frac{64}{4}\right)} = 4$$



$$\sqrt{\left(\frac{7^2+1^2+(-6)^2+(-2)^2}{4}\right)} = \sqrt{\left(\frac{90}{4}\right)} = 4.74...$$

Ref: https://www.mathsisfun.com/data/standard-deviation.html

### **Median and Mode**






- The median of a finite list of numbers is the "middle" number when those numbers are listed in order from smallest to greatest.
- Mode is the most frequent value in the data set

#### **Examples:**


- Test Scores out of 20: (0, 11, 15, 8, 18, 19, 7, 8, 9, 12)
- Test Scores out of 20: (0, 11, 15, 8, 18, 7, 8, 9, 12)
- Test Scores out of 20: (0, 11, 15, 8, 18, 19, 7, 8, 9, 15)

### **Data Characterization**





- It has been shown that the mean (average) can be used to describe the data characteristics.
- However, it is possible to find two sets of data to have equal averages, but different degrees of scatter
- It has been a common mistake in many cases of applications to put all emphasis on the average but overlook the scatter of the data
- Such a mistake usually leads to unnecessary erroneous conclusions which could have been easily avoided if the scatter of the data had been considered.





Group C Group D

Table 1. Modified Table in Lee and Kim's Research (Adapted from Korean J Anesthesiol 2017; 70: 39-45)

Variable	Group	Baseline	After drug	1 min	3 min	5 min
SBP	С	135.1 ± 13.4	139.2 ± 17.1	186.0 ± 26.6*	160.1 ± 23.2*	140.7 ± 18.3
	D	135.4 ± 23.8	131.9 ± 13.5	165.2 ± 16.2*.†	127.9 ± 17.5 <sup>†</sup>	$108.4 \pm 12.6^{+,+}$
DBP	C	$79.7 \pm 9.8$	$79.4 \pm 15.8$	$104.8 \pm 14.9^*$	87.9 ± 15.5*	$78.9 \pm 11.6$
	D	$76.7 \pm 8.3$	$78.4 \pm 6.3$	97.0 ± 14.5*	$74.1 \pm 8.3^{\dagger}$	$66.5 \pm 7.2^{+.+}$
MBP	C	$100.3 \pm 11.9$	$103.5 \pm 16.8$	137.2 ± 18.3*	116.9 ± 16.2*	$103.9 \pm 13.3$
	D	97.7 ± 14.9	$98.1 \pm 8.7$	123.4 ± 13.8*.†	$95.4 \pm 11.7^{\dagger}$	$83.4 \pm 8.4^{+,+}$

Values are expressed as mean ± SD. Group C: normal saline, Group D: dexmedetomidine. SBP: systolic blood pressure, DBP: diastolic blood pressure MBP: mean blood pressure. HR: heart rate.

Table 2. Difference between a Regular Table and a Heat Map

Exa	mple of a	regular t	able	Example of a heat map			
SBP	DBP	MBP	HR	SBP	DBP	MBP	HR
128	66	87	87	128	66	87	87
125	43	70	85	125	43	70	85
114	52	68	103	114	52	68	103
111	44	66	79	111	44	66	79
139	61	81	90	139	61	81	90
103	44	61	96	103	44	61	96
94	47	61	83	94	47	61	83

All numbers were created by the author. SBP: systolic blood pressure, DBP: diastolic blood pressure, MBP: mean blood pressure, HR: heart rate.





Fig. 1. Line graph with whiskers. Changes in systolic blood pressure (SBP) in the two groups. Group C: normal saline, Group D: dexmedetomidine.

**Reference:** In, Junyong, and Sangseok Lee. "Statistical data presentation." *Korean journal of anesthesiology* 70.3 (2017): 267.


250



#### **Scatter Plot**





Scatter plots are used to investigate association between two variables





### **Bar Graph**







Bar graphs are used to indicate and compare values in discrete category or groups

Reference: In, Junyong, and Sangseok Lee. "Statistical data presentation." Korean journal of anesthesiology 70.3 (2017): 267.




### **Box and Whisker Graph**






Reference: In, Junyong, and Sangseok Lee. "Statistical data presentation." Korean journal of anesthesiology 70.3 (2017): 267.







**Misleading Plot** 

Reference: In, Junyong, and Sangseok Lee. "Statistical data presentation." Korean journal of anesthesiology 70.3 (2017): 267.



Table 3. Types of Charts depending on the Method of Analysis of the Data

Analysis	Subgroup	Number of variables	Туре
Comparison	Among items	Two per items	Variable width column chart
_	-	One per item	Bar/column chart
	Over time	Many periods	Circular area/line chart
		Few periods	Column/line chart
Relationship		Two	Scatter chart
_		Three	Bubble chart
Distribution		Single	Column/line histogram
		Two	Scatter chart
		Three	Three-dimensional area chart
Comparison	Changing over time	Only relative differences matter	Stacked 100% column chart
-		Relative and absolute differences matter	Stacked column chart
	Static	Simple share of total	Pie chart
		Accumulation	Waterfall chart
		Components of components	Stacked 100% column chart with subcomponents

Reference: In, Junyong, and Sangseok Lee. "Statistical data presentation." Korean journal of anesthesiology 70.3 (2017): 267.