


WWU  
MÜNSTER


Institut für  
Landschaftsökologie



# Machine learning for earth observation: Mapping the „Area of Applicability“ of spatial prediction models

**Hanna Meyer**

Remote Sensing & Spatial Modelling,  
Institute of Landscape Ecology, WWU Münster



# Problem: Moving from field observations to maps of ecosystem variables



# Problem: Moving from field observations to maps of ecosystem variables




# Problem: Moving from field observations to maps of ecosystem variables




How do we fill the gaps between sampling locations?


# Remote Sensing of landscapes




# Remote Sensing of landscapes




# Remote Sensing of landscapes




# Remote Sensing of landscapes



# Remote Sensing of landscapes




# Remote Sensing of landscapes



# Predictive modelling of the environment


e.g. vegetation cover from satellite (VIS/NIR)



Remote sensing data

# Predictive modelling of the environment


e.g. vegetation cover from satellite (VIS/NIR)



# Predictive modelling of the environment


e.g. vegetation cover from satellite (VIS/NIR)

Field data




Remote sensing data

Typical ecological variables from satellite?




# Predictive modelling of the environment

e.g. vegetation cover from satellite (VIS/NIR)




Typical ecological variables from satellite?




Models that can deal with complex nonlinear relationships are required!


# Predictive modelling of the environment: The machine learning way




# Predictive modelling of the environment: The machine learning way




# Predictive modelling of the environment: The machine learning way



# Predictive modelling of the environment: The machine learning way




# Predictive modelling of the environment: The machine learning way




# Global maps of ecosystem variables based on machine learning

Proportion of publications that use machine learning in environmental remote sensing



# Global maps of ecosystem variables based on machine learning

Proportion of publications that use machine learning in environmental remote sensing




Including **global** datasets on

- soil properties
- abundances of microorganisms
- Biodiversity
- tree restoration potential
- ...and many more

# Global maps of ecosystem variables based on machine learning

Proportion of publications that use machine learning in environmental remote sensing



- Including **global** datasets on
- soil properties
  - abundances of microorganisms
  - Biodiversity
  - tree restoration potential
  - ...and many more

Machine learning as a „magic tool“ to map basically everything ?

# ...but there are increasingly doubts about the methods

Wissenschaft

## Wenn die KI daneben liegt

Welche Fehler drohen, wenn Forscher Wissenslücken per Computer schließen wollen, zeigen zwei aktuelle Klimastudien.

Von Tin Fischer

6. November 2019, 16:44 Uhr / Editiert am 9. November 2019, 17:42 Uhr / DIE ZEIT  
Nr. 46/2019, 7. November 2019 / [9 Kommentare](#)



# DEEP TROUBLE FOR DEEP LEARNING

BY DOUGLAS HEAVEN

Nature 574, 163-166 (2019)

The screenshot shows a news article from 'The Scientist' website. At the top left, there is a navigation bar with 'Home' and 'News & Opinion'. The main title of the article is 'Researchers Find Flaws in High-Profile Study on Trees and Climate'. The background of the article section is dark, showing a landscape image.

Four independent groups say the work overestimates the carbon-absorbing benefits of global forest restoration, but the authors insist their original estimates are accurate.

Oct 17, 2019  
KATARINA ZIMMER

[www.the-scientist.com](http://www.the-scientist.com)

# ...but there are increasingly doubts about the methods

Wissenschaft

## Wenn die KI daneben liegt

Welche Fehler drohen, wenn Forscher Wissenslücken per Computer schließen wollen, zeigen zwei aktuelle Klimastudien.

Von Tin Fischer

6. November 2019, 16:44 Uhr / Editiert am 9. November 2019, 17:42 Uhr / DIE ZEIT  
Nr. 46/2019, 7. November 2019 / [9 Kommentare](#)



# DEEP TROUBLE FOR DEEP LEARNING

BY DOUGLAS HEAVEN

Nature 574, 163-166 (2019)

The screenshot shows a news article from 'The Scientist' website. At the top left, there's a navigation bar with 'Home' and 'News & Opinion'. The main title of the article is 'Researchers Find Flaws in High-Profile Study on Trees and Climate'. The background of the article section is dark, showing a landscape image of trees.

Four independent groups say the work overestimates the carbon-absorbing benefits of global forest restoration, but the authors insist their original estimates are accurate.

Oct 17, 2019  
KATARINA ZIMMER

[www.the-scientist.com](http://www.the-scientist.com)

## Have we been too ambitious? Why might the models fail?

# What we have learned so far...

See e.g. workshop at previous OpenGeoHub Summer School

Spatial dependencies need to be taken into account for machine learning applications!

# What we have learned so far...

See e.g. workshop at previous OpenGeoHub Summer School

Spatial dependencies need to be taken into account for machine learning applications!

- Standard validation procedures lead to an overoptimistic view on prediction performance  
→ Spatial validation is essential

# What we have learned so far...

See e.g. workshop at previous OpenGeoHub Summer School

Spatial dependencies need to be taken into account for machine learning applications!

- Standard validation procedures lead to an overoptimistic view on prediction performance
  - Spatial validation is essential
- Spatial dependencies can lead to misinterpretation of predictor variables
  - Spatial variable selection is required

# What we have learned so far...



See e.g. workshop at previous OpenGeoHub Summer School

Spatial dependencies need to be taken into account for machine learning applications!


- Standard validation procedures lead to an overoptimistic view on prediction performance
  - Spatial validation is essential
- Spatial dependencies can lead to misinterpretation of predictor variables
  - Spatial variable selection is required

...but is this sufficient for reliable (global) mapping?


# Largest challenge: predictions far beyond training samples




# Largest challenge: predictions far beyond training samples




# Largest challenge: predictions far beyond training samples




- Transfer to new space required





# Largest challenge: predictions far beyond training samples



- Transfer to new space required
- New space might differ in environmental properties




# Largest challenge: predictions far beyond training samples




- Transfer to new space required
- New space might differ in environmental properties
- But what if the algorithm has never seen such properties?

# Machine learning models are weak in extrapolations




- Machine learning can fit very complex relationships.

# Machine learning models are weak in extrapolations




- Machine learning can fit very complex relationships.
- But gaps in predictor space are problematic (the model has no knowledge about these areas!)

# Machine learning models are weak in extrapolations



- Machine learning can fit very complex relationships.
- But gaps in predictor space are problematic (the model has no knowledge about these areas!)
- **A measure for “unknown space” is needed**

# Distances in feature space as a measure for “unknown space”




- Unknown space: Environmental conditions that are very different from the training locations

\*More details: <https://arxiv.org/abs/2005.07939>

# Distances in feature space as a measure for “unknown space”

Is this environment known to the model?




- Unknown space: Environmental conditions that are very different from the training locations

\*More details: <https://arxiv.org/abs/2005.07939>

# Distances in feature space as a measure for “unknown space”

Is this environment known to the model?




- Unknown space: Environmental conditions that are very different from the training locations
- Suggestion: Dissimilarity Index based on distances in the (weighted) predictor space\*

\*More details: <https://arxiv.org/abs/2005.07939>

# Distances in feature space as a measure for “unknown space”

Is this environment known to the model?




- Unknown space: Environmental conditions that are very different from the training locations
- Suggestion: Dissimilarity Index based on distances in the (weighted) predictor space\*

For each new location/pixel: how distant is it from what the algorithm has seen?

\*More details: <https://arxiv.org/abs/2005.07939>

# Mapping the area of applicability - Example



Predictors



# Mapping the area of applicability - Example


Virtual Response and simulated samples

Predictors




# Mapping the area of applicability - Example


Predictors



Virtual Response and simulated samples




Prediction




# Mapping the area of applicability - Example


Predictors



Virtual Response and simulated samples




Prediction




Where can we trust the predictions  
and where should we better not?






# Mapping the area of applicability - Example



# Mapping the area of applicability - Example



# Mapping the area of applicability - Example



Threshold = DI of cross-validated training data

DI < threshold = inside AOA


DI > threshold = outside AOA

Outside AOA

# Why is it relevant to map the area of applicability?

Results are not just nice maps but used for...


- subsequent modeling
- nature conservation
- risk assessment
- ...



# Why is it relevant to map the area of applicability?


Results are not just nice maps but used for...

- subsequent modeling
- nature conservation
- risk assessment
- ...




**Predictions should only be presented for the area of applicability to avoid error propagation or misplanning (and to keep trust in the methods)!**

# Today's hands-on exercise




- Machine learning for remote sensing based land cover classification

# Today's hands-on exercise



- Machine learning for remote sensing based land cover classification
- Analysing the area of applicability

# Today's hands-on exercise



- Machine learning for remote sensing based land cover classification
- Analysing the area of applicability
- Assessing the transferability of a model

