

Disk Granulometries on the Digital Grid: Why a Single Structuring Element Is Not Enough

Peter Zsoldos

Tampere University, Tampere, Finland
TODO@tuni.fi

Abstract. The usual way to measure size distributions is to open an image by the digital disks of increasing radius that common libraries provide. We show that this is not a granulometry: these disks violate Matheron’s absorption axiom for most pairs of radii, and the resulting size distributions can increase with the radius. Two remedies exist. The first is classical: replacing each opening by the union of openings by all larger disks yields a correct granulometry that is as round as the disks themselves and costs the same. We recommend it in place of single openings. The second keeps a *single* structuring element per size, as in octagons, periodic lines or Bresenham decompositions, and it forces polygonal shapes. Our main result proves that this is unavoidable. If the mean radius of a Minkowski chain grows by at most δ per element, every element of radius ρ deviates from every disk by at least about $\rho/(16\pi^2\delta^2)$, in any dimension. An explicit chain of periodic lines attains this rate within a factor of 4, and for a fixed library the exact slope is attained by octagons. Experiments on synthetic and real images confirm the diagnosis, the remedy, and when the anisotropy of single-element families matters.

Keywords: Granulometry · Digital disk · Minkowski sum · Lattice polygon · Width function · Pattern spectrum.

1 Introduction

Granulometries, introduced by Matheron [8], formalise sieving: a family of openings $(\gamma_\lambda)_{\lambda \geq 0}$ that are decreasing in λ and satisfy the absorption law $\gamma_\lambda \gamma_\mu = \gamma_\mu \gamma_\lambda = \gamma_{\max(\lambda, \mu)}$. The size distribution and pattern spectrum derived from them [7,12,13] are among the most widely used morphological descriptors. The textbook construction uses a *single structuring element per size*: $\gamma_\lambda(X) = X \circ B_\lambda$. Absorption then holds if and only if B_μ is B_λ -open, i.e. a union of translates of B_λ , for all $\lambda \leq \mu$. Euclidean disks satisfy this condition.

On the grid $\mathbb{Z}^2$ they do not. The digital disks $\{p \in \mathbb{Z}^2 : \|p\| \leq r\}$ of common software are not open with respect to each other (Sect. 3): with `scikit-image` [17], 73% of the pairs of radii up to 40 violate the condition. Size distributions computed as $r \mapsto |X \circ D_r|$, the most direct use of these libraries, then increase with the radius on most random test images. There are two ways out.

- **Several structuring elements per size.** The sup-closure $\gamma'_r = \bigvee_{s \geq r} \gamma_{D_s}$, i.e. the union of the openings by all disks of radius at least r , is a granulometry for any

family (D_s) (Proposition 1). With digital disks it is correct, as round as the disks, and computable with one opening per radius. The construction is a classical consequence of Matheron’s theory [8,12], and opening transforms [16] and local thickness [5] implicitly rely on it. We do not claim it as new. Our point is that it, rather than single disk openings, should be used to compute size distributions with disks.

- **One structuring element per size.** The classical alternative builds the elements as *Minkowski chains* $D_k = D_{k-1} \oplus C_k$, which are automatically granulometric [1,6,13,16]. Examples are octagons from neighbourhood sequences [10,4], periodic lines [6] and Bresenham segments [1]. These families are polygonal. We prove that this is unavoidable and determine the price.

Our contributions are:

- **Diagnosis** (Sect. 3). We quantify the violation of the axiom by standard digital disks and its effect on size distributions.
- **Lower bound for single-element chains** (Theorem 1, Corollaries 1–3). If the mean radius of a Minkowski chain grows by at most δ per element, then *every* element of radius ρ has error $\varepsilon \geq (1 - o(1)) \rho / (16\pi^2 \delta^2)$, where the error is the Hausdorff distance to the nearest Euclidean disk. If the fitted radius grows by bounded steps, the maximal error is at least $c \rho^{1/3}$. For a fixed library the slope $\kappa / (1 + \kappa)$ is given by a linear program, and octagons attain it.
- **Matching upper bound and higher dimensions** (Theorems 2 and 3). An explicit chain of periodic lines achieves $\Theta(\rho / \delta^2)$, within a factor of 4 asymptotically, and $\Theta(\sqrt{\rho})$ when $\delta \sim \rho^{1/4}$. In $\mathbb{Z}^n$, $\varepsilon \geq c_n \rho \delta^{-2n}$.
- **Evidence and recommendations** (Sect. 6). On synthetic and real images, we compare single disk openings, their sup-closure and single-element families, on whole size distributions and on per-object sieve sizes. The comparison confirms that the classical sup-closure should replace single openings. When a single element per size is required, our LP-weighted periodic-line chains halve the error of octagons; this is a niche, and they do not compete with the sup-closure.

In short, round and correct disk granulometries on $\mathbb{Z}^2$ exist, but they need several structuring elements per size. Our contribution is to show that the common practice fails, to explain why the single-element route cannot be repaired, and to state what to use instead; we do not propose a new granulometry. Related work: chamfer norms [2,15] approximate Euclidean balls by polygons with fixed edge directions, and Corollary 3 applies when their balls form Minkowski chains (e.g. 3×3 masks). Neighbourhood sequences [4] optimise the roundness of a fixed library. Soille and Talbot [14] noted that radial decompositions by Bresenham lines are not granulometric. Normand [9] builds convex structuring elements by unions of translates, which lies outside the scope of our bound. To our knowledge, no lower bound of this kind has been published.

2 Preliminaries

For a finite set $X \subset \mathbb{Z}^n$ and a unit vector u , let $h_X(u) = \max_{p \in X} \langle p, u \rangle$ be the support function and $w_X(u) = h_X(u) + h_X(-u)$ the *width function*, i.e. the width of $\text{conv } X$ in

direction u . Both are additive under Minkowski addition [11]:

$$h_{X \oplus Y} = h_X + h_Y, \quad w_{X \oplus Y} = w_X + w_Y. \quad (1)$$

The *anisotropy* of X is $A(X) = \max_u w_X - \min_u w_X$, and its *mean radius* is $\bar{r}(X) = \frac{1}{2} \int w_X d\sigma$, where σ is the uniform probability measure on the sphere; $\bar{r}$ is additive as well. In the plane, Cauchy's formula gives a concrete meaning: $\bar{r}(X)$ is the perimeter of $\text{conv } X$ divided by 2π , i.e. the radius of the circle with the same perimeter. For a segment of length L it equals L/π . We say that X is ε -round with radius ρ if $d_H(\text{conv } X, B(c, \rho)) \leq \varepsilon$ for some centre c . Then $|h_X(u) - \langle c, u \rangle - \rho| \leq \varepsilon$, and hence

$$2\rho - 2\varepsilon \leq w_X \leq 2\rho + 2\varepsilon, \quad A(X) \leq 4\varepsilon, \quad |\bar{r}(X) - \rho| \leq \varepsilon. \quad (2)$$

The opening of Y by a structuring element D is $Y \circ D = \bigcup \{D + t : D + t \subseteq Y\}$. A family (D_k) generates a *single-element granulometry* $\gamma_k(Y) = Y \circ D_k$ if and only if $D_l \circ D_k = D_l$ for all $k \leq l$ [8,12]; since the relation is transitive, consecutive elements suffice. A *Minkowski chain* is a family with $D_k = D_{k-1} \oplus C_k$, and every Minkowski chain generates a single-element granulometry. Its *increments* are the C_k , its *steps* are $\bar{r}(D_k) - \bar{r}(D_{k-1}) = \bar{r}(C_k)$, and, if all C_k belong to a finite set $\mathcal{L}$ (up to translation), $\mathcal{L}$ is its *library*. Without the single-element restriction, any family of structuring elements can be turned into a granulometry.

The following fact is classical [8,12]; we include a short proof for completeness.


Proposition 1 (sup-closure). *For any finite sets $D_1, \dots, D_R \subset \mathbb{Z}^n$, the operators $\gamma'_r = \bigvee_{s=r}^R \gamma_{D_s}$, $r = 1, \dots, R$, where $\gamma_D(Y) = Y \circ D$, form a granulometry.*

Proof. A supremum of openings is an opening [12], and γ'_r is decreasing in r . Let $r \leq s$. Since γ'_r and γ'_s are anti-extensive and increasing, $\gamma'_r \gamma'_s \leq \gamma'_s$ and $\gamma'_s \gamma'_r \leq \gamma'_s$. Conversely, for every $t \geq s$ we have $\gamma'_r \gamma'_s \geq \gamma_{D_t} \gamma_{D_t} = \gamma_{D_t}$ and $\gamma'_s \gamma'_r \geq \gamma_{D_t} \gamma_{D_t} = \gamma_{D_t}$, and taking the supremum over $t \geq s$ gives $\geq \gamma'_s$. $\square$

Computing γ'_r from $r = R$ down to 1 needs one opening per radius, the same cost as the single-element family γ_{D_r} . The structuring elements are unchanged, so with digital disks the sup-closure is exactly as round as the disks. The rest of the paper explains why the single-element route cannot achieve this.

3 Digital Disks Are Not Granulometric

Figure 1(a) shows the smallest instance of the problem. We tested the condition “ D_r is D_s -open” for all pairs $1 \leq s < r \leq 40$, for the digital disks of `scikit-image` [17] (`disk`, the Gauss digitisation $\{p : \|p\| \leq r\}$) and of `OpenCV` [3] (`MORPH_ELLIPSE`). As Table 1 shows, most pairs violate it, including most consecutive pairs. When D_r is not D_s -open, $X \circ D_r \not\subseteq X \circ D_s$ for some X , and the size distribution $r \mapsto |X \circ D_r|$ can *increase*, so the pattern spectrum takes negative values. On random 120×120 images made of 3 to 11 disks, rectangles or digital disks ($r \leq 15$), this happens for 63% (`scikit-image`) and 60% (`OpenCV`) of the images. For example, in one image the opened

(a) $\text{disk}(5) \circ \text{disk}(4) \neq \text{disk}(5)$ 

red: pixels not covered by
any translate of $\text{disk}(4)$ inside



(b) directions, $\|v\| \leq 3$ (c) gap $\Rightarrow$ not round

Fig. 1. (a) The digital disk of radius 5 is not open with respect to that of radius 4: the red pixels (marked with crosses) lie in no translate of the smaller disk that fits inside. (b) Idea of Theorem 1: lattice vectors of norm at most ℓ (here 3) leave gaps of $\arctan(1/\ell)$ next to the axes. (c) A polygon whose edges come from such vectors has no edge with a direction in the gap. Over the corresponding range of normals a single vertex carries the support and sticks out of the circle (schematic, gap exaggerated).

Table 1. Violations of the absorption condition by standard digital disks, $1 \leq s < r \leq 40$. Last row: the sup-closure (several elements per size).

family	violating pairs (s, r)	violating pairs $(r - 1, r)$	images with non- monotone size distr.
scikit-image disk	571/780 (73%)	30/39	253/400
OpenCV MORPH_ELLIPSE	628/780 (81%)	36/39	241/400
sup-closure of disk (Prop. 1)	not applicable		0/400

area grows from 2053 to 2089 pixels between $r = 8$ and $r = 9$, and `scipy.ndimage`, `scikit-image` and `OpenCV` agree up to border conventions.

A granulometric *subsequence* of digital disks $\{p : \|p\|^2 \leq n\}$ ($n \leq 900$) with radius gaps at most G reaches radius 2.0 for $G = 1$, 9.9 for $G = 2$ and 21.8 for $G = 3$, and reaches 30 only with $G \geq 5$: such granulometries are very coarse.

4 Lower Bounds for Single-Element Chains

Additivity (1) turns the roundness of two elements of a chain into a constraint on the increments between them. All edges of the increments are lattice vectors. If the increments are short, these vectors leave a gap of directions around each axis (Fig. 1b), and a polygon whose edges avoid an angular interval cannot be round (Fig. 1c). Adding more elements to the chain makes the polygon larger but does not fill the gap, so the deviation grows with the radius.

Lemma 1 (direction gap). *Let $\ell \geq 1$. No nonzero $(a, b) \in \mathbb{Z}^2$ with $a^2 + b^2 \leq \ell^2$ has a direction angle in $(0, \arctan(1/\ell))$.*

Proof. A direction in that interval requires $a > 0$ and $b \geq 1$, hence $b/a \geq 1/a \geq 1/\ell$. $\square$

Theorem 1. Let $(D_k)_{k=0}^K$ be a Minkowski chain in $\mathbb{Z}^2$ whose increments have diameter at most $\ell \geq 1$. If D_0 is ε_0 -round with radius ρ_0 and D_K is ε_K -round with radius ρ_K , then, with $c = 1 - \cos(\frac{1}{2} \arctan \frac{1}{\ell})$,

$$4(\varepsilon_0 + \varepsilon_K) \geq 2(\rho_K - \rho_0 - \varepsilon_0 - \varepsilon_K) c. \quad (3)$$

Proof. *Step 1.* Every edge of $\text{conv } C_k$ joins two lattice points, so it is a lattice vector of norm at most ℓ . *Step 2.* Let $P = \sum_k \text{conv } C_k$ and $Q = \frac{1}{2}(P - P)$. By (1), $W := w_{D_K} - w_{D_0} = w_P = 2h_Q$. The edges of the centrally symmetric polygon Q are parallel to edges of the $\text{conv } C_k$, so by Lemma 1 (applied to $\pm e$) none has a direction in $(0, \alpha)$, where $\alpha = \arctan(1/\ell)$. The normal angles in $(\frac{\pi}{2}, \frac{\pi}{2} + \alpha)$ therefore belong to the normal cone of a single vertex q of Q : $h_Q(\theta) = |q| \cos(\theta - \theta_q)$ on an interval I of length at least α . *Step 3.* We may assume $s := \rho_K - \rho_0 - \varepsilon_0 - \varepsilon_K > 0$, since otherwise (3) is trivial. By (2), $\min W \geq 2s$, so $h_Q > 0$ and $|\theta - \theta_q| < \frac{\pi}{2}$ on I . On an interval of length α within $(\theta_q - \frac{\pi}{2}, \theta_q + \frac{\pi}{2})$, the function $\cos(\theta - \theta_q)$ varies by at least $1 - \cos(\alpha/2)$: by at least that much if $\theta_q \in I$, and by at least $1 - \cos \alpha$ otherwise, by monotonicity. With $|q| \geq \min h_Q \geq s$ this gives $\max W - \min W \geq 2sc$. *Step 4.* Finally $\max W - \min W \leq A(D_K) + A(D_0) \leq 4\varepsilon_K + 4\varepsilon_0$ by (2). $\square$

Only the first and the last element enter the bound. For a chain starting from a single point ($\varepsilon_0 = \rho_0 = 0$), (3) gives $\varepsilon_K \geq \rho_K c / (2 + c)$, where $c = \frac{1}{8\ell^2}(1 + O(\ell^{-2}))$: every element is at least $(1 - o(1))\rho / (16\ell^2)$ away from a disk. Different constraints on the steps bound ℓ in different ways.

Corollary 1 (mean-radius steps). If a Minkowski chain starting from a point has mean-radius steps $\bar{r}(C_k) \leq \delta$, then $\ell \leq \pi\delta$, and every element of radius ρ has error $\varepsilon \geq \rho c / (2 + c) = (1 - o(1))\rho / (16\pi^2\delta^2)$, with $c = 1 - \cos(\frac{1}{2} \arctan \frac{1}{\pi\delta})$. In particular, granulometric chains with bounded steps have error growing linearly with the radius.

Proof. By Cauchy's formula the perimeter of a planar convex set is π times its mean width, and it is at least twice its diameter. Hence $\text{diam conv } C_k \leq \frac{\pi}{2} \cdot 2\bar{r}(C_k) \leq \pi\delta$. $\square$

Corollary 2 (fitted-radius steps). If all elements of a Minkowski chain are ε -round with radii satisfying $\rho_k - \rho_{k-1} \leq \Delta$, then $\varepsilon \geq (1 - o(1))((\rho_K - \rho_0)/512)^{1/3}$ as $\rho_K \rightarrow \infty$.

Proof. By (1) and (2), $w_{C_k} = w_{D_k} - w_{D_{k-1}} \leq 2\Delta + 4\varepsilon$, so $\ell \leq 2\Delta + 4\varepsilon$. If ε stayed bounded, then so would ℓ , and (3) would force $\varepsilon \rightarrow \infty$. Hence $\ell = 4\varepsilon(1 + o(1))$, and (3) with $\varepsilon_0 = \varepsilon_K = \varepsilon$ becomes $8\varepsilon \geq (\rho_K - \rho_0)/(64\varepsilon^2)(1 + o(1))$. $\square$

Corollary 3 (fixed library). Let $\mathcal{L}$ be a finite library containing a set with more than one point, and let

$$\kappa(\mathcal{L}) = \inf \left\{ \frac{1}{4} A(\sum_C \lambda_C w_C) : \lambda \geq 0, \min_u \sum_C \lambda_C w_C(u) = 2 \right\}.$$

Then $\kappa := \kappa(\mathcal{L}) > 0$, and the last element of every chain with library $\mathcal{L}$ satisfies $\varepsilon_K \geq (\kappa(\rho_K - \rho_0 - \varepsilon_0) - \frac{1}{4} A(D_0)) / (1 + \kappa)$, i.e. $\varepsilon \geq \frac{\kappa}{1+\kappa} \rho - O(1)$.

Proof. For λ in the simplex, $f(\lambda) = A(\sum \lambda_C w_C) / \min \sum \lambda_C w_C$ takes values in $(0, \infty]$: $\sum \lambda_C w_C$ is the width function of the polygon $\sum \lambda_C \text{conv } C$, which does not have constant width, because bodies of constant width are strictly convex [11]. Moreover f is continuous where finite and tends to ∞ where the minimum vanishes, so it attains a positive minimum, and $\kappa = \frac{1}{2} \min f > 0$. For the chain, write $W = w_{D_K} - w_{D_0} = \sum_C n_C w_C$ and $s = \frac{1}{2} \min W \geq \rho_K - \rho_0 - \varepsilon_0 - \varepsilon_K$. Then $A(W) \geq 4\kappa s$ and $A(W) \leq 4\varepsilon_K + A(D_0)$. $\square$

We compute κ by linear programming on a grid of 3600 directions, with all D_4 -images included. It is 0.2071 for $\{N_4\}$ or $\{N_8\}$ (squares) and 0.0412 for octagons. For libraries of the 8-connected digital straight segments (DSS) of all primitive v with $\|v\| \leq m$, it is 0.0176 for $m = \sqrt{5}$ and 0.0028 for $m = 10$. For the corresponding periodic lines $\{0, v\}$ it is 0.0137 and, for $m = 5$, 0.0038. Since the grid underestimates the anisotropy, the computed values are lower approximations, and the bounds remain valid. For example, $\kappa(\{N_4\}) = (\sqrt{2} - 1)/2$ is the anisotropy of the square. Corollary 3 is tight for octagons: the element of fitted radius 961.5 of the best octagon chain has $\varepsilon/\rho = 0.039571$, while $\kappa/(1 + \kappa) = 0.039570$.

5 Matching Upper Bound and Higher Dimensions

The lower bound is attained, up to constants, by a simple explicit chain. For $m \geq 1$, let V_m contain one vector from each pair $\pm v$ of integer points on the boundary of $[-m, m]^2$. These are $4m$ vectors of norm at most $\sqrt{2}m$, and consecutive directions differ by at most $\arctan(1/m) \leq 1/m$, the largest gap being between $(m, 0)$ and $(m, 1)$. Let Z be the polygon circumscribed to the unit disk whose edges have the directions of $\pm v_m$; it is centrally symmetric, hence $Z = \sum_{v \in V_m} [-\frac{\ell_v}{2} \hat{v}, \frac{\ell_v}{2} \hat{v}]$ with $\hat{v} = v/\|v\|$, and $1 \leq h_Z \leq \sec(\frac{1}{2m})$. For $t \geq 0$ set $n_v(t) = \lfloor t\ell_v/\|v\| \rfloor$ and

$$D(t) = \bigoplus_{v \in V_m} n_v(t) \{0, v\},$$

the Minkowski sum of $n_v(t)$ copies of the periodic line $\{0, v\}$ [6]. As t grows, the counts increase. When several counts increase at the same t , we add the corresponding lines one at a time, so the elements form a Minkowski chain of periodic lines starting from $\{0\}$.

Theorem 2. *This chain has mean-radius steps at most $\sqrt{2}m/\pi$, and every element is ε -round with some radius ρ , with $\varepsilon \leq \rho/(8m^2) + 2m^2$.*

Proof. A step adds one periodic line, whose mean radius is $\|v\|/\pi \leq \sqrt{2}m/\pi$. Every element, including those created when ties are broken, satisfies $n_v = t\ell_v/\|v\| - \varphi_v$ for some t and $\varphi_v \in [0, 1]$. The centred support function of $\text{conv } D = \sum_v n_v [0, v]$ is therefore $H = \sum_v n_v |\langle v, u \rangle|/2 = t h_Z - \sum_v \varphi_v |\langle v, u \rangle|/2$. Since $\sum_v \|v\| \leq 4\sqrt{2}m^2$ and $\sec y - 1 \leq y^2$ for $y \leq \frac{1}{2}$, we get $t - 2\sqrt{2}m^2 \leq H \leq t + t/(4m^2)$. The disk of radius $\rho = (\max H + \min H)/2 \geq t - 2\sqrt{2}m^2$ is thus within $\varepsilon \leq t/(8m^2) + \sqrt{2}m^2 \leq \rho/(8m^2) + \sqrt{2}m^2 + \frac{\sqrt{2}}{4} \leq \rho/(8m^2) + 2m^2$. $\square$




Fig. 2. (a) Error of practical families for $\rho \leq 48$; mean-radius steps are below 2.2. (b) Chains of Theorem 2 for $m = 2, 4, 8$, with the upper bound of Theorem 2 (dashed) and the lower bound of Corollary 1 (dotted), and the multiscale chain against $0.3\sqrt{\rho}$.

With $\delta = \sqrt{2}m/\pi$, Theorem 2 gives $\varepsilon \leq \rho/(4\pi^2\delta^2) + \pi^2\delta^2$, while Corollary 1 gives $\varepsilon \geq (1 - o(1))\rho/(16\pi^2\delta^2)$. For $\rho \geq 4\pi^4\delta^4$, where the constant term is dominated, the best achievable error is therefore $\Theta(\rho/\delta^2)$, with constants within a factor of 8, and of 4 asymptotically. We can also concatenate chains with $m_j = 2^j$, each run up to $t_j \propto 16^j$ on top of the frozen previous ones; anisotropies and radii add. The result is a single chain with steps $O(\rho^{1/4})$ and error $O(\sqrt{\rho})$, which is optimal for steps $O(\rho^{1/4})$ by Corollary 1. Numerically (Fig. 2b), the fixed- m chains have slope about $1/(18m^2)$, between the bounds $1/(32m^2)$ and $1/(8m^2)$, and the multiscale chain satisfies $\varepsilon \leq 0.28\sqrt{\rho}$ for $20 \leq \rho \leq 5 \cdot 10^4$.

Like all our bounds, Theorem 2 concerns convex hulls. Sums of periodic lines can leave isolated holes, as in [6]. Adding the 2×2 square reduces them to 0.2–2% of the hull's lattice points, and holes do not affect the granulometric property.

Higher dimensions. In $\mathbb{Z}^n$ the directions are not ordered, but a measure argument replaces Lemma 1. We state the result with unspecified constants; its exponent is not sharp, since for $n = 2$ it gives ℓ^{-4} instead of ℓ^{-2} .

Theorem 3. Let (D_k) be a Minkowski chain in $\mathbb{Z}^n$, $n \geq 2$, whose increments have diameter at most $\ell \geq 1$, and let D_0 and D_K be ε_0 - and ε_K -round (with respect to Euclidean balls). There are constants $b_n > 0$ such that (3) holds with c replaced by $1 - \cos \alpha_n$, where $\sin \alpha_n = 1/(4b_n(2\ell + 1)^n)$. Hence $\varepsilon_K \geq c_n(\rho_K - \rho_0 - \varepsilon_0)\ell^{-2n} - \varepsilon_0$. If the mean-radius steps are at most δ , then $\ell \leq 2\delta/\mu_n$ with $\mu_n = \Gamma(\frac{n}{2})/(\sqrt{\pi}\Gamma(\frac{n+1}{2}))$, and bounded steps again force linear growth.

Table 2. Families for $\rho \leq 48$: largest mean-radius step δ , maximum and mean error, and the largest radius reached with $\max \varepsilon \leq 1$. The sup-closure uses several elements per size.

family	gran.	max step δ	max ε	mean ε	ρ at $\varepsilon \leq 1$
Gauss disks (disk)	no	1.19	0.35	0.21	–
sup-closure of Gauss disks	yes	1.19	0.35	0.21	≥ 48
octagons (neighbourhood seq.)	yes	2.17	1.85	0.98	25
Bresenham/DSS, 8 directions	yes	1.48	3.27	1.75	12
periodic lines, 8 dir., LP weights	yes	1.42	1.09	0.60	41
periodic lines, 13 dir., LP weights	yes	1.46	1.04	0.48	46
DSS, $\ v\ \leq 6$, MILP	yes	2.09	1.00	0.57	44

Proof (sketch). The directions u split into regions in each of which a single vertex of $Q = \frac{1}{2}(P - P)$ attains $h_Q(u)$ (the normal fan of Q). The boundaries between these regions lie in the hyperplanes $e^\perp$ for the edges e of the conv C_k , i.e. lattice vectors with $\|e\| \leq \ell$. There are at most $(2\ell + 1)^n$ of them, and each band $\{u : |\langle u, e/\|e\| \rangle| < s\}$ of the sphere has measure at most $b_n s$. So some u_0 satisfies $|\langle u_0, e \rangle| \geq 2 \sin \alpha_n \|e\|$ for all e , and the geodesic cap of radius α_n around u_0 meets no boundary. On this cap $h_Q(u) = \langle q, u \rangle$ is linear. Restricting it to the great circle through u_0 and q gives a variation of at least $|q|(1 - \cos \alpha_n)$, and Steps 3 and 4 are as before. Finally, a segment of length L has mean width $\mu_n L$, and mean width is monotone under inclusion. $\square$

6 Experiments

Families. We compare Gauss disks (disk), their sup-closure (Proposition 1), and five single-element families with mean-radius steps between 1.2 and 2.2, i.e. about one element per integer radius. These are octagons; Bresenham/DSS segments as in [1]; periodic lines as in [6] in 8 and in 13 directions (the D_4 -images of $(1, 0)$, $(1, 1)$, $(2, 1)$ and, for 13, also $(3, 1)$); and a chain of individual DSS computed by mixed-integer programming (HiGHS solver), which minimises the maximal error for $r \leq 48$. Segment libraries use the optimal weights λ^* of Corollary 3. We move towards the multiplicities $\text{round}(t\lambda^*)$ one set at a time, greedily keeping the element roundest. These chains contain the 2×2 square, and all their elements are hole-free.

Table 2 and Fig. 2(a) summarise roundness. Among single-element families, the optimised chains halve the error of octagons and reach a third of that of Bresenham decompositions, but they remain three times less round than the disks used by the sup-closure. At these radii Corollary 1 only forces 0.06–0.14 for the steps in the table. The lower bound is far from binding for small disks; its force lies in the growth rate (Fig. 2b). As Theorem 1 predicts through ℓ , larger steps help: with radius steps up to 4.1, the 13-direction chain reaches 0.41. With real multiplicities, a linear program finds monotone DSS families with error at most 0.43 up to radius 2000 (numerically; not proven), which points to integrality, used in Step 1 of the proof, as the obstruction.

Size distributions under rotation. For each scene X , rotated by every multiple of 5° up to 45° , we compute $F_\theta(r) = |\gamma_r(X_\theta)|/|X_\theta|$, implementing openings by chain elements

Table 3. Top: rotation sensitivity S ($\times 10^3$) of size distributions; in parentheses, the number of increases when nonzero. Bottom: per-object sieve size on coins (21 objects, inscribed radius 18–30 pixels; one merged object larger than the tested radii is excluded), error and rotation standard deviation in % of the inscribed radius.

	Gauss	sup-cl.	octag.	DSS 8	per. 8	per. 13	MILP
squares (synthetic)	4.0 (17)	4.0	30.2	13.3	11.8	9.5	17.6
nuclei (microscopy)	1.7	1.7	1.9	1.9	2.0	1.6	1.4
coins	5.0	5.0	5.0	6.4	4.2	5.0	6.2
gravel	1.4	1.4	1.3	1.6	1.1	1.6	1.2
coins: size error (%)	2.8	2.8	4.8	5.7	4.6	4.1	3.8
coins: rotation std (%)	0.65	0.65	1.13	1.20	1.06	0.54	1.00




Fig. 3. Size distributions of the rotated synthetic scene, one curve per angle. Single openings by Gauss disks are nearly isotropic but non-monotone. Their sup-closure removes the increases and keeps the isotropy. Octagons are monotone but orientation dependent, and the 13-direction periodic chain reduces this dependence by a factor of three.

as cascades of erosions and dilations by their elementary sets. We report the rotation sensitivity $S = \text{mean}_r \text{std}_\theta F_\theta(r)$ and the number of increases $F_\theta(r+1) > F_\theta(r)$, which are impossible for a granulometry. The scenes are nine squares of sides 17 to 81 pixels, rasterised exactly after rotation ($r \leq 40$), and three real images from `scikit-image`: fluorescence microscopy of nuclei (`human_mitosis`, $r \leq 15$), coins ($r \leq 30$), and a material texture (`gravel`, $r \leq 20$). The real images are smoothed, rotated with bilinear interpolation, and thresholded at the Otsu level of the unrotated image.

Per-object sieve size. Sieving assigns to each object O the size $s(O) = \max\{r : \gamma_r(X) \cap O \neq \emptyset\}$ of the largest element that fits inside it, which is the same for Gauss disks and their sup-closure. We compare $s(O)$ with the Euclidean inscribed radius, the maximum of the distance transform in O . We report the mean relative error and the mean standard deviation of $s(O)$ under rotation, relative to the inscribed radius, on objects matched by centroid.

Results. Table 3 and Fig. 3 confirm the picture from the theory.

- *The classical sup-closure is the best choice, as expected.* On every scene it has exactly the rotation sensitivity of Gauss disks, it never increases, and it removes the 17 increases that single Gauss-disk openings produce on the synthetic scene; on the random images of Sect. 3, it removes all increases (0 of the 400 images, against 253 of 400 for single openings).
- *Single-element families pay for their polygonal shape.* On the synthetic scene, octagons are 7.5 times more orientation dependent than the disks, and the 13-direction periodic chain reduces this to 2.4 times.
- *The cost shows on large objects.* Per object, on coins (inscribed radii 18–30 pixels), octagons measure sizes with 4.8% error and 1.1% rotation scatter, against 2.8% and 0.65% for the disks. The 13-direction chain has the smallest scatter (0.54%) and a 4.1% error.
- *It does not show on small objects or on whole images.* On nuclei and gravel, with inscribed radii of 3–10 pixels, discretisation dominates: every family errs by 13–24% and none is consistently best. On whole size distributions of the real images, all families agree within the effect of interpolation and thresholding.

The anisotropy of single-element families matters most for large or polygonal objects. The failure of single disk openings is a correctness issue on any image.

Recommendations.

- To compute size distributions or pattern spectra with disks, do not open by each library disk separately. Use the sup-closure of Proposition 1: open by every radius from the largest down and accumulate the unions. This costs nothing extra and yields a correct and round granulometry.
- Only if a single structuring element per size is required, e.g. to implement each opening as a fixed cascade of two-point erosions and dilations whose cost grows linearly rather than quadratically with the radius, use a Minkowski chain with LP weights. It improves on octagons and Bresenham decompositions, but not on the sup-closure: up to radius 50, the 13-direction periodic chain has error at most 1.04 pixels, and larger radii need larger libraries (Fig. 2b).

Our code and the computed chains are available¹.

7 Conclusion

Opening by each digital disk of a library is not a granulometry, and its size distributions can increase. The classical sup-closure fixes this at no cost and should be used instead. Single-element families cannot do as well: with mean-radius steps δ , their error is $\Theta(\rho/\delta^2)$, and bounded steps force linear growth in every dimension. Open questions: the optimal rate when only the fitted radius has bounded steps, between $\rho^{1/3}$ and $\rho^{1/2}$; whether single-element granulometries that are not Minkowski chains, built from unions of translates as in [9], can escape the bound; and matching upper bounds in $\mathbb{Z}^3$.

¹ Repository link withheld for review.

Acknowledgments. This work was supported by an Erasmus Mundus Joint Master (EMJM) scholarship in Imaging, funded by the European Union.

Use of AI Tools. Claude (Anthropic), a large language model, was used to assist with the literature search, the implementation and execution of the experiments, the derivation and checking of the proofs, and the drafting and editing of the manuscript. The author reviewed all content and takes full responsibility for it.

Disclosure of Interests. The author has no competing interests to declare that are relevant to the content of this article.

References

1. Adams, R.: Radial decomposition of disks and spheres. *CVGIP: Graphical Models and Image Processing* **55**(5), 325–332 (1993). <https://doi.org/10.1006/cgip.1993.1024>
2. Borgefors, G.: Distance transformations in digital images. *Computer Vision, Graphics, and Image Processing* **34**(3), 344–371 (1986). [https://doi.org/10.1016/S0734-189X\(86\)80047-0](https://doi.org/10.1016/S0734-189X(86)80047-0)
3. Bradski, G.: The OpenCV library. *Dr. Dobb's Journal of Software Tools* (2000)
4. Farkas, J., Baják, S., Nagy, B.: Approximating the Euclidean circle in the square grid using neighbourhood sequences. *Pure Mathematics and Applications* **17**, 309–322 (2006)
5. Hildebrand, T., Rügsegger, P.: A new method for the model-independent assessment of thickness in three-dimensional images. *Journal of Microscopy* **185**(1), 67–75 (1997). <https://doi.org/10.1046/j.1365-2818.1997.1340694.x>
6. Jones, R., Soille, P.: Periodic lines: Definition, cascades, and application to granulometries. *Pattern Recognition Letters* **17**(10), 1057–1063 (1996). [https://doi.org/10.1016/0167-8655\(96\)00066-9](https://doi.org/10.1016/0167-8655(96)00066-9)
7. Maragos, P.: Pattern spectrum and multiscale shape representation. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **11**(7), 701–716 (1989). <https://doi.org/10.1109/34.192465>
8. Matheron, G.: *Random Sets and Integral Geometry*. Wiley, New York (1975)
9. Normand, N.: Convex structuring element decomposition for single scan binary mathematical morphology. In: *Discrete Geometry for Computer Imagery (DGCI 2003)*. LNCS, vol. 2886, pp. 154–163. Springer (2003). https://doi.org/10.1007/978-3-540-39966-7_14
10. Rosenfeld, A., Pfaltz, J.L.: Distance functions on digital pictures. *Pattern Recognition* **1**(1), 33–61 (1968). [https://doi.org/10.1016/0031-3203\(68\)90013-7](https://doi.org/10.1016/0031-3203(68)90013-7)
11. Schneider, R.: *Convex Bodies: The Brunn–Minkowski Theory*. Cambridge University Press, 2nd edn. (2014). <https://doi.org/10.1017/CBO9781139003858>
12. Serra, J.: *Image Analysis and Mathematical Morphology*. Academic Press, London (1982)
13. Soille, P.: *Morphological Image Analysis: Principles and Applications*. Springer, Berlin, 2nd edn. (2003). <https://doi.org/10.1007/978-3-662-05088-0>
14. Soille, P., Talbot, H.: Directional morphological filtering. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **23**(11), 1313–1329 (2001). <https://doi.org/10.1109/34.969120>
15. Thiel, É.: *Géométrie des distances de chanfrein*. Habilitation à diriger des recherches, Université de la Méditerranée, Aix-Marseille II (2001)
16. Vincent, L.: Fast grayscale granulometry algorithms. In: *Mathematical Morphology and Its Applications to Image Processing (ISMM 1994)*. pp. 265–272. Kluwer (1994). https://doi.org/10.1007/978-94-011-1040-2_34
17. van der Walt, S., Schönberger, J.L., Nunez-Iglesias, J., Boulogne, F., Warner, J.D., Yager, N., Gouillart, E., Yu, T.: scikit-image: Image processing in Python. *PeerJ* **2**, e453 (2014). <https://doi.org/10.7717/peerj.453>