

Correct or Round? Granulometries by Digital Disks

Anonymous Author(s)

Anonymous Institution

Abstract. Size distributions by disk openings are a standard tool, but on $\mathbb{Z}^2$ disk granulometries face a dilemma. The digital disks of common libraries are round, but they violate Matheron's absorption axiom for most pairs of radii, so the size distributions they produce can increase with the radius. The classical correct alternatives, Minkowski chains of small structuring elements such as octagons, periodic lines or Bresenham decompositions, are polygonal. We prove that this dilemma is unavoidable and determine its price. If the mean radius of a Minkowski chain grows by at most δ per element, every element of radius ρ deviates from every disk by at least about $\rho/(16\pi^2\delta^2)$, so fine granulometric chains are linearly anisotropic, in any dimension. An explicit chain of periodic lines attains this rate within a factor of 4, and for a fixed library the exact slope, computed by linear programming, is attained by octagons. Experiments on synthetic and real images illustrate the violation of the axiom and the anisotropy of correct chains, and lead to practical recommendations.

Keywords: Granulometry · Digital disk · Minkowski sum · Lattice polygon · Width function · Pattern spectrum

1 Introduction

Granulometries, introduced by Matheron [9], formalise sieving: a family of openings $(\gamma_\lambda)_{\lambda \geq 0}$ that are decreasing in λ and satisfy the absorption law $\gamma_\lambda \gamma_\mu = \gamma_\mu \gamma_\lambda = \gamma_{\max(\lambda, \mu)}$. The size distribution and pattern spectrum derived from them [8, 13, 14] are among the most widely used morphological descriptors. For openings by structuring elements B_λ , absorption holds if and only if B_μ is B_λ -open, i.e. a union of translates of B_λ , for all $\lambda \leq \mu$. Euclidean disks satisfy this condition, and they form the canonical isotropic granulometry.

On the grid $\mathbb{Z}^2$, practitioners face a choice that is rarely made explicit. The digital disks $\{p \in \mathbb{Z}^2 : \|p\| \leq r\}$ of common software are round, but they are *not* open with respect to each other (Sect. 3). With `scikit-image` [18], 73% of the pairs of radii up to 40 violate the condition, and size distributions computed with these disks increase with the radius on most random test images. The classical remedy [1, 7, 14, 17] builds the structuring elements as *Minkowski chains* $D_k = D_{k-1} \oplus C_k$, which are automatically granulometric. Their elementary sets

C_k come from small libraries: the 4- and 8-neighbourhoods give octagons (neighbourhood sequences [11,4]), and there are also periodic lines [7] and Bresenham segments [1]. These families are correct but polygonal.

We show that this trade-off is a theorem, and we determine its rate.

- **Lower bound** (Theorem 1, Corollaries 1–3). If the mean radius of a Minkowski chain grows by at most δ per element, then *every* element of radius ρ has error $\varepsilon \geq (1 - o(1)) \rho / (16\pi^2 \delta^2)$, where the error is the Hausdorff distance to the nearest Euclidean disk. Every fine chain is therefore *linearly* anisotropic. If instead the fitted radius grows by bounded steps, then the maximal error is at least $c\rho^{1/3}$. For a fixed finite library the slope $\kappa/(1 + \kappa)$ is given by a linear program, and octagons attain it.
- **Matching upper bound** (Theorem 2). An explicit chain of periodic lines along the boundary of the square $[-m, m]^2$ has steps $\delta \leq \sqrt{2}m/\pi$ and error $\varepsilon \leq \rho/(8m^2) + 2m^2$. The optimal error is thus $\Theta(\rho/\delta^2)$ for $\rho \gtrsim \delta^4$, with constants within a factor of 4 asymptotically, and $\Theta(\sqrt{\rho})$ when $\delta \sim \rho^{1/4}$.
- **Higher dimensions** (Theorem 3). In $\mathbb{Z}^n$, $\varepsilon \geq c_n \rho \delta^{-2n}$: bounded steps force linear anisotropy in every dimension.
- **Evidence**. We quantify the violation of the axiom by standard digital disks (Table 1), compare practical families (Table 3), and test size distributions on rotated synthetic and real images (Table 4).

The message for practice is that one can be correct (granulometric) or round, but not both at all scales, and our bounds quantify the price. Several lines of related work bear on this problem.

- Chamfer norms [2,16] approximate Euclidean balls by polygons with a fixed set of edge directions. When their balls form Minkowski chains, as for 3×3 masks, Corollary 3 applies to them.
- Neighbourhood sequences [4] and zonohedra [6] optimise the roundness of a fixed library.
- Soille and Talbot [15] noted that radial decompositions by Bresenham lines do not generate granulometries.
- Normand [10] builds convex structuring elements by unions of translates instead of Minkowski sums, a construction outside the scope of our bound.

To our knowledge, no lower bound of this kind has been published.

2 Preliminaries

For a finite set $X \subset \mathbb{Z}^n$ and a unit vector u , let $h_X(u) = \max_{p \in X} \langle p, u \rangle$ be the support function and $w_X(u) = h_X(u) + h_X(-u)$ the *width function*, i.e. the width of $\text{conv } X$ in direction u . Both are additive under Minkowski addition [12]:

$$h_{X \oplus Y} = h_X + h_Y, \quad w_{X \oplus Y} = w_X + w_Y. \quad (1)$$

The *anisotropy* of X is $A(X) = \max_u w_X - \min_u w_X$, and its *mean radius* is $\bar{r}(X) = \frac{1}{2} \int w_X d\sigma$, where σ is the uniform probability measure on the sphere;




Fig. 1. (a) The digital disk of radius 5 is not open with respect to that of radius 4: the red pixels lie in no translate of the smaller disk that fits inside. (b) Idea of Theorem 1: lattice vectors of norm at most ℓ (here 3) leave gaps of $\arctan(1/\ell)$ next to the axes. (c) A polygon whose edges come from such vectors has no edge with a direction in the gap. Over the corresponding range of normals a single vertex carries the support and sticks out of the circle (schematic, gap exaggerated).

$\bar{r}$ is additive as well. In the plane, Cauchy’s formula gives a concrete meaning: $\bar{r}(X)$ is the perimeter of $\text{conv } X$ divided by 2π , i.e. the radius of the circle with the same perimeter. For a segment of length L it equals L/π . We say that X is ε -round with radius ρ if $d_H(\text{conv } X, B(c, \rho)) \leq \varepsilon$ for some centre c . Then $|h_X(u) - \langle c, u \rangle - \rho| \leq \varepsilon$, and hence

$$2\rho - 2\varepsilon \leq w_X \leq 2\rho + 2\varepsilon, \quad A(X) \leq 4\varepsilon, \quad |\bar{r}(X) - \rho| \leq \varepsilon. \quad (2)$$

The opening of Y by a structuring element D is $Y \circ D = \bigcup \{D + t : D + t \subseteq Y\}$. A family (D_k) generates a granulometry if and only if $D_l \circ D_k = D_l$ for all $k \leq l$ [9,13]; since the relation is transitive, consecutive elements suffice. A *Minkowski chain* is a family with $D_k = D_{k-1} \oplus C_k$, and every Minkowski chain is granulometric. Its *increments* are the C_k , its *steps* are $\bar{r}(D_k) - \bar{r}(D_{k-1}) = \bar{r}(C_k)$, and, if all C_k belong to a finite set $\mathcal{L}$ (up to translation), $\mathcal{L}$ is its *library*.

3 Digital Disks Are Not Granulometric

Figure 1(a) shows the smallest instance of the problem. We tested the condition “ D_r is D_s -open” for all pairs $1 \leq s < r \leq 40$, for the digital disks of `scikit-image` [18] (`disk`, the Gauss digitisation $\{p : \|p\| \leq r\}$) and of `OpenCV` [3] (`MORPH_ELLIPSE`). As Table 1 shows, most pairs violate it, including most consecutive pairs. When D_r is not D_s -open, $X \circ D_r \not\subseteq X \circ D_s$ for some X , and the size distribution $r \mapsto |X \circ D_r|$ can *increase*, so the pattern spectrum takes negative values. On random 120×120 images made of 3 to 11 disks, rectangles or digital disks ($r \leq 15$), this happens for 63% (`scikit-image`) and 60% (`OpenCV`) of the images. For example, in one image the opened area

Table 1. Violations of the absorption condition by standard digital disks, $1 \leq s < r \leq 40$.

family	violating pairs (s, r)	violating pairs $(r - 1, r)$	images with non- monotone size distr.
<code>scikit-image disk</code>	571/780 (73%)	30/39	253/400
<code>OpenCV MORPH_ELLIPSE</code>	628/780 (81%)	36/39	241/400

grows from 2053 to 2089 pixels between $r = 8$ and $r = 9$, and `scipy.ndimage`, `scikit-image` and `OpenCV` agree up to border conventions.

One could try to extract a granulometric *subsequence* of digital disks. Over all origin-centred disks $\{p : \|p\|^2 \leq n\}$ with $n \leq 900$, the largest radius reachable by a chain of consecutively open disks with radius gaps at most G is 2.0 for $G = 1$, 9.9 for $G = 2$ and 21.8 for $G = 3$, and radius 30 is reached only with $G \geq 5$. Digital disks thus give at best very coarse granulometries.

4 Lower Bounds

Additivity (1) turns the roundness of two elements of a chain into a constraint on the increments between them. All edges of the increments are lattice vectors. If the increments are short, these vectors leave a gap of directions around each axis (Fig. 1b), and a polygon whose edges avoid an angular interval cannot be round (Fig. 1c). Adding more elements to the chain makes the polygon larger but does not fill the gap, so the deviation grows with the radius.

Lemma 1 (direction gap). *Let $\ell \geq 1$. No nonzero $(a, b) \in \mathbb{Z}^2$ with $a^2 + b^2 \leq \ell^2$ has a direction angle in $(0, \arctan(1/\ell))$.*

Proof. A direction in that interval requires $a > 0$ and $b \geq 1$, hence $b/a \geq 1/a \geq 1/\ell$. $\square$

Theorem 1. *Let $(D_k)_{k=0}^K$ be a Minkowski chain in $\mathbb{Z}^2$ whose increments have diameter at most $\ell \geq 1$. If D_0 is ε_0 -round with radius ρ_0 and D_K is ε_K -round with radius ρ_K , then, with $c = 1 - \cos(\frac{1}{2} \arctan \frac{1}{\ell})$,*

$$4(\varepsilon_0 + \varepsilon_K) \geq 2(\rho_K - \rho_0 - \varepsilon_0 - \varepsilon_K)c. \quad (3)$$

Proof. Step 1. Every edge of $\text{conv } C_k$ joins two lattice points, so it is a lattice vector of norm at most ℓ . *Step 2.* Let $P = \sum_k \text{conv } C_k$ and $Q = \frac{1}{2}(P - P)$. By (1), $W := w_{D_K} - w_{D_0} = w_P = 2h_Q$. The edges of the centrally symmetric polygon Q are parallel to edges of the $\text{conv } C_k$, so by Lemma 1 (applied to $\pm e$) none has a direction in $(0, \alpha)$, where $\alpha = \arctan(1/\ell)$. The normal angles in $(\frac{\pi}{2}, \frac{\pi}{2} + \alpha)$ therefore belong to the normal cone of a single vertex q of Q : $h_Q(\theta) = |q| \cos(\theta - \theta_q)$ on an interval I of length at least α . *Step 3.* We may assume $s := \rho_K - \rho_0 - \varepsilon_0 - \varepsilon_K > 0$, since otherwise (3) is trivial. By (2), $\min W \geq 2s$, so $h_Q > 0$ and $|\theta - \theta_q| < \frac{\pi}{2}$ on I . On an interval of length α within

$(\theta_q - \frac{\pi}{2}, \theta_q + \frac{\pi}{2})$, the function $\cos(\theta - \theta_q)$ varies by at least $1 - \cos(\alpha/2)$: by at least that much if $\theta_q \in I$, and by at least $1 - \cos \alpha$ otherwise, by monotonicity. With $|q| \geq \min h_Q \geq s$ this gives $\max W - \min W \geq 2sc$. *Step 4.* Finally $\max W - \min W \leq A(D_K) + A(D_0) \leq 4\varepsilon_K + 4\varepsilon_0$ by (2). $\square$

Only the first and the last element enter the bound. For a chain starting from a single point ($\varepsilon_0 = \rho_0 = 0$), (3) gives $\varepsilon_K \geq \rho_K c/(2 + c)$, where $c = \frac{1}{8\ell^2}(1 + O(\ell^{-2}))$: every element is at least $(1 - o(1))\rho/(16\ell^2)$ away from a disk. Different constraints on the steps bound ℓ in different ways.

Corollary 1 (mean-radius steps). *If a Minkowski chain starting from a point has mean-radius steps $\bar{r}(C_k) \leq \delta$, then $\ell \leq \pi\delta$, and every element of radius ρ has error $\varepsilon \geq \rho c/(2 + c) = (1 - o(1))\rho/(16\pi^2\delta^2)$, with $c = 1 - \cos(\frac{1}{2} \arctan \frac{1}{\pi\delta})$. In particular, granulometric chains with bounded steps have error growing linearly with the radius.*

Proof. By Cauchy's formula the perimeter of a planar convex set is π times its mean width, and it is at least twice its diameter. Hence $\text{diam conv } C_k \leq \frac{\pi}{2} \cdot 2\bar{r}(C_k) \leq \pi\delta$. $\square$

Corollary 2 (fitted-radius steps). *If all elements of a Minkowski chain are ε -round with radii satisfying $\rho_k - \rho_{k-1} \leq \Delta$, then $\varepsilon \geq (1 - o(1))((\rho_K - \rho_0)/512)^{1/3}$ as $\rho_K \rightarrow \infty$.*

Proof. By (1) and (2), $w_{C_k} = w_{D_k} - w_{D_{k-1}} \leq 2\Delta + 4\varepsilon$, so $\ell \leq 2\Delta + 4\varepsilon$. If ε stayed bounded, then so would ℓ , and (3) would force $\varepsilon \rightarrow \infty$. Hence $\ell = 4\varepsilon(1 + o(1))$, and (3) with $\varepsilon_0 = \varepsilon_K = \varepsilon$ becomes $8\varepsilon \geq (\rho_K - \rho_0)/(64\varepsilon^2)(1 + o(1))$. $\square$

Corollary 3 (fixed library). *Let $\mathcal{L}$ be a finite library containing a set with more than one point, and let*

$$\kappa(\mathcal{L}) = \inf \left\{ \frac{1}{4} A(\sum_C \lambda_C w_C) : \lambda \geq 0, \min_u \sum_C \lambda_C w_C(u) = 2 \right\}.$$

Then $\kappa := \kappa(\mathcal{L}) > 0$, and the last element of every chain with library $\mathcal{L}$ satisfies $\varepsilon_K \geq (\kappa(\rho_K - \rho_0 - \varepsilon_0) - \frac{1}{4} A(D_0))/(1 + \kappa)$, i.e. $\varepsilon \geq \frac{\kappa}{1 + \kappa} \rho - O(1)$.

Proof. For λ in the simplex, $f(\lambda) = A(\sum \lambda_C w_C)/\min \sum \lambda_C w_C$ takes values in $(0, \infty]$: $\sum \lambda_C w_C$ is the width function of the polygon $\sum \lambda_C \text{conv } C$, which does not have constant width, because bodies of constant width are strictly convex [12]. Moreover f is continuous where finite and tends to ∞ where the minimum vanishes, so it attains a positive minimum, and $\kappa = \frac{1}{2} \min f > 0$. For the chain, write $W = w_{D_K} - w_{D_0} = \sum_C \lambda_C w_C$ and $s = \frac{1}{2} \min W \geq \rho_K - \rho_0 - \varepsilon_0 - \varepsilon_K$. Then $A(W) \geq 4\kappa s$ and $A(W) \leq 4\varepsilon_K + A(D_0)$. $\square$

We compute κ by linear programming on a grid of 3600 directions (Table 2). Since the grid underestimates the anisotropy, the computed values are lower approximations, and the bounds remain valid. For example, $\kappa(\{N_4\}) = (\sqrt{2} - 1)/2$ is the anisotropy of the square. Corollary 3 is tight for octagons: the element of fitted radius 961.5 of the best octagon chain has $\varepsilon/\rho = 0.039571$, while $\kappa/(1 + \kappa) = 0.039570$.

Table 2. Slope κ of Corollary 3 for classical libraries (D_4 -images included). Segment libraries contain the 8-connected digital straight segments (DSS) or the periodic lines $\{0, v\}$ of all primitive v with $\|v\| \leq m$.

library	κ	library	κ	library	κ
$\{N_4\}$ or $\{N_8\}$	0.2071	DSS, $m = \sqrt{5}$	0.0176	periodic, $m = \sqrt{5}$	0.0137
$\{N_4, N_8\}$ (octagons)	0.0412	DSS, $m = 10$	0.0028	periodic, $m = 5$	0.0038

5 Matching Upper Bound and Higher Dimensions

The lower bound is attained, up to constants, by a simple explicit chain. For $m \geq 1$, let V_m contain one vector from each pair $\pm v$ of integer points on the boundary of $[-m, m]^2$. These are $4m$ vectors of norm at most $\sqrt{2}m$, and consecutive directions differ by at most $\arctan(1/m) \leq 1/m$, the largest gap being between $(m, 0)$ and $(m, 1)$. Let Z be the polygon circumscribed to the unit disk whose edges have the directions of $\pm V_m$; it is centrally symmetric, hence $Z = \sum_{v \in V_m} [-\frac{\ell_v}{2}\hat{v}, \frac{\ell_v}{2}\hat{v}]$ with $\hat{v} = v/\|v\|$, and $1 \leq h_Z \leq \sec(\frac{1}{2m})$. For $t \geq 0$ set $n_v(t) = \lfloor t\ell_v/\|v\| \rfloor$ and

$$D(t) = \bigoplus_{v \in V_m} n_v(t) \{0, v\},$$

the Minkowski sum of $n_v(t)$ copies of the periodic line $\{0, v\}$ [7]. As t grows, the counts increase. When several counts increase at the same t , we add the corresponding lines one at a time, so the elements form a Minkowski chain of periodic lines starting from $\{0\}$.

Theorem 2. *This chain has mean-radius steps at most $\sqrt{2}m/\pi$, and every element is ε -round with some radius ρ , with $\varepsilon \leq \rho/(8m^2) + 2m^2$.*

Proof. A step adds one periodic line, whose mean radius is $\|v\|/\pi \leq \sqrt{2}m/\pi$. Every element, including those created when ties are broken, satisfies $n_v = t\ell_v/\|v\| - \varphi_v$ for some t and $\varphi_v \in [0, 1]$. The centred support function of $\text{conv } D = \sum_v n_v[0, v]$ is therefore $H = \sum_v n_v |\langle v, u \rangle|/2 = t h_Z - \sum_v \varphi_v |\langle v, u \rangle|/2$. Since $\sum_v \|v\| \leq 4\sqrt{2}m^2$ and $\sec y - 1 \leq y^2$ for $y \leq \frac{1}{2}$, we get $t - 2\sqrt{2}m^2 \leq H \leq t + t/(4m^2)$. The disk of radius $\rho = (\max H + \min H)/2 \geq t - 2\sqrt{2}m^2$ is thus within $\varepsilon \leq t/(8m^2) + \sqrt{2}m^2 \leq \rho/(8m^2) + \sqrt{2}m^2 + \frac{\sqrt{2}}{4} \leq \rho/(8m^2) + 2m^2$. $\square$

With $\delta = \sqrt{2}m/\pi$, Theorem 2 gives $\varepsilon \leq \rho/(4\pi^2\delta^2) + \pi^2\delta^2$, while Corollary 1 gives $\varepsilon \geq (1 - o(1))\rho/(16\pi^2\delta^2)$. For $\rho \geq 4\pi^4\delta^4$, where the constant term is dominated, the best achievable error is therefore $\Theta(\rho/\delta^2)$, with constants within a factor of 8, and of 4 asymptotically. We can also concatenate chains with $m_j = 2^j$, each run up to $t_j \propto 16^j$ on top of the frozen previous ones; anisotropies and radii add. The result is a single chain with steps $O(\rho^{1/4})$ and error $O(\sqrt{\rho})$, which is optimal for steps $O(\rho^{1/4})$ by Corollary 1. Numerically (Fig. 2b), the fixed- m chains have slope about $1/(18m^2)$, between the bounds $1/(32m^2)$ and $1/(8m^2)$, and the multiscale chain satisfies $\varepsilon \leq 0.28\sqrt{\rho}$ for $20 \leq \rho \leq 5 \cdot 10^4$.




Fig. 2. (a) Error of practical families for $\rho \leq 48$; mean-radius steps are below 2.2. (b) Chains of Theorem 2 for $m = 2, 4, 8$, with the upper bound of Theorem 2 (dashed) and the lower bound of Corollary 1 (dotted), and the multiscale chain against $0.3\sqrt{\rho}$.

Like all our bounds, Theorem 2 concerns convex hulls. Minkowski sums of periodic lines can leave isolated holes, as Jones and Soille’s decompositions also do [7]. Adding the 2×2 square to the chain reduces them to 0.2–2% of the hull’s lattice points in our tests, and holes do not affect the granulometric property.

Higher dimensions. In $\mathbb{Z}^n$ the directions are not ordered, but a measure argument replaces Lemma 1. We state the result with unspecified constants; its exponent is not sharp, since for $n = 2$ it gives ℓ^{-4} instead of ℓ^{-2} .

Theorem 3. *Let (D_k) be a Minkowski chain in $\mathbb{Z}^n$, $n \geq 2$, whose increments have diameter at most $\ell \geq 1$, and let D_0 and D_K be ε_0 - and ε_K -round (with respect to Euclidean balls). There are constants $b_n > 0$ such that (3) holds with c replaced by $1 - \cos \alpha_n$, where $\sin \alpha_n = 1/(4b_n(2\ell + 1)^n)$. Hence $\varepsilon_K \geq c_n(\rho_K - \rho_0 - \varepsilon_0)\ell^{-2n} - \varepsilon_0$. If the mean-radius steps are at most δ , then $\ell \leq 2\delta/\mu_n$ with $\mu_n = \Gamma(\frac{n}{2})/(\sqrt{\pi}\Gamma(\frac{n+1}{2}))$, and bounded steps again force linear growth.*

Proof (sketch). The directions u split into regions in each of which a single vertex of $Q = \frac{1}{2}(P - P)$ attains $h_Q(u)$ (the normal fan of Q). The boundaries between these regions lie in the hyperplanes $e^\perp$ for the edges e of the $\text{conv } C_k$, i.e. lattice vectors with $\|e\| \leq \ell$. There are at most $(2\ell + 1)^n$ of them, and each band $\{u : |\langle u, e/\|e\| \rangle| < s\}$ of the sphere has measure at most $b_n s$. So some u_0 satisfies $|\langle u_0, e \rangle| \geq 2\sin \alpha_n \|e\|$ for all e , and the geodesic cap of radius α_n around u_0 meets no boundary. On this cap $h_Q(u) = \langle q, u \rangle$ is linear. Restricting it to the great circle through u_0 and q gives a variation of at least $|q|(1 - \cos \alpha_n)$, and Steps 3 and 4 are as before. Finally, a segment of length L has mean width $\mu_n L$, and mean width is monotone under inclusion. $\square$

6 Practical Families and Experiments

Families. Table 3 compares families for $\rho \leq 48$ with mean-radius steps between 1.2 and 2.2, i.e. about one element per integer radius. For the fixed libraries

Table 3. Error ε (pixels) of families for $\rho \leq 48$: largest mean-radius step δ , maximum and mean error, and the largest radius reached with $\max \varepsilon \leq 1$.

family	gran.	max step δ	max ε	mean ε	ρ at $\varepsilon \leq 1$
Gauss disks (disk)	no	1.19	0.35	0.21	–
octagons (neighbourhood seq.)	yes	2.17	1.85	0.98	25
Bresenham/DSS, 8 directions	yes	1.48	3.27	1.75	12
periodic lines, 8 dir., LP weights	yes	1.42	1.09	0.60	41
periodic lines, 13 dir., LP weights	yes	1.46	1.04	0.48	46
DSS, $\ v\ \leq 6$, MILP	yes	2.09	1.00	0.57	44


(Bresenham/DSS segments as in [1], periodic lines as in [7]), we use the optimal weights λ^* of Corollary 3. We move towards the multiplicities $\text{round}(t\lambda^*)$ one elementary set at a time, choosing at each step the set that keeps the element roundest; the chains contain the 2×2 square. We also include a chain of individual DSS computed by mixed-integer programming (HiGHS [5]), which minimises the maximal error over $r \leq 48$. All elements of these granulometric families were verified to be hole-free. The optimised chains halve the error of octagons and reach about a third of that of Bresenham decompositions, but they remain three times less round than Gauss disks. At these radii Corollary 1 only forces 0.06–0.14 for the steps in the table. The lower bound is far from binding for small disks, and its force lies in the growth rate at large radii (Fig. 2b). Relaxing the step size helps, as Theorem 1 predicts through ℓ : with radius steps up to 4.1, the 13-direction periodic chain reaches 0.41. Conversely, if multiplicities could be real, a monotone family of DSS with $\|v\| \leq 45$ would have error at most 0.43 on a grid of radii up to 2000, as we found by solving a linear program; we do not claim that this persists for all radii. Integrality is what Step 1 of the proof exploits.

Size distributions. For each scene X and angle $\theta \in \{0^\circ, 5^\circ, \dots, 45^\circ\}$ we compute $F_\theta(r) = |X_\theta \circ D_r|/|X_\theta|$, implementing openings by chain elements as cascades of erosions and dilations by their elementary sets. We report the rotation sensitivity $S = \text{mean}_r \text{std}_\theta F_\theta(r)$ and the number of increases $F_\theta(r+1) > F_\theta(r)$, which are impossible for a granulometry. The scenes are nine squares of sides 17 to 81 pixels, rasterised exactly after rotation ($r \leq 40$), and three real images from `scikit-image`: fluorescence microscopy of nuclei (`human_mitosis`, $r \leq 15$), `coins` ($r \leq 30$) and a material texture (`gravel`, $r \leq 20$). The real images are smoothed, rotated with bilinear interpolation and thresholded at the Otsu level of the unrotated image.

Table 4 and Fig. 3 illustrate the dilemma of the title. On the synthetic scene, Gauss disks are the least orientation dependent but produce 17 increases, i.e. negative pattern-spectrum values. Octagons are correct but strongly orientation dependent, and the periodic-line chain divides their sensitivity by three. On the three real images, the families differ by at most $0.6 \cdot 10^{-3}$ (nuclei, gravel) and $2.2 \cdot 10^{-3}$ (coins), with no consistent ranking, i.e. within the effect of interpolation

Table 4. Rotation sensitivity $S (\times 10^3)$ of size distributions; in parentheses, the number of increases of F_θ when nonzero.

scene	Gauss	octagons	DSS 8 dir.	periodic 8	periodic 13	DSS MILP
squares (synthetic)	4.0 (17)	30.2	13.3	11.8	9.5	17.6
nuclei (microscopy)	1.7	1.9	1.9	2.0	1.6	1.4
coins	5.0	5.0	6.4	4.2	5.0	6.2
gravel	1.4	1.3	1.6	1.1	1.6	1.2

**Fig. 3.** Size distributions of the rotated synthetic scene, one curve per angle: Gauss disks are nearly isotropic but non-monotone; octagons are monotone but orientation dependent; the periodic-line chain is monotone and three times less orientation dependent.

and thresholding; no increases occur there. Rounded, randomly oriented objects average out the anisotropy of the structuring element. The choice of family therefore matters most for polygonal or aligned structures. The violation of the axiom by Gauss disks, in contrast, is a correctness issue that can surface on any image (Table 1).

Recommendations. If size distributions must be correct (monotone, with non-negative pattern spectra), use a Minkowski chain, not library disks. For radii up to about 50, the periodic-line chain with the 13 directions of $(1,0), (1,1), (2,1), (3,1)$ and their D_4 -images, together with the 2×2 square and LP weights, has error at most 1.04 pixels (mean 0.48). Openings by it are cascades of two-point erosions and dilations, whose cost grows linearly with the radius instead of quadratically. For larger radii, use larger libraries: by Corollary 3 a library with slope κ loses $\frac{\kappa}{1+\kappa}\rho + O(1)$, and the best choice grows with the range of radii (Fig. 2b). If only roundness matters and absorption is not needed, Gauss disks remain the roundest option. Our code and the computed chains are available¹.

¹ Repository link withheld for review.

7 Conclusion

Standard digital disks do not generate granulometries, and granulometric Minkowski chains cannot be uniformly round. With mean-radius steps δ their error is $\Theta(\rho/\delta^2)$, and $\Theta(\sqrt{\rho})$ when $\delta \sim \rho^{1/4}$; bounded steps force linear growth in every dimension. For practice, one must choose between correct but anisotropic chains, whose loss the constant κ quantifies, and round but incorrect digital disks. Our periodic-line chains halve the loss of octagons. Three questions remain open: the optimal rate when only the fitted radius has bounded steps, between $\rho^{1/3}$ and $\rho^{1/2}$; whether granulometries that are not Minkowski chains, where D_l is merely a union of translates of D_k , can escape the bound; and matching upper bounds in $\mathbb{Z}^3$.

References

1. Adams, R.: Radial decomposition of discs and spheres. *CVGIP: Graphical Models and Image Processing* **55**(5), 325–332 (1993)
2. Borgefors, G.: Distance transformations in digital images. *Computer Vision, Graphics, and Image Processing* **34**(3), 344–371 (1986)
3. Bradski, G.: The OpenCV library. *Dr. Dobb's Journal of Software Tools* (2000)
4. Farkas, J., Baják, S., Nagy, B.: Approximating the Euclidean circle in the square grid using neighbourhood sequences. *Pure Mathematics and Applications* **17**, 309–322 (2006)
5. Huangfu, Q., Hall, J.A.J.: Parallelizing the dual revised simplex method. *Mathematical Programming Computation* **10**(1), 119–142 (2018)
6. Jensen, P.M., Trinderup, C.H., Dahl, A.B., Dahl, V.A.: Zonohedral approximation of spherical structuring element for volumetric morphology. In: *Image Analysis (SCIA 2019)*. LNCS, vol. 11482, pp. 128–139. Springer (2019)
7. Jones, R., Soille, P.: Periodic lines: Definition, cascades, and application to granulometries. *Pattern Recognition Letters* **17**(10), 1057–1063 (1996)
8. Maragos, P.: Pattern spectrum and multiscale shape representation. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **11**(7), 701–716 (1989)
9. Matheron, G.: *Random Sets and Integral Geometry*. Wiley, New York (1975)
10. Normand, N.: Convex structuring element decomposition for single scan binary mathematical morphology. In: *Discrete Geometry for Computer Imagery (DGCI 2003)*. LNCS, vol. 2886, pp. 154–163. Springer (2003)
11. Rosenfeld, A., Pfaltz, J.L.: Distance functions on digital pictures. *Pattern Recognition* **1**(1), 33–61 (1968)
12. Schneider, R.: *Convex Bodies: The Brunn–Minkowski Theory*. Cambridge University Press, 2nd edn. (2014)
13. Serra, J.: *Image Analysis and Mathematical Morphology*. Academic Press, London (1982)
14. Soille, P.: *Morphological Image Analysis: Principles and Applications*. Springer, Berlin, 2nd edn. (2003)
15. Soille, P., Talbot, H.: Directional morphological filtering. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **23**(11), 1313–1329 (2001)
16. Thiel, É.: *Géométrie des distances de chanfrein*. Habilitation à diriger des recherches, Université de la Méditerranée, Aix-Marseille II (2001)

17. Vincent, L.: Fast grayscale granulometry algorithms. In: Mathematical Morphology and Its Applications to Image Processing (ISMM 1994). pp. 265–272. Kluwer (1994)
18. van der Walt, S., Schönberger, J.L., Nunez-Iglesias, J., Boulogne, F., Warner, J.D., Yager, N., Gouillart, E., Yu, T.: scikit-image: Image processing in Python. *PeerJ* **2**, e453 (2014)