

Math 4B: Differential Equations

Lecture 04: Separable ODEs

- Separable First-order ODEs,
- Those equations from last time,
- & More!

© 2021 Peter M. Garfield

Please do not distribute outside of this course.

Basic Example

Two of our basic examples of ODEs was

$$\frac{dy}{dt} = ky \quad \text{and} \quad \frac{dy}{dt} = p(t)y$$

where k was a constant and $p(t)$ was a function of t . We found the answers, but the question today is...

HOW?

The idea is today's topic: separation of variables.

Separation of Variables

The idea:

Move all the “ y ”s to one side and the “ t ”s to the other:

$$\frac{1}{y} \frac{dy}{dt} = k \quad \text{or} \quad \frac{1}{y} dy = k dt.$$

Which one is better?

We integrate and both are the same:

$$\int \frac{1}{y} \frac{dy}{dt} dt = \int k dt \quad \text{or} \quad \int \frac{1}{y} dy = \int k dt.$$

$$\text{So } \ln |y| = kt + C_1 \quad \implies \quad |y| = e^{C_1} e^{kt} = C_2 e^{kt} \quad \implies \quad y = C e^{kt}.$$

The General Idea

Last Time we dealt with solving the general first order linear ODE

$$y' + p(t)y = q(t),$$

where $p(t)$ and $q(t)$ are arbitrary functions of t .

Today we'll look at separable first order ODEs; that is, ones that can be written as

$$M(y)y' = N(t) \quad \text{or, equivalently} \quad M(y) dy = N(t) dt.$$

Idea: Just integrate (both are the same):

$$\int M(y) \frac{dy}{dt} dt = \int N(t) dt \quad \text{or} \quad \int M(y) dy = \int N(t) dt.$$

Example 1

1. Find the most general solution to the initial value problem (IVP)

$$\begin{cases} y \frac{dy}{dt} = 4t \\ y(0) = 3. \end{cases}$$

Solution: The equation arrives separated, so integrate:

$$\int y \frac{dy}{dt} dt = \int 4t dt \quad \text{or} \quad \int y dy = \int 4t dt$$


Then

$$\frac{1}{2}y^2 = 2t^2 + C \quad \implies \quad y = \pm\sqrt{4t^2 + 2C}.$$

When $t = 0$, $y = 3$ means $3 = +\sqrt{4(0)^2 + 2C}$, so $2C = 9$.

Thus $y = \sqrt{4t^2 + 9}$.

Here's a picture of the direction field and a few solutions:



Example 2

2. Solve the ODE

$$\frac{dy}{dt} = te^{t^2+y}.$$

That is, find the general solution to this ODE.

Solution: We separate to get

$$e^{-y} \frac{dy}{dt} = te^{t^2} \quad \text{or} \quad e^{-y} dy = te^{t^2} dt.$$

We integrate to get

$$\int e^{-y} \frac{dy}{dt} dt = \int te^{t^2} dt \quad \text{or} \quad \int e^{-y} dy = \int te^{t^2} dt.$$

Then $-e^{-y} = \frac{1}{2}e^{t^2} + C$, so

$$e^{-y} = K - \frac{1}{2}e^{t^2} \quad \implies \quad y = -\ln\left(K - e^{t^2}/2\right).$$

Example 3

3. Solve the ODE

$$\frac{dy}{dx} = \frac{x}{1+y^2}$$

That is, find the general solution to this ODE.

Solution: We separate to get

$$(1+y^2) \frac{dy}{dx} = x \quad \text{or} \quad (1+y^2) dy = x dx.$$


We integrate to get

$$\int (1+y^2) \frac{dy}{dx} dx = \int x dx \quad \text{or} \quad \int (1+y^2) dy = \int x dx.$$

Then

$$y + \frac{1}{3}y^3 = \frac{1}{2}x^2 + C \quad \implies \quad y = ??? .$$

Here's a picture of the direction field and some solutions:



$$y + \frac{1}{3}y^3 = \frac{1}{2}x^2 + C$$

Example 4

4. Find the most general solution to the ODE

$$\frac{dy}{dt} = \frac{1}{4}y(5 - y).$$

How does the behavior of the solution change based on the initial condition $y(0)$?

Solution: The equation is not hard to separate:

$$\frac{1}{y(5 - y)} \frac{dy}{dt} = \frac{1}{4} \quad \text{or} \quad \frac{1}{y(5 - y)} dy = \frac{1}{4} dt.$$

As usual, we integrate:

$$\int \frac{1}{y(5 - y)} \frac{dy}{dt} dt = \int \frac{1}{4} dt \quad \text{or} \quad \int \frac{1}{y(5 - y)} dy = \int \frac{1}{4} dt.$$

We use partial fractions to write $\frac{1}{y(5 - y)} = \frac{1}{5} \left(\frac{1}{y} + \frac{1}{5 - y} \right)$.

Example 4 (cont'd)

Integrating, we get

$$\frac{1}{5} (\ln |y| - \ln |5 - y|) = \frac{1}{4}t + C_1 \quad \implies \quad \frac{y}{5 - y} = Ke^{5t/4}.$$

Thus

$$y = \frac{5Ke^{5t/4}}{1 + Ke^{5t/4}}$$

or, equivalently,

$$y = \frac{5}{1 + Ce^{-5t/4}}.$$

Here's a picture of the direction field and a few solutions:

