

CH_4

CH_4

4.1 Fourier Series Representation of Continuous-time Periodic Signals

Trigonometric Fourier Series

An Alternative Form for the Fourier Series

Complex Exponentially Fourier Series

Names of the Equations and Coefficients

The Effect of Symmetry

Even Function

Odd Function

Odd Harmonic Function

4.2 Convergence of the Fourier Series

Dirichlet Conditions

Gibbs Phenomenon

4.3 Parseval's Relation

4.1 Fourier Series Representation of Continuous-time Periodic Signals

Trigonometric Fourier Series

Periodic Signal $x(t)$, with period T_1

$$x(t) = a_0 + \sum_{n=1}^{\infty} a_n \cos(n\omega_1 t) + b_n \sin(n\omega_1 t)$$


- Constant Component: $a_0 = \frac{1}{T_1} \int_{t_0}^{t_0+T_1} x(t) dt$
- Cosine Coefficients: $a_n = \frac{2}{T_1} \int_{t_0}^{t_0+T_1} x(t) \cos(n\omega_1 t) dt$
- Sine Coefficients: $b_n = \frac{2}{T_1} \int_{t_0}^{t_0+T_1} x(t) \sin(n\omega_1 t) dt$

An Alternative Form for the Fourier Series

$$x(t) = c_0 + \sum_{n=0}^{\infty} c_n \cos(n\omega_1 t + \phi_n)$$

$$c_0 = a_0 \quad c_n = \sqrt{a_n^2 + b_n^2} \quad \phi_n = \arctan\left(-\frac{b_n}{a_n}\right)$$

$$a_n = c_n \cos(\phi) \quad b_n = -c_n \sin(\phi)$$



- $x(t)$ is real and even

$$b_n = 0 \quad c_n = a_n$$

$$\phi_n = \begin{cases} 0 & a_n > 0 \\ \pm\pi & a_n < 0 \end{cases}$$

- $x(t)$ is real and odd

$$a_n = 0 \quad \phi_n = \pm\frac{\pi}{2}$$


Complex Exponentially Fourier Series

$$\begin{aligned} x(t) &= a_0 + \sum_{n=1}^{\infty} a_n \cos(n\omega_1 t) + b_n \sin(n\omega_1 t) \\ &= a_0 + \sum_{n=1}^{\infty} \left[\frac{a_n - jb_n}{2} e^{jn\omega_1 t} + \frac{a_n + jb_n}{2} e^{-jn\omega_1 t} \right] \end{aligned}$$

Let

$$\begin{aligned} F(n\omega_1) &= \frac{a_n - jb_n}{2} \\ F(-n\omega_1) &= \frac{a_n + jb_n}{2} \\ \therefore \sum_{n=1}^{\infty} F(-n\omega_1) e^{-jn\omega_1 t} &= \sum_{n=-1}^{-\infty} F(n\omega_1) e^{jn\omega_1 t} \\ \therefore x(t) &= \sum_{n=-\infty}^{\infty} F(n\omega_1) e^{jn\omega_1 t} \end{aligned}$$

$$F(n\omega_1) = \frac{1}{T_1} \int_0^{T_1} x(t) e^{-jn\omega_1 t} dt$$



$$F(n\omega_1) = |F(n\omega_1)|e^{j\phi_n}$$

$$|F(n\omega_1)| = \frac{1}{2}\sqrt{a_n^2 + b_n^2} = \frac{1}{2}c_n \quad \phi_n = \arctan\left(-\frac{b_n}{a_n}\right)$$

$$|F_n| = |F_{-n}| = \frac{1}{2}c_n \quad |F_n| + |F_{-n}| = c_n \quad F_n + F_{-n} = a_n$$

since the complex exponential Fourier series is

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t} \quad \omega_0 = \frac{2\pi}{T}$$

- $a_1 e^{\pm j\omega_0 t}$: fundamental components (first harmonic components)
- $a_n e^{\pm jn\omega_0 t}$: Nth harmonic components

Names of the Equations and Coefficients

Given that $\omega_0 = \frac{2\pi}{T}$

- Synthesis Equation:

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t}$$

- Analysis Equation:

$$a_k = \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt$$

- Fourier Series Coefficients (Spectral Coefficients): $\{a_k\}$
- Magnitude Spectrum: $|a_k|$
- Phase Spectrum: $\angle a_k$

the magnitude spectrum and phase spectrum all have **discrete property**, **harmonic property** and **convergence property**

- Constant Component:

$$a_0 = \frac{1}{T} \int_T x(t) dt$$

The Effect of Symmetry

Even Function

The Fourier Series only contains **constant coefficients** and **cosine coefficients**, and $F(n\omega)$ is the **real function**

$$b_n = 0 \quad a_n \neq 0$$
$$F_n = \frac{1}{2}a_n \quad \phi_n = 0$$

Odd Function

The Fourier Series function only contains **sine coefficients**, and $F(n\omega_1)$ is the **virtual function**

$$a_0 = 0 \quad a_n = 0$$
$$b_n \neq 0 \quad F_n = -\frac{1}{2}jb_n$$

Odd Harmonic Function

$$x(t) = -x(t \pm \frac{T}{2})$$
$$\begin{cases} a_0 = 0 \\ a_n = b_n = 0 \end{cases} \quad n = 2k$$
$$\begin{cases} a_n = \frac{4}{T} \int_0^{\frac{T}{2}} x(t) \cos(n\omega_1 t) \\ b_n = \frac{4}{T} \int_0^{\frac{T}{2}} x(t) \sin(n\omega_1 t) \end{cases} \quad n = 2k - 1$$






4.2 Convergence of the Fourier Series

Dirichlet Conditions

- must be integrable
- bounded variation
- finite number of discontinuities

Gibbs Phenomenon

the truncated Fourier series approximation $x_N(t)$ of a discontinuous signal $x(t)$ will in general exhibit high-frequency ripples and overshoot $x(t)$ near the discontinuities



4.3 Parseval's Relation

the **total average power** in a periodic signal equals the sum of the average powers in **all of its harmonic components**