

a Design of the what-if validation
Cut
1 min before recorded start
Response window, 6 h
Real ICU record up to the cut
Vitals
Labs
Medications
Procedures
Give: the recorded start
Withhold
Null arm: second withhold
another seed
b Contrast, observational and simulated
Obs. / Didymos
P (Holm)
Norepinephrine, MAP (mmHg)
−3.3 / −3.6
1.00
Fluid bolus, MAP (mmHg)
−5.8 / −4.3
1.00
Fluid bolus, urine (mL)
−8.2 / −108.4
1.00
Furosemide, urine (mL)
337 / 403
1.00
IMV, FiO2 records (n)
2.8 / 3.1
1.00
IMV, SpO2 (%)
1.1 / 1.2
1.00
CRRT, potassium (mmol/L)
0.24 / 0.44
1.00
CRRT, creatinine 24 h (mg/dL)
0.50 / 1.04
1.00
Insulin, glucose (mg/dL)
38.0 / −0.3
0.31
Observational contrast
Didymos, simulated factual contrast
c Null arm
d Factual accuracy
0.3
0.5
0.7
Expected sign, share
0
0.5
1
CRPS / persistence
0.5
0.8
1
80% coverage
e Timing of the cut, MAP
Norepinephrine
Fluid bolus
−2
−1
0
1
2
Give minus withhold, starters (mmHg)
Cut 0.5–6 h before (pre-specified)
Cut 1 min before