

Rolling anchors
24-hour landmark
48-hour landmark
Mortality
Death, 24 h
ICU death
In-hospital death
Organ failure
Circulatory failure, 8 h
Circulatory failure, 12 h
Hypoxaemia, 4 h
Hypoxaemia, 12 h
AKI stage ≥1, 24 h
AKI stage ≥1, 48 h
AKI stage ≥2, 24 h
AKI stage ≥2, 48 h
AKI stage 3, 48 h
AKI ≥2 (creatinine), 24 h
AKI ≥2 (creatinine), 48 h
AKI ≥2 (urine), 12 h
AKI ≥2 (urine), 24 h
Liver dysfunction, 48 h
Hyperglycaemia, 12 h
Hypoglycaemia, 12 h
Hypoglycaemia, 24 h
Sepsis
Sepsis, 12 h
Sepsis, 24 h
Sepsis without lactate, 24 h
Intervention onsets
New IMV, 4 h
New IMV, 12 h
New IMV, 24 h
New vasopressor, 4 h
New vasopressor, 12 h
New vasopressor, 24 h
New RRT, 24 h
New RRT, 48 h
New NIV, 12 h
New NIV, 24 h
New IMV or NIV, 12 h
Discharge
Discharge alive, 24 h
Discharge alive, 48 h
Threshold crossings
Hypotension, 1 h
Hypotension, 4 h
Hypotension, 12 h
Tachycardia, 4 h
Tachycardia, 12 h
Tachypnoea, 4 h
Tachypnoea, 12 h
Fever ≥38.0 °C, 12 h
Fever ≥38.3 °C, 12 h
Lactate above 4, 12 h
Acidaemia, 12 h
Hyperkalaemia, 12 h
Hyperkalaemia, 24 h
Potassium above 5.5, 24 h
Hyponatraemia, 24 h
Thrombocytopenia, 24 h
Thrombocytopenia, 48 h
Anaemia, 24 h
Leukocytosis, 24 h
Remaining stay
ICU stay >7 days
−0.2
0
0.2
Difference, AUROC
Favours Didymos (FDR 5%)
Favours LightGBM (FDR 5%)
Not significant