
a Events without a vocabulary token
Continuous infusions, RICD
85.1% to 0.0%
Intermittent doses, RICD
31.6% to 2.9%
Locations, RICD
12.7% to 0.0%
Outputs, RICD
14.5% to 0.0%
Continuous infusions, zigongICU
55.5% to 0.6%
Intermittent doses, zigongICU
6.5% to 1.2%
Locations, zigongICU
0.1% to 0.0%
0
25
50
75
100
Share of the domain's events (%)
Original encoding
Mapped codes
b Mapped codes minus original, zero-shot
Death, 24 h, RICD
AKI stage ≥2, 48 h, RICD
Hyperkalaemia, 24 h, RICD
Death, 24 h, zigongICU
New vasopressor, 24 h, zigongICU
Thrombocytopenia, 48 h, zigongICU
ICU death, RICD (L)
In-hospital death, RICD (L)
ICU stay >7 days, RICD (L)
Death, 24 h, zigongICU (L)
ICU death, zigongICU (L)
In-hospital death, zigongICU (L)
ICU stay >7 days, zigongICU (L)
−0.1
0
0.1
Difference in AUROC (95% CI)
c The training grid
Before adaptation
Held-out FT loss (nats per token)
2.2
2.0
1.8
1.6
1.4
1.2
1.0
3 × 10−5, 1 pass
3 × 10−5, 3 passes
1 × 10−4, 1 pass
1 × 10−4, 3 passes (chosen)
0
0.01
0.02
0.03
Change of the seven databases' loss (mean, nats per token)
Learning rate 3 × 10−5
1 × 10−4
1 pass over FT
3 passes
d Membership signal of the FT stays
RICD
zigongICU
Pooled
0.45
0.50
0.55
0.60
AUC, FT stays against TEST stays (95% CI)
Didymos, adapted
Didymos, zero-shot