

Intro to ML

October 13th, 2021

CHAPTER 5:

Multivariate Methods

Multivariate Data

- Multiple measurements (sensors)
- d inputs/features/attributes: d -variate
- N instances/observations/examples

$$\mathbf{X} = \begin{bmatrix} X_1^1 & X_2^1 & \dots & X_d^1 \\ X_1^2 & X_2^2 & \dots & X_d^2 \\ \vdots & & & \\ X_1^N & X_2^N & \dots & X_d^N \end{bmatrix}$$


Multivariate Parameters

Mean: $E[\mathbf{x}] = \boldsymbol{\mu} = [\mu_1, \dots, \mu_d]^T$

Covariance: $\sigma_{ij} \equiv \text{Cov}(x_i, x_j)$

Correlation: $\text{Corr}(x_i, x_j) \equiv \rho_{ij} = \frac{\sigma_{ij}}{\sigma_i \sigma_j}$

Multivariate Gaussian
Distribution parameters



$$\Sigma \equiv \text{Cov}(\mathbf{X}) = E[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T] = \begin{bmatrix} \sigma_1^2 & \sigma_{12} & \cdots & \sigma_{1d} \\ \sigma_{21} & \sigma_2^2 & \cdots & \sigma_{2d} \\ \vdots & & & \\ \sigma_{d1} & \sigma_{d2} & \cdots & \sigma_d^2 \end{bmatrix}$$


Parameter Estimation from data (training)

Sample mean \mathbf{m} : $m_i = \frac{\sum_{t=1}^N x_i^t}{N}, i = 1, \dots, d$

Covariance matrix \mathbf{S} : $s_{ij} = \frac{\sum_{t=1}^N (x_i^t - m_i)(x_j^t - m_j)}{N}$

Correlation matrix \mathbf{R} : $r_{ij} = \frac{s_{ij}}{s_i s_j}$

Multivariate Normal Distribution (your input feature)



$$\mathbf{x} \sim \mathcal{N}_d(\boldsymbol{\mu}, \Sigma)$$

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right]$$

Multivariate Normal Distribution

Use of inverse variance

- Larger variance adds less distance
- Correlated variable contribute less





- Mahalanobis distance: $(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})$
measures the distance from \mathbf{x} to $\boldsymbol{\mu}$ in terms of $\boldsymbol{\Sigma}$ (normalizes for difference in variances and correlations)

- Bivariate: $d = 2$


$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{bmatrix}$$

$$p(x_1, x_2) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \exp\left[-\frac{1}{2(1-\rho^2)}(z_1^2 - 2\rho z_1 z_2 + z_2^2)\right]$$
$$z_i = (x_i - \mu_i) / \sigma_i$$


Bivariate Normal




$\text{Cov}(x_1, x_2)=0, \text{Var}(x_1)=\text{Var}(x_2)$




$\text{Cov}(x_1, x_2)>0$



$\text{Cov}(x_1, x_2)=0, \text{Var}(x_1)>\text{Var}(x_2)$



$\text{Cov}(x_1, x_2)<0$



Parametric Classification

- If $p(\mathbf{x} | C_i) \sim N(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$


$$p(\mathbf{x} | C_i) = \frac{1}{(2\pi)^{d/2} |\boldsymbol{\Sigma}_i|^{1/2}} \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu}_i)^T \boldsymbol{\Sigma}_i^{-1} (\mathbf{x} - \boldsymbol{\mu}_i) \right]$$

- Discriminant functions

$$\begin{aligned} g_i(\mathbf{x}) &= \log p(\mathbf{x} | C_i) + \log P(C_i) \\ &= -\frac{d}{2} \log 2\pi - \frac{1}{2} \log |\boldsymbol{\Sigma}_i| - \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu}_i)^T \boldsymbol{\Sigma}_i^{-1} (\mathbf{x} - \boldsymbol{\mu}_i) + \log P(C_i) \end{aligned}$$

Repeat: Estimation of Parameters

$r_i = 1$ if label = 1, otherwise 0


$$\hat{P}(C_i) = \frac{\sum_t r_i^t}{N}$$
$$\mathbf{m}_i = \frac{\sum_t r_i^t \mathbf{x}^t}{\sum_t r_i^t}$$
$$\mathbf{S}_i = \frac{\sum_t r_i^t (\mathbf{x}^t - \mathbf{m}_i)(\mathbf{x}^t - \mathbf{m}_i)^T}{\sum_t r_i^t}$$

$$g_i(\mathbf{x}) = -\frac{1}{2} \log |\mathbf{S}_i| - \frac{1}{2} (\mathbf{x} - \mathbf{m}_i)^T \mathbf{S}_i^{-1} (\mathbf{x} - \mathbf{m}_i) + \log \hat{P}(C_i)$$

Different \mathbf{S}_i for each class

Quadratic discriminant

Quadratic form

$$g_i(\mathbf{x}) = -\frac{1}{2} \log |\mathbf{S}_i| - \frac{1}{2} (\mathbf{x}^T \mathbf{S}_i^{-1} \mathbf{x} - 2 \mathbf{x}^T \mathbf{S}_i^{-1} \mathbf{m}_i + \mathbf{m}_i^T \mathbf{S}_i^{-1} \mathbf{m}_i) + \log \hat{P}(C_i)$$

$$= \mathbf{x}^T \mathbf{W}_i \mathbf{x} + \mathbf{w}_i^T \mathbf{x} + w_{i0}$$

where

$$\mathbf{W}_i = -\frac{1}{2} \mathbf{S}_i^{-1}$$

$$\mathbf{w}_i = \mathbf{S}_i^{-1} \mathbf{m}_i$$

$$w_{i0} = -\frac{1}{2} \mathbf{m}_i^T \mathbf{S}_i^{-1} \mathbf{m}_i - \frac{1}{2} \log |\mathbf{S}_i| + \log \hat{P}(C_i)$$

Total number of parameters:

- Mean: Kd (k is the number of classes)
- Variance: $K(d(d+1))/2$

Variance is the problem here

Common Covariance Matrix \mathbf{S}

- Shared common sample covariance \mathbf{S} for all class

$$\mathbf{S} = \sum_i \hat{P}(C_i) \mathbf{S}_i \quad \leftarrow \text{Weighted variance}$$

- Discriminant reduces to

$$g_i(\mathbf{x}) = -\frac{1}{2}(\mathbf{x} - \mathbf{m}_i)^T \mathbf{S}^{-1}(\mathbf{x} - \mathbf{m}_i) + \log \hat{P}(C_i)$$


which is a linear discriminant

$$g_i(\mathbf{x}) = \mathbf{w}_i^T \mathbf{x} + w_{i0}$$

where

$$\mathbf{w}_i = \mathbf{S}^{-1} \mathbf{m}_i \quad w_{i0} = -\frac{1}{2} \mathbf{m}_i^T \mathbf{S}^{-1} \mathbf{m}_i + \log \hat{P}(C_i)$$

Common Covariance Matrix S




Diagonal S

- When $x_j, j = 1, \dots, d$, are independent, Σ is diagonal
 $p(x|C_i) = \prod_j p(x_j|C_i)$ (Naive Bayes' assumption)

$$g_i(\mathbf{x}) = -\frac{1}{2} \sum_{j=1}^d \left(\frac{x_j^t - m_{ij}}{s_j} \right)^2 + \log \hat{P}(C_i)$$

Classify based on weighted Euclidean distance (in s_j units) to the nearest mean

Diagonal S



*variances may be
different*

Independent Inputs: Naive Bayes Classifier

- If x_i are independent, off diagonals of Σ are 0, **Mahalanobis distance** reduces to weighted (by $1/\sigma_i$) **Euclidean distance**:

$$p(\mathbf{x}) = \prod_{i=1}^d p_i(x_i) = \frac{1}{(2\pi)^{d/2} \prod_{i=1}^d \sigma_i} \exp \left[-\frac{1}{2} \sum_{i=1}^d \left(\frac{x_i - \mu_i}{\sigma_i} \right)^2 \right]$$

- If variances are also equal, reduces to Euclidean distance


Diagonal S, equal variances

- **Nearest mean classifier:** Classify based on Euclidean distance to the nearest mean

$$\begin{aligned} g_i(\mathbf{x}) &= -\frac{\|\mathbf{x} - \mathbf{m}_i\|^2}{2s^2} + \log \hat{P}(C_i) \\ &= -\frac{1}{2s^2} \sum_{j=1}^d (x_j^t - m_{ij})^2 + \log \hat{P}(C_i) \end{aligned}$$

- Each mean can be considered a prototype or template and this is template matching

Diagonal S , equal variances




Model Selection


<i>Assumption</i>	<i>Covariance matrix</i>	<i>No of parameters</i>
Shared, Hyperspheric	$\mathbf{S}_i = \mathbf{S} = s^2 \mathbf{I}$	1
Shared, Axis-aligned	$\mathbf{S}_i = \mathbf{S}$, with $s_{ij} = 0$	d
Shared, Hyperellipsoidal	$\mathbf{S}_i = \mathbf{S}$	$d(d+1)/2$
Different, Hyperellipsoidal	\mathbf{S}_i	$K d(d+1)/2$

- As we increase complexity (less restricted \mathbf{S}), bias decreases and variance increases
- Assume simple models (allow some bias) to control variance (regularization)


Population likelihoods and posteriors




Arbitrary covar.




Diag. covar.



Shared covar.



Equal var.



Discrete Features

$$p_{ij} \equiv p(x_j=1|C_i)$$

Binary features:

if x_j are independent (Naive Bayes')

$$p(\mathbf{x} | C_i) = \prod_{j=1}^d p_{ij}^{x_j} (1 - p_{ij})^{(1-x_j)}$$

the discriminant is linear

$$\begin{aligned} g_i(\mathbf{x}) &= \log p(\mathbf{x} | C_i) + \log P(C_i) \\ &= \sum_j [x_j \log p_{ij} + (1 - x_j) \log (1 - p_{ij})] + \log P(C_i) \end{aligned}$$

Estimated parameters

$$\hat{p}_{ij} = \frac{\sum_t x_j^t r_i^t}{\sum_t r_i^t}$$

Discrete Features

Multinomial (1-of- n_j) features: $x_j \in \{v_1, v_2, \dots, v_{n_j}\}$

$$p_{ijk} \equiv p(z_{jk}=1 | C_i) = p(x_j = v_k | C_i)$$

if x_j are independent $p(\mathbf{x} | C_i) = \prod_{j=1}^d \prod_{k=1}^{n_j} p_{ijk}^{z_{jk}}$

$$g_i(\mathbf{x}) = \sum_j \sum_k z_{jk} \log p_{ijk} + \log P(C_i)$$

$$\hat{p}_{ijk} = \frac{\sum_t z_{jk}^t r_i^t}{\sum_t r_i^t}$$